



CHEMISTRY


BOOKS - MS CHOUHAN

ALCOHOL, ETHERS AND EPOXIDES

Level 1

1. The following transformation involves a carbocation rearrangement. The carbocation is generated by protonation of the hydroxyl group, followed by the loss of water. Which bond has to migrate in the

carbocation to yield the product indicated (after the deprotonation)



A. a

B. b

C. c



D. d

Answer:




[Watch Video Solution](#)


2. Identify the major product.




A.



B.



C.





D.




Answer:

 [Watch Video Solution](#)

3. Complete the following reaction



A.








Answer:



Watch Video Solution

4. Predict the product when given compound reacts with $LiAlH_4$.






D.


Answer:

 [Watch Video Solution](#)


5. Predict the product when given compound (A, in the above question 4) reacts with $NaBH_4$




A.



B.




C.



D.

Answer:

[▶ Watch Video Solution](#)








The labelled $-O^{18}$ will be in :

- A. H_2O
- B. Methyl benzoate
- C. Both (a) and (b)
- D. Benzoic acid

Answer:




[▶ Watch Video Solution](#)



Answer:


 Watch Video Solution

8. Which is oxidized most easily?







Answer:

 Watch Video Solution





Product (P) is :




Answer:

10. Which one of the following reacts with HBr at the fastest rate?



Answer:

 [Watch Video Solution](#)



Above conversion can be done by :

A. NaBH_4


B. LiAlH_4

C. PCC

D. KMnO_4

Answer:




 Watch Video Solution




12.

(A) ,

product (A) is :







D.




Answer:

 Watch Video Solution



(P) of the reaction is :







Answer:


Watch Video Solution

14. Predict the major product of the given reaction.







A.



B.



C.




D.


Answer:

 [Watch Video Solution](#)


15. Identify the major product ,




A.



B.



C.




D.



Answer:




Watch Video Solution




(A) ,


Product (A) is :




A.



B.




C.



D.


Answer:

[▶ Watch Video Solution](#)




product (A) is :


(A) Major ,




A.



B.



C.








D.

Answer:

 [Watch Video Solution](#)



18. Complete the following reaction



Answer:


 Watch Video Solution

19. Complete the following reaction




(3° alcohol)


A. No reaction



B.






C.




D.


Answer:

 Watch Video Solution






C.



D.

Answer:


[View Text Solution](#)




21.

Major


product obtained in the reaction is :




A.



B.



C.







D.





Answer:

 [Watch Video Solution](#)

22. Consider the following alcohols,




The order of decreasing reactivities of these alcohols towards nucleophilic substitution with HBr is:



Answer:

 [Watch Video Solution](#)



23.

Sum of number of alcoholic groups in product (P) and (Q) is

A. 1

B. 2



C. 3

D. 4



Answer:

 [Watch Video Solution](#)


24. In presence of dil. HCl, compound A is converted to a constitutional isomer (B), compound B is:



A.



C.








D.

Answer:



Watch Video Solution

25. Complete the following reaction



Answer:

 Watch Video Solution



26.


Product of

the reaction is :

- A. Racemic
- B. Diastereomer
- C. Meso
- D. Optically pure

Answer:





[▶ Watch Video Solution](#)



27.


Chemist

added extra 0.5 mole of Et-Li in above reaction to obtain product (A), which is ?



Answer:





Watch Video Solution



28.


Major

product of above esterification reaction is:



Answer:

Watch Video Solution



products are obtained by the mechanism

A. S_N1

B. S_N2

The

C. $SN - NGP$

D. $SN - Ar$

Answer:


 Watch Video Solution




Product


(A) of reaction is :

A.






B.



C.






D.


Answer:




Watch Video Solution

31. Complete the following reaction





C.




D.

Answer:

[▶ Watch Video Solution](#)

32. Which is the best reagent to convert isopropyl alcohol to isopropyl bromide?



A. HBr


B. $SOBr_2$

C. Br_2

D. CH_3MgBr


Answer:


 Watch Video Solution



33.

Major product obtained in the above reaction is :






B.

C. Racemic

D. Diastereomers

Answer:

 [Watch Video Solution](#)



34.

Total number of stereoisomers of product (B) will be:

A. 2

B. 3





C. 4


D. 5

Answer:

 Watch Video Solution

35. Complete the following reaction







D.


Answer:

 [Watch Video Solution](#)

36. Complete the following reaction



A.



B.



C.




D.

Answer:



Watch Video Solution



(1)




(2)

37.





For the above conversion the correct order of reagents used is :

 Watch Video Solution



38.

Find missing reagents .



Answer:

 [Watch Video Solution](#)

39. In solvolysis of 1, 2-dimethyl propyl p-toluene sulfonate in acetic acid at 75°C , (alkene + substitution products) will be formed by mechanism?

A. S_N2, E_2


B. S_N2, E_1

C. S_N1, E_2

D. S_N1, E_1

Answer:

 [Watch Video Solution](#)



40.

x = moles of anhydride consumed

A. 1

B. 2

C. 3

D. 4

Answer:



Watch Video Solution



41. Identify product when (R) - and (S) - 2 - butanol reacts with (R,R) tartaric acid in acidic medium.



- A. Racemic
- B. Diastereomer
- C. Meso
- D. Pure enantiomer

Answer:

 [Watch Video Solution](#)

42. An alcohol of formula $C_9H_{12}O$ reacts with $Na_2Cr_2O_7$ to form a compound having formula $C_9H_{10}O$. The original alcohol might be







Answer:

 [Watch Video Solution](#)


43. An optically active alcohol of formula $C_9H_{12}O_2$ produced the following compound when refluxed with $KMnO_4$




The original compound showed these properties also :




What is structure of (A) ?



A.



B.



C.

D. both (a) and (b)

Answer:

 [Watch Video Solution](#)

44. Which are not cleaved by HIO_4 ?

I : glycerol

II : glycol

III : 1,3 - propenediol

IV : methoxy - 2- propanol

A. I,II,III,IV

B. I,II

C. II,III


D. III,IV




Answer:



[Watch Video Solution](#)

45. Which of the following reactions require an oxidising agent?








Answer:


 [Watch Video Solution](#)

46. What is the major product of the following reaction ?






B.



C.






D.

Answer:

[▶ Watch Video Solution](#)

47. Which of the esters shown, after reduction with $LiAlH_4$ and aqueous workup, will yield two molecules of only a single alcohol ?






D. None of these

Answer:

 [Watch Video Solution](#)

48. For the following reaction, select the statement that best describes the change.




- A. The alcohol is oxidized to an acid, and the Cr(VI) is reduced
- B. The alcohol is oxidized to an aldehyde, and the Cr(VI) is reduced
- C. The alcohol is reduced to an aldehyde, and the Cr(VI) is oxidized
- D. The alcohol is oxidized to a ketone, and the Cr(VI) is reduced

Answer:

 [Watch Video Solution](#)

49. What is the product of the following reaction ?






- A. Only 1
- B. 1:1 mixture of 2 and 3
- C. Only 2
- D. 1:1: 1 mixture of 1,2, and 3

Answer:

 [Watch Video Solution](#)


50. An organic compound B is formed by the reaction of ethylmagnesium iodide ($\text{CH}_3\text{CH}_2\text{MgI}$) with a substance A, followed by treatment with dilute aqueous acid. Compound B does not react with PCC in dichloromethane. Identify A ?



Answer:

 [Watch Video Solution](#)

51. Which of the following reagents would carry out the following transformation? (D = ^2H)



A. NaBD_4 in CH_3OH

B. $LiAlH_4$, then D_2O


C. $NaBD_4$ in CH_3OD

D. $LiAlD_4$, then D_2O

Answer:

 [Watch Video Solution](#)

52. Which sequence of steps describes the best synthesis of 2-methyl-3-



pentanone ?

A. (1) 1 - Propanol + $(CH_3)_2CHMgBr$, diethyl ether

(2) H_3O^+

(3) PCC, CH_2Cl_2

B. (1) 1-Propanol + $Na_2Cr_2O_7$, H_2SO_4 , H_2O heat

(2) $SOCl_2$

(3) $(CH_3)_2CHCl$, $AlCl_3$

C. (1) 1-Propanol + PCC, CH_2Cl_2

(2) $(CH_3)_2CHLi$, diethyl ether

(3) H_3O^+

(4) $Na_2Cr_2O_7$, H_2SO_4 , H_2O heat

D. (1) 2-Propanol + $Na_2Cr_2O_7$, H_2SO_4 , H_2O , Heat

(2) $CH_3CH_2CH_2Li$, diethyl ether




(3) H_3O^+

(4) PCC, CH_2Cl_2

Answer:

 [Watch Video Solution](#)

53. Diols (I-IV) which react with CrO_3 in aqueous H_2SO_4 and yield products that readily undergo decarboxylation on heating, are :










- A. I and II
- B. II and III
- C. II and IV
- D. I and IV

Answer:


 [Watch Video Solution](#)

54. Which of following compounds are not oxidized by HIO_4 ?





Watch Video Solution

55. Complete the following reaction




A.






B.



C.




D.




Answer:

[▶ Watch Video Solution](#)

56. Unknown (A) in the reaction (given in Q. 55) is :



A.



Answer:


 [Watch Video Solution](#)

57. In the given table, identify the incorrect option. The digit in box indicate the moles of that substance.


Reactant	HIO_4 consumed	HCO_2H formed	HCHO formed
(a) $\text{HO}-\text{CH}_2-\overset{\text{OH}}{\text{CH}}-\text{CH}_2-\text{OH}$	2	1	2
(b) $\text{R}-\overset{\text{OH}}{\text{CH}}-\overset{\text{OH}}{\text{CH}}-\overset{\text{OH}}{\text{CH}}-\text{CH}_2-\text{OH}$	3	2	1
(c) $\text{HO}-\text{CH}_2-\overset{\text{OCH}_3}{\text{CH}}-\text{CH}_2\text{OH}$	0	0	0
(d) $\text{HO}-\text{CH}_2-\overset{\text{OH}}{\text{CH}}-\overset{\text{OCH}_3}{\text{CH}_2}$	1	1	1

 **Watch Video Solution**


58. Succinic acid $\xrightarrow{\Delta}$ (A) $\xrightarrow[\Delta]{\text{NH}_2}$ (B) $\xrightarrow[\text{KOH}]{\text{Br}_2}$ (C), Product (C) will be :




A. |




B. |



C. |




D. |




Answer:

59. Given are the structures of cyclic D-glucoside. Moles of HIO_4 consumed with X and Y are respectively:



(X)



(Y)

- A. 2,2
- B. 3,3
- C. 2,3
- D. 3,2

Answer:

60. Moles of formic acid formed in X and Y respectively are:

A. 1,2

B. 2,1

C. 2,3

D. 3,2

Answer:



Watch Video Solution

61. Fructose is oxidised by periodic acid [HIO_4]. The number of moles of

HCOOH formed from each mole of fructose are

A. 1,1

B. 2,2,

C. 1,2

D. 2,1

Answer:



 [Watch Video Solution](#)

62. Which among the following gives positive iodoform test as well as positive Fehling test ?





- A. methanol, ethanol, propanone
- B. ethanol, isopropanal, methanal
- C. ethanol, ethanal, isopropyl alcohol
- D. propanal, propanol-2, propanone

Answer:

 [Watch Video Solution](#)




Product (C) of the reactions is:






Answer:

 [Watch Video Solution](#)



64. Product (B) of the reaction is :







Answer:

 [Watch Video Solution](#)

65. Two unknown compounds X and Y, both having molecular formula C_4H_8O , give following results with four chemical tests.

	Bromine	Na metal	Chromic acid	Lucas reagent
Compound X	decolourises	bubbles	Orange to Green	No reaction
Compound Y	No reaction	No reaction	No reaction	No reaction



Compound X and Y respectively are :




Answer:

 [Watch Video Solution](#)


66. Complete the following reaction




A.



B.



C.








D.

Answer:



Watch Video Solution


67. Assign the structure of major product (X) of the reaction given below.



Answer:







Watch Video Solution



68.


Product (A) and (B) respectively in the above reaction are







Answer:

 Watch Video Solution

69.



Product (A) of above reaction is :






Answer:

 [Watch Video Solution](#)



In above reaction molecular formula of glycerol increases by :






Answer:

 Watch Video Solution

71. Give the best conditions for this transformation:



A. $CH_3OH, H^+ (cat.)$ heat

B. $H_2O, H^+ (cat.)$ heat



C. Mg, ether, CH_3OH

D. $SOCl_2, CH_3OH$


Answer:

 Watch Video Solution


72. Give the major organic product of the following reaction.




A.



B.




C.




D.

Answer:


[▶ Watch Video Solution](#)




Product (D) in above reaction is :




A.



B.



C.




D.

Answer:




[Watch Video Solution](#)

74. Select the best method for the preparation of the following compounds :



and



(MCPBA = Metachloro per benzoic acid)


- A. reaction of cyclohexanone with CH_3Li
- B. reaction of 1-methylcyclohexene with $Hg(OAc)_2$ followed by $NaBH_4$
- C. reaction of cyclohexene with $BH_3:NaOH/H_2O_2$, following by CH_3Br
- D. reaction of cyclohexene with MCPBA, followed by CH_3MgBr

Answer:



[Watch Video Solution](#)

75. Identify the reagents (1-4), required for the transformations shown and arrange them in correct order.



(1) $LAH(LiAlH_4)$ (2) OSO_4

(3) $NaIO_4$ (4) $NaBH_4$

A. 1 → 3 → 4 → 2

B. 2 → 3 → 1 → 4


C. 2 → 1 → 3 → 4

D. 1 → 2 → 3 → 4

Answer:

 [Watch Video Solution](#)

76. Which describes the best stereochemical aspects of the following reaction ?







- A. Inversion of configuration occurs at the carbon undergoing substitution.
- B. Retention of configuration occurs at the carbon undergoing substitution.
- C. Racemization (loss of configuration) occurs at the carbon undergoing substitution.
- D. The carbon undergoing substitution is not stereogenic

Answer:

[Watch Video Solution](#)

77. Which of following is an example of Pinacol-Diazotization ?




Answer:




Watch Video Solution

78. (A) $\xrightarrow{\text{H}_2\text{O}^*}$ B + C, (B) and (C) both give +ve iodoform test. Compound

(A) is









D. both (b) and (c)

Answer:

 [Watch Video Solution](#)






79. A solution of Ph_3CCO_2H in conc. H_2SO_4 gives (X) when poured into methanol X is :



Answer:

 [Watch Video Solution](#)

80. Complete the following reaction





Answer:





Watch Video Solution

81. In the given pair of alcohols, in which pair second alcohol is more reactive than first towards hydrogen bromide?






C.



Answer:

 [Watch Video Solution](#)

82. Rank the transition states that occur during the following reaction steps in order of increasing stability (least \rightarrow most stable):



A. $1 < 2 < 3$

B. $2 < 3 < 1$



C. $1 < 3 < 2$




D. $2 < 21 < 3$

Answer:

 [Watch Video Solution](#)

83. Complete the following reaction









Answer:


Watch Video Solution

84. In which of the following reactions hydrogen gas will not be evolved ?



Answer:


Watch Video Solution



85.

When Grignard reagent reacts with (B) product (D) will be obtained.


Reactant (A) of the above reaction is




A.



B.



C.







D.

Answer:


[Watch Video Solution](#)

86. A $\xrightarrow{\text{LiAlH}_4}$ 2(B), structure of (A) is :
(Chiral alcohol only)




Answer:

[Watch Video Solution](#)





87.

Above conversion can be achieved by :



Answer:


 Watch Video Solution



88. (Mandelic acid)

Identify product of above Fischer esterification reaction:

A. $\text{Ph} - \overset{\text{O-Et}}{\text{CH}} - \text{CO}_2\text{H}$



B.


C. $\text{Ph} - \underset{\text{OH}}{\text{C}}\text{H} - \text{CO}_2\text{Et}$

D. $\text{Ph} - \underset{\text{OH}}{\text{C}}\text{H} - \overset{\text{O}}{\parallel} \text{C} - \text{Et}$





Answer:



Watch Video Solution




Product (C) of the above reaction is :







Answer:

 [Watch Video Solution](#)

90. What is the major product of the following reaction?





Answer:

 [Watch Video Solution](#)

91. The major reason that phenol is a better Bronsted acid than cyclohexanol is that:


- A. it is a better proton donor.
- B. the cyclohexyl group is an electron donating group by induction, which destabilizes the anion formed in the reaction by resonance.
- C. phenol is able to stabilize the anion formed in the reaction.

D. the phenyl group is an electron withdrawing group by induction, which stabilizes the anion formed in the reaction.

Answer:

 [Watch Video Solution](#)

92. Which of these reagents would accomplish the following reduction?



A. $NaBH_4$

B. $LiAlH_4$


C. 1 mole H_2 , poisoned catalyst, low pressure

D. H_3O^+





Answer:

 [Watch Video Solution](#)

93. Complete the following reaction




Identify A and B :







Answer:



Watch Video Solution








Product (A) in above reaction is :



Answer:


 [Watch Video Solution](#)








Answer:

 Watch Video Solution



Fastest of reaction will be when R is :



Answer:



Watch Video Solution


97. Select the correct statement.

- A. Solvolysis of $(CH_3)_2C = CH - CH_2 - Cl$ in ethanol is over 6000 times greater than alkyl chloride ($25^\circ C$)
- B. $CH_3 - CH = CH - CH_2 - OH$ when reacts with HBr give a mixture of 1-bromo-2-butene and 3-bromo 1-butene
- C. When solution of 3-buten-2-ol in aqueous sulphuric acid is allowed to stand for one week, it was found to contain both 3-buten-2-ol and 2-buten-1-ol
- D. All of these

Answer:






Watch Video Solution



98.

Above


transformation can be carried out by :







Answer:

 Watch Video Solution

99.



Compound (C) is :






Answer:



Watch Video Solution


100. Iodoform can be obtained on warming NaOH and iodine with:



Answer:

 [Watch Video Solution](#)


101. Which of these is a reducing agent ?




Answer:


 [Watch Video Solution](#)

102. Complete the following reaction







A.



B.





C.





D.

Answer:

 [Watch Video Solution](#)





D. No reaction

Answer:

 [Watch Video Solution](#)

104. 1, 2, 3 - butanetriol undergoes oxidative cleavage of HIO_4 During this process

A. 1 equivalent of HIO_4 consumed & HCO_2H & $H_3C - \underset{\underset{O}{||}}{C} - CO_2H$

are formed

B. 2 equivalents of HIO_4 consumed &

HCO_2H , $HCH = O$ & $CH_3 - CH - O$ are formed

C. 3 equivalents of HIO_4 consumed & $HCO_2H(2eq.)$ & 1 eq. of

CH_3CO_2H are formed



D. 2 equivalents of HIO_4 consumed & 2 e.q . Of HCO_2H & 1 eq. of

$CH_3CH = O$ is formed


Answer:

 Watch Video Solution


105. Complete the following reaction




A.



B.




C.



D.

Answer:

 [Watch Video Solution](#)



Above reaction is/an example of

A. esterification

B. saponification



C. trans-esterification

D. hydrolysis

Answer:

 Watch Video Solution


107. What is the major organic product of the following sequence of reactions ?




Answer:

 Watch Video Solution

108. The structure of the product formed in the reaction given below is :




A.




B.



C.






D.

Answer:

 Watch Video Solution



A. $HCHO, HCO_2H$






B. $HCHO, 2HCO_2H$

C. $CO_2, 2HCO_2H$

D. $CO_2, HCHO, HCO_2H$



Answer:

 Watch Video Solution



Answer:

 [Watch Video Solution](#)



Ratio of moles of formic acid obtained in reaction (i) and reaction (ii) is :

A. $3/4$

B. $4/5$

C. 1


D. $5/4$

Answer:




Watch Video Solution


112. Which of the following compound gives $HCHO$, CO_2 , $2HCO_2H$ when oxidised by periodic acid ?




A.



B.



C.



D.

Answer:



[Watch Video Solution](#)

113. Hydration of 3- phenylbut -1- ene with dil , H_2SO_4 mainly gives

A. 3-Phenylbutan-1-ol

B. 3-Phenylbutan-2-ol

C. 2-Phenylbutan-2-ol

D. 2-Phenylbutan-1-ol

Answer:

 [Watch Video Solution](#)

114. Decarboxylation of sodium salicylate with soda lime forms

A. Salicylic acid

B. Phenol


C. Benzene

D. None of these

Answer:

 [Watch Video Solution](#)

115. Complete the following reaction








- A. 1
- B. 2
- C. 3
- D. 4

Answer:


 [Watch Video Solution](#)

116. Complete the following reaction






Answer:

Watch Video Solution



117.


Reactant A of the above reaction is :



Answer:

 [Watch Video Solution](#)

118. Which statement is true



- A. (A) - $\begin{matrix} CH_2 - OH \\ | \\ CH_2 - SH \end{matrix}$
- B. (B) = $NaBH_4$
- C. (C) = $KMnO_4$
- D. (D) = H_3O^{\oplus}

Answer:

 Watch Video Solution

119. In the Libermann's nitroso reaction, sequential changes in the colour of phenol occurs as :

- A. Brown or red \rightarrow green \rightarrow deep blue

B. Red \rightarrow deep blue \rightarrow green

C. Red \rightarrow green \rightarrow white

D. White \rightarrow red \rightarrow green

Answer:

 [Watch Video Solution](#)

120. Ethanol when reacts with PCl_5 gives A, $POCl_3$ and HCl. A reacts with dry Ag_2O to form B (major product) and AgCl. A and B respectively are:

A. C_2H_2Cl and $C_2H_5OC_2H_5$

B. C_2H_4 and $C_2H_4OC_2H_5$


C. C_2H_6 and $C_2H_5OC_2H_5$

D. C_2H_6 and $C_2H_5NO_2$

Answer:

 [Watch Video Solution](#)

121. Complete the following reaction




- A. $\text{CH}_3(\text{CH}_2)_3 - \text{CO}_2\text{H}$
- B. $\text{CH}_3(\text{CH}_2)_3 - ^{14}\text{CO}_2\text{H}$
- C. $\text{CH}_3 - (\text{CH}_2)_3\text{CO}_2\text{H}$
- D. $\text{CH}_3 - \overset{\text{O}}{\underset{^{14}}{\text{C}}} - \text{O} - \text{H}$


Answer:

 Watch Video Solution


122. Choose the best synthesis of phenyl n-propyl ether.




A.



B.



C.





D.


Answer:

 [Watch Video Solution](#)


123. Complete the following reaction



A.



B.




C.

D. None of these

Answer:

 [Watch Video Solution](#)

124. What is true for the equilibrium reaction ?




- A. The use of equimolar quantities of CH_3OH and CH_3COOH will give the greatest yield of the ester at equilibrium
- B. Removal of water will increase the amount of ester at equilibrium
- C. Addition of CH_3COOCH_3 will cause the formation of equal an equal number of moles of water
- D. Application of pressure increases the amount of ester at equilibrium




Answer:

 [Watch Video Solution](#)

125. PCC (Pyridiniumchloro chromate) is a good reagent for which of the following transformations ?



A.



Answer:

 [Watch Video Solution](#)

126. How many primary alcohols (including stereoisomers) are possible with formula $C_5H_{12}O$?

- A. Two
- B. Three
- C. Four
- D. Five

Answer:

 [Watch Video Solution](#)

127. 1-Phenylethanol can be prepared by the reaction of benzaldehyde with the product obtained in the reaction between

- A. CH_3I and Mg
- B. C_2H_5I and Mg
- C. CH_3Br and $AlCl_3$
- D. CH_3OH and $ZnCl_2$

Answer:

 [Watch Video Solution](#)

128. 0.092 g of a compound with the molecular formula $C_3H_8O_3$ on reaction with an excess of CH_3MgI gives 67.00 mL of methane at STP. The

number of active hydrogen atoms present in a molecule of the compound is :

- A. one
- B. two
- C. three
- D. four




Answer:

 [Watch Video Solution](#)

129. Migratory aptitude of the following in decreasing order is :






A. $a > c > b > d$





Answer:

 Watch Video Solution

130. The major product formed in the reaction is :





Answer:

 [Watch Video Solution](#)

131. Reaction of R-2-butanol with p-toluenesulphonyl chloride in pyridine then LiBr gives:

- A. R-2-butyl bromide
- B. S-2-butyl tosylate
- C. R-2-butyl tosylate
- D. S-2-butyl bromide

Answer:

 [Watch Video Solution](#)

132. Optically active 2-octanol rapidly loses its optical activity when exposed to :

- A. dilute acid
- B. dilute base
- C. light
- D. humidity

Answer:



Watch Video Solution

133. If (\pm) 2-methyl butanoic acid were esterified by reaction with (\pm) 2-butanol, how many optically active compounds would be present in the final equilibrium reaction mixture?

- A. 2

B. 3





C. 4


D. 6

Answer:

 Watch Video Solution

134. Complete the following reaction






D.


Answer:

[▶ Watch Video Solution](#)


135. Complete the following reaction




A. No reaction



B.




C.







Answer:

 Watch Video Solution





Product obtained in the above reaction is :



Answer:

 Watch Video Solution

137. Reaction - (1) :




Product (B) and (C) respectively are :

- A. CH_2CHO, CH_3CO_2H
- B. CH_3CO_2H, CH_3CHO
- C. CH_3CHO in both reaction
- D. CH_3CO_2H in both reaction

Answer:

 [Watch Video Solution](#)

138. Complete the following reaction






A.



B.



C.




D.

Answer:



Watch Video Solution



139.

this




conversion can be achieved by

- A. O_3, Zn , then LiAlH_4
- B. $\text{O}_3 / \text{H}_2\text{O}_2$, then LiAlH_4
- C. cold dil . $\text{KMnO}_4 / \text{HIO}_4$, then LiAlH_4
- D. All of these

Answer:

 [Watch Video Solution](#)

140. Which of the following alcohol on treatment with HCl give 3-chloro-3-methyl cyclohexene as a product?







D. All of these

Answer:

 [Watch Video Solution](#)

141. Complete the following reaction






D.

Answer:

 [Watch Video Solution](#)

142. Esterification (shown below) is a reaction converting a carboxylic acid to its ester. It involves only the carbonyl carbon. Esterification of (-)-lactic acid with methanol yields (+)-methyl lactate. Assuming that there are no side reactions, what is true about this reaction ?








- A. anS_{N2} process has occurred, inverting the absolute configuration of the chiral center
- B. anS_{N1} reaction at the chiral center has inverted the optical rotation
- C. A diastereomer has been produced, diastereomers have different physical properties including optical rotation
- D. Optical rotation is not directly related to absolute configuration, so the change in sign of rotation is merely a coincidence

Answer:





[Watch Video Solution](#)


143. Which of the following sets of reagents, used in the order shown, would successfully accomplish the conversion shown ?




Answer:

 Watch Video Solution






B.



C.







D.

Answer:

 [Watch Video Solution](#)

145. Which of the following compound on hydrolysis followed by heating gives a product, which gives positive iodoform test?



Answer:

 [Watch Video Solution](#)

146. Treatment of a 2° OH with $\text{CrO}_3 / \text{H}_2\text{SO}_4$ yields an/a :

A. aldehyde

B. carboxylic acid


C. ester

D. ketone

Answer:

 [Watch Video Solution](#)

147. Esterification of the acid P with the alcohols Q will give :



A. only one enantiomer


B. a mixture of diastereomers

C. a mixture of enantiomers

D. only one diastereomer


Answer:

 Watch Video Solution




148.

Identify major product of the reaction, when the given compound is hydrolysed and heated strongly:




A.



B.



C.




D.

Answer:



[Watch Video Solution](#)

149. Complete the following reaction



find out the reactivity order with LiAlH_4 :

A. $A > B > C > D$

B. $B > C > D > A$


C. $D > C > B > A$




D. $B > D > C > A$

Answer:

 [Watch Video Solution](#)


150. Find out the reaction in which obtained product give positive isocyanide test:





Answer:

 [Watch Video Solution](#)



151.

In the above given compound how many functional group reduced by LAH (Lithium aluminium hydride) and SBH (sodium borohydride) respectively?

A. 4,4

B. 4,3

C. 3,4

D. 4,2

Answer:



Watch Video Solution

152. An unknown compound (A) (molar mass = 180) on acylation gives a product (molar mass = 390) than find the number of hydroxyl group present in compound (A).

A. 5

B. 6


C. 10

D. 1


Answer:

 Watch Video Solution


153. Which of the following compound is differentiated by NaHCO_3 as well as by NaOH ?




A.



B.



C.




D.

Answer:

 Watch Video Solution

154. Arrange the following compounds in order of their reactivity towards



hydration

A. $i < ii < iii$

B. $i < iii < ii$

C. $ii < i < iii$


D. $ii < iii < i$

Answer:



Watch Video Solution

155. Choose the statement that is true about A, B and C in the following separation scheme.



A. A = tetrahydroisoquinoline, B = naphthalene and C = inorganic ions

such as Na^+ and Cl^-

B. A = naphthalene, B = tetrahydroisoquinoline and C = inorganic ions

such as Na^+ and Cl^-

C. A = inorganic ions such as Na^+ and Cl^- , B = naphthalene and C

=tetrahydroisoquinoline

D. A = inorganic ions such as Na^+ and Cl^- , B = naphthalene and C =


tetrahydroisoquinoline

Answer:



Watch Video Solution

156. Complete the following reaction







- A. 2
- B. 3
- C. 4
- D. 5

Answer:



Watch Video Solution

157. Complete the following reaction



D. (a) and (c) both

Answer:

 Watch Video Solution

158. 1-Phenoxypropane is treated with excess of conc. HI at 0° C and the mixture of products is treated with thionyl chloride. The products formed are

- A. n-propanol + Chlorobenzene
- B. Phenol + n-propyl iodide
- C. n-propyl chloride + Chlorobenzene
- D. n-propyl chloride + Phenol

Answer:

 [Watch Video Solution](#)


159. Amongst the following compounds, the compound having the lowest boiling point is




B.



C.





D.



Answer:

 [Watch Video Solution](#)


160. Which statement/s are true about products P & Q.




A. P & Q are position isomers

B. P show geometrical isomerism but Q not

C.




D.




Answer:

 [Watch Video Solution](#)




161.

Which of the following is/are possible end product of the above reaction?



A.



B.



C.

D. both (a) and (c)












Answer:

 [Watch Video Solution](#)

1. Consider the pairs of ethers, numbered I through V, shown below. To the right of each pair is a description of reaction conditions to be applied to each. One compound of the pair will react more rapidly than the other.









Which ether of the two will be more rapidly cleaved?

Write your answer in box.

	(A)	Ether Pairs	(B)	Cleavage Conditions
I.		$\text{O}-\text{CH}(\text{CH}_3)_2$ &		Treated with HBr in CH_3CN , 40°C
II.		$\text{O}-\text{C}(\text{CH}_3)_3$ &		Treated with H_2SO_4 in CH_3CN , 40°C
III.			& 	Treated with H_2SO_4 in CH_3CN , 40°C
IV.		&		Treated with 5% aqueous H_2SO_4 , 25°C
V.		&		Treated with 5% aqueous H_2SO_4 , 25°C



Watch Video Solution



Above compounds (a) to (h) are isomers of $\text{C}_5\text{H}_{12}\text{O}$.









2.

Based on the above isomer answer the following (A to F).

Which isomer is most reactive towards dehydration by conc. H_2SO_4 ?



Watch Video Solution



3. Above compounds (a) to (h) are isomers of $\text{C}_5\text{H}_{12}\text{O}$.









Based on the above isomer answer the following (A to F).

Which isomer will undergo rearrangement when treated with conc.

H_2SO_4 ?



Watch Video Solution











4. Above compounds (a) to (h) are isomers of $\text{C}_5\text{H}_{12}\text{O}$.

Based on the above isomer answer the following (A to F).

Which isomers on dehydration with conc. H_2SO_4 give alkene which is capable to show geometrical isomerism ?



Watch Video Solution




5. Above compounds (a) to (h) are isomers of $\text{C}_5\text{H}_{12}\text{O}$.

Based on the above isomer answer the following (A to F).

Which isomer is least acidic ?

 [Watch Video Solution](#)




6. Above compounds (a) to (h) are isomers of $\text{C}_5\text{H}_{12}\text{O}$.

Based on the above isomer answer the following (A to F).

Which isomers on dehydration give most stable alkene?

 [Watch Video Solution](#)




Above compounds (a) to (h) are isomers of $\text{C}_5\text{H}_{12}\text{O}$.

7.

Based on the above isomer answer the following (A to F).

Which isomer on dehydration with conc. H_3PO_4 undergo maximum rearrangement?

 Watch Video Solution



[O] = Oxidation [H] = Reduction

8. Consider the above sequence and answer A to F.

Conversion ($\text{CH}_3 - \text{CH}_3 \rightarrow \text{CH}_3 - \text{CH}_2 - \text{OH}$) alkane \rightarrow alcohol is achieved by :

A. $Br_2/h\nu$, a/cKOH


B. $Br_2/h\nu$, aq. KOH

C. Br_2/CCl_4 , $LiAlH_4$

D. Br_2/CCl_4 , $NaBH_4$

Answer: A-b ;

 Watch Video Solution



9. Consider the above sequence and answer A to F.

Conversion $R-CH_2-OH \rightarrow R-CHO$ can be done by :

A. PCC/CH_2Cl_2


B. $Cu, 300^\circ C$

C. CrO_3

D. All of these

Answer: B::D

 Watch Video Solution




10. Consider the above sequence and answer A to F.

Conversion $R - CHO \rightarrow R - CO_2H$ can be done by

- A. $KMnO_4$
- B. H_2CrO_4
- C. $K_2Cr_2O_7$
- D. All of these

Answer: C::D

 Watch Video Solution



11. Consider the above sequence and answer A to F.

Conversion $R - CO_2H \rightarrow R - CHO$ can be done by :

A. $LiAlH_4$


B. $NaBH_4$

C. $DIBAL - H$

D. All of these




Answer: C::D

 Watch Video Solution



12. Consider the above sequence and answer A to F.


Conversion $R - CHO \rightarrow R - CH_2 - OH$ can be done by :



D. All of these

Answer: D


 Watch Video Solution



[O] = Oxidation [H] = Reduction

13. Consider the above sequence and answer A to F.

Reduction $R-CH_2-OH \rightarrow R-CH_3$ can be done by :



Answer: D

 [Watch Video Solution](#)

14. Which of the following is true for 3- methylbutanal ?

a.	This compound may be classified as an aldehyde.	
b.	This compound may be classified as a ketone	
c.	An aldol reaction takes place on treatment with NaOH solution.	
d.	There is no reaction with LiAlH_4 in ether solution.	
e.	An excess of CH_3MgBr in ether reacts to give 4-methyl-2-pentanol.	
f.	Wolff-Kishner reduction gives butane.	
g.	This compound is an isomer of 3-pentanone.	

 [Watch Video Solution](#)




15. This problem is an introduction to the planning of multistep syntheses

For use, you have six reactant compounds (A through F) , and eight reagents (1 through 8), shown below

Following these lists, five multistep syntheses are outlined. For each of


these, certain reactants or reagents must be identified by writing an appropriate letter or number in designated answer boxes. Write a single letter or number, indicating your choice of the best reactant or reagent, in each answer box.

Reactant Compounds :



Reagents :

- | | |
|---|---|
| (1) Jones' reagent [$\text{Na}_2\text{Cr}_2\text{O}_7$ in H_3O^+] | (2) PCC [CrO_3 in pyridine + HCl] |
| (3) Sodium hydride NaH | (4) Sodium borohydride NaBH_4 |
| (5) Thionyl chloride SOCl_2 | (6) Phosphorus tribromide PBr_3 |
| (7) Aluminium trichloride AlCl_3 | (8) Magnesium turnings in ether |



Watch Video Solution

16. Which of the following is true for 3-methyl-2-butanone?

a.	It may be prepared by CrO_3 oxidation of 2-methyl-2-butanol.	
b.	Its reaction with NaBH_4 gives a secondary alcohol.	
c.	It may be prepared by acidic Hg^{2+} catalyzed hydration of 3-methyl-1-butyne.	
d.	It forms a silver mirror on treatment with $[\text{Ag}(\text{NH}_3)_2]^+$.	
e.	This compound is an isomer of 4-penten-1-ol.	

 [Watch Video Solution](#)

17. Which of these methods would serve to prepare 1-phenyl-2-propanol?

a.	Addition of benzyl Grignard reagent to acetaldehyde (ethanal).	
b.	Addition of phenyl lithium to propylene oxide (methyloxirane).	
c.	Addition of phenyl Grignard reagent to acetone (2-propanone).	
d.	Acid-catalyzed hydration (addition of water to) of 2-phenyl-1-propene.	
e.	Addition of methyl Grignard reagent to acetophenone (methyl phenyl ketone).	
f.	Addition of methyl Grignard reagent to phenylacetaldehyde.	

 [Watch Video Solution](#)

18. Match the Column (I) and (II) .

Column (I)		Column (II)	
Reaction		Name of Reaction	
(a)	$\begin{array}{c} & \\ \text{---} & \text{---} \\ & \\ \text{OH} & \text{OH} \end{array} \xrightarrow[\Delta]{\text{H}^+}$	(p)	Pinacol-Pinacolone rearrangement

(b)	$\begin{array}{c} & \\ \text{---} & \text{---} \\ & \\ \text{OH} & \text{NH}_2 \end{array} \xrightarrow[2\text{HCl}]{\text{NaNO}_2}$	(q)	Semi-Pinacol reaction
(c)	$\text{Cyclic Pinacolone} \xrightarrow[\Delta]{\text{H}^+}$	(r)	Pinacolic-Diazotization
(d)	$\begin{array}{c} & \\ \text{---} & \text{---} \\ & \\ \text{OH} & \text{OH} \end{array} \xrightarrow[(2) \text{Et}_3\text{N}, \Delta]{(1) \text{TiCl}_4}$	(s)	Pinacol fischer reaction

Watch Video Solution

19. Match the Column (I) and (II) .

Column (I)		Column (II)	
Reactant		Products	
(a)		(p)	
(b)		(q)	
(c)		(r)	
(d)		(s)	

Watch Video Solution


Reaction 1.	$ \begin{array}{c} \text{CH}_2\text{-OH} \\ \\ \text{C} = \text{O} \\ \\ \text{HO} - \text{C} - \text{H} \\ \\ \text{H} - \text{C} - \text{OH} \\ \\ \text{H} - \text{C} - \text{OH} \\ \\ \text{CH}_2\text{OH} \end{array} \xrightarrow{5\text{HIO}_4} $
Reaction 2.	$ \begin{array}{c} \text{CHO} \\ \\ \text{HO} - \text{C} - \text{H} \\ \\ \text{H} - \text{C} - \text{OH} \\ \\ \text{H} - \text{C} - \text{OH} \\ \\ \text{H} - \text{C} - \text{OH} \\ \\ \text{CH}_2\text{OH} \end{array} \xrightarrow{5\text{HIO}_4} $

20.

Ratio of moles of formaldehyde obtained in the reaction (1) and reaction (2) ?

Watch Video Solution

21. Di-tert-glycols rearrange in the presence of acid to give *alpha*-tertiary ketones. The trivial name of the simplest glycol of this type is pinacol, and this type of reaction therefore is named pinacol rearrangement (in this specific case, the reaction is called a pinacol-pinacolone rearrangement). The rearrangement involves 4 steps. one of the hydroxyl groups is protonated in the first step. A molecule of water is eliminated in the second step and a tertiary carbocation is formed. The carbocation rearranges in the third step into a more stable carboxonium ion via a [1,2] rearrangement. In the last step, the carboxonium ion is deprotonated and the product ketone is obtained.



What is R.D.S. of pinacol-pinacolone rearrangement?

A. I step

B. II step

C. III step



D. IV step

Answer: B



[Watch Video Solution](#)

22. Di-tert-glycols rearrange in the presence of acid to give *alpha*-tertiary ketones. The trivial name of the simplest glycol of this type is pinacol, and this type of reaction therefore is named pinacol rearrangement (in this specific case, the reaction is called a pinacol-pinacolone rearrangement). The rearrangement involves 4 steps. one of the hydroxyl groups is protonated in the first step. A molecule of water is eliminated in the second step and a tertiary carbocation is formed. The carbocation rearranges in the third step into a more stable carboxonium ion via a [1,2] rearrangement. In the last step, the carboxonium ion is deprotonated and the product ketone is obtained.





How many products obtained in above reaction?

- A. 1
- B. 2
- C. 3
- D. 4

Answer: B


[▶ Watch Video Solution](#)

23. Di-tert-glycols rearrange in the presence of acid to give *alpha*-tertiary ketones. The trivial name of the simplest glycol of this type is pinacol, and this type of reaction therefore is named pinacol rearrangement (in this specific case, the reaction is called a pinacol-pinacolone rearrangement). The rearrangement involves 4 steps. one of the hydroxyl groups is protonated in the first step. A molecule of water is eliminated in the second step and a tertiary carbocation is formed. The carbocation rearranges in the third step into a more stable carboxonium ion via a [1,2] rearrangement. In the last step, the carboxonium ion is deprotonated and the product ketone is obtained.





A.



B.



C.







D.

Answer: C

 [Watch Video Solution](#)

24. Di-tert-glycols rearrange in the presence of acid to give *alpha*-tertiary ketones. The trivial name of the simplest glycol of this type is pinacol, and this type of reaction therefore is named pinacol rearrangement (in this specific case, the reaction is called a pinacol-pinacolone rearrangement). The rearrangement involves 4 steps. one of the hydroxyl groups is protonated in the first step. A molecule of water is eliminated in the second step and a tertiary carbocation is formed. The carbocation rearranges in the third step into a more stable carboxonium ion via a [1,2] rearrangement. In the last step, the carboxonium ion is deprotonated

and the product ketone is obtained.



B.




C.

D. None of these

Answer: A:D

 Watch Video Solution

Subjective Problems



1.

Maximum number of moles of Ac_2O consumed by reactant (A) is :

 Watch Video Solution