



# BIOLOGY

## BOOKS - AAKASH SERIES

# EXCRETORY PRODUCTS AND THEIR ELIMINATION

## Exercise | Excretory Products And Excretory Organs

1. least toxic form of excretory product is

A. ammonia

B. urea

C. uric acid

D. carbon dioxide

**Answer: A**



**Watch Video Solution**

**2. Different types of excretory structures and animals are given below. Match them**

appropriately and mark the correct answer

**Column-I**

**Column-II**

A) Protonephridia

1) Prawn

B) Nephridia

2) Cockroach

C) Malpighian tubules

3) Earthworm

D) Antennal glands/  
green glands

4) Flat worm

1) A - 4, B - 3, C - 2, D - 1

2) A - 3, B - 2, C - 1, D - 4

3) A - 2, B - 1, C - 3, D - 4

4) A - 2, B - 1, C - 4, D - 3



**Watch Video Solution**

**3. Read the following and choose the correct combination**

<b>Ammonotelic</b>	<b>Ureotelic</b>	<b>ricotelic</b>
1) Bony fish	Shark fish	Silver fish
2) Shark fish	Bony fish	Cuttlefish
3) Bony fish	Shark fish	Crayfish
4) Crayfish	Bony fish	Silverfish



**Watch Video Solution**

**4. Protonephridia of flatworms are primarily concerned with**

**A. ionic and fluid volume regulation**

**B. excretion of ammonium ions with their excretory organs**

C. egestion of undigested waste

D. expulsion of gametes during copulation

**Answer: A**



**Watch Video Solution**

5. To minimize the loss of water from the body  
the suitable condition is

A. Ammonotelism

B. Ureotelism

C. Uricotelism

D. Both (1) and (2)

**Answer: C**



**Watch Video Solution**

6. An advantage of excreting nitrogenous wastes in the form of uric acid is that :

A. Uric acid can be excreted in almost solid form

B. The formation of uric acid requires a great deal of energy

C. Uric acid is the first metabolic breakdown product

D. Uric acid may be excreted through the lungs

**Answer: A**



**Watch Video Solution**

7. Ammonia is the main nitrogenous excretory material in

A. Amphibians

B. Aves

C. fresh water fishes

D. Reptiles

**Answer: C**



**Watch Video Solution**



8. Animals which excrete large amount of ammonia are

A. Terrestrial

B. Amphibians

C. Egg Laying

D. Aquatic

**Answer: D**



**Watch Video Solution**

9. Uric acid is nitrogenous waste in

A. Mammals and molluses

B. Birds and lizards

C. Frog and cartilaginous fishes

D. Insects and bony fishes

**Answer: B**



**Watch Video Solution**

**10.** Excretion of nitrogenous waste product mainly as uric acid by birds is helpful in

- A. Conserving body heat
- B. Eliminating excess water
- C. Conserving body water
- D. Eliminating body water

**Answer: C**



**Watch Video Solution**

11. Which one of the following simplest excretory organs ?

- A. Alveoli
- B. Flame cells
- C. Nephridia
- D. Kidney

**Answer: B**



**Watch Video Solution**

12. Elimination of which substance requires large amount of water ?

A. Urea

B. Creatinine

C. Ammonia

D. Uric acid

**Answer: C**



**Watch Video Solution**

13. Which of the following are ammono-telic?

A. Terrestrial amphibians

B. Land snails

C. Marine fishes

D. Aquatic insects

**Answer: D**



**Watch Video Solution**

**14.** In some animals desired osmolarity is maintained in the kidney matrix by retaining

A. creatinine

B. urea

C. ammonia

D. uric acid

**Answer: B**



**Watch Video Solution**

## 15. Excretory structures of rotifers

A. renette glands

B. solenocytes

C. flame cells

D. nephridia

**Answer: C**



**Watch Video Solution**



**16.** The following substances are the excretory products in animals. Choose the least toxic from among them

A. Urea

B. Uric acid

C. Ammonia

D. Carbon dioxide

**Answer: B**



**Watch Video Solution**

17. Which one of the following statements is incorrect?

A. Birds and land snails are uricotelic animals

B. Mammals and frogs are ureotelic animals

C. Aquatic amphibians and aquatic insects are ammonotelic animals

D. Birds and reptiles are ureotelic

**Answer: D**



**Watch Video Solution**

**18. Which of the following pairs is wrong?**

A. Uricotelic - birds

B. Ureotelic - insects

C. Ammonotelic - Tadpole

D. Ureotelic - Elephant

**Answer: B**



Watch Video Solution

## Exercise I Human Excretory System

1. Part of human kidney that has projections called calyces is

A. hilum

B. renal capsule

C. cortex

D. renal pelvis

**Answer: D**



**Watch Video Solution**

**2. Columns of Bertini are present between**

A. renal cortex and medulla

B. Bowman's capsules

C. renal pyramids

D. kidney and urinary bladder

**Answer: C**



[Watch Video Solution](#)

3. Glomerulus in kidney is formed by

- A. renal artery and renal vein
- B. afferent arteriole and renal venule
- C. efferent arteriole and renal portal vein
- D. afferent renal arteriole

**Answer: D**



[Watch Video Solution](#)

4. Medullary nephrons are

A. without Henle loop

B. without vasa recta

C. with long Henle loop

D. with less blood supply

**Answer: C**



**Watch Video Solution**

5. Vasa recta run parallel to

A. PCT

B. DCT

C. loop of Henle

D. all of these

**Answer: C**



**Watch Video Solution**



## 6. In Bowman's capsule

A. afferent arteriole is narrower whereas efferent arteriole is wider

B. afferent arteriole is wider whereas efferent arteriole is narrow

C. afferent capillary is wider and efferent capillary is narrow

D. afferent capillary is narrow and efferent capillary is wide

**Answer: B**



**Watch Video Solution**

7. Through the hilum of a kidney these emerge out

- A. renal artery and ureter
- B. renal vein and ureter
- C. nerve and renal artery
- D. renal artery and renal vein

**Answer: B**



**Watch Video Solution**

**8. Juxtamedullary nephrons are much efficient in reabsorption of components because**

A. they are found adjacent area of cortex with medulla

B. they have larger Bowman's capsules

C. they have long loop of Henle and well

developed vasa recta

D. they are present in desert animals only

**Answer: C**



**Watch Video Solution**

**9. Vasa recta is the continuation of**

A. afferent arteriole

B. efferent arteriole

C. renal vein


D. renal nerve

**Answer: B**



**Watch Video Solution**

**10.** Read the following and choose the correct pertaining to Malpighian body



A. 'A' has lesser diameter than that of 'B'

B. 'B' continues as vasa recta in all vertebrates

C. 'C' is formed by two layers of simple squamous epithelium

D. 'D' continues as loop of Henle and then  
as convoluted tubules

**Answer: C**



**Watch Video Solution**

**11.** Which of the following are found in cortical region of kidney of mammals?

A. PCT and DCT

B. PCT and loop of Henle

C. Loop of Henle and collecting duct

D. DCT and loop of Henle

**Answer: A**



**Watch Video Solution**

**12.** In human kidney, the blood vessel which is not a portal vein but starts with blood capillaries and end with blood capillaries is

A. afferent arteriole



B. renal vein

C. renal artery

D. efferent arteriole

**Answer: D**



**Watch Video Solution**

**13.** In humans, kidneys are located between the levels of

A. last cervical and 3rd thoracic vertebrae

B. last thoracic and 3rd lumbar vertebrae

C. last lumbar and 3rd sacral vertebrae

D. 3rd thoracic and last lumbar vertebrae

**Answer: B**



**Watch Video Solution**

**14.** Broad funnel shaped space in the kidney,  
inner to hilum is

A. renal column

B. renal pelvis

C. renal calyx

D. renal sinus

**Answer: B**



**Watch Video Solution**

**15.** extensions of cortex between the medullary pyramids of kidney and known as

A. columns of Bertini

B. ducts of Bellini

C. renal calyces

D. renal pyramids

**Answer: A**



**Watch Video Solution**

**16.** Which of the following is located in the medulla of the kidney?

A. Malpighian body

B. PCT

C. Henle's loop

D. DCT

**Answer: C**



**Watch Video Solution**

**17.** Majority of the nephrons in the human kidney are

A. Juxta medullary - with long loop of Henle

B. Juxta medullary - with short loop of

Henle

C. Cortical - with long loop of Henle

D. Cortical - with short loop of Henle

**Answer: D**



**Watch Video Solution**

**18.** Which one of the following statement is incorrect?

A. The medullary zone of kidney is divided into a few conical masses called medullary pyramids projecting into the calyces

B. Inside the kidney the cortical region extends in between the medullary pyramids as renal pelvis

C. Glomerulus along with Bowman's capsule is called the renal corpuscle

D. Renal corpuscle, proximal convoluted tubule (PCT) and distal convoluted tubule (DCT) of the nephron are situated in the cortical region of kidney

**Answer: B**



**Watch Video Solution**

**Exercise I Urine Formation**



1. The net pressure gradient that causes the fluid to filter out of the glomeruli into the capsule is

A. 20 mm Hg

B. 50 mm Hg

C. 75 mm Hg

D. 30 mm Hg

**Answer: A**



**Watch Video Solution**

2. The Glomerular capillary blood pressure causes filtration of blood through ?

A. 2 layers

B. 1 layer

C. 3 layers

D. 5 layers

**Answer: C**



**Watch Video Solution**

3. Which of the following occur in epithelium of Bowman's capsule?

- A. Nephrocytes
- B. Podocytes
- C. Neural cells
- D. Protonephridia

**Answer: B**



**Watch Video Solution**

4. Materials that are not filtered in Bowman's capsule are

A. simple sugars

B. water

C. amino acids

D. proteins

**Answer: D**



**Watch Video Solution**

5. On their activation JG cells secrete

- A. erythropoietin
- B. carbonic anhydrase
- C. renin
- D. angiotensin

**Answer: C**



**Watch Video Solution**

6. JG cells get activated when there is

A. more GFR

B. normal GFR

C. fall in GFR

D. both (1) and (3)

**Answer: C**



**Watch Video Solution**

**7. Filtration of the blood takes place at**

A. Proximal convoluted tubules

B. Distal convoluted tubules

C. Malpighian body

D. collecting ducts

**Answer: C**



**Watch Video Solution**

**8. In kidney ultrafiltration occurs due to**

A. osmotic concentration

B. glomerular hydrostatic pressure

C. deoxygenated blood

D. exocytosis

**Answer: B**



**Watch Video Solution**

**9. Glomerular filtrate contains**

A. all the constituents of plasma

B. all the constituents of blood

C. similar to the features of serum



D. all the constituents of plasma excluding  
larger proteins

**Answer: D**



**Watch Video Solution**

**10.** Glomerular filtration per minute is equal to

A.  $1/2$  of blood pumped by each ventricle  
per minute

B.  $\frac{1}{5}$  of blood pumped by each ventricle  
per beat

C.  $\frac{1}{2}$  of blood pumped by each ventricle  
per beat


D.  $\frac{1}{5}$  of blood pumped by each ventricle  
per minute

**Answer: D**



**Watch Video Solution**

11. Identify the correct set of parts in the diagram given below.



A. A - Fenestration, B - Filtration slit, C - Podocyte

B. A - Podocyte, B - Filtration slit, C - Fenestration

C. A - Filtration slit, B - Fenestration, C -  
Podocyte

D. A - Filtration slit, B - Podocyte, C -  
Fenestration

**Answer: A**



**Watch Video Solution**

**12.** The movement of ions against the concentration gradient will be:

A. Active transport

B. Osmosis

C. Diffusion

D. All

**Answer: A**



**Watch Video Solution**

**13.** Volume of the blood filtered by the kidneys per minute is

A. 180ml

B. 125ml

C. 1100-1200ml

D. 5000ml

**Answer: C**



**Watch Video Solution**

**14. Filtration slits are formed by**

A. endothelium of glomerular capillaries

B. podocytes of glomerular capillaries

C. podocytes of Bowman's capsule

D. podocytes of renal capsule

**Answer: C**



**Watch Video Solution**

**15.** Amount of filtrate formed by kidneys per minute is approximately

A. 125ml/min

B. 180 ml/min

C. 1100-1200 ml/min

D. 1500 ml/min

**Answer: A**



**Watch Video Solution**

## Exercise I Function Of The Tubules

1. The ascending limb of loop of Henle



A. permeable to water but not for electrolytes

B. permeable to both water and electrolytes

C. permeable to electrolytes but not for water

D. impermeable for both electrolytes and water

**Answer: C**



**Watch Video Solution**

2. The concentration of filtrate will \_\_\_\_\_ as it moves down in the descending limb of loop of Henle

A. increase

B. decrease

C. remain unchanged

D. first increase and then decrease

**Answer: A**



**Watch Video Solution**

3. From this part urea can be reabsorbed into interstitium to maintain osmolarity of that region

A. PCT

B. DCT

C. loop of Henle

D. collecting duct

**Answer: D**



[Watch Video Solution](#)

4. Urine becomes more and more hypertonic as it passes through

- A. descending limb of loop of Henle
- B. proximal convoluted tubule
- C. ascending limb of loop of Henle
- D. distal convoluted tubule

**Answer: A**



[Watch Video Solution](#)

5. Match the following regarding excretory system and choose correct one.

Column-I	Column-II
A) Proximal convoluted tubule	1) Formation of concentrated urine
B) Distal convoluted tubule	2) Filtration of blood
C) Henle's loop	3) Reabsorption of 70-80% of electrolytes
D) Counter current	4) Ionic balance mechanism
E) Renal corpuscle	5) Maintenance of concentration gradient in renal medulla

1) A - 3, B - 5, C - 4, D - 2, E - 1

2) A - 3, B - 4, C - 1, D - 5, E - 2

3) A - 1, B - 3, C - 2, D - 5, E - 4

4) A - 3, B - 1, C - 4, D - 5, E - 2



Watch Video Solution

6. Major portion of glomerular filtrate is reabsorbed in

- A. proximal convoluted tubule
- B. distal convoluted tubule
- C. ascending limb of loop of Henle
- D. descending limb of loop of Henle

**Answer: A**



**Watch Video Solution**

7. Essential nutrients like glucose, amino acids and vitamins reabsorbed in

A. DCT

B. PCT

C. Loop of Henle

D. Collecting duct

**Answer: B**



**Watch Video Solution**

8. Under normal conditions which one is completely reabsorbed in the renal tubule?

A. urea

B. uric acid

C. glucose

D. creatinine

**Answer: C**



**Watch Video Solution**



9. Which one is impermeable to reabsorption of electrolytes in nephron?

- A. proximal convoluted tubule
- B. distal convoluted tubule
- C. ascending limb of loop of Henle
- D. descending limb of loop of Henle

**Answer: D**



**Watch Video Solution**

**10.** Maintenance of pH and ionic balance of blood takes place by the selective secretion of certain ions takes place in

PCT

DCT

Collecting duct

All the above

A. PCT

B. DCT

C. Collecting duct


D. All the above

**Answer: D**



**Watch Video Solution**

**11.** Read the following diagram and choose the correct statements.



I). In part 'A', the water reabsorption takes place II) In part 'B', the water and electrolytes reabsorption take place III) In part 'C', the water reabsorption takes place IV) In part 'D', the water is reabsorbed but electrolytes are not reabsorbed

A. I and II

B. I and III

C. III and IV

D. I and IV

**Answer: B**



**Watch Video Solution**

**12.** Identify the correct statement from the following

- A. 70-80% of electrolytes and water are reabsorbed in PCT
- B. Volume of the glomerular filtrate is 180 liters
- C. Volume of urine per day is 1.5 liters
- D. All the above

**Answer: D**



**Watch Video Solution**

**13.** Which of the following is incorrect?

A. Primary urine is hypotonic to cortical fluid

B. Urine in CD is hypertonic to plasma of blood

C. Urine in CD is hypertonic to medullary fluid

D. Urine in CD is equal to the concentration of medullary fluid

**Answer: C**



**Watch Video Solution**

**14. Which is mismatched.**

A. Bowman's capsule - Glomerular filtration

B. PCT - Absorption of  $Na^+$  and  $K^+$

C. DCT - Absorption of glucose

D. None of these

**Answer: C**





Watch Video Solution

15. Maximum absorption of  $Na^+$  and  $K^+$  ions occur in

- A. Bowman's capsule
- B. Loop of Henle
- C. Distal convoluted tubule
- D. Proximal convoluted tubule

**Answer: A**



**16.** Renal fluid present in the part of renal tubule which is impermeable to water but permeable to salts is

- A. Hypertonic to the medullary fluid
- B. Hypotonic to the medullary fluid
- C. Isotonic to the cortical fluid
- D. Hypertonic to blood and isotonic to the cortical fluid

**Answer: B**



**Watch Video Solution**

**17.** During formation of urine NaCl is returned to the medullary interstitium from the

- A. Ascending limb of Henle's loop
- B. Descending portion of the vasa recta
- C. Ascending portion of the vasa recta
- D. Descending limb of Henle's loop

**Answer: C**



**Watch Video Solution**

**18.** Which process plays important role in maintenance of ionic and acid base balance of body fluids

- A. ultrafiltration
- B. tubular secretion
- C. reabsorption
- D. glomerular filtration

**Answer: B**



**Watch Video Solution**

## **Exercise I Mechanism Of Concentration Of The Filtrate**

**1. Counter current mechanism observed in the renal medulla helps in formation of concentrated urine**

dilute urine

reabsorption of nutrients

reabsorption of creatinine

A. concentrated urine

B. dilute urine

C. reabsorption of nutrients

D. reabsorption of creatinine

**Answer: A**



**Watch Video Solution**

2. Statement (A): Humans have the ability to produce concentrated urine.

Statement (B) : Majority of the nephrons in the human kidney have very long loops of Henle and well-developed vasa recta which are involved in countercurrent mechanism.

A. Both A and B are correct

B. Both A and B are false

C. A is true, but B is false

D. A is false, but B is true

**Answer: C**



**Watch Video Solution**

**3.** Loop of Henle plays a significant role in the maintenance of

A. low osmolarity in medullary interstitial fluid

B. high osmolarity in cortical interstitial fluid



C. low osmolarity in cortical interstitial fluid

D. high osmolarity in medullary interstitial fluid

**Answer: D**



**Watch Video Solution**

4. Osmolarity close to cortex is 300 mOsmol/L and 1200 mOsmol/L in the inner medulla, the gradient mainly caused by

A. bicarbonates and glucose

B. glucose and amino acids

C. NaCl and urea

D. NaCl only

**Answer: C**



**Watch Video Solution**

**5. Human urine as compared to human blood is normally : —**

A. Hypotonic

B. Hypertonic

C. Isotonic

D. All of these

**Answer: B**



**Watch Video Solution**

**6.** We can produce concentrated / dilute urine.

This is facilitated by a special mechanism.

Identify the mechanism.

- A. Reabsorption from PCT
- B. Reabsorption from collecting Duct
- C. Reabsorption/Secretion in DCT
- D. Counter current mechanism in Henle's loop/Vasa recta

**Answer: D**



**Watch Video Solution**

**Exercise I Regulation Of Kidney Function**

1. Hormone released by hypothalamus, which is related with the functioning of kidney is

A. renin

B. ANF

C. vasopressin

D. angiotensin

**Answer: C**



**Watch Video Solution**

2. Angiotensin II is

- A. a vasoconstrictor
- B. a vasodilator
- C. activating adrenal medulla
- D. decreasing GFR

**Answer: A**



**Watch Video Solution**

### 3. Atrial Natriuretic Factor (ANF)

- A. causes vasodilation
- B. increases blood pressure
- C. decreases GFR
- D. activates renin

**Answer: A**



**Watch Video Solution**

4. Match the following regarding excretory system and choose correct one.

**Column-I**

- A) JG cells
- B) Hypothalamus
- C) Angiotensin II
- D) ANF

**Column-II**

- 1) Vasodilator
- 2) Vasoconstrictor
- 3) Vasopressin
- 4) Renin

- 1) A - 4, B - 3, C - 2, D - 1
- 2) A - 3, B - 2, C - 1, D - 4
- 3) A - 2, B - 1, C - 3, D - 4
- 4) A - 2, B - 1, C - 4, D - 3



**Watch Video Solution**

5. Statement (A): Atrial natriuretic peptide (ANP) mechanism acts as a counter check on



the RAAS (Renin-Angiotensin-Aldosterone system).

Statement (B) : ANP causes contraction of vascular smooth muscles.

A. Both A and B are correct

B. Both A and B are false

C. A is true, but B is false

D. A is false, but B is true

**Answer: C**



**Watch Video Solution**

6. Which of the following statement is correct?

A. ANF and renin-angiotensin mechanism  
are similar in their function

B. ADH and ANF are vasodilators

C. Vasopressin prevents more loss of water  
through urine

D. JGA releases renin when glomerular  
blood pressure increases

**Answer: C**



**Watch Video Solution**

7. Enzyme released from kidney is

A. renin

B. uricase

C. pepsin

D. none of these

**Answer: A**



Watch Video Solution

8. Read the following

Angiotensinogen  $\xrightarrow{A}$  Angiotensin - I  $\xrightarrow{B}$

Angiotensin- II  $\xrightarrow{C}$  Aldosterone Identify A, B

and C

A. A-Renin, B-Angiotensin converting

enzyme and C-Adrenal medulla

B. A-Renin, B-Angiotensin converting

enzyme and C-Adrenal cortex

C. A-Angiotensin converting enzyme, B-

Renin and C-Adrenal medulla

D. A-Angiotensin converting enzyme, B-

Renin and C-Adrenal cortex

**Answer: B**



**Watch Video Solution**

**9. Role of Angiotensin II is**

A. Decreases the GFR

B. Enhances reabsorption of  $Na^+$

C. Stimulates the secretion of ANP

D. Decreases BP

**Answer: B**



**Watch Video Solution**

**10. Increase in body fluid volume**

A. 1. activates the osmoreceptors

B. 2. suppresses release of ADH

C. 3. increases ADH secretion

D. 4, prevents diuresis

**Answer: B**



**Watch Video Solution**

**11.** Which one of the following is also known as antidiuretic hormone?

A. Oxytocin

B. Vasopressin

C. Adrenaline

D. Calcitonin

**Answer: B**



**Watch Video Solution**

## Exercise I Micturition

1. Micturition occurs due to

A. contraction of bladder muscles



B. contraction of urethral sphincter

C. relaxation of urethral sphincter

D. Both (1) and (3)

**Answer: D**



**Watch Video Solution**

2. Amount of urea excreted out per day is

A. 1 - 1.5 lt

B. 25-30 gm

C. 180 lt

D. 10-15 gm

**Answer: B**



**Watch Video Solution**

**3. Urine of a diabetes mellitus patient consists of**

A. no glucose

B. glucose

C. ketone bodies

D. both (2) and (3)

**Answer: D**



**Watch Video Solution**

**4. The pH of urine is approximately**

A. 6.5

B. 7

C. 6

D. 7.5

**Answer: C**



**Watch Video Solution**

## Exercise I Role Of Other Organs In Excretion

1. Sweat glands help in elimination of

A. NaCl

B. Urea

C. Lactic acid

D. All the above

**Answer: D**



**Watch Video Solution**

**2. Substances eliminated along with sebum**

A. hydrocarbons

B. sterols

C. waxes

D. all the above

**Answer: D**



**Watch Video Solution**

**3.** A large quantity of one of the following is removed from our body by lungs.

A.  $CO_2$  only

B.  $H_2O$  only

C.  $CO_2$  and  $H_2O$

D. ammonia

**Answer: C**



**Watch Video Solution**

**4. Sweating is meant for**

A. removal of excess salt

B. regulation of body temperature

C. killing of skin bacteria

D. removal of excess water

**Answer: B**



**Watch Video Solution**

**5. Lactic acid is excreted by**

A. sebaceous glands

B. sudoriferous glands

C. liver

D. kidneys

**Answer: B**





Watch Video Solution

## Exercise I Disorders Of The Excretory System

1. Haemodialysis is helpful for the patients who are suffering from

- A. diabetes insipidus
- B. uremia
- C. glomerulonephritis
- D. diabetes mellitus

**Answer: B**



**Watch Video Solution**

2. Formation of insoluble mass of crystallised salts in the kidney is

A. glomeulonephritis

B. uremia

C. renal calculi

D. renal failure

**Answer: C**



**Watch Video Solution**

**3. Dialysis fluid contains all the constituents as in plasma except**

A. water

B. amino acids

C. nitrogenous wastes

D. sugars

**Answer: C**



**Watch Video Solution**

4. The condition of accumulation of urea in the blood is termed as

- A. renal calculi
- B. glomerulonephritis
- C. uremia
- D. ketonuria

**Answer: C**



**Watch Video Solution**

5. Dialysing unit (artificial kidney) contains a fluid which is almost same as plasma except that it has

A. high glucose

B. no urea

C. high urea

D. high uric acid

**Answer: B**



**Watch Video Solution**

6. After the process of dialysis, the cleared blood is pumped back to the body through a vein by adding this

- A. anti-heparin
- B. heparin
- C. dialyzing fluid
- D. nitrogenous wastes

**Answer: A**



**Watch Video Solution**

7. Match the abnormal conditions given in Column A with their explanations, given in Column B and Choose the correct option

	<b>Column A</b>		<b>Column B</b>
A.	Glycosurea	i.	Accumulation of uric acid in joints
B.	Renal calculi	ii.	Inflammation in glomeruli
C.	Glomerular nephritis	iii.	Mass of crystallised salts within the kidney
D.	Gout	iv.	Presence of glucose in urine



**Watch Video Solution**

## Exercise II Excretory Products And Excretory Organs

### 1. Match the following.

#### List - I

- A) Renette glands
- B) Antennary glands
- C) Pericardial glands
- D) Uricose glands
- E) Coxal glands

#### List - II

- I) *Palamnaeus*
- II) *Wuchereria*
- III) *Periplaneta*
- IV) *Pila*
- V) *Carcinus*

- 1) A-III, B-V, C-IV, D-I, E-II
- 2) A-II, B-V, C-IV, D-III, E-I
- 3) A-III, B-IV, C-V, D-II, E-I
- 4) A-II, B-III, C-IV, D-V, E-I



Watch Video Solution



## 2. Choose the correct statements

A) Uric acid is converted into allantoin by the action of uricase enzyme, in humans

B) Most of the ammonia is lost as ammonium ions across the epithelium of gills, in bony fishes

C) Trimethylamine oxide protects body proteins from the damaging effect of uric acid in cartilaginous fishes

A. A, B only

B. B, C only

C. A, C only

D. All

**Answer: B**



**Watch Video Solution**

**3. Nitrogenous waste products are eliminated mainly as**

A. Urea in tadpole & ammonia in adult frog

B. Ammonia in tadpole and urea in adult  
frog

C. Urea in both tadpole & adult frog

D. Urea in tadpole and uric acid in adult  
frog

**Answer: B**



**Watch Video Solution**

4. In ornithine cycle, enzyme arginase breaks down arginine into

- A. Citrulline and ammonia
- B. Ornithine and ammonia
- C. Ornithine and urea
- D. Citrulline and urea

**Answer: C**



**Watch Video Solution**

5. Trimethylamine is the excretory product in

- A. Marine teleosts
- B. Fresh water fishes
- C. Terrestrial Molluscs
- D. Amphibians

**Answer: A**



**Watch Video Solution**

6. Urea is derived from

A. Facts

B. Amino acids

C. Carbohydrates

D. Uric acid

**Answer: B**



**Watch Video Solution**

7. Ornithine cycle operates in

A. Stomach

B. Pancreas

C. Liver

D. Oral cavity

**Answer: C**



**Watch Video Solution**

**8.** Physiologically urea is produced by the action of an enzyme

A. Uricase

B. Urease

C. Arginase

D. None

**Answer: C**



**Watch Video Solution**

**9. Identify the correct matching pair**

A. fish - uricotelic

B. man - ureotelic



C. bird - ammonotelic

D. frog - uricotelic

**Answer: B**



**Watch Video Solution**

**10.** Correct order of excretory organs in Cockroach, Earthworm and Rabbit respectively

: —

A. Skin, Malpighian tubules, kidneys

B. Malpighian tubules, nephridia, kidneys

C. Nephridia, Malpighian tubules, kidneys

D. Nephridia, kidneys, green glands

**Answer: B**



**Watch Video Solution**

**11.** Which one of the following pair of waste substances is removed from blood in ornithine cycle :

A.  $CO_2$  and urea

B. Ammonia and urea

C.  $CO_2$  and ammonia

D. Urea and sodium salt

**Answer: C**



**Watch Video Solution**

**12.** The presence of arginase confirms that

A. Arginine is synthesised

B. Ammonia is formed

C. Arginine is being converted into  
ornithine

D. Arginine is being converted into  
citrulline

**Answer: C**



**Watch Video Solution**

13. Kidney of a mammal resembles to the contractile vacuole of Amoeba in expelling out

A. Glucose

B. Excess water

C. Urea

D. Ammonia

**Answer: B**



**Watch Video Solution**

14. In man purine metabolism results in the formation of

A. Urea

B. Uric acid

C. Ammonia

D. Allantoin

**Answer: B**



**Watch Video Solution**

15. Metanephridia, Malpighian tubules and antennal glands are the excretory organs of the following respectively

A. Hydras, sponges and planarians

B. Prawns, cockroaches and crabs

C. Earthworms, insects and prawns

D. Starfishes, cyclostomes and bony fishes

**Answer: C**



**Watch Video Solution**

**16.** Earthworms are

A. Ureotelic when plenty of water is available

B. Uricotelic when plenty of water is available

C. Uricotelic under conditions of water scarcity

D. Ammonotelic when plenty of water is available



**Answer: D**



**Watch Video Solution**

## Exercise II Human Excretory System

1. The functional adult kidney of human being is

A. archinephros

B. pronephros

C. mesonephros

D. Metanephros

**Answer: D**



**Watch Video Solution**

2. Which blood vessel carries least percentage of urea?

A. Hepatic vein

B. Renal vein

C. Hepatic portal vein

D. Renal artery

**Answer: B**



**Watch Video Solution**

**3. The afferent and efferent vessels are**

A. Arterial in nature

B. Venous in nature

C. One is arterial and the other is venous

D. None of the above

**Answer: A**



**Watch Video Solution**

4. All Bowman's capsules of the kidney are found in

A. Pelvis

B. Medulla

C. Cortex

D. None

**Answer: C**



**Watch Video Solution**

5. Complete loop of Henle is found in

A. Amphibia

B. Reptilia

C. Birds

D. Mammals

**Answer: D**



Watch Video Solution

6. Which of the following is not a function of kidneys?

A. Regulation of blood pressure

B. Removal of urea

C. Regulation of acidity of fluids

D. Secretion of antibiotics

**Answer: D**



7. Which type of kidneys are found in amphibians?

- A. Holonephric
- B. Mesonephric
- C. Pronephric
- D. Metanephric

**Answer: B**



## Exercise II Urine Formation

1. One is found in blood not in nephric filtrate

A. Urea

B. Glucose

C. Amino acids

D. Globulin

**Answer: D**



**Watch Video Solution**



2. The effective filtration pressure in uriniferous tubule is

A. 75 mm Hg

B. 10 mm Hg

C. 100 mm Hg

D. 125 mm Hg

**Answer: B**



**Watch Video Solution**

3. Blood fraction remaining unchanged after circulation through kidney is

A. Urea and uric acid

B. Urea and proteins

C. Urea and glucose

D. Glucose and proteins

**Answer: A**



**Watch Video Solution**

4. Changeable threshold material in Renal tubules

A. Water and Glucose

B. Urea and uric acid

C. Glucose and amino acids

D. Water and salts

**Answer: D**



**Watch Video Solution**

5. Due to insufficient filtration in the Bowman's capsule, all are likely to happen except :

- A. Accumulation of fluid in the body
- B. Increase in blood pressure
- C. Increase in blood urea level
- D. Loss of glucose through urine

**Answer: D**



**Watch Video Solution**

## Exercise II Function Of The Tubules

1. Match the following regarding excretory system and choose correct one.

Column-I	Column-II
A) Proximal convoluted tubule	1) Simple epithelium
B) Collecting duct	2) Simple cuboidal epithelium
C) Bowman's capsule	3) Simple squamous epithelium
D) Urinary bladder	4) Transitional epithelium

- 1) A - 4, B - 3, C - 2, D - 1
- 2) A - 3, B - 2, C - 1, D - 4
- 3) A - 2, B - 1, C - 3, D - 4
- 4) A - 2, B - 1, C - 4, D - 3



[Watch Video Solution](#)

## 2. Route of tubular secretion is

A. Blood → Interstitial fluid → Renal fluid

B. Renal fluid → Interstitial fluid → Blood

C. Blood → Renal fluid → Interstitial fluid

D. Interstitial fluid → Renal fluid → Blood

**Answer: A**



**Watch Video Solution**

**3. Which of the following statement is correct pertaining to the functioning of nephron?**

A. Conditional reabsorption of water takes place in proximal convoluted tubule

B. Concentrated filtrate when passes upward in ascending limb of loop of

Henle dilutes due to passage of electrolytes into medullary fluid

C. More glucose is reabsorbed in distal convoluted tubule

D. There is no hormonal influence on the reabsorption of filtration load in collecting duct

**Answer: B**



**Watch Video Solution**



# Exercise II Mechanism Of Concentration Of The Filtrate

1. Concentration of urine depends upon which organ:

A. Bowman's capsule

B. length of Henle's loop

C. P.C.T

D. network of capillaries arising from  
glomerulus

**Answer: B**



**Watch Video Solution**

**2. Counter current exchange occurs in**

A. PCT

B. DCT

C. Vasa recta

D. Loop of Henle

**Answer: C**



Watch Video Solution

## Exercise II Regulation Of Kidney Function

1. Which regulates reabsorption of salts from Glomerular filtrate ?

- A. Oxytocin
- B. Vasopressin
- C. Glucocorticoids
- D. Mineralocorticoids

**Answer: D**



**Watch Video Solution**

## Exercise II Role Of Other Organs In Excretion

1. Workers in deep mines usually suffer from dehydration because:-

- A. Water is lost due to evaporation
- B. Water is lost due to defaecation
- C. Water is lost in the form of urine

D. Water is lost along with salts in the form  
of sweat

**Answer: D**



**Watch Video Solution**

2. Why do we pass more urine in wet and cold  
season?

A. Impairment of water absorption by  
nephrons

B. Kidney becomes more active

C. Sweating is much decreased

D. ADH secretion is increased

**Answer: C**



**Watch Video Solution**

## **Exercise II Disorders Of The Excretory System**

**1. A condition of failure of kidney to form urine is called**

A. Creatinine

B. Haematuria

C. Anuria

D. Ketonuria

**Answer: C**



**Watch Video Solution**

## 2. Match the following.

### Column-I

- A) Renal calculi
- B) Gout
- C) Uraemia
- D) Glucosuria

### Column-II

- 1) Uric acid deposition
- 2) Hyperparathyroidism
- 3) Diabetes insipidus
- 4) Kidney failure
- 5) Hypoinsulinism

- 1) A - 2, B - 1, C - 4, D - 5
- 2) A - 5, B - 2, C - 3, D - 1
- 3) A - 3, B - 2, C - 4, D - 1
- 4) A - 3, B - 4, C - 2, D - 5



**Watch Video Solution**

## 3. Presence of RBC in urine called : —

A. Anuria



B. Haematuria

C. Glycosuria

D. Ketonuria

**Answer: B**



**Watch Video Solution**

4. Urine of a person who takes a protein-deficient diet will have

A. Little glucose

B. Less urea

C. Excess urea

D. Little fat

**Answer: B**



**Watch Video Solution**

5. Urine of a human being suffering from diabetes inspidus is

A. Tasteless and thick

B. Sweet and thick

C. Tasteless and watery

D. Sweet and watery

**Answer: C**



**Watch Video Solution**

**6.** A person is undergoing prolonged fasting.

His urine will be found to contain abnormal quantities of –

A. Fats

B. Amino acids

C. Ketones

D. Glucose

**Answer: C**



**Watch Video Solution**

7. What will happen if one kidney is removed from the body of a human

A. Death due to poisoning

B. Uremia and death

C. Stoppage of urination

D. Nothing, the person will survive and remain normal, kidney will become hypertrophied

**Answer: D**



**Watch Video Solution**

8. In diabetes mellitus the patient drink more water as there is urinary loss of

A. Salt

B. Insulin

C. Protein

D. Glucose

**Answer: D**



**Watch Video Solution**

9. If kidneys fail to reabsorb water the effect on tissue would

- A. Remain unaffected
- B. Shrink and shrivel
- C. Absorb water from blood plasma
- D. Take more  $O_2$  from blood

**Answer: B**



**Watch Video Solution**

**10. Ketonuria is**

A. Albumin in urine

B. Globulin in urine

C. Ketone bodies in urine

D. None of the above

**Answer: C**



**Watch Video Solution**



**11.** Which of the following statements are correct?

(i). Glucose has high threshold value.

(ii). Urine is concentrated in Henle's loop

(iii). Haemodialyser removes urea, uric acid, glucose and plasma proteins

(iv). In glomerulus, urea, uric acid, water, glucose and plasma proteins are filtered out.

A. A, C, D

B. B, C, D

C. A, B

D. A, C

**Answer: C**



**Watch Video Solution**

**12. Match the following regarding excretory system and choose correct one.**

**Column-I**

- A) JG cells
- B) Hypothalamus
- C) Angiotensin II
- D) ANF

**Column-II**

- 1) Vasodilator
- 2) Vasoconstrictor
- 3) Vasopressin
- 4) Renin

- 1) A - 4, B - 3, C - 2, D - 1
- 2) A - 3, B - 2, C - 1, D - 4
- 3) A - 2, B - 1, C - 3, D - 4
- 4) A - 2, B - 1, C - 4, D - 3



[Watch Video Solution](#)

**13.** Match the following regarding excretory system and choose correct one.

**Column-I**

- A) Urochrome
- B) Ornithine cycle
- C) Angiotensinogen
- D) Hilum

**Column-II**

- 1) Urea forms in liver
- 2) Colour of urine
- 3) Concave border of kidney
- 4) Produced by liver

- 1) A - 4, B - 3, C - 2, D - 1
- 2) A - 3, B - 2, C - 1, D - 4
- 3) A - 2, B - 1, C - 3, D - 4
- 4) A - 2, B - 1, C - 4, D - 3



[Watch Video Solution](#)

1. The part of nephron involved in active reabsorption of sodium is

A. Bowman's capsule

B. Descending limb of Henle's loop

C. Distal convoluted tubule

D. Proximal convoluted tubule

**Answer: D**



**Watch Video Solution**

2. In mammals, which blood vessel would normally carry largest amount of urea

- A. Renal vein
- B. Dorsal aorta
- C. Hepatic vein
- D. Hepatic portal vein

**Answer: C**



**Watch Video Solution**

3. Human urine is usually acidic because

A. Hydrogen ions are actively secreted into the filtrate

B. The sodium transporter exchanges one hydrogen ion for each sodium ion, in peritubular capillaries

C. Excreted plasma proteins are acidic

D. Potassium and sodium exchange generates acidity

**Answer: A**



**Watch Video Solution**

**4. Removal of proximal convoluted tubule from the nephron will result in**

- A. No urine formation
- B. More diluted urine
- C. More concentrated urine

D. No change in quality and quantity of urine

**Answer: B**



**Watch Video Solution**

5. Which of the following does not favour the formation of large quantities of dilute urine?

A. Atrial - natriuretic factor

B. Alcohol



C. Caffeine

D. Renin

**Answer: D**



**Watch Video Solution**

**6.** Which of the following causes an increase in sodium reabsorption in the distal convoluted tubule?

A. Increase in aldosterone levels

B. Decrease in antidiuretic hormone levels

C. Decrease in aldosterone levels

D. Decrease in antidiuretic hormone levels


**Answer: A**



**Watch Video Solution**

7. Figure shows human urinary system with structures labelled A to D. Select option which correctly identifies them and gives their

## characteristics and/or functions



A. A-Adrenal gland-located at the anterior part of Kidney. Secrete catecholamines which stimulate glycogen breakdown

B. B-Pelvis-broad funnel shaped space inner to hilum, directly connected to loop of Henle.

C. C-Medulla-inner zone of kidney and contains complete nephrons.

D. D-Cortex-outer part of kidney and do not contain any part of nephrons.

**Answer: A**



**Watch Video Solution**

8. Which one of the following options gives the correct categorization of animals

according to the type of nitrogenous waste they give out?



**Watch Video Solution**

**9.** Which one of the following characteristics is common both in human and adult frogs?

- A. Internal fertilization
- B. Nucleated RBCs
- C. Ureotelic mode of excretion
- D. Four-chambered heart

**Answer: C**



**Watch Video Solution**

**10.** A fall in glomerular filtration rate (GFR) activates

A. Adrenal cortex to release aldosterone

B. Adrenal medulla to release adrenaline

C. Posterior pituitary to release vasopressin

D. Juxtaglomerular cells to release renin

**Answer: D**



**Watch Video Solution**

**11.** The maximum amount of electrolytes and water (70 - 80 percent) from the glomerular filtrate is reabsorbed in which part of the nephron?

A. Proximal convoluted tubule

B. Descending limb of loop of Henle

C. Ascending limb of loop of Henle

D. Distal convoluted tubule

**Answer: A**



**Watch Video Solution**

**12.** Uricotelic mode of passing out nitrogenous wastes is found in

A. Insects and Amphibians



B. Reptiles and Birds

C. Birds and Annelids

D. Amphibians and Reptiles

**Answer: B**



**Watch Video Solution**

**13.** which of the following statement is correct with respect to kidney function regulation?

- A. During summer when body loses a lot of water by evaporation, the release of ADH is suppressed
- B. When someone drinks a lot of water, ADH release is suppressed
- C. Exposure to cold temperature stimulates ADH release
- D. An increase in glomerular blood flow stimulates formation of Angiotensin II.

**Answer: B**



Watch Video Solution

**14.** Which one of the following is a correct pair showing the function of a specific part of the human nephron?

A. Afferent arteriole: carries the blood away from the glomerulus towards renal vein

B. Podocytes: Create minute spaces (slitpores) for the filtration of blood into the Bowman's capsule

C. Henle's loop: most reabsorption of the major substances from the glomerular filtrate

D. Distal convoluted tubule: reabsorption of  $K^+$  ions into the surrounding blood capillaries

**Answer: B**



**Watch Video Solution**

15. Which one of the following statements in regard to the excretion by the human kidneys is correct?

A. Descending limb of loop of Henle is impermeable to water

B. Distal convoluted tubules is incapable in reabsorbing  $HCO_3$

C. Nearly 99 percent of the glomerular filtrate is reabsorbed by the renal tube

D. Ascending limb of loop of Henle is impermeable to electrolytes

**Answer: C**



**Watch Video Solution**

**16.** The principal nitrogenous excretory compound in humans is synthesized

A. In kidneys but eliminated mostly through liver

B. In kidneys as well as eliminated by  
kidneys

C. In liver and also eliminated by the same  
through bile

D. In the liver, but eliminated mostly  
through kidneys

**Answer: D**



**Watch Video Solution**

17. What will happen if the stretch receptors of the urinary bladder wall are totally removed?

A. There will be no micturition

B. Urine will continue to collect normally in bladder

C. Micturition will continue

D. Urine will not collect in the bladder

**Answer: C**



**Watch Video Solution**



**18.** Uric acid is the chief nitrogenous component of the excretory products of

A. Frog

B. Man

C. Earthworm

D. Cockroach

**Answer: D**



**Watch Video Solution**

**19.** Consider the following four statements (i)-(iv) about certain desert animals such as kangaroo rat.

i) They have dark colour and high rate of reproduction and excrete solid urine ii) They do not drink water, breathe at a slow rate to conserve water and have their body covered with thick hairs iii) They feed on dry seeds and do not required drinking water iv)They excrete very concentrated urine and do not use water to regulate body temperature. Out of these four, which two are correct :

A. (iii) and (i)

B. (i) and (ii)

C. (iii) and (iv)

D. (ii) and (iii)

**Answer: C**



**Watch Video Solution**

**20.** A person who is on a long hunger strike and is surviving only on water, will have

A. Less urea in his urine

B. More sodium in his urine

C. Less amino acids in his urine

D. More glucose in his blood

**Answer: A**



**Watch Video Solution**