

PHYSICS

BOOKS - HC VERMA

OPTICAL INSTRUMENTS

Example

1. Two boys one 52 inches tall and the other 55

inches tall, are standing at distances 4.0 m

and 5.0 m respectivley from an eye. Which boy will taller?



Watch Video Solution

2. A compound microscope has an objective of focal length 1 cm and an eyepiece of focal length 2.5 cm. An object has to be placed at a distance of 1.2 cm away from the objective for normal adjustment. a.Find the angular magnification. b.Find the length of the microscope tube.



3. A nearsighted man can clearly see objects up to a distance of 1.5m. Calculate the power of the lens of the spectacles necessary for the remedy of this defect.



Watch Video Solution

Worked Out Examples

1. An object is seen through a simple microscope of focal length 12 cm. Find the angular magnification produced if the image is formed at the near point of the eye which is 25 cm away from it.



Watch Video Solution

2. An object is placed at a distance of 10 cm from a convex lens of focal length 12 cm. Find the position and nature of the image.



Watch Video Solution

3. A small object is placed at distance of 3.6 cm from a magnifier of focal length 4.0 cm. a. find the position of theimage. B. find the linear magnification c. Find the angular magnification.



4. A compound microscope consists of an objective lens of focal length 2.0 cm and an

eyepiece of focal length 6.25 cm separated by a distance of 15 cm. How far from the objective should an object be placed in order to obtain the final image at (a) the least distance of distinct vision (25cm), and (b) at infinity? What is the magnifying power of the microscope in each case?



Watch Video Solution

5. The separation L between the objective (f=0.5cm) and the eyepiece (f=5cm) of a

compound microscope is 7 cm. Where should a small object be placed so that the eye is least strained to see the image? Find the angular magnification produced by the microscope.



Watch Video Solution

6. A astronmical telescope has objective and eyepiece of focal lengths 40 cm and 4 cm respectively To view an object 200 cm away

from the objective ,the lenses must be separated by a distance:



Watch Video Solution

7. A Galilean telescope is constructed by an objective of focal length 50 cm and a eyepiece of focal length 5.0 cm a. find the tube length and magnifying power when it is used to see n object at a large distancei normal adjustment b. if the telescope is to focus an object 2.0 m away from the objective what should be the tube length and angular magnification, the image again forming at infinity?



Watch Video Solution

8. The image of the moon is focused by a converging lens of focal length 50 cm on a plane screen. The image is seen by an unaided eye from a distance of 25 cm. Find the angular magnification achieved due to the converging lens.



9. The near and far points of a person are at 40cm and 250cm respectively. Find the power of the lens he/she should use while reading at 25cm. With this lens on the eye, what maximum distance is clearly visible?



Watch Video Solution

10. A young boy can adjust the power of his eye lens between 50 D and 60 D. His far point

is infinity. A. What is the distance of his retina

from the eye-lens? b. What is the near point?



Watch Video Solution

Question For Short Answer

1. Can virtual image be formed on the retina in a seeing process?



2. Can the image formed by a simple microscope be projected on a screen without using any additional lens or mirror?



Watch Video Solution

3. The angular magnification of a system is less than one. Does it mean that the image formed is inverted?



4. A simple microscope using a single lens often shows coloured image of a white source. Why?



Watch Video Solution

5. A magnifying glass is a converging lens placed close to the eye. A farsighted person uses spectacles having converging lenses. Compare the functions of a converging lens used as a magnifying glass and as spectacles.



6. A person is viewing an extended object. If a converging lens is placed in front of his eyes, will he feel that the size has increased?



Watch Video Solution

7. The magnifying power of a converging lens used as a simple microscope is $\left(1+\frac{D}{f}\right)$. A compound microscope is a combination of two such converging lenses. Why don't we

. In other words, why can the objective not be treated as a simple microscope but the eyepiece can?

have magnifying power $\left(1+\frac{D}{f_0}\right)\left(1+\frac{D}{f_e}\right)$



Watch Video Solution

8. By mistake, an eye surgeon puts a concave lens in place of the lens in the eye after a cataract operation. Will the patient be able to see clearly any object placed at any distance?



9. The magnifying power of a simple microscope is given by $1+\frac{D}{f}$, where D is the least distance for clear vision. For farsighted persons, D is greater than the usual. Does it mean that the magnifying power of a simple microscope is greater for a farsighted person as compared to a normal person? Does it mean that a farsighted person can see an insect more clearly under a microscope than a normal person?



10. Why are the magnification properties of microscopes and telescopes defined in terms of the ratio of angles and not in terms of the ratio of sizes of objects and images?



Watch Video Solution

11. An object is placed at a distance of 30 cm from a converging lens of focal length 15 cm. A normal eye (near point 25 cm, far point

infinity) is placed close to the lens on the other side. (a) Can the eye see the object clearly? (b) What should be the minimum separation between the lens and the eye so that the eye can clearly see the object?



Watch Video Solution

12. A compound microscope forms an inverted image of an object. In which of the following cases it is likely to create difficulties? a. Looking at small germs. b.Looking at circular

spots. c. Looking at a vertical tube containing some water.



Watch Video Solution

Objective 1

1. The size of an object as perceived by an eye depends primarily on

A. A. actual of the object

B. B. distances of the object from the eye

- C. C. aperture of the pupil
- D. D. size of the image formed on the retina

Answer: D



Watch Video Solution

2. The muscles of a normal eye are least

strained when the eye is focused on an object

- A. far away from the eyes
- B. very close to the eye

C. at abot 25 cm from the eye

D. at about 1 m from the eye

Answer: A



Watch Video Solution

3. Why is a normal eye not able to see clearly the objects placed closer than 25 cm?

A. the focal length of the eye is 25 cm

- B. the distance of the retina from the eye lens is 25 cm
- C. the eye is not able to decreae the distance between the eye lens and the retina beyond a limit
- D. the eye is not able to decrease the focal length beyond a limit.

Answer: D



4. When objects at different distances are seen by the eye, which of the following remain constant?

A. the focal length of the eye lens

B. the object distance from the eye lens

C. the radius of curvature of the eye lens

D. the image distance from the the eye lens

Answer: D



5. A person can clearly see objects between 25 cm and 200 cm. Which of the following may represent the range of clear vision for a person B having muscles stronger than A but all other parameters of eye identical to that of A

- A. 25 cm to 200 cm
- B. 18 cm to 200 cm
- C. 25 cm to 300 cm
- D. 1 cm to 300 cm

Answer: B



Watch Video Solution

6. The focal length of a normal eye lens is about

A. 1 mm

B. 2 cm

C. 25 cm to 300 cm

D. 1 m

Answer: B



Watch Video Solution

7. The distance of the eye lens from the retina is x. For a normal eye, the maximum focal length of the eye lens

$$A. = x$$

B.
$$< x$$

$$\mathsf{C.} > x$$

D.
$$=2x$$

Answer: A



Watch Video Solution

8. A man wearing glasses of focal length +1 m cannot clearly see beyond 1 m. What does it signify?

A. if he is farsighted

B. if he is nearsighted

C. if his vision is normal

D. in each of these cases

Answer: D



Watch Video Solution

9. An object is placed at a distance u from a simple microscope of focal length f. The angular magnification obtained depends

A. on f but not on u

B. on u but not on f

C. on f as well as u

D. neither on f nor on u

Answer: C



- **10.** To increase the angular magnification of a simple microscope, one should increase
 - A. the focal length of the lens
 - B. the power of the lens
 - C. the apeture of the lens
 - D. the object size

Answer: B



Watch Video Solution

11. A man is looking at a small object placed at his near point. Without altering the position of lhis eye or the object, he puts a simple microscope of magnifying power 5x before his eyes. The angular magnification achieved is

A. 5

B. 2.5

C. 1

D. 0.2

Answer: C



Watch Video Solution

Objective 2

1. When we see an object the image formed on the retina is

B. virtual

C. erect

D. inverted.

Answer: A::D



Watch Video Solution

2. Compound microscope, astronomical telescope, simple microscope, flood light.

- A. simple microscope
- B. compound microscope
- C. Astronomical telescope
- D. Galilean telescope

Answer: A::D



Watch Video Solution

3. The maximum focal length of the eye lens of a person is greater than its distance from the retina. The eye is

A. always strained in looking at an object

B. strained for objects at large distances only

C. strained for objects at short distances only

D. unstrained for all distances

Answer: A



- 4. Mark the correct options.
 - A. A) If the far point goes ahead, the power of the divergent lens should be reduced
 - B. B) If the near point goes ahead, the power of the convergent lens should be reduced.
 - C. C) If the far point 1 m away form the eye, divergent lens should be used.

D. D) If the near point is 1 m away from the eye, divergent lens should be used

Answer: A::C



Watch Video Solution

5. The focal length of the objective of a compound microscope is f_0 and its distance from the eyepiece is L. The object is placed at a distance u from the objective. For proper working of the instrument.,

A.
$$L < u$$

$$\mathtt{B.}\,L>u$$

C.
$$f_0 < L < 2f_0$$

D.
$$L>2f_0$$

Answer: B::D



Watch Video Solution

Exercises

1. An object is to be seen through a simple microscope of focal length 12 cm. Where should the object be placed so as to produce maximum angular magnification? The least distance for clear vision is 25 cm.



Watch Video Solution

2. A simple microscope has a magnifying power of 3.0 when the image is formed at the near point (25 cm) of a normal eye. (a) What is

its focal length? (b) What will be its magnifying power if the image is formed at infinity?



Watch Video Solution

3. A child has near point at 10 cm. What is the maximum angular magnification the child can have with a convex lens of focal length 10 cm?



4. A simple microscope is rated 5 X for a normal relaxed eye. What Will be its magnifying power for a relaxed farsighted eye whose near point is 40 cm?



Watch Video Solution

5. Find the maximum magnifying power of a compound microscope having a 25 diopter lens as the objective, a 5 diopter lens as the eyepiece and the separation 30 cm between

the two lenses. The least distance for clear vision is 25 cm.



Watch Video Solution

6. The separation between the objective and the eyepiece of a compound microscope can be adjusted between 9.8 cm to 11.8 cm. If the focal lengths of the objective and the eyepiece are 1.0 cm and 6 cm respectively, find the range of the magnifying power if the image is always needed at 24 cm from the eye.

7. In a compound microscope, the objective and eyepiece have focal lengths 0.95 cm and 5 cm respectively and are kept at a distance of 20 cm .the final image is formed at a distance of 25 cm from the eyepiece. Calculate the position of the object and the total magnification .



8. A compound microscope has a magnifying power of 100 when the image is formed at infinity. The objective has a focal length of 0.5 cm and the tube length is 6.5 cm. What is the focal length of the eyepiece.



Watch Video Solution

9. A compound microphone has an objective of focal length 1 cm and an eyepiece of focal length 2.5 cm. An object has to be placed at a

distance of 1.2 cm away from the objective for normal adjustment. a.Find the angular magnification. b.Find the length of the microscope tube.



Watch Video Solution

10. An optical instrument used for angular magnification has a 25 D objective and a 20D eyepiece. The tube length is 25 cm when the eye is least strained . Whether it is a microscope or a telescope?

11. An astronimical telescope is to be designed to have a magnifying power of 50 in normal adjustment. If the length of the tube is 102 cm, find the powers of the objective and the eyepiece.



12. The eyepice of an astronomicasl telescope has a focasl length of 10 cm. The telescope is

focussed for normal vision of distant objects when the tube length is 1.0 m. find the focal length of the bjective and the magnifying power of the telescope.



Watch Video Solution

13. A Galilean telescope is 27 cm long when focussed to form an image at infinity. If the objective has a focal length of 30 cm, what is the focal length of the eyepiece?



14. A farsighted person cannot see objects placed closer to 50 cm. Find the power of the lens needed to see the objects at 20 cm.



Watch Video Solution

15. A nearsighted person cannot clearly see beyond 200 cm. Find the power of the lens needed to see objects at large distances.



16. A person wears glasses of power -2.5D. Is the person short sighted or long sighted ? What is the far point of the person without glasses ?



Watch Video Solution

17. A professor reads a greeting card received on his 50th birthday with + 2.5 D glasses keeping the card 25 cm away. Ten years later, he reads his farewell letter with the same

glasses but he has to keep the letter 50 cm away. What power of lens should he now use?



Watch Video Solution

18. A normal eye has retina 2 cm behind the eye-lens. What is the power of the eye-lens when the eye is (a) fully relaxed, (b) most strained?



19. The near point and the far point of a child are at 10 cm and 100 cm. If the retina is 2.0 cm behind the eye-lens, what is the range of the power of the eye-lens?



Watch Video Solution

20. A nearsighted person cannot see beyond 25 cm. Assuming that the separation of the glass from the eye is 1 cm, find the power of lens needed to see distant objects.

Watch Video Solution

21. A person has near point at 100 cm. What power of lens is needed to read at 20 cm if he/she uses (a) contact lens, (b) spectacles having glasses 2.0 cm separated from the eyes?



22. A lady uses + 1.5 D glasses to have normal vision from 25 cm onwards. She uses a 20 D

lens as a simple microscope to see an object. Find the maximum magnifying power if she uses the microscope (a) together with her glass (b) without the glass. Do the answers suggest that an object can be more clearly seen through a microscope without using the correcting glasses?



Watch Video Solution

23. A lady cannot see objects closer than 40cm from the left eye and closer than 100cm from

the right eye. While on a mountaining trip, she is lose from her team. She tries to make an astronomical trip, from her reading glasses to look from her teammates. (a) Which glass should she use as the eyepiece? (b) What magnification can she get with relaxed eye?

