





PHYSICS

BOOKS - HC VERMA

THERMAL AND CHEMICAL EFFECT OF ELECTRIC CURRENT



1. Find the heat developed in each of the three resistors shown in figure in 1 minute.





2. Calculate the electric current required to deposit 0.972g of chromium in three hours. ECE of chromium is $0.00018gC^{-1}$



1. A current of 30A is registered when the terminals of a dry cell of EMF 1.5 VOLTS are connected through an ammeter. Neglecting the meter resistance, find the amount of heat produced in the battery in 10 seconds.

Watch Video Solution

2. A room heater is rated 500W, 220V. (a) Find the resistance of its coil. (b) If the supply voltage drops to 200V, what will be the power consumed? (c) If an electric bulb rated 100W, 220V is connected in series with this heater, what will be the power consumed by the heater and by the bulb when the supply is at 220V?



3. A battery of ε and internal resistance r is used in a circuit with a variable external resistance R. Find the value of R for which the power consumed in R is maximum.

Watch Video Solution

4. The junctions of a Ni - Cu thermocouple are maintained at $0^{\circ}C$ and $100^{\circ}C$. Calculate the Seebeck emf produced in the loop. $a_{Ni,Cu} = 16.3 \times 10^{-6} V^{\circ} C^{-1}$ and $b_{Ni,Cu} = -0.042 \times 10^{-6} V^{\circ} C^{-3}$

Watch Video Solution

5. Find the neutral and inversion temperatures for Ni - Cu thermocople with the cold junction at $0^{\circ}C$. Use data from previous example.

Watch Video Solution

6. An electric current of 0.4A is passed through a silver voltameter fro half an hour.Find the amount of silver deposited on the cathode. ECE of silver $= 1.12 \times 10^{-4} kgC^{-1}$

Watch Video Solution

7. A silver and a copper voltameter are connected in series with a 12.0V battery of negligible resistance. It is found that 0.806g of silver is deposited in half an hour.Find (a) the mass of the copper deposited and (b) the energy supplied by the battery. *ECE* of silver $= 1.12 \times 10^{-6} kgC^{-1}$ and that of copper $= 6.6 \times 10^{-7} kgC^{-1}$

Watch Video Solution

8. A current of 1A is possed through a dilute solution of sulphuric acid for some time to liberate 1g of oxygen. How much hydrogen is liberated during this period? How long

 $= 96500 Cmol^{-1}.$

Watch Video Solution



1. If a constant potential difference is applied across a bulb, the current slightly decreases as time passes and then becomes constant. Explain.



2. Two unequal resistances R_1 and R_2 ae connected across two identical batteries of $emf \varepsilon$ and internal

resistance r(figure). Can the thermal energies developed in R_1 and R_2 be equal in a given time. If yes, what will be the condition? (Figure Question)



Watch Video Solution

3. When a current passes through a resistor, its

temperature increases. Is it an adiabatic process?



4. Apply the first law of thermodynamics to a resistor carrying a current *i*. Identify which of the quantities ΔQ , ΔU and ΔW are zero, which are positive and which are negative.



5. Do all the thermocouples have a neutral temperature?

Watch Video Solution

6. Is inversion temperature always double of the neutral temperature? Does the unit of temperature have an effect in deciding this question?



7. Is neutral temperature always the arithmetic mean of the inversion temperature and the temperature of the cold junction? Does the unit of temperature have an effect in deciding this question?

Watch Video Solution

8. Do the electrodes in an electrolytic cell have fixed

polarity like a battery?



9. As temperature increases, the vicosity of liquids decrease considerably. Will this decrease the resistance of an electrolyte as the temperature increases?

Watch Video Solution			
Objective 1			

1. Which of the following plotes may represent the thermal energy produced in a resistor in a given time as a

function of the electric current? Figure





2. A constant current i is passed through a resistor. Taking the temperature coefficient of resistance into account, indicate which of the plots shown in Figure best represents the rate of production of thermal energy in

the resistor



3. Consider the following statements regarding a thermocouple. (A) The neutral temperature does not depend on the temperature of the cold junction. (B) The inversion temperature does not depend on the temperature of the cold junction.

A. Both A and B are correct.

B. A is correct but B is wrong

C. B is correct but A is wrong

D. Both A and B are wrong

Answer: B

Watch Video Solution

4. The heat developed in a system is proportional to the current through it.

A. It cannot be Thomson heat.

B. It cannot be Peltier heat.

C. It cannot be Joule heat.

D. It can be any of the three heats mentioned above.

Answer: C



5. Consider the following two statements (A) Free-electron density is different in different metals. (B) Free-electron density in a metal depends on temperature.

Seebeck effect is caused

A. due to both A and B

B. due to A but not due to B

C. due to B but not due to A

D. neither due to A nor due to B

Answer: A



6. Consider the statements A and B in the previous queston. Peltier effect is caused by

A. Free electron density is different in different metals.

B. Free electron density in a metal depends on temperature.

A. due to both A and B

B. due to A but not due to B

C. due to B but not due to A

D. neither due to A nor due to B

Answer: B



7. Consider the statement A and B . Thomson effect is caused by

(A) Free-electron density is different in different metals.

(B) Free electron density in a metal depends on temperature.

A. due to both A and B

B. due to A but not due to B

C. due to B but not due to A

D. neither due to A nor due to B

Answer: C



- 8. Faraday constant
 - A. depends on the amount of the electrolyte
 - B. depends on the current in the electrolyte
 - C. is a universal constant
 - D. depends on the amount of charge passed through

the electrolyte

Answer:

Objective 2

 Two resistors having equal resistances are joined in series and a current is passed through the combination.
 Neglect any variation in resistance as the temperature changes. In a given time interval,

A. equal amounts of thermal energy must be produced

in the resistors

B. unequal amount of thermal energy may be produced

C. the temperature must rise equally in the resistors

D. the temperature may rise equally in the resistors.

Answer: A::D



2. A copper strip AB and an iron strip AC are joined at A. The junction A is maintained at $0^{\circ}C$ and the free ends B and C are maintained at $100^{\circ}C$. There is a potential difference between

A. the two ends of the copper strip

B. the copper end and the iron end at the junction

C. the two ends of the iron strip

D. the free ends B and C.

Answer: A::B::C::D



3. The constants a and b for the pair silver-lead are $2.50\mu V^{\circ}C^{-1}$ and $0.012\mu V^{\circ}C^{-2}$ respectively. For a silver-lead thermocouple with colder junction at $0^{\circ}C$,

A. there will be no neutral temperature

B. there will be no inversion temperature

C. there will not be any thermo-emf even if the

junctions are kept at different temperatures

D. there will be no current in the thermocouple even if

the junctions are kept at different temperature.

Answer: A::B



4. An electrolysis experiment is stopped and the battery terminals are reversed.

A. The electrolysis will stop

B. The rate of liberation of material at the electrodes

will increase.

C. The rate of liberation of material will remain the same.

D. Heat will be produced at a greater rate.

Answer: C

Watch Video Solution

5. The electrochemical equivalent of a material depends on

A. the nature of the material

B. the current through the electrolyte containing the material

- C. the amount of charge passed through the electrolyte
- D. the amount of this material pressent in the electrolyte.



2. A coil of resistance 100Ω is connected across a battery of emf6.0V. Assume that the heat developed in the coil

is used to raise its temperature. If the heat capacity of the coil is $4.0JK^{-1}$, how long will it take to raise the temperature of the coil by $15^{\circ}C$?



3. The specification on a heater coil is 250V, 500W. Calculate the resistance of the coil. What will be the resistance of a coil of 1000W to operate at the same voltage?



4. A heater coil is to be constructed with a nichrome wire

 $\left(
ho=1.0 imes10^{-6}\Omega m
ight)$ which can operate at 500W when

connected to a 250V supply. (a) What would be the resistance of the coil? (b) If the cross-sectional area of the wire is $0.5mm^2$, what length of the wire will be needed? (c) If the radius of each turn is 4.0mm, how many turns will be there in the coil?



5. A bulb with rating 250V, 100W is connected to a power supply of 220V situatd 10m away using a copper wire of area of cross section $5mm^2$. How much power will be consumed by the connecting wire? Resistivity of copper $= 1.7 \times 10^{-8}\Omega m$. **6.** An electric bulb, when connected across a power supply of 220V, consumes a power of 60W. If the supply drops supply is suddently increased to 240V, what will be the power consumed?

Watch Video Solution

7. A servo voltage stabiliser restricts the voltage output to $220V \pm 1 \%$. If an electric bulb rated at 220V, 100W is connected to it, what will be the minimum and maximum power consumed by it?

Watch Video Solution

8. An electric bulb marked 220V, 100W will get fused if it is made to consume 150W or more. What voltage fluctuation will the bulb withstand?

Watch Video Solution

9. An immersion heater rated 1000W, 220V is used to heat $0.01m^3$ of water. Assuming that the power is supplied at 220V and 60% of the power supplied is used to heat the water, how long will it take to increase the temperature of the water from $15^{\circ}C$ to $40^{\circ}C$?

Watch Video Solution

10. An electric kettle used to prepare tea, takes 2 minutes to boil 4 cups of water (1 cup contains 200 cc of water) if the room temperature is $25^{\circ}C$, (a) If the cost of power consumption is Re1.00 per unit (1unit = 1000wa - hour), calculate the cost of boiling 4 cups of water. (b) What will be the corresponding cost if the room temprature drops to $5^{\circ}C$?



Watch Video Solution

11. The coil of an electric bulb takes 40 watts to start glowing. If more than 40 W is supplied, 60% of the extra power is converted into light and the remaining into heat. The bulb consumes 100 W at 220 V. Find the percentage

drop in the light intensity at a point if the supply voltage

changes from $220V \rightarrow 200V$.



12. The 2.0 Ω resistor show in figure is dipped into a calorimeter containing water. The heat capacity of the calorimeter together with water is $2000JK^{-1}$. (a) If the circuit is active for 15 minutes, what would be the rise in the temperature of the water? (b) Suppose the 6.0 Ω resistor gets burnt. What would be the rise in the

temperature of the water in the next 15 minutes? (Figure)



13. The temperatures of the junctions of a bismuth-silver thermocouple are maintained at $0^{\circ}C$ and $0.001^{\circ}C$. Find the thermo-emf (Seebeck emf) developed. For bismuth-

silver,

$$a = -46 imes 10^{-6} V^{\,\circ} C^{\,-1} \, ext{ and } \, b = -0.48 imes 10^{-6} V^{\,\circ} C^{\,-2}$$



14. Find the thermo-emf developed in a copper-silver thermocopule when the junctions are kept at $0^{\circ}C$ and $40^{\circ}C$. Use the data in table.

Watch Video Solution

15. Find the neutral temperature and inversion temperature of copper-iron thermocouple if the reference junction is kept at $0^{\circ}C$. Use the data in table.



16. Find the charge required to flow through an electrolyte to liberate one atom of (a) a monovalent material and (b) a divalent material.

Watch Video Solution

17. Find the amount of silver liberated at cathode if

0.500A of current is passed through $AgNO_3$ electrolyte

for 1 hour. Atomic weight of silver s $107.9 gmol^{-1}$



18. An electroplating unit plates 3.0g of silver on a brass plate in 3.0 minutes. Find the current used by the unit. The electrochemical equivalent of silver is $1.12 \times 10^{-6} kg C^{-1}$.

Watch Video Solution

19. Find the time requried to liberate 1.0 litre of hydrogen

at STP in an electrolytic cell by a current of 5.0A.



20. Two voltameters, one having a solution of silver salt and the other of a trivalent-metal salt, are connected in

series and a current of 2A is maintained for 1.50hours. It is found that 1.00g of the trivalent metal is deposited. (a) What is the atomic weight of the trivalent metal? (b) How much silver is deposited during this period? Atomic weight of silver is $109.9gmol^{-1}$

C

Watch Video Solution

21. A brass plate having surface area $200cm^2$ on one side is electroplated with 0.10mm thick silver layers on both sides using a 15A current. Find the time taken to do the job. The specific gravity of silver is 10.5 and its atomic weight is $107.9gmol^{-1}$

Watch Video Solution

22. Figure, shows an electrolyte of AgCI through which a current is passed. It is observed that 2.68g of silver is deposited in 10 minutes on the cathode. Find the heat developed in the 20Ω resistor during this period. Atomic weight of silver is $107.9gmol^{-1}$. (Figure)



Watch Video Solution

23. The potential difference across the terminals of a battery of emf12V and internal resistance 2Ω drops to 10V when it is connected to a silver voltameter. Find the silver deposited at the cathode in half an hour. Atomic weight of silver is $107.9gmol^{-1}$.



24. A plate of area $10cm^2$ is to be electroplated with copper ($density9000kgm^{-3}$ to a thickness of 10 micrometres on both sides, using a cell of 12V. Calculate the energy spend by the cell in the process of deposition. If this energy is used to heat 100g of water, calculate the rise in the temperature of the water. ECE of copper

 $= 3 imes 10^{-7} kg C^{-1}$ and specific heat capacity of water

$$= 4200 J k g^{-1} K^{-1}.$$

Watch Video Solution