




CHEMISTRY

BOOKS - MS CHOUHAN





ALDEHYDES AND KETONES

Level 1



, Product A


is :



Answer: A



Watch Video Solution



Above conversion can be achieved by :

A. A) Wolf-Kishner reduction


B. B) Clemmensen reduction

C. C) $LiAlH_4$

D. D) $NaBH_4$

Answer: B

 [Watch Video Solution](#)





3.

Above conversion can be achieved by :

- A. Wolf-Kishner reduction
- B. Clemmensen reduction
- C. $\text{HS} - \text{CH}_2 - \text{CH}_2 - \text{SH}_2$ following by Raney Ni
- D. None of these

Answer: D


 [Watch Video Solution](#)




4.

, Product


of the Clemmensen reduction is :




A. A)



B. B)




C. C)



D. D)

Answer: C

 Watch Video Solution



Above conversion can be achieved by :

A. Wolf-Kishner reduction


B. Clemmensen reduction

C. $LiAlH_4$

D. $NaBH_4$

Answer: A

 Watch Video Solution



Identify product (A) & (B) from the given product P, Q, R:

A. A = P, B = Q




B. A = Q, B = R

C. A = Q, B = P


D. A = R, B = P

Answer: C


 Watch Video Solution



C.





D.



Answer: B


 Watch Video Solution

8.



, Product


(A) is :




A.



B.



C.



D.

Answer: B

 Watch Video Solution

9.



, Above

conversion can be carried out by :

A. Clemmensen reduction

B. Wolf-Kishner reduction


C. $LiAlH_4$

D. $NaBH_4$

Answer: B

 [Watch Video Solution](#)

10. Increasing order of equilibrium constants for the formation of a hydrate:



A. $IV < III < II < I$


B. $IV < III < I < II$

C. $I < II < III < IV$





D. $II < III < I < IV$

Answer: C

 Watch Video Solution




(A) is :




Answer: C

 Watch Video Solution




Predict the product of hydrolysis of the above molecule.


A.




B.



C.




D.







Answer: C

 [Watch Video Solution](#)




13. _____, This conversion can be achieved by :



Answer: B

 [Watch Video Solution](#)




Compound (A) & (B) can be differentiated by :


- A. 2-4-DNP
- B. Fehling solution
- C. Lucas reagent
- D. $NaHSO_3$

Answer: B


 [Watch Video Solution](#)




(A) is :




A.



B.




C.



D.

Answer: B

 Watch Video Solution




Reagents to carry out above conversion, P, Q, R respectively are :


A. $H_2C = CH - CH_2 - Br$, (HO^\ominus) , $[HO^\ominus, \Delta]$, Wacker-process

B. $H_2C = CH - CH_2 - Br$, (HO^\ominus) , Wacker-process, HO^\ominus, Δ

C. Wacker process,




D. Wacker process,



Answer: B





 Watch Video Solution

17.



Above reaction is a Baeyer Villiger rearrangement of an asymmetric

ketone with magnesium mono peroxy phthalate hexahydrate (in the drawing, Mg^{+2} is omitted for clarity) Identify major product.



Answer: B

 Watch Video Solution



18.


Above compounds can be differentiated by following reagent:

- A. 2-4 DNP (Brady reagent)
- B. Tollen's reagent
- C. Lucas reagent
- D. $NaHSO_3$

Answer: B







[Watch Video Solution](#)




19.

Product (D) will be :



Answer: B

20.



Relationship between products (Q) and (S) is:

- A. Positional isomer
- B. Chain isomer
- C. Stereoisomer
- D. Functional isomer





Answer: D

21. In the reaction,



, the product


(X) is :



Answer: B

Watch Video Solution

22. Rank the following in order of increasing value of the equilibrium constant for hydration, K_{hyd} . (smallest value first).



A. $1 < 2 < 3$


B. $3 < 1 < 2$

C. $2 < 1 < 3$

D. $2 < 3 < 1$

Answer: B

 [Watch Video Solution](#)



23.

Above conversion can be achieved by :

A. $Zn(Hg), HCl$

B. $NH_2 - NH_2 / KOH / \Delta$








C. $LiAlH_4$

D. H_2 / Ni


Answer: B

 [Watch Video Solution](#)

24. Which sequence represents the best synthesis of hexanal ?




 Watch Video Solution




25. _____, Product

(B) in this reaction is :

A. A)






B. B)




C. C)







D. D)

Answer: C

 Watch Video Solution




Compound (x) in the above reaction is :







Answer: B

 Watch Video Solution




Product (C) of the above reaction is :




Answer: B

 [Watch Video Solution](#)




(B) of the reaction is :







Answer: A

 [Watch Video Solution](#)





29.

Product (A) of the above reaction is (bromination occur not in the benzene ring) :




Answer: B




30.

Positive


Tollens test Compound (A) is :




A. A)



B. B)




C. C)






D. D)

Answer: C

 Watch Video Solution




(A) in this reaction is :







D. both (b) and ©

Answer: D

32.






Compound (P) is :



Answer: C

33. Correct order of reactivity of following compounds towards Grignard reagent?



A. A) $I > II > III$


B. B) $II > I > III$

C. C) $II > III > I$

D. D) $I > III > II$


Answer: B

 Watch Video Solution




34. _____, Product


(A) is :




A.



B.



C.




D.

Answer: B

 Watch Video Solution

35.



The number of products obtained in above reaction is :

A. 2

B. 3


C. 4

D. 5

Answer: A

 Watch Video Solution


36. What reagent and/or reaction conditions would you choose to bring about the following conversion?



- A. 1. LiAlH_4 , 2. H_2O
- B. H_2O , H_2SO_4 , heat
- C. H_2O , NaOH , heat
- D. PCC, CH_2Cl_2

Answer: B

 Watch Video Solution




Mel excess
KH excess

→ 81% yield ;


37.

, Product


of the reaction is :




A.



B.




C.



D.

Answer: C

 [Watch Video Solution](#)



38.

The above reduction can be best carried out by :

A. A) Clemmensen reduction



B. B) Wolf-Kishner reduction

C. C) NaBH_4

D. D) None of these

Answer: D

 [Watch Video Solution](#)



Product (A) and (B) is differentiated by :

A. 2-4-DNP


B. NaOI

C. Na-metal




D. $NaHSO_3$


Answer: B

 [Watch Video Solution](#)



End product (C) in above reaction is :








D.


Answer: A

 Watch Video Solution

41. Compound (X) C_4H_8O , which reacts with 2,4-DNP derivative and gives negative haloform test is :



C. C)



Answer: B

 [Watch Video Solution](#)


42. When a nucleophile encounters a ketone, the site of attack is :

- A. A) the carbon atom of the carbonyl
- B. B) the oxygen atom of the carbonyl
- C. C) both the carbon and oxygen atoms, with equal probability
- D. D) no attack occurs as ketones do not react with nucleophiles


Answer: A

 [Watch Video Solution](#)


43. The correct order of rate of reaction toward nucleophilic addition reaction:




(a)




(b)



(c)



(d)



(e)

A. A) $a > b > c > d > e$

B. B) $a > b > d > c > e$

C. C) $a > d > e > b > c$

D. D) $a > b > e > d > c$

Answer: A




Watch Video Solution

44.

The

structure



would be




best classified as a (an) :

- A. A) Acetal
- B. B) Hemiacetal
- C. C) Hydrate
- D. D) Cyanohydrin

Answer: B




45. Which of the following pairs of reactants is most effective in forming an enamine ?






D. None of these form an enamine.

Answer: B


46. The reaction of $\text{C}_6\text{H}_5\text{CH} = \text{CHCHO}$ with LiAlH_4 gives :







Answer: A

 Watch Video Solution



(B) of the reaction is :






B.



C.




D.


Answer: B

 [Watch Video Solution](#)


48. Which of following compound is hemiacetal ?



A.



B.






C.

D. all of these

Answer: D


 [Watch Video Solution](#)

49. $Ph - CH_2 - C \equiv N \xrightarrow[THF]{LDA} \xrightarrow{CH_3I} 71\%$, End product of the reaction will be :





Answer: C

 Watch Video Solution




Above conversion can be achieved by :




Answer: B

 Watch Video Solution




Products,

Product of the reaction is/are :



A.




B.

C. HCHO

D. Both (a) and (c)

Answer: D


 Watch Video Solution




52.

, Product


(A) of the reaction is :




A.



B.




C.



D.


Answer: B

 [Watch Video Solution](#)




, Product


(C) of the reaction is :




A.



B.




C.



D.

Answer: B

 Watch Video Solution



Product (A) of the reaction will be :



Answer: C

 [Watch Video Solution](#)

55. $R - \overset{\overset{O}{||}}{C} - H \xrightarrow{R-NH_2} R - CH = N - R$. This reaction gives best yield at :





- A. A) pH 1 - 2
- B. B) pH 4 - 5
- C. C) pH 10 - 11
- D. D) pH 13 - 14

Answer: B

 [Watch Video Solution](#)

56. An aromatic compound A of the molecular formula $C_8H_{10}O$ on reaction with iodine and dilute NaOH gives a yellow precipitate. The

structure of the compound is expected to be :







Answer: B

 [Watch Video Solution](#)

57. Compound 'A' (molecular formula C_3H_8O) is treated with acidified potassium dichromate to form a product 'B' (molecular formula C_3H_6O). 'B' forms a shining silver mirror on warming with ammoniacal silver nitrate. 'B' when treated with an aqueous solution of

$H_2NCONHNH_2$, HCl and sodium acetate gives a product 'C'.


Identify the structure of C :



Answer: A

 Watch Video Solution

58. In the reaction, the acid obtained will be :



A. A) D-isomer

B. B) L-isomer


C. C) (80%D + 20%L) mixture

D. D) (50%D + 50%L) mixture

Answer: D

 [Watch Video Solution](#)

59. In the following sequence :



A. A) $CH_3CH_2CH_2NH_2$


B. B) $CH_3CH_2CH_2CONHCH_3$

C. C) $CH_3CH_2CH_2NHCOCCH_3$





D. D) $CH_3CH_2CH_2CONHCOCCH_3$

Answer: C

 [Watch Video Solution](#)






Product (G) is :



Answer: C


 Watch Video Solution

61. Carbonyl compounds can generally be converted to hydrocarbons by:



Answer: C


 Watch Video Solution




62.

, Product


(A) is :




A.



B.



C.



D.

Answer: A

 [Watch Video Solution](#)




63. Which statement about the aldol condensation is correct ?

- A. A Lewis acid is commonly used as a catalyst
- B. The initial step is probably the formation of a carbanion
- C. A Lewis base is employed to induce carbocation formation
- D. The carbon chain is lengthened through the elimination of 1 mole of water.

Answer: B

 [Watch Video Solution](#)





64. A compound gives a positive test with $I_2 / NaOH$ and is extracted from benzene by saturated $NaHSO_3$. It may be :



Answer: B

 [Watch Video Solution](#)


65. Which of the following compounds on reaction with excess CH_3MgBr and subsequent hydrolysis will give a tertiary alcohol?



D.


Answer: B




 Watch Video Solution



66.


Reactant (A) and (B) is :





Answer: D


 Watch Video Solution



Product (A) is :




A.



B.

C. 





D.

Answer: B

 [Watch Video Solution](#)

68. Which of the following does not form a stable hydrate on addition of H_2O ?






B.



C.




D.

Answer: D

 Watch Video Solution


69. Consider the following sequence of reactions.




The ketone (A) is :




A.



B.



C.




D.

Answer: B




[Watch Video Solution](#)

70. In the reaction,




, the product


(X) is :




A.



B.



C.



D.

Answer: C

 [Watch Video Solution](#)



71. The conversion of acetophenone into benzoic acid can be achieved by its reaction with :

- A. sodium hydroxide followed by acidification
- B. iodine and sodium hydroxide, followed by acidification
- C. hydroxylamine followed by reaction with H_2SO_4
- D. m-chloroperoxobenzoic acid

Answer: B

 [Watch Video Solution](#)


72. In which of the following compounds the methylenic hydrogens are the most acidic?







Answer: D

 Watch Video Solution


73. Which is the major product of the following reaction ?







Answer: D

 Watch Video Solution



Product (B) is :



Answer: A

 [Watch Video Solution](#)

75. The presence of unsaturation in organic compounds can be tested with :

A. Schiff's reagent

B. Tollen's reagent




C. Fehling's reagent

D. Baeyer's reagent

Answer: D

 Watch Video Solution

76. Which of the following gives iodoform test?





D. None of these

Answer: A

 Watch Video Solution


77. Which of the following *beta*-keto carboxylic acid does not undergo decarboxylation on heating?



D. None of these


Answer: A




 Watch Video Solution



78.

Product (D) is :







Answer: A

 [View Text Solution](#)


79. The compound shown in the below undergoes racemization on reaction with aqueous acid.




Which of the following structures best represents the intermediate responsible for this process ?




A.



B.



C.








D.

Answer: C

 Watch Video Solution

80. The final product of the following sequence of reaction is :







Answer: C


 Watch Video Solution

81. The amino ketone shown below undergoes a spontaneous cyclization on standing. What is the major product of this intramolecular reaction ?







A.



B.



C.







D.

Answer: D

 [Watch Video Solution](#)


82. Compound (A) $C_6H_{12}O$ is optically active. Compound (A) give negative Tollens test and positive test with 2-4-di-nitro phenyl

hydrazine. Identify A.



Answer: B





 [Watch Video Solution](#)



83.


, Product

(A) of the reaction is :




Answer: B


 Watch Video Solution



Product (Q) will be:




A.



B.




C.







D.

Answer: B






Product (B) of above the reaction is :



Answer: B




86. _____, Product

obtained in the reaction is :

- A. Diastereomer
- B. Racemic
- C. Meso
- D. Optically pure enantiomer


Answer: A

 [Watch Video Solution](#)




87. , Product (


C) of the reaction is :



B.




C.



D.


Answer: B



 Watch Video Solution




88.

Product (A) and (C) is :






C.







Answer: C

 Watch Video Solution




Product (B) is :





Answer: B

 Watch Video Solution



90.

Product (P) is :

A. Hemiacetal


B. Acetal

C. Alcohol

D. Alkane

Answer: B

 [Watch Video Solution](#)







(Oxallyl cation)

91.


, Product


of rearrangement is :



Answer: C

 [Watch Video Solution](#)



Reactant	K_{eq}
PhCHO	a
	b
$\text{Ph}-\overset{\text{O}}{\parallel}{C}-\text{CH}_3$	c
$\text{CH}_3-\overset{\text{O}}{\parallel}{C}-\text{H}$	d

92.

The correct order of decreasing value of K_{eq} is :

A. $a > b > c > d$

B. $d > a > b > c$






C. $d > b > a > c$

D. $d > a > c > d$

Answer: B

 Watch Video Solution





93. Product (B) of the given reaction is :




Answer: B

 Watch Video Solution

94. End product (C) of the reaction is :







D.

Answer: B


 Watch Video Solution



Compound (A) will be :







Answer: C



96.

, Product


of the reaction is :







Answer: C

 Watch Video Solution


97. Which pair of reactants compounds may be used to make given acetal ?









Answer: D

 Watch Video Solution




98. (A) , (A) & (B)

are isomers , Isomer (B) is :



Answer: B

 Watch Video Solution



99.

(A) and (B) is differentiated by:

A. NaH

B. 2-4 DNA

C. Tollen's reagent

D. $NaHSO_3$

Answer: C

 Watch Video Solution

100. Which of the following pairs cannot be differentiated by Tollen's reagent ?

A. Benzaldehyde and benzyl alcohol

B. Hexanal and 2-hexanone


C. 2-Hexanol and 2-hexanone




D. Pentanal and diethyl ether

Answer: C

 [Watch Video Solution](#)

101. An optically active compound $C_6H_{12}O$ gives positive test with 2, 4-dinitrophenyl hydrazine, but negative with Tollens' reagent, what is the structure of the compound ?









Answer: C

 Watch Video Solution

102. Compound (A) $\text{C}_6\text{H}_{12}\text{O}_3$, when treated with I_2 in aqueous sodium hydroxide gives yellow precipitate. When A is treated with Tollens reagent no reaction occur. When A is hydrolysed and then treated with Tollens reagent, a silver mirror is formed in test tube.


Compound (A) will be :







Answer: C

 Watch Video Solution




(C) obtained is :





C.







D.

Answer: C

 Watch Video Solution


104. Which of following does not react with $NaHSO_3$ (sodium bisulphite)?








Answer: C

 Watch Video Solution




(A) is :





C.





D.

Answer: C


 Watch Video Solution

106.






, Product

(A) is :




A.




Answer: B


Watch Video Solution




(B) in this reaction is :




A.



B.



C.






D.

Answer: A

 [Watch Video Solution](#)




Arrange the following reagent in the correct order in which above transformation is carried out :






Answer: C

 Watch Video Solution




Compound (C) can show geometrical isomerism. Product (E) of the reaction will be:



Answer: C

 [Watch Video Solution](#)

110. Arrange in their increasing order of equilibrium constants for hydration ?



A. $A < B < C < D < E$

B. $A < C < B < E < D$


C. $A < C < E < B < D$





D. $C < A < B < E < D$

Answer: B

 [Watch Video Solution](#)


111. End products of the following sequence of reactions are :






Answer: C




 Watch Video Solution



112. _____, Product


(P) of the reaction will be :





Answer: B

 Watch Video Solution



113. . Products

of the reaction are :

A. Racemic mixture

B. Diastereomers

C. Meso

D. Mixture of meso compound and optically active compound




Answer: B

 Watch Video Solution



114.




∴ reactant (A) is :




Answer: C



115. $CH_3CH_2 - \overset{O}{\parallel} C - CH_3 \xrightarrow[HCl]{NaNO_2}$, Major product of this reaction is :





C.







D.

Answer: C




116. . Product

(A) & (B) are :




Answer: A

Watch Video Solution




3, 4-Methylene
dioxybenzaldehyde




117. . Product


(B) is :




A.



B.





C.



D.

Answer: A


Watch Video Solution




. Product

118.


(A) is :



A.



B.




C.



D.




Answer: A


 Watch Video Solution



119. _____ . Product


(A) is :








Answer: B

Watch Video Solution




Which of following can be used as a catalyst in the above reaction?



Answer: C

Watch Video Solution

121. Arrange the following carbonyl compounds in decreasing order of their reactivity in nucleophilic addition reaction.



A. $ii > iii > i > iv$

B. $ii > i > iv > iii$


C. $iii > ii > i > iv$

D. $iii > i > iv > ii$





Answer: B

 Watch Video Solution

122. The following reaction were carried out.




The final product formed in the above reaction sequence is :



Answer: B

 Watch Video Solution




Yield of each step as actually carried out in the laboratory is given above. What is overall yield of reaction?

- A. 0.42
- B. 0.31
- C. 0.21
- D. 0.6

Answer: C

 Watch Video Solution




124.

What is the appropriate reagent to carry out above synthesis, i.e., A, B, C respectively are :

- A. Br_2 / H^+ , LiAlH_4 , H^{\oplus}
- B. Br_2 / H^+ , NaBH_4 , HO^-
- C. NBS , AlCl_3 , HO^-
- D. $\text{Br}_2 / \text{HO}^-$, BF_3 , HO^-

Answer: B

 Watch Video Solution



125. Product (E) is : _____, Product




(E) is :

- A. Nylon 66
- B. Nylon 6
- C. Styrene
- D. Polystyrene

Answer: B

 [Watch Video Solution](#)


126. Methyl vinyl ketone on reaction with LiCuMe_2 gives a major product, whose structure is :





Answer: A


 Watch Video Solution

127. Which of following is in capable to show iodoform test ?





C.




D.

Answer: C


 Watch Video Solution


128.




Product (B) of above reaction is :

A.






B.




C.



D.



Answer: A


 [Watch Video Solution](#)



129. _____, Product

obtained is :






C.




D. None of these


Answer: B

 Watch Video Solution




Reactent (A) is :





Answer: C

 Watch Video Solution



131.

Value of x in above reaction is :

A. 1



B. 2

C. 3

D. 4

Answer: A

 Watch Video Solution




132.

Molecular weight of compound (A) is :





- A. 58
- B. 120
- C. 60
- D. 182

Answer: A

 Watch Video Solution



Product (A) of above reaction will be :




Answer: C

 **Watch Video Solution**

134. Which of the following will form stable hydrate ?

A. CCl_3CHO (Chloral)



B.


C. All of these

D.

Answer: D

 Watch Video Solution

135. The pH at which maximum hydrate is present in an solution of oxaloacetic acid :




- A. pH = 0
- B. pH = 12
- C. pH = 4
- D. pH = 6

Answer: A

 Watch Video Solution

136. Arrange their stabilities of given gem-diols in decreasing order.



A. $I > II > III$

B. $III > II > I$



C. $I > III > II$


D. $III > I > II$

Answer: A

 Watch Video Solution

137. Maximum hydration takes place of :





Answer: A


 Watch Video Solution

138. The conversion, $\text{PhCN} \rightarrow \text{PhCOCH}_3$, can be achieved most conveniently by reaction with:

- A. CH_3MgBr followed by hydrolysis
- B. $\text{I}_2 - \text{NaOH}$, CH_3I
- C. dil. H_2SO_4 followed by reaction with CH_2N_2
- D. LAH followed by reaction with CH_3I


Answer: A

 Watch Video Solution




139.


In the above reaction, product (B) is :




A.



B.




C.



D.


Answer: B

 Watch Video Solution




140.


Structure of A is :




A.



B.



C.




D.





Answer: D

[▶ Watch Video Solution](#)

141.




Product (B) is :



Answer: B






 Watch Video Solution

142.




To carry out about conversion, arrange the following reagents in

correct order.




Answer: D




 Watch Video Solution



143.

Product (B) is :










Answer: A

Watch Video Solution

144.




Identify appropriate reagents for the above reaction :







Answer: C

 Watch Video Solution



145. _____, Product




(X) of this reaction is :

- A. 
- B. 
- C. 
- D. 

Answer: B

 Watch Video Solution

146. The K_{eq} values in HCN addition to following aldehydes are in the order :



A. $I > II > III$


B. $II > III > I$

C. $III > I > II$

D. $II > I > III$

Answer: D

 Watch Video Solution



147.

relation between K_1 and K_2 is :

A. $K_1 = K_2$

B. $K_1 > K_2$

C. $K_2 > K_1$

D. $K_1 = K_2 = 1$

Answer: B

 [Watch Video Solution](#)

148. Which of the following is correct for the reaction ?




- A. A is cyanohydrin
- B. Nucleophilic-addition reaction
- C. The above reaction is not shown by alkenes
- D. All of these

Answer: D

 [Watch Video Solution](#)

149. Which of the following compounds (*i* through *v*) should not be classified as an acetal?



A. ii and iii

B. iv




C. i

D. none (they are all acetals)

Answer: D

[Watch Video Solution](#)

150. In which of reactions final product is NOT a ketone :






D. 1-butyne $\xrightarrow{\text{NaNH}_2}$ (A) $\xrightarrow{\text{CH}_3\text{I}}$ (B) $\xrightarrow[\text{CH}_3\text{CO}_2\text{H}]{\text{O}_3, \text{H}_2\text{O}, \text{THF}}$ (C)

Answer: C

 Watch Video Solution


151. The reaction of ethyl methyl ketone with $\text{Cl}_2 / \text{excess OH}^-$ gives the following major product



Answer: B

 Watch Video Solution

152. The product obtained from the following sequence of reactions is



- A. propanol
- B. 2-propanol
- C. 1-propanol
- D. propane

Answer: B

 Watch Video Solution

153. Which of the following shows oxidation of reactant.



A. 1

B. 1, 2



C. 1, 2, 3

D. All of these


Answer: B

 [Watch Video Solution](#)


154. The enolate ion that reacts with




A.



B.




C.







D.

Answer: C

 [Watch Video Solution](#)



What is the final product T.













Answer: D

 Watch Video Solution

1. Select the best choice for example (A to L) from the examples (a to n) given below. Write your choice in the box given.

A.	An acetal derivative of a ketone.	
B.	A chiral ketone.	
C.	An aldehyde that gives a aldol condensation with itself.	
D.	An oxime derivative	
E.	A reagent that reduces aldehydes to 1°- alcohols.	
F.	An α , β -unsaturated ketone.	
G.	A reagent that oxidizes aldehydes to carboxylic acids.	
H.	A reagent that reduces ketones to alkanes.	
I.	An enamine derivative of a ketone.	
J.	An intermediate in imine formation.	
K.	A cyclic hemiacetal.	
L.	A cyanohydrin derivative.	









(a)		(b)		(c)	
(d)		(e)		(f)	$\text{Zn(Hg)H}_3\text{O}^{(+)}$
(g)		(h)	NaBH_4 aq. alcohol	(i)	
(j)	$\text{Ag(NH}_3)_2^{(+)}\text{OH}^{(-)}$	(k)		(l)	

(m)		(n)	$\text{CH}_3-\text{CH}_2-\text{C}(=\text{O})-\text{H}$
------------	---	------------	--

 [View Text Solution](#)





2. The following questions refer to the compounds (A to G) shown below :

i. Which compounds are reduced by sodium borohydride ?		ii. Which compounds are hydrolyzed by hot aqueous acid ?		iii. Which compound are oxidized by $\text{CrO}_3/\text{pyridine}$?	
A	E	A	E	A	E
B	F	B	F	B	F
C	G	C	G	C	G
D	H	D	H	D	H






A.		B.		C.		D.	
E.		F.		G.		H.	

 Watch Video Solution


3. Match of the column :

Column (I)		Column (II)	
(a)		(p)	racemic mixture
(b)		(q)	Diastereomers
(c)		(r)	Nu-addition reaction
(d)		(s)	Nu-Substitutions reaction

4. Complete the following table.

	REACTANT	REAGENT(S)/ CONDITIONS	MAJOR ORGANIC PRODUCTS
a.		H ₂ /Pd - C in ethanol (solvent)	A
b.		H ⁺ /H ₂ O/Δ	B
c.		(CH ₃) ₂ C ⁻ —P ⁺ (C ₆ H ₅) ₃	C
d.		1. Li ⁺ [(CH ₃) ₂ Cu] ⁻ in dry ether 2. H ⁺ /H ₂ O	D
e.	E	OH ⁻ /ethanol/Δ	

5. Consider the following reactions and answer A and B.



Suggest a reagent appropriate step (a0 the synthesis.

A. HO^- / Br_2 (1 mole)

B. H^+ / Br_2 (1 mole)


C. both (a) and (b)

D. None of these

Answer: C

 Watch Video Solution

6. Consider the following reactions and answer A and B.



Yield of each step as actually carried out in laboratory is given above each arrow. What is overall yield of the reaction ?

A. 0.6


B. 0.21

C. 0.4

D. 0.68

Answer: B

 Watch Video Solution











7.

Degree of unsaturation present in compound (A + B + C) is ?



 Watch Video Solution


8. Within each set, which compound should be more reactive toward carbonyl addition reaction ?

	A	B
Set (1)	$\text{CH}_3-\overset{\text{O}}{\parallel}{\text{C}}-\text{CH}_2-\text{Br}$	$\text{CH}_3-\overset{\text{O}}{\parallel}{\text{C}}-\text{CH}_2-\text{CH}_2-\text{Br}$
Set (2)	$\text{CH}_3-\overset{\text{O}}{\parallel}{\text{C}}-\overset{\text{O}}{\parallel}{\text{C}}-\text{CH}_3$	$\text{CH}_3-\overset{\text{O}}{\parallel}{\text{C}}-\text{CH}_2-\text{CH}_3$
Set (3)	$\text{CH}_3\text{O}-\text{C}_6\text{H}_4-\text{CH}=\text{O}$	$\text{O}_2\text{N}-\text{C}_6\text{H}_4-\text{CH}=\text{O}$
Set (4)	$\text{C}_6\text{H}_5-\text{CH}=\text{O}$	$\text{CH}_3-\overset{\text{O}}{\parallel}{\text{C}}-\text{H}$
Set (5)		

Set (6)		
Set (7)		
Set (8)		
Set (9)	$\text{C}_6\text{H}_5-\overset{\text{O}}{\parallel}{\text{C}}-\text{CH}_3$	$\text{C}_6\text{H}_5-\overset{\text{O}}{\parallel}{\text{C}}-\text{C}_6\text{H}_5$
Set (10)	$\text{CH}_3-\overset{\text{O}}{\parallel}{\text{C}}-\text{CH}_2-\text{CH}_3$	$\text{C}_6\text{H}_5-\overset{\text{O}}{\parallel}{\text{C}}-\text{H}$

9. Match the Column (I) and Column (II). (Matrix)

Column (I)		Column (II)
(A)		(p) Formation of six member ring takes place
(B)		(q) Final product is Ketone

(C)	$\text{CH}_3 - \overset{\text{O}}{\parallel}{\text{C}} - \text{CH}_2 - \text{CH}_2 - \text{CH}_2 - \overset{\text{O}}{\parallel}{\text{C}} - \text{H} \xrightarrow[\Delta]{\text{HO}^-} (\text{A})$	(r) Final product formed will give positive Tollens test
(D)		(s) Final product formed will react with 2,4-DNP. (2,4-di-nitrophenyl hydrazine)

10. Consider reactions A through F. Those carbon atoms undergoing change, as part of a functional group, are marked as C^{12} , C^{14} or starred. In the cases shown, each carbon atom has either been reduced or oxidized. Your job is to identify the change in oxidation

state that has occurred for each of the marked carbon.











Reaction		C ¹²	C ¹⁴
A.	$\text{CH}_3\underset{12}{\text{C}}\text{H}=\underset{14}{\text{C}}\text{H}_2 \xrightarrow{\text{Br}_2} \text{CH}_3\text{CHBrCH}_2\text{Br}$	Reduced	Reduced
		Oxidized	Oxidized
B.	$\text{CH}_3\underset{12}{\text{C}}\text{H}=\underset{14}{\text{C}}\text{H}_2 \xrightarrow[\text{(ii) H}_2\text{O}_2, \text{NaOH}]{\text{(i) B}_2\text{H}_6} \text{CH}_3\text{CH}_2\text{CH}_2\text{OH}$	Reduced	Reduced
		Oxidized	Oxidized
C.	$\text{CH}_3\text{CH}_2\overset{*}{\text{C}}\text{H}=\text{O} \xrightarrow{\text{NaBH}_4} \text{CH}_3\text{CH}_2\text{CH}_2\text{OH}$	Reduced	
		Oxidized	
D.	$\text{CH}_3\text{CH}_2\overset{*}{\text{C}}\text{H}=\text{O} \xrightarrow[\text{H}_2\text{O, pH} > 8]{\text{Ag}^{(+)}} \text{CH}_3\text{CH}_2\text{CO}_2\text{H}$	Reduced	
		Oxidized	
E.	$\text{CH}_3\text{CO}\underset{12}{\text{C}}\underset{14}{\text{C}}\text{H}_2\text{CO}_2\text{H} \xrightarrow{\text{Heat}} \begin{matrix} \text{CH}_3\text{COCH}_3 \\ + \\ \text{O}=\text{C}=\text{O} \end{matrix}$	Reduced	Reduced
		Oxidized	Oxidized
F.	$\text{H}_2\underset{12}{\text{C}}=\underset{14}{\text{C}}(\text{OH})\text{C}_2\text{H}_5 \xrightarrow{\text{tautomerization}} \text{H}_3\text{CCOC}_2\text{H}_5$	Reduced	Reduced
		Oxidized	Oxidized

 Watch Video Solution

11. Consider the possible formation of an aldehyde or ketone product when each of the ten compounds in the column on the left is treated with each of the reagents shown in the top row. Check the designated answer box if you believe an aldehyde or ketone will be formed.

Assume that the reagents may be present in excess. For each checked

reaction, try to draw the structure of the major product (s).

Starting	PCC $C_2H_5NHCrO_3Cl$	Jone's Reagent CrO_3 in aq. acid	$Pb(OAc)_4$ [or HIO_4]	(i) O_3 , (ii) Zn dust	H_3O^+	(i) BH_3 in THF (ii) H_2O_2 + NaOH
						
						
						
						
						
						
						
						
						
						




[View Text Solution](#)


12. Wittig reaction :

The reaction of a phosphorus ylide with an aldehyde (or) ketone

introduces a carbon-carbon double bond in place of the carbonyl bond.



Mechanism :





Driving force of the reaction is high bond energy of ($\text{P}=\text{O}$).

($\Delta H = -ve$).



, Major

product (A) is :




D.

Answer: A


 Watch Video Solution

13. Wittig reaction :

The reaction of a phosphorus ylide with an aldehyde (or) ketone introduces a carbon-carbon double bond in place of the carbonyl bond.




Mechanism :





Driving force of the reaction is high bond energy of ($\text{P}=\text{O}$).

($\Delta H = -ve$).



product (A) is :






D.

Answer: A


 [Watch Video Solution](#)

14. Wittig reaction :

The reaction of a phosphorus ylide with an aldehyde (or) ketone introduces a carbon-carbon double bond in place of the carbonyl bond.



Mechanism :



Driving force of the reaction is high bond energy of (P = O).

($\Delta H = -ve$).



, Major

product (A) is


- A. cis-2-butene
- B. trans-2-butene
- C. iso-butene
- D. 1-butene

Answer: B


 Watch Video Solution

15. Wittig reaction :

The reaction of a phosphorus ylide with an aldehyde (or) ketone introduces a carbon-carbon double bond in place of the carbonyl bond.




Mechanism :







Driving force of the reaction is high bond energy of ($\text{P}=\text{O}$).

($\Delta H = -ve$).



is :






D.

Answer: B


 [Watch Video Solution](#)

16. Wittig reaction :

The reaction of a phosphorus ylide with an aldehyde (or) ketone introduces a carbon-carbon double bond in place of the carbonyl bond.





Mechanism :





Driving force of the reaction is high bond energy of (P = O).

($\Delta H = -ve$).

Identify major product in given intramolecular wittig reaction :



Product (A) and (B) respectively are :




Answer: A


 Watch Video Solution

17. Wittig reaction :

The reaction of a phosphorus ylide with an aldehyde (or) ketone introduces a carbon-carbon double bond in place of the carbonyl bond.




Mechanism :







Driving force of the reaction is high bond energy of ($\text{P}=\text{O}$).

($\Delta H = -ve$).



, Product (A)





is :




Answer: B

 Watch Video Solution

18. Match the column :





Column (I)		Column (II)	
Conversion		Reagent	
(a)		(p)	$\text{NH}_2/\text{NH}_2/\text{HO}^\ominus, \Delta$ (Wolff-Kishner reduction)
(b)		(q)	Zn(Hg), HCl (Clemmensen reduction)
(c)		(r)	LiAlH_4
(d)		(s)	None

 Watch Video Solution



19.


Reactant (A) is :



Answer: C







View Text Solution



20.

Product (B) is :



Answer: D



View Text Solution

