



CHEMISTRY

BOOKS - MS CHOUHAN

BIOMOLECULES

Level 1

1. Which statement correctly completes the statement ?

Except for glycine, which is achiral, all the amino acids present in proteins

A. are chiral, but racemic

B. have the L configuration at their α carbon

C. have the R configuration at their α carbon





D. have the configuration at their α carbon

Answer: B



Watch Video Solution

2. Assume that a particular amino acid has an isoelectric point of 6.0. In a solution at pH 1.0, which of the following species will predominate ?




Answer: A



Watch Video Solution

3. The pK_a values for the three ionizable groups X, Y and Z of glutamic acid are 4.3, 9.7 and 2.2 respectively



The isoelectric point for the amino acid is :

A. 7

B. 3.25

C. 4.95

D. 5.95

Answer: B



Watch Video Solution

4. An amino acid may be represented by




general formula $H_2N - \overset{R}{\underset{|}{C}H} - COOH$. If

$R = -CH_2C_6H_5$ then it is phenylalanine

(Phe) and if $R = CH_3$ then it is alanine (Ala).

Find the sequence of reagents from those

given below to synthesize Phe- Ala.



A. 1 and 2

B. 1 and 4

C. 2 and 3

D. 3 and 4

Answer: B



Watch Video Solution

5. Iso-electric point of alanine is ($\text{pH} = 6$). At which pH , maximum concentration of zwitterion of alanine will be present?

A. $\text{pH} > 6$

B. $\text{pH} < 6$


C. $\text{pH} = 6$

D. $\text{pH} = 7$

Answer: C



Watch Video Solution



Identify the amino acid obtained by hydrolysis of the above compound:

A. Glycine

B. Alanine

C. Both (a) and (b)

D. None of these

Answer: C



Watch Video Solution

7. At iso-electric point:

A. conc. of cation is equal to conc of anion

B. Net charge is zero.

C. Maximum conc. of di-polar ion (Zwitter ion) will be present

D. All of the above

Answer: D



Watch Video Solution

8. Which of following amino acid has lowest iso-electric point ?

A. Glycine

B. Alanine

C. Aspartic acid


D. Lysine

Answer: C



Watch Video Solution

9. Find iso-electric point of given amino acid



$$pK_b = 4.4$$

A. 3.3

B. 5.9

C. 9.6


D. 11.8

Answer: B



Watch Video Solution

10. Find iso-electric point of the given amino acid



A. A) 5.5

B. B) 6.5

C. C) 3


D. D) 5

Answer: C



Watch Video Solution

11.



, Product (B) of given reaction is :

A. Glycine

B. Alanine

C. valine

D. Leucine

Answer: B



Watch Video Solution

12. Which amino acid does not contain chiral centre ?

A. Valine

B. Leucine

C. Glycine

D. Iso-leucine

Answer: C



Watch Video Solution

13. Which of the following is Sanger reagent ?

A. 2,4-Di-nitro fluobenzene

B. Phenyl isocyanate

C. 2,4-Di-nitro chlorobenzene

D. 2,4-Di-nitro iodobenzene

Answer: A



Watch Video Solution

14. A D-carbohydrate is :

A. Always dextrorotatory

B. Always laevorotatory

C. Always the mirror of the corresponding

L-carbohydrate




D. None of these

Answer: D




Watch Video Solution

15. Which L-sugar on oxidation gives an optically active dibasic acid (2 COOH groups)?




D.



Answer: A



Watch Video Solution



given osazone can be obtained by :

A. D-glucose

B. D-mannose

C. D-Idose

D. Both (a) & (b)

Answer: D



Watch Video Solution

17. Which of the following pair gives same phenyl osazone?


- A. D-Glucose and D-Allose
- B. D-Glucose and D-Alfrose
- C. D-Glucose and D-Mannose

D. D-Glucose and D-Talose


Answer: C

 [Watch Video Solution](#)


18. Which of the following is the Fischer projection of L-threonine (also known as (2S, 3R)2-amino-3-hydroxybutanoic acid) ?




A.



B.



C.






D.

Answer: B



Watch Video Solution

19. Among the three compounds shown below, two yield the same product on reaction with warm HNO_3 . The exception is :



D. None of these

Answer: B



Watch Video Solution

20. The optical rotation of the α form of a pyranose is $+150.7^\circ$, that of the β -form is $+52.8^\circ$. In solution an equilibrium mixture of these anomers has an optical rotation of $+80.2^\circ$. The percentage of the α -form in equilibrium mixture is :

A. 0.28

B. 0.32

C. 0.68




D. 0.72

Answer: A



Watch Video Solution

21. Which of the following represents the anomer of the compound shown?



C. both

D. None of these


Answer: B



Watch Video Solution

22. Which set of terms correctly identifies the carbohydrate shown? Which set of terms correctly identifies the carbohydrate shown?

(1) Pentose (2) Pentulose (3) Hexulose (4) Hexose (5) Aldose (6) Ketose (7) Pyranose (8) Furanose



A. 2,6,8

B. 2,6,7

C. 1,5,8

D. A set of terms other than these

Answer: A



Watch Video Solution

23. For the complex conversion of D-glucose into the corresponding osazone, the minimum

number of equivalents of phenyl hydrazine
required is :

A. two

B. three

C. four


D. five

Answer: B



Watch Video Solution

24. Which one of the following compounds will form an osazone derivative ?



Answer: A




Watch Video Solution

25. Which of the following structure is L-arabinose?

A. 

B. 

C. 



Answer: C



Watch Video Solution

26. Which one of the statements concerning the equilibrium shown is true ?



- A. The two structures are enantiomers of each other. They have equal but opposite optical rotations and racemize slowly at room temperature
- B. The two structures are enantiomers of each other. They racemize too rapidly at

room temperature for their optical rotations to be measured

C. The two structures are diastereomers of each other. Their interconversion is called mutarotation

D. The two structures are diastereomers of each other. Their interconversion does not require breaking and making bonds, only a change in conformation

Answer: C



Watch Video Solution

27. The configurations of the chirality centres in D-threose (shown) are :



A. 2R, 3R

B. 2R, 3S

C. 2S, 3R

D. 2S, 3S

Answer: C



Watch Video Solution


28. Rapid interconversion of α -D-glucose and β -D-glucose in solution is known as:

- A. racemization
- B. asymmetric induction
- C. fluxional isomerization
- D. mutarotation

Answer: D



29. Identify the correct set of stereochemical relationships amongst the following monosaccharides I-IV



A. I and II are anomers, III and IV are epimers

B. I and III are epimers, II and IV are anomers

C. I and II are epimers, III and IV are anomers

D. I and III are anomers, I and II are epimers

Answer: D




Watch Video Solution

30. (i) Deficiency of which vitamin cause night blindness ?


(ii) Name the base that is found in nucleotide of RNA only.

(iii) Glucose on reaction with HI gives n - hexane . What does it suggest about the structure of glucose ?

A. 



B.



C.




D. None of these

Answer: A



Watch Video Solution

31. What is the structure of L-glyceraldehyde?




D. Both (a) & (b)

Answer: D



Watch Video Solution



32.

the

given is enol form of:

A. D-glucose

B. D-mannose

C. D-fructose

D. All of these

Answer: D



Watch Video Solution

33. D-glucose $\xrightleftharpoons{HO^-} A + B$, A and B are :

A. D-mannose & D-mannitol

B. D-mannose & D-Fructose

C. D-Allose & D-Altrose

D. D-Glucose & D-Idose

Answer: B



Watch Video Solution

34. Stereoisomers of aldoheptose is (a) and stereoisomers of ketoheptose is (b). Ratio of a/b is:

A. $\frac{1}{2}$

B. $\frac{2}{1}$

C. $\frac{4}{1}$

D. $\frac{1}{4}$

Answer: B



Watch Video Solution

35. D-Glucose $\xrightarrow[\Delta]{HNO_3}$ to (A)`, Product (A) is :

A. D-Gluconic acid

B. D-Glucitol

C. D-Fructose

D. D-Glucaric acid

Answer: D



Watch Video Solution

36. D-glucose & D-Fructose can be differentiated by :

A. Fehling solution

B. Tollens reagent

C. Benedict test

D. Br_2 / H_2O

Answer: D



Watch Video Solution

37. D-Glucose exist in x different forms. The value of x (stereoisomer) is :

A. 2

B. 3

C. 4

D. 5

Answer: B



Watch Video Solution

38. D-Mannose $\xrightleftharpoons{HO^-}$ D - Glucose $\xrightleftharpoons{HO^-}$ (A),

Product (A) of above reaction is

A. D-glucose

B. D-fructose


C. D-talose

D. D-idose

Answer: B

 Watch Video Solution

39. Which of the molecules below will react with Ag^+ ?



A. (i), (iii) and (v)

B. (ii) and (iv)


C. (iv) and (vi)

D. (i), (ii), (iii) and (vi)

Answer: C



Watch Video Solution



40.

Which of the compounds (A-D) depicted above is NOT a hemiacetal linkage ?

A. Compound A

B. Compound B

C. Compound C





D. None of the above (they are all hemiacetals)

Answer: D



Watch Video Solution

41. Which of the following Fischer projection formula is same as D-Glyceraldehyde ?





Answer: C



Watch Video Solution

42. What is the structure of the monomer from which the following polymer was made ?




Answer: B





Watch Video Solution

43. The following structure represents a subunit of a hydrocarbon polymer that may be prepared by a radical polymerization method. Identify the monomer that has been polymerized to make this polymer chain.



A.






Answer: C



Watch Video Solution

44. Choose the answer that has correctly identified the number of acetals and

hemiacetals in isomaltose.




Acetal	Hemiacetal	Acetal	Hemiacetal
(a) 0	0	(b) 1	0
(c) 0	1	(d) 1	1




Watch Video Solution


45. Predict the product of the following reaction.




A.




B.



C.



D.



Answer: A



Watch Video Solution

46. Which reagent/s can be used to distinguish glucose and fructose ?

(I) Bromine water (II) Tollen's reagent (III)

Schiff's reagent

A. (I), (II) and (III)

B. (II) and (III)

C. Only (I)





D. Only (III)

Answer: C



Watch Video Solution

47. Choose the peptide that matches the abbreviation Phe-Val-Ala.




Answer: A



Watch Video Solution

48. Which of the following carbohydrate(s) would not undergo mutarotation in aqueous solution?



A. II only

B. I, III and IV only

C. II and IV only


D. I and III only

Answer: C



Watch Video Solution

49. The number of peptide bonds in the compound.



A. 1

B. 2

C. 3


D. 4

Answer: A




Watch Video Solution


50. Serine ($HOCH_2CH(NH_2)COOH$) is an essential amino acid. The correct Fisher projection of serine is




B.



C.



D.



Answer: D



Watch Video Solution

Level 2


1. Match the column (I) and Column (II).

(Matrix)

	Column (I) Molecule	Column (II) Configuration
(a)	$\begin{array}{c} \text{CHO} \\ \\ \text{H} - \text{C} - \text{OH} \\ \\ \text{CH}_2\text{OH} \end{array}$	(p) R- (Rectus)
(b)	$\begin{array}{c} \text{CHO} \\ \\ \text{HO} - \text{C} - \text{H} \\ \\ \text{CH}_2\text{OH} \end{array}$	(q) S- (Sinister)
(c)	$\begin{array}{c} \text{NH}_2 \\ \\ \text{H} - \text{C} - \text{CO}_2\text{H} \\ \\ \text{CH}_3 \end{array}$	(r) D
(d)	$\begin{array}{c} \text{NH}_2 \\ \\ \text{H} - \text{C} - \text{CH}_3 \\ \\ \text{CO}_2\text{H} \end{array}$	(s) L



Watch Video Solution



2.


One cyclic acetal form of D - galactose is shown above.

Which atom is the anomeric carbon ?

- (a) Atom A (b) Atom B (c) Atom C (d) Atom D
(e) Atom E (f) Atom F



[Watch Video Solution](#)



3.

One cyclic acetal form of D - galactose is shown above.

Which name most completely describes this cyclic acetal form?

(a) α - *D*- Galactofuranose


(b) β - *P* - Galactofuranose

(c) α - D- Galactopyranose

(d) β - D- Galactopyranose



Watch Video Solution




4.

One cyclic acetal form of D - galactose is shown above.

How many compound which is given below is isomer of D-Glucose ?

D-Mannose, D-Fructose, D-Gulose, D-Idose, D-Galactose, D-Arabinose, D-Ribose.

 [Watch Video Solution](#)




5.

One cyclic acetal form of D - galactose is

shown above.

How many acidic group is present in given amino acid ?



Watch Video Solution