

CHEMISTRY




BOOKS - MS CHOUHAN

HYDROCARBONS (ALKENES)

Level 1

1. (R)-3-bromocyclopentene (shown below) reacts with Br_2 / CCl_4 to form two products, Y and Z, Y is not optically active (does not rotate plane-polarized light). What is the structure of Y?






Answer: C






Watch Video Solution



2.

reactant (A)

can be :



D. All of these

Answer: D


Watch Video Solution




3.

major


product of the reaction is :




A.



B.




C.



D.

Answer: C

 Watch Video Solution






4.

Which of the following products cannot be obtained in ozonolysis of o-xylene?

CHO


A. |

CHO



Answer: D


 Watch Video Solution




5.

major


product of the reaction is :




A.



B.




C.






D.

Answer: B

 Watch Video Solution




(A) of the reaction is :



D. None of these




Answer: B

 Watch Video Solution



products (C






) of the reaction is :



Answer: C

 Watch Video Solution





8. Complete the following reaction



Answer: C

 Watch Video Solution






9. The reaction of propene with H_3O^+ will proceed with which of the following intermediates ?



Answer: C

 [Watch Video Solution](#)





10. Which of the following bromides is the major product of the reaction shown below, assuming that there are no carbocation rearrangement ?



Answer: D

 Watch Video Solution


11. Which of the following reactions results in the formation of a pair of diastereomers ?




Answer: B

 Watch Video Solution


12. What is a likely product of the reaction shown ?




A.




B.



C.




D.



Answer: D

 Watch Video Solution


13. Which of the following, when undergoing addition of HBr, will form ONLY a pair of diastereomers ?



A.



B.



C.



D.

Answer: C

 Watch Video Solution

14. How many transition states and intermediates will be formed during the course of following reaction ?







- A. 3 transition states and 3 intermediates
- B. 4 transition states and 3 intermediates
- C. 3 transition states and 2 intermediates
- D. 5 transition states and 4 intermediates

Answer: B

 [Watch Video Solution](#)

15. Product of which of the following reactions, is racemic mixture ?




Answer: B



Watch Video Solution

16. The product(s) of the following reaction can best be described as :






- A. a racemic mixture
- B. a single enantiomer
- C. a pair of diastereomers
- D. an achiral molecule

Answer: C



Watch Video Solution

17. Taking into account the stability of various carbocations and, as well as the rules governing mechanisms of carbocation rearrangements, which reaction is most likely to occur during the given reaction ?



D. None





Answer: D

Watch Video Solution

18. Consider the following reaction in which the intermediate carbocation loses H^+ to give the final product







Which of the following energy profiles best represents the overall reaction?



Answer: D

 Watch Video Solution


19. Methyl vinyl ether, $H_2C = CH-OCH_3$, reacts with Br_2/CH_3OH . If methanol is reacting as water would, and if this reaction follows a typical mechanism of electrophilic addition, what would be the expected product ?



Answer: B

 Watch Video Solution

20. 2, 4-hexadiyne (C_6H_6) is allowed to react with Li in $NH_3(liq)$. The product obtained is treated with 1 equivalent of Cl_2 in CCl_4 . Which of the following constitutional isomers are possible products ?



A. I and II

B. II and III


C. I and V

D. I and III


Answer: D

 Watch Video Solution


21. Which of the following is the best stereochemical representation when reaction between 1-methylcyclohexene and NBS react in aqueous dimethyl sulfoxide ?



A.



B.



C.


D. None of these

Answer: B




[Watch Video Solution](#)


22. Which of the following is among the major products of the reaction of (E)-3-methyl-2-pentene with BH_3 in THF followed by the addition of H_2O_2/HO^- ?




A.



B.




C.



D.

Answer: A

23. Compare rate of dehydration of (i), (ii) and (iii) by conc. H_2SO_4



A. (i) > (iii) > (ii)


B. (i) > (ii) > (iii)

C. (ii) > (i) > (iii)

D. (ii) > (iii) > (i)

Answer: B

24. How many products will be formed in this reaction ?



A. 10


B. 2

C. 3

D. 4

Answer: B


 Watch Video Solution




25.

Product (A) of the reaction is:


A.



B.



C.




D.




Answer: B



Watch Video Solution





A.



B.




C.



Answer: B


 Watch Video Solution

27. Predict the product (A) of the following reaction







A.



B.



C.




D.

Answer: D







Watch Video Solution



28.

Major-




product (A) is:



Answer: B




29. Di-imide (N_2H_4) is used to reduce double bond of:






Answer: D

Watch Video Solution



End product of the reaction is :







Answer: B

 Watch Video Solution




Product (A) is :






Answer: C


 Watch Video Solution



32.

Product (A) is :







D.

Answer: B

 Watch Video Solution




Product (X) will be :




A.

B.



C.




D.




Answer: A

 Watch Video Solution

34.



Product (C) is :




A.



B.



C.




Answer: B

 [Watch Video Solution](#)




MMPP → Magnesium mono peroxy phthalate. Product (X) is :




A.



B.




C.



D.




Answer: B

 Watch Video Solution



Product


(P) is :



D. None of these

Answer: B




 Watch Video Solution




37.

Product

(A) is :






D.



Answer: B

 [Watch Video Solution](#)



(no ring





substitution) product (A) is :



Answer: B






39. metachloroperbenzoic acid Product (A) of the above reaction is :




Answer: B

 Watch Video Solution

40. The major product of the following reaction sequence is :





D.

Answer: B

 Watch Video Solution

41. Which one of the following compounds gives acetone ($(CH_3)_2C = O$) as one of the product of its ozonolysis ?



A.




B.



C.

D.




Answer: D




Watch Video Solution


42. Addition of HCl to 3,3-dimethyl-1-butene yields two products, one of which has a rearranged carbon skeleton. Among the following carbocations, select the possible intermediates in that reaction ?




1



2



3



4

A. 1,2

B. 1,3


C. 2,4

D. 2,3

Answer:

 [Watch Video Solution](#)

43. Conversion of cyclohexene to cyclohexanol can be conveniently achieved by :



C. hydroboration, oxidation


D. hydroboration hydrolysis

Answer: C

 [Watch Video Solution](#)

44. Trans-cyclohexane-1,2-diol can be obtained by the reaction of cyclohexene with:






Answer: C

 [Watch Video Solution](#)


45. Bromination of (E)-2-butenedioic acid gives



Answer: A

 [Watch Video Solution](#)


46. The major product formed during the reaction of 1-methylcyclopentene with CH_3CO_3H is




A.



B.



C.




D.





Answer: C



Watch Video Solution




reaction is :




Answer: C


 Watch Video Solution

48. Complete the following reaction







A.



B.



C.




D.

Answer: B







Watch Video Solution



49.

Product






(P) is :



Answer: B

Watch Video Solution




50. What is the major product expected from the following reaction ?



Answer: B


 Watch Video Solution

51. $CH_3 - CH = CH_2 \xrightarrow[\text{(low conc.)}]{Br_2 / h\nu}$ (A), Product (A) of the reaction is :



Answer: B

 Watch Video Solution



Y is

A. $O_3 / Zn(H_2O)$


B. HIO_4

C. CrO_3

D. Cold dil $KmnO_4$


Answer: A

 Watch Video Solution




53.


Product of the reaction is:




A.



B.



C.





D.

Answer: B

 [Watch Video Solution](#)

54. Which compound is a possible product from addition of Br_2 to 1-butene?



Answer: D

 [Watch Video Solution](#)

55. Addition of Br_2 to cis-2-butene would give a product which is:

- A. achiral
- B. racemic
- C. meso
- D. optically active

Answer: B

 [Watch Video Solution](#)

56. Addition of Br_2 to trans-2-butene would give a product which is

- A. achiral
- B. racemic
- C. meso
- D. optically active

Answer: C

 [Watch Video Solution](#)

57. Addition of OsO_4 to cyclopentene would give a product which is:

- A. achiral

B. racemic

C. meso

D. optically active

Answer: C



[Watch Video Solution](#)

58. Addition of BH_3 followed by H_2O_2 to trans-2-butene would give a product which is:

A. achiral

B. racemic


C. meso

D. optically active

Answer: B






[Watch Video Solution](#)



59.

..., Reagent A

may be :








D. All are possible

Answer: C

 Watch Video Solution

60. The major product of the following reaction is :






D.



Answer: A

 Watch Video Solution




61.


Identify (B):



C.




D.



Answer: B

 [View Text Solution](#)

62. Which of the following is a major product of the reaction shown below?



 [Watch Video Solution](#)


63. In methyl alcohol solution, bromine reacts with ethylene (ethene) to yield $BrCH_2$, CH_2OCH_3 in addition to 1,2-dibromoethane because





- A. the methyl alcohol solvates the bromine
- B. the ion formed initially may react with Br^- or CH_3OH
- C. this is a free radical reaction
- D. the reaction follows Markovnikov's rule

Answer: A

 Watch Video Solution

64. Which of the following compound was the starting material for the oxidation shown below?









Answer: B

 Watch Video Solution

65. Which series of reactions will achieve the following transformation ?






Answer: D


 Watch Video Solution

66. Taking into account the stability of various cycloalkanes and carbocations, as well as the rules governing mechanisms of carbocation rearrangements, what is the most likely product of this reaction ?







A.



B.



C.






D.

Answer: B




Watch Video Solution


67. A triene is treated with ozone followed by zinc in acetic acid to give the following three products. What is the structure of the triene ?



A.



B.



C.



D.




Answer: D



Watch Video Solution

68. Which of the following compound would yield trialkylborane shown below when treated with BH_3/THF ?




- A. 2-methylbut-1-ene
- B. 2-methylbut-2-ene
- C. 3-methylbut-1-ene
- D. 3-methylbut-1-yne

Answer: A



[Watch Video Solution](#)

69. If the following compound is treated with Pd/C in excess of hydrogen gas, how many stereoisomers of the product will be obtained ?



A. 1

B. 2


C. 3

D. 4


Answer: C

 Watch Video Solution


70. Which is the most precise designation of stereochemistry for the products formed in the electrophilic addition of DBr to 1-methylcyclohexene? ($D = {}^2H$, an isotope of hydrogen)



A.



B.




C.

D. Both (a) and (b)

Answer: D

 Watch Video Solution

71. Consider the addition of HBr to 3,3-Dimethyl-1-butene shown below. What is the best mechanistic explanation for the formation of the observed product?



- A. Protonation of the alkene followed by a hydride shift and addition of bromide to the carbocation
- B. Double bond shift in the alkene following by the protonation and addition of bromide to the carbocation
- C. Addition of bromide to the alkene followed by a double bond shift and protonation

D. Protonation of the alkene followed by a methyl shift and addition of bromide to the carbocation

Answer: D

 [Watch Video Solution](#)

72. Propene $CH_3CH = CH_2$ can be converted into 1-propanol by oxidation. Indicate which sets of reagents amongst the following is ideal to effect the above conversion ?

A. $KMnO_4$ (alkaline)

B. Osmium tetroxide (OsO_4 / CH_2Cl_2)




C. B_2H_6 and alk. H_2O_2

D. O_3 / Zn

Answer: C

 [Watch Video Solution](#)

73. Which is the most suitable reagent among the following distinguish compound (3) from the others ?



A. Bromine in carbon tetrachloride

B. Bromine in acetic acid solution

C. Alk. $KMnO_4$

D. Ammonical silver nitrate

Answer: D



Watch Video Solution

74. The principal organic product formed in the reaction given below is :



- A. $CH_3 - CHBr(CH_2)_8COOH$
- B. $CH_2 = CH(CH_2)_8COBr$
- C. $CH_2BrCH_2(CH_2)_8COOH$
- D. $CH_2 = CH(CH_2)_7CHBrCOOH$

Answer: C



[Watch Video Solution](#)


75. When 2-butyne is treated with $H_2 / Pd - BaSO_4$, the product formed will be :

- A. cis-2-butene
- B. trans-2-butene
- C. 1-butene

D. 2-hydroxy butane

Answer: A

 Watch Video Solution



In the above reaction X is

A. HNO_3

B. O_2




C. O_3

D. KMnO_4

Answer: A

 Watch Video Solution

77. Which of the following alkene on catalytic hydrogenation given cis and trans-isomer ?



D. all of these

Answer: D

 [Watch Video Solution](#)




78. In the reaction of hydrogen bromide with an alkene (in the absence of peroxides), the first step of the reaction is the to the alkene.

- A. fast addition of an electrophilic
- B. slow addition of an electrophile
- C. fast addition of a nucleophilic
- D. slow addition of a nucleophile

Answer: B

 [Watch Video Solution](#)

79. Which of the following alcohols cannot be prepared from hydration of an alkene ?



D.




Answer: D


 [Watch Video Solution](#)


80. Which of the species shown below is the most stable form of the intermediate in the electrophilic addition of Cl_2 in water to cyclohexene to form a halohydrin ?

A.




B.





C.



D.

Answer: D

 [Watch Video Solution](#)

81. The reaction, $(CH_3)_2C = CH_2 + Br\cdot \rightarrow (CH_3)_2\dot{C} - CH_2Br$ is an example of a/an step in a radical chain reaction.

A. initiation

B. termination





C. propagation

D. heterolytic cleavage

Answer: C

 Watch Video Solution





82. Which of the following most accurately describes the first step in the reaction of hydrogen chloride with 1-butene ?



Answer: B

 Watch Video Solution

83. Which of the following best describes the flow of electrons in the acid-catalyzed dimerization of $(CH_3)_2C = CH_2$?




Answer: A




Watch Video Solution


84. Hydroboration of 1-methylcyclopentene using B_2D_6 , followed by treatment with alkaline hydrogen peroxide, gives




A.



B.



C.




D.

Answer: A



Watch Video Solution



85.






The correct statements with respect to the above pair of reactions are that (I) the reactions are stereospecific (II) (X) is erythro and (Y) is threo isomer (III) (X) is threo and (Y) is erythro isomer (IV) each of (P) and (Q) gives a mixture of (x) and (Y)

- A. I and II
- B. I and III
- C. I and IV
- D. II and IV

Answer: A

 [Watch Video Solution](#)

86. Identify .Q. in the following sequence of reactions :



Answer: D


Watch Video Solution

87. 4-Pentenoic acid when treated with I_2 and NaHCO_3 gives :



- A. 4,5-diiodopentanoic acid
- B. 5-iodomethyl-dihydrofuran-2-one
- C. 5-iodo-tetrahydropyran
- D. 4-pentenolyl iodide

Answer: B


 Watch Video Solution




(B) of the reaction is:



C.




D.




Answer: B

 Watch Video Solution







89. (Styrene)



Product


(C) is :




Answer: C

 Watch Video Solution


90. Which of the following will give a mixture of cis and trans-1,4-dimethyl cyclohexane, when undergo catalytic hydrogenation ?



A.



B.



Cis-3-6 dimethyl
cyclohexene


C.

D. Both (a) and (b)


Answer: D

 [Watch Video Solution](#)


91. An optically active compound A with molecular formula C_8H_{14} undergoes catalytic hydrogenation to give meso compound, the structure of (A) is :




A.



B.




C.



D.

Answer: B

Watch Video Solution



How many products will be formed in above reaction ?

A. 2


B. 4

C. 3

D. 6

Answer: B

 Watch Video Solution



93.

Product of the reaction is :

A. Racemic

B. Diastereomers

C. Meso

D. Pure enantiomers

Answer: A

 [Watch Video Solution](#)

94. cis-2-butene $\xrightarrow[\text{Peroxide}]{\text{HBr}}$ product , Product of the reaction is :

A. Racemic


B. Diastereomers

C. Meso


D. E and Z isomer

Answer: A

 [Watch Video Solution](#)



(a)



(b)

95.

Rate of reaction towards reduction using (H_2 / Pt):

A. a b

B. a = b


C. b a

D. Reduction of given molecule is not possible



Answer: A



Watch Video Solution



of the above reaction is :




D. Both (a) and (b)





Answer: D

 [Watch Video Solution](#)

97.





MCPBA → Metachloroperbenzoic acid



Answer: B






Watch Video Solution




98.

Product of

the reaction is :





D.

Answer: A

 Watch Video Solution





99. $CH_3 - CH = CH_2 \xrightarrow[(2) CH_3CO_2T]{(1) THF:BD_3} (A)$ Product (A) of the above reaction is:



Answer: C

 Watch Video Solution


100. Optically active isomer (A) of (C_5H_9Cl) on treatment with one mole of H_2 gives an optically inactive compound (B) compound (A) will be :






Answer: D

 Watch Video Solution

101. An organic compound C_4H_6 on ozonolysis give $HCHO$, CO_2 , CH_3CHO . Compound will be:






D.




Answer: B

 Watch Video Solution




102. major
product of this reaction is :








Answer: B

 Watch Video Solution



Product (C) in the above reactions is :






D.

Answer: D

 [Watch Video Solution](#)



in the above reaction is :

A. 2, 2, 3-trimethyl pentane

B. 2, 2, 4-trimethyl pentane

C. 2,2-dimethyl hexane





D. n-octane

Answer: B

 Watch Video Solution




(Q) is :




Answer: B

 Watch Video Solution




106.


Product (C) of the reaction is:




A.



B.



C.





D.


Answer: C




107. What is the major product expected from the following reaction ?




A.



B.



C.





D.


Answer: B

 Watch Video Solution


108. Choose the correct product of this reaction :



A.



B.





C.

D. 

Answer: B

 Watch Video Solution




109.

A, Product

A is:


A. 



B.



C.



D.

Answer: D

 Watch Video Solution




110.


Product,

Product is :

A. 



B.




C.





D. 

Answer: D

 [View Text Solution](#)

111. Choose the correct product of the following reactions :





Answer: C

 Watch Video Solution

112. How many stereoisomeric tetrabromides will be formed in the following reaction ?



A. 2

B. 3


C. 4

D. 6

Answer: B

 [Watch Video Solution](#)


113. How many stereoisomeric pentabromides will be formed in the following reaction ?



- A. 2
- B. 3
- C. 4
- D. None of these

Answer: A

 Watch Video Solution



Identify

(Z) in the above sequence of reactions :



B.



C.




D.



Answer: B

 Watch Video Solution

115.




major product (A) of the above reaction :

A.




B.






C.



D.

Answer: C

 [Watch Video Solution](#)



116.





(only one enantiomer is taken) Which of the following statement is correct about A and B ?

- A. A and B are mixture of diastereomers
- B. A and B are mixture of enantiomer
- C. A and B are optically active
- D. B is racemic mixture

Answer: A

 Watch Video Solution

117. Complete the following reaction






D.

Answer: B

 Watch Video Solution



118.



Product (Y) of the above reaction is :



A.



B.



Answer: B


 [Watch Video Solution](#)

119. In the reaction $Me-C \equiv C - Et \xrightarrow{Na / lig. NH_3} P \xrightarrow[CCl_4]{Br_2} (Q)$, then Q is :

- A. A pure compound which is optically inactive due to internal compensation
- B. A binary mixture which is optically inactive due to external compensation
- C. A binary mixture which is optically active
- D. A pure compound which is optically inactive due to absence of chiral centre

Answer: B

 Watch Video Solution





120.
Which (π -bond) will reduce first, when above compound undergoes catalytic hydrogenation ?

- A. a
- B. b
- C. c
- D. d


Answer: D



121. Compound A, which is a degradation product of the antibiotic vermiculine has following structure





▶ View Text Solution




122.

Major product (A) is :






B.



C.




D.

Answer: C



Watch Video Solution

123. In the reaction given below, the product would be :



A. a mixture of diastereomers

B. optically active

C. optically pure enantiomer

D. a racemic mixture

Answer: A

 [Watch Video Solution](#)

124. Surprisingly, the reaction shown below goes through classical carbocation. What is the major product of this reaction ?



A. trans-1, 3-dibromocyclohexane

B. cis-1, 3-dibromocyclohexane








C. trans-1, 2-dibromocyclohexane

D. cis-1, 2-dibromocyclohexane

Answer: B

 Watch Video Solution

125. The major product of the reaction given below is :



A. (i) and (ii)





B. (iii) and (iv)

C. (v) and (vi)


D. none of these

Answer: C

126. Which reaction will occur at the fastest rate ?



Answer: D



127.


Above

reaction is known as:

- A. Wurtz reaction
- B. Fittig reaction
- C. Wurtz fittig reaction
- D. Kolbe electrolysis

Answer: C

 [Watch Video Solution](#)



Product A is :

- A. propane
- B. propanol
- C. propanoic acid
- D. propene

Answer: A

 [Watch Video Solution](#)


129. Which of the following compound give diastereomers when treated with Br_2 in CCl_4 ?




A. Methylcyclopentane



B. 1-Methylcyclopentene



C. 3-Methylcyclopentene



D. 4-Methylcyclopentene

Answer: D




Watch Video Solution

130. A mixture of C_2H_6 , C_2H_4 and C_2H_2 is bubbled through alkaline solution of copper (I) chloride, contained in Woulf's bottle. The gas coming out is

- A. original mixture
- B. C_2H_6
- C. C_2H_6 and C_2H_4 mixture
- D. C_2H_4 and C_2H_2

Answer: C

 Watch Video Solution



The

number of possible products for x and y is :

- A. 2,4

B. 3,5

C. 3,6

D. 3,4

Answer: B



[Watch Video Solution](#)

132. Select the incorrect statement :

A. Bromine is more selective and less reactive

B. Chlorine is less selective and more reactive

C. Benzyl free radical is more stable than 2° free radical

D. Vinyl free radical more stable than allyl free radical

Answer: D




[Watch Video Solution](#)


133. Which of the following compound does not evolve CO_2 gas, when undergo oxidative ozonolysis ?



A.



B.



D.

Answer: D

 Watch Video Solution

134. cis-3-hexene $\xrightarrow{(a)}$ meso 3,4-hexanediol trans-3-hexene $\xrightarrow{(b)}$ meso 3,4-hexanediol. Choose pair of reagent (a, b) for above conversions.

A. Cold $KmNO_4$, OsO_4


B. cold $KmnO_4$, RCO_3H / H_3O^+

C. RCO_3H / H_3O^+ cold $KmnO_4$

D. None of these

Answer: B

 Watch Video Solution



Product (C) of the above reaction is :

A. 1,3-hexadiene

B. 1,4-pentadiene

C. 1,3-butadiene

D. 1,3-heptadiene

Answer: B

 [Watch Video Solution](#)

136. How many carbon-hydrogen bond orbitals are available for overlap with the vacant p-orbital in ethyl carbocation ?

A. 0


B. 3

C. 5

D. 6

Answer: B

 [Watch Video Solution](#)




137.

To achieve above conversion, the reagents used will be :


- A. $O_3 / H_2O_2, HO^- / \Delta$
- B. $HBr, alcKOH, O_3, LiAlH_4, H^+ / \Delta$
- C. $HBr, t - buOK, O_3, KMnO_4, \Delta$
- D. $HCl, KMnO_3(cold) , H^+ / \Delta$

Answer: B


 Watch Video Solution




138.




A.



B.



C.




D.

Answer: B

 Watch Video Solution

139. Decreasing order of heat evolved upon catalytic hydrogenation of given reactants with a H_2 (Pd/C) is :



A. $b > c > a > d$

B. $d > a > c > b$





C. $d > c > a > b$

$$D. c > b > c > d$$

Answer: B

 Watch Video Solution

140.



The correct order of heat of hydrogenation of given molecules is :

A. $d > c > a > b$

B. $d > c > b > a$


C. $b > a > c > d$

D. $d > a > c > b$





Answer: C

 Watch Video Solution

141.




Product (A) of the above reaction is :



D.

Answer: B




142.


Product (A) is :




A.



B.



C.





D.


Answer: A

 Watch Video Solution


143. What is the product of 1, 4-addition in the reaction shown below?



C.




D.



Answer: D

 [Watch Video Solution](#)

144.




Dehydration of the above compound will give :

- A. meso product
- B. racemic mixture
- C. diastereomer
- D. optically pure enantiomer

Answer: B

 Watch Video Solution




145.

What is stereochemistry of product ?

- A. Racemic mixture
- B. Optically inactive
- C. Diastereomers
- D. Meso product

Answer: A

 Watch Video Solution



146.

End product formed in the above reaction is :

- A. Optically active
- B. Racemic
- C. Meso
- D. Diastereomer

Answer: D

 [Watch Video Solution](#)

147. How many moles of BH_3 are needed to react completely with 2 mole of 1-pentene in hydroboration-oxidation reaction ?

- A. 2 mole

B. 3 mole

C. 2/3 mole

D. 3/2 mole


Answer: C

 Watch Video Solution


148.




Product (B) in the above reaction is :




A.



B.




C.






D.

Answer: C

 Watch Video Solution



Product of the above reaction is :





D. Both (a) and (b)

Answer: B

 Watch Video Solution


150. In which of the following reactions 1,3-butadiene will be obtained as a major product ?






D. All of these


Answer: B

 Watch Video Solution




Identify A.







Answer: B


 Watch Video Solution




152.

Product (A) is :






C.



D.

Answer: B

 Watch Video Solution



Bromination take place at :

A. a

B. b


C. c

D. d

Answer: A

 [Watch Video Solution](#)

154. Which is incorrect statement about heats of combustion ?








C. *Iso - bute* \neq $>$ *trans - 2 - bute* \neq $>$ *1 - bute* \neq

D. *n - Hexa* \neq $<$ *n - Heptane* $<$ *n - Octane*

Answer: C

 [Watch Video Solution](#)


155. Predict the major product of the reaction.



Answer: B

 Watch Video Solution

156.



→ cold dil. KMnO_4 → Product of the reaction is:

product


of the reaction is :

- A. Meso compound
- B. Enantiomeric pair
- C. Diastereomers
- D. Optically pure enantiomer

Answer: B







Watch Video Solution




157.

Product (A) of above reaction is:



Answer: D

[Watch Video Solution](#)




158. Products,

Comment up on optical activity of products.





- A. Diastereomers
- B. Racemic mixture
- C. Meso
- D. Optically pure enantiomer

Answer: B


 [Watch Video Solution](#)



Addition of a mineral acid to an olefin bond leads to major product, Identify it:





Answer: C




In polyenes that contain differently substituted ($C=C$) double bonds, it is possible to hydrogenate chemoselectively one ($C=C$) double bond.

Product is :






B.



C.



D.

Answer: B

 [Watch Video Solution](#)



MCPBA \rightarrow meta-chloro perbenzoic acid) Stereochemistry of the product of above reaction is :

A. Meso

B. Racemic

C. Diastereomers

D. Optically inactive due to absence of chiral center.


Answer: B

 [Watch Video Solution](#)




162.


Identify product (P).




A.



B.




C.



D.


Answer: B







163.

A isomerise to B on addition of traces of acid H_2SO_4 . Compound (B) is :






B.



C.




D.

Answer: C




Watch Video Solution




164.


Product (A) of the reaction is :



A.



B.




C.

D. None of these

Answer: B

 [Watch Video Solution](#)




165.

Product


(A) is :




A.



B.



C.



D.


Answer: C



Watch Video Solution

166. Which of the following reactions do not represent the major product

of given Birch reductions ?



A. (i), (iii), (vi)


B. (iv), (vi), (vii)

C. (iv), (v), (vi)

D. (i), (ii), (v), (vii)




Answer: B


 Watch Video Solution



167. Product (A) is:


Product (A) is:





Answer: B

 [Watch Video Solution](#)




168.

Correct statement about above reaction is:

- A. A = cis-2-chlorocyclohexanol,
- B. A = trans-2-chloro cyclohexanol,
- C. A = trans-2-chlorocyclohexanol,
- D. A = cis-2-chlorocyclohexanol,

Answer: C



 Watch Video Solution




169.

Predict


the major product:



B.




C.



D.


Answer: C

 Watch Video Solution




170.


(A) is :




A.



B.




C.



D.

Answer: C


 Watch Video Solution




171.

Major


product of the reaction is :




A.



B.



C.



D.

Answer: B



Watch Video Solution



172.


stereochemistry of the product is:

- A. Diastereomers
- B. Racemic mixture
- C. Meso
- D. Pure Enantiomers

Answer: A



[Watch Video Solution](#)




obtained is/are :

- A. Diastereomers
- B. Meso
- C. Racemic
- D. Optically pure enantiomers


Answer: B

 [Watch Video Solution](#)



174. Product


(x) is :




A.



B.



C.




D.




Answer: B



Watch Video Solution



product (A) is :




D. none

Answer: B



Watch Video Solution



Comment on optical activity of the products:

A. Racemic



B. Diastereomer

C. Meso

D. Optically pure enantiomer

Answer: A


 Watch Video Solution



177.

Product


(al is :




A.



B.



C.




D.

Answer: B



Watch Video Solution



178. Alkene (A) will be :

Alkene (A) will be :

A. cis-2-pentene


B. cis-2-hexene

C. cis-4-octene

D. trans-2-hexene

Answer: C

 [Watch Video Solution](#)



179. Product (A) is





Product (A) is

- A. trans-2-butane
- B. cis-2-butene
- C. 1-butene
- D. Iso-butene

Answer: B

 [Watch Video Solution](#)


180. In which of the following reactions, two products will be formed other than phosphonium ylide (POPh_3)



Answer: C

 Watch Video Solution

181. To carry out the given conversions, select the correct option:



A. $a = \text{Ag}_2\text{O}$, $b = \text{Zn} / \text{CH}_3\text{CO}_2\text{H}$, $c = \text{LiAlH}_4$

B. $a = \text{H}_2\text{O}_2$, $b = \text{CH}_3 - \text{S} - \text{CH}_3$, $c = \text{NaBH}_4$




C. Both (a) and (b)


D. None of these

Answer: C


 Watch Video Solution

182. The product (A) of given alkoxymercuration de-mercuration is :






C.





D.


Answer: B

 Watch Video Solution



End product of the reaction is :









Answer: A

 [Watch Video Solution](#)

184. Major product of the given reaction is :



Answer: B

 [Watch Video Solution](#)


185. The rate constant for a reaction can be increased by a the stability of the reactant or by b the stability of the transition state. Select the correct choice for a and b.


- A. decreasing, decreasing
- B. increasing, decreasing
- C. decreasing, increasing
- D. increasing, increasing

Answer: C

 [Watch Video Solution](#)

186. Major product of the given reaction is :






A.



B.




C.



Answer: C





Watch Video Solution



Major product (A) is :


187.

Major product (A) is :





C.








D.

Answer: C




[Watch Video Solution](#)

188. In the given reaction, only one alkene undergo preferential oxidation by electrophilic ozone. Identify product (P) of the given reaction:



Answer: B

Watch Video Solution




189. Product


(P) is:




A.



B.



C.






D.


Answer: D

 Watch Video Solution



190. Major product of the reaction is :






D.

Answer: B

 Watch Video Solution







Product (B) is :

191.

Product





(B) is :



Answer: C

 Watch Video Solution

192. Isobutene, in the presence of H_2SO_4 , forms a mixture of two isomeric alkene (C_8H_{16}). The major alkene is :




Answer: B



 Watch Video Solution

193. An unknown alkene (A) reacts with 3 mole of H_2 gas in presence of platinum catalyst to form 1-isopropyl-4-methyl cyclohexane. When


unknown alkene (A) is ozonized and reduced, following product are obtained




The alkene (A) is :



B.




C.






D.

Answer: B

 [Watch Video Solution](#)



Product (C) is :



D. Both (a) and (b)


Answer: B

 Watch Video Solution


195. The following reaction take place in high yields.




Use your knowledge of alkene chemistry to predict a product even though you have never seen this reaction before




A.



B.




C.



Answer: B

 Watch Video Solution



What is the ratio of glyoxal to pyruvaldehyde obtained in the above reaction ?


A. 1 : 3

B. 3 : 1




C. 1 : 2

D. 2 : 3

Answer: C




Which of the following product cannot be obtained in above reaction ?







D. None of these

Answer: C




Product (A) is :






Answer: B

 [Watch Video Solution](#)



Product (A) of the reaction is :



D. None of these

Answer: A




Watch Video Solution




200.


Product (A) is :




A.



B.




C.




D.

Answer: B


[▶ Watch Video Solution](#)



(a)



(b)



(c)

201.

Arrange the above in the decreasing order of reactivity towards HBr :

A. a gtbgtc

B. b gtagtc





C. b gtcgta

D. a gtcgtb

Answer: B

 Watch Video Solution


202. Which reaction has the lowest ΔG^\ddagger or (Activation-Energy)?



Answer: D

 Watch Video Solution

203. Which of the following will rearrange ?



A. 1

B. 1 and 3


C. All

D. 1,2,4


Answer: C

 Watch Video Solution

204. Which of the following is most likely to undergo a favorable hydride shift ?




A.



B.



C.




D.

Answer: A

 Watch Video Solution

205. Energy profile diagram for dehydration of 2-butanol using conc.

H_2SO_4 is given below:



Product (b) of above reaction is :

- A. 1-butene
- B. cis-2-butene
- C. trans-2-butene
- D. iso-butene

Answer: B



Watch Video Solution

206. How many alkene on catalytic hydrogenation given isopentane as a product ?

A. A) 2

B. B) 3


C. C) 4

D. D) 5

Answer: B


 Watch Video Solution

207. Which of the following would not rearrange to a more stable form?






B.



C.




D.

Answer: C

Watch Video Solution

208. Consider the following reaction.



In this reaction SbF_5 acts as:

A. an acid


B. a base

C. a nucleophile


D. an electrophile


Answer: D

 [Watch Video Solution](#)




Product (Z) is:






B.



C.




D.

Answer: C



Watch Video Solution



210.


Relation between (B) and (C) is:





- A. Enantiomer
- B. Diastereomer
- C. Geometrical isomer
- D. Meso

Answer: B::C

 Watch Video Solution

211. The reaction of HBr with the following compound would produce :





Answer: B

 Watch Video Solution

212.  is an example of: _____ is an example of:

A. Nucleophilic addition

B. Nucleophilic substitution

C. Electrophilic substitution

D. Electrophilic addition

Answer: C



[Watch Video Solution](#)

213. Olefins can be hydrogenated by :

A. A) Zinc and HCl

B. B) Nascent hydrogen

C. C) Raney Ni and H


D. D) Lithium hydride in ether

Answer: C




[Watch Video Solution](#)

214. What are the products obtained on hydroboration-oxidation of the



given alkene



A. I and III

B. II and IV


C. II and VI

D. III and V

Answer: D

 Watch Video Solution

215.







Relation between A and B, C and D are :

- A. Position, chain
- B. Position, Functional
- C. Chain, Identical
- D. Metamer, Functional

Answer: B

 Watch Video Solution

216. In which reaction syn addition doesn't take place.



Answer: D

 [Watch Video Solution](#)

217. Number of olefin of $C_4H_8 \xrightarrow{HBr}$ Number of Markonikow product
(X) (y)

(including stereo) x+y is :

A. 5

B. 6

C. 7

Answer: C

 Watch Video Solution

Level 2

Reagents

| | | | |
|----------------------------------|--|---|--|
| A. HCl | B. Br ₂ | C. Hg(OAc) ₂ in H ₂ O | D. B ₂ H ₆ (BH ₃) in ether |
| E. H ₂ O ₂ | F. KMnO ₄ in H ₂ O | G. HOBr | H. NaBH ₄ |

1.

In each reagent box write a letter designating the best reagent and condition selected from the above list of reagents.

| Reactant | Reagent | | Product |
|---|---------|--------------------------|---|
| $(\text{CH}_3)_2\text{CHCH}=\text{CH}_2$ 3-methyl-1-butene | (i) | <input type="checkbox"/> | $(\text{CH}_3)_2\text{CHCH}(\text{Cl})\text{CH}_3$ 2-Chloro-3-methyl butane |
| | (ii) | <input type="checkbox"/> | $(\text{CH}_3)_2\text{CHCHBrCH}_2\text{Br}$ 1,2-dibromo-3-methyl butane |
| | (iii) | <input type="checkbox"/> | $(\text{CH}_3)_2\text{CHCHOHCH}_2\text{Br}$ 1,bromo-3-methyl 2-butanol |
| | (iv) | <input type="checkbox"/> | $(\text{CH}_3)_2\text{CHCH}(\text{OH})\text{CH}_3$ 3-methyl-2-butanol |
| | (v) | <input type="checkbox"/> | $(\text{CH}_3)_2\text{CHCH}(\text{OH})\text{CH}_2\text{OH}$ 3-methyl-1, 2 butanediol |

 Watch Video Solution

2. Propene ($CH_3-CH=CH_2$) can be transformed to compounds (a to j) listed in the left-hand column. Write letter designating the reagent, you believe will achieve desired transformation. In the case of a multi step sequence write the reagent in the order they are to be used.











| Desired Product | No. of Steps | Write options | Reagent List |
|---------------------|--------------|---------------|--------------------------|
| a. $CH_3CHBrCH_2Br$ | one | | A. $Hg(OAc)_2$ in H_2O |
| b. $(CH_3)_2CHOH$ | two | | B. B_2H_6 in THF |

| | | | |
|--------------------------|-------|--|----------------------------------|
| c. $CH_3CH_2CH_2OH$ | two | | C. $NaBH_4$ in alcohol |
| d. CH_3COCH_3 | three | | D. Br_2 in CH_2Cl_2 |
| e. CH_3CH_2CHO | three | | E. H_2O_2 in aqueous base |
| f. $CH_3CH(OH)CH_2Br$ | one | | F. HOBr (NBS in aqueous acetone) |
| g. $(CH_3)_2CHBr$ | one | | G. HBr in CH_2Cl_2 |
| h, k. $CH_3CH(OH)CH_2OH$ | two | | H. OsO_4 in ether |
| i. $CH_3-CH_2-CH_2-Cl$ | three | | I. Thionyl chloride ($SOCl_2$) |
| j. $CH_3-C \equiv CH$ | two | | J. $NaHSO_3$ in aqueous acetone |
| | | | K. NaOH in alcohol and reflux |
| | | | L. $NaNH_2$ (strong base) |



Watch Video Solution


3. In each reaction box write a single letter designating the best reagent and condition selected from the list at bottom of the page. (F.S., → first step, S.S. → second step, T.S. → third step)

| Reaction | Reactant | Options | Product |
|----------|--|--|--|
| 1. |  | $\xrightarrow[\text{S.S. } \square]{\text{F.S. } \square}$ |  |
| 2. |  | $\xrightarrow[\text{T.S. } \square]{\text{S.S. } \square}$ |  |
| 3. |  | $\xrightarrow[\text{S.S. } \square]{\text{F.S. } \square}$ |  |
| 4. |  | $\xrightarrow[\text{S.S. } \square]{\text{F.S. } \square}$ |  |
| 5. |  | $\xrightarrow[\text{T.S. } \square]{\text{S.S. } \square}$ |  |

| | | | |
|--|--|--|--|
| A. $\text{NaBH}_4/\text{alcohol}$ | B. $\text{Ph} - \text{CO}_3\text{H}/\text{CH}_2\text{Cl}_2$ | C. PCC | D. $\text{CH}_3\text{ONa}/\text{CH}_3\text{OH}$ |
| E. B_2H_6 in THF | F. $\text{H}_2\text{O}_2/\text{aq. NaOH}$ | G. H_3PO_4 & heat | H. $\text{AlCl}_3/\text{C}_6\text{H}_6$ |
| I. O_3 in CH_2Cl_2 | J. Br_2 in CH_2Cl_2 | K. 20% KOH & heat | L. $\text{Ph} - \text{Li}/\text{ether}$ |

 [View Text Solution](#)






















4. Match the reagents a-j with products A-J. There is one best product for each reaction.



(x)

The molecule (x) is the starting material for all reactions in problem. Do

the ones you know first and then tackle the rest by deductive reasoning

| Products | | Reagents | Option |
|--|---|---|--------|
|  |  | (a) H_2O heat, pH 7 | |
|  |  | (b)  | |
|  |  | (c) tBuOK , polar aprotic solvent | |
|  |  | (d) (1) O_3 , ether (2) H_2O , NaOH , H_2O_2 | |
|  |  | (e) $\text{Br}_2, \text{CCl}_4$ | |
|  |  | (f) NBS , $\text{h}\nu$, CCl_4 | |
|  |  | (g) (1) H_3O^+ (2) NaOH , H_2O | |
|  |  | (h) (1) BH_3 , ether (2) H_2O_2 | |
|  |  | (i) (1) OsO_4 (2) NaOH , H_2O | |
|  |  | (j) $\text{H}_2/\text{Pd}/\text{C}(\text{EtOH})$ | |







View Text Solution

5. Match the column :

| Column (I) | | Column (II) | |
|------------|---|-------------|---|
| (a) | $\text{CH}_3-\text{C}\equiv\text{C}-\text{CH}_3$ | (p) | cis-product with $\text{H}_2/\text{Pd} \cdot \text{BaSO}_4$ |
| (b) | $\text{CH}_3-\text{CH}_2-\text{C}\equiv\text{CH}$ | (q) | Trans-product with $\text{Na}/\text{liq. NH}_3$ |
| (c) | $\text{CH}_3-\text{C}\equiv\text{CH}$ | (r) | White with amm. AgNO_3 |
| (d) | $\text{CH}_3-\text{C}\equiv\text{C}-\text{Et}$ | (s) | H_2 gas with Na |

 Watch Video Solution









6. Match the column I with column II and with column III (Matrix).

| Column-I | | Column- II | | Column- III | |
|----------|---|--------------------------|------------------|---|---|
| Reaction | | Nature of product formed | | Number of chiral center present in product. (Consider only one isomer in case of racemic mixture or Diastereomer) | |
| (a) |  | (p) | Racemic mixture | (w) | 0 |
| (b) |  | (q) | Meso | (x) | 1 |
| (c) |  | (r) | Diastereomer | (y) | 2 |
| (d) |  | (s) | Vicinal dihalide | (z) | 3 |



Watch Video Solution




7. Match the column I and II.

| Column (I) | | Column (II) | |
|------------|--|-------------|--|
| | Reaction | | Product |
| (a) |  | (p) |  |
| (b) |  | (q) |  |
| (c) |  | (r) |  |
| (d) |  | (s) |  |



Watch Video Solution

8. Sum of molecular mass of A, B, C, D (i.e. A+B + C + D) is equal to :




 [Watch Video Solution](#)

9. $C_2FClBrI \xrightarrow[Ni]{H_2} (A)$ (exclude stereoisomer)
(all isomers)

(2) C_4H_8 (alkene) $\xrightarrow[Ni]{H_2} (B)$ (exclude stereoisomer) Total number of
(all isomers)

products (A) and (B) is equal to:

 [Watch Video Solution](#)



10.

How many

products are formed by sum of P,Q,R,S ?





Watch Video Solution





11. Vladimir Markovnikov rule : Alkenes undergo electrophilic addition reactions. It is triggered by the acid acting as a electrophile toward π -electrons of the double bond. Markovnikov's rule states that when an unsymmetrically substituted alkene reacts with a hydrogen halide, the hydrogen atom adds to the carbon that has the greater number of hydrogen, e.g.,



Mechanism :



Which of the following is most reactive toward Markovnikov addition ?




D.



Answer: B

 Watch Video Solution


12. Vladimir Markovnikov rule : Alkenes undergo electrophilic addition reactions. It is triggered by the acid acting as a electrophile toward π -electrons of the double bond. Markovnikov's rule states that when an unsymmetrically substituted alkene reacts with a hydrogen halide, the hydrogen atom adds to the carbon that has the greater number of hydrogen, e.g.,






Mechanism :



What is the energy profile for the given reaction ?



A.




Answer: C



 [Watch Video Solution](#)

13. Vladimir Markovnikov rule : Alkenes undergo electrophilic addition reactions. It is triggered by the acid acting as a electrophile toward π -electrons of the double bond. Markovnikov's rule states that when an unsymmetrically substituted alkene reacts with a hydrogen halide, the




hydrogen atom adds to the carbon that has the greater number of hydrogen, e.g.,



Mechanism :



In which of following reactions carbocation rearrangement is possible ?



D. All of these

Answer: D



[Watch Video Solution](#)

14. Vladimir Markovnikov rule : Alkenes undergo electrophilic addition reactions. It is triggered by the acid acting as a electrophile toward t-




electrons of the double bond. Markovnikov's rule states that when an unsymmetrically substituted alkene reacts with a hydrogen halide, the hydrogen atom adds to the carbon that has the greater number of hydrogen, e.g.,





Mechanism :



Identify the major products r_1 , r_2 , and r_3 in the given reactions.









Answer: B

 Watch Video Solution

15. Vladimir Markovnikov rule : Alkenes undergo electrophilic addition reactions. It is triggered by the acid acting as a electrophile toward π -electrons of the double bond. Markovnikov's rule states that when an unsymmetrically substituted alkene reacts with a hydrogen halide, the hydrogen atom adds to the carbon that has the greater number of hydrogen, e.g.,



In which of the following reactions, product is racemic mixture ?



D. All of these



Answer: D

 [Watch Video Solution](#)




16. Vladimir Markovnikov rule : Alkenes undergo electrophilic addition reactions. It is triggered by the acid acting as a electrophile toward π -electrons of the double bond. Markovnikov's rule states that when an unsymmetrically substituted alkene reacts with a hydrogen halide, the hydrogen atom adds to the carbon that has the greater number of hydrogen, e.g.,



Mechanism :



In which of the following reactions, diastereomers will be formed ?




D. All of these

Answer: D




Watch Video Solution

17.



What is electrophile in first step?




Answer: B




Watch Video Solution


18.



What is nucleophile in first step?



B. 1-butene




Answer: B






Watch Video Solution

19.




What is electrophile in second step ?






Answer: C

 Watch Video Solution

20.




What is nucleophile in second step ?



Answer: B

 Watch Video Solution

21.



Which step is rate determining step ?

A. attack of nucleophile CH_3OH

B. attack of electrophile H^{\oplus}

C. attack of nucleophile H_2O

D. attack of electrophile $\overset{\oplus}{C}H_3$

Answer: B

 Watch Video Solution

22. Match the column I and II :

| Column (I) | | Column (II) |
|------------|--|--|
| Conversion | | Reagent |
| (a) | | (p) $\text{SO}_2\text{Cl}_2 / h\nu$ (2 equivalent) |
| (b) | | (q) NBS (2 equivalent) |
| (c) | | (r) NBS then $\text{SO}_2\text{Cl}_2/h\nu$ |
| (d) | | (s) $\text{SO}_2\text{Cl}_2 / h\nu$ then NBS |



Watch Video Solution