



MATHS

BOOKS - ARIHANT MATHS

COORDINATE SYSTEM AND COORDINATES

Example

1. Draw the polar coordinates

$$\left(2, \frac{\pi}{3}\right), \left(-2, \frac{\pi}{3}\right), \left(-2, -\frac{\pi}{3}\right) \text{ and } \left(2, -\frac{\pi}{3}\right)$$

on the plane.



Watch Video Solution

2. Draw the polar coordinate $\left(3, \frac{5\pi}{4}\right)$ on the plane.



[Watch Video Solution](#)

3. Find the cartesian coordinates of the points whose polar coordinates are

$$\left(5, \pi - \tan^{-1}\left(\frac{4}{3}\right)\right)$$



[Watch Video Solution](#)

4. Find the cartesian coordinates of the points whose polar coordinates are

$$\left(5\sqrt{2}, \frac{\pi}{4}\right)$$



[Watch Video Solution](#)

5. Find the polar coordinates of the points whose cartesian coordinates are

$$(-2, -2)$$



[Watch Video Solution](#)

6. Find the polar coordinates of the points whose cartesian coordinates are

$$(-3, 4)$$



[Watch Video Solution](#)

 Watch Video Solution

7. Transform to Cartesian coordinates the equations:

$$r^2 = a^2 \cos 2\theta$$



Watch Video Solution

8. Transform the equation $x^2 + y^2 = ax$ into polar form.



Watch Video Solution

9. Prove that the distance of the point $(a \cos \alpha, a \sin \alpha)$ from the origin is independent of α

 [Watch Video Solution](#)

10. The distance between the points $(a \cos \alpha, a \sin \alpha)$ and $(a \cos \beta, a \sin \beta)$ where $a > 0$

 [Watch Video Solution](#)

11. If $P(x, y)$ is a point equidistant from the points $A(6, -1)$ and $B(2, 3)$, show that $x - y = 3$.

 [Watch Video Solution](#)

12. Using distance formula, show that the points $(1, 5)$, $(2, 4)$ and $(3, 3)$ are collinear.



Watch Video Solution

13. An equilateral triangle has one vertex at $(0, 0)$ and another at $(3, \sqrt{3})$. What are the coordinates of the third vertex ?



Watch Video Solution

14. By using the concept of slope, show that the points $(-2,-1)$, $(4,0)$, $(3,3)$ and $(-3,2)$ are the vertices of a parallelogram.



Watch Video Solution

15. Let the opposite angular points of a square be $(3, 4)$ and $(1, -1)$. Find the coordinates of the remaining angular points.



Watch Video Solution

16. Find the circumcentre of the triangle whose vertices are $(-2, -3)$, $(-1, 0)$ and $(7, -6)$. Also find the radius of the circumcircle.



[Watch Video Solution](#)

17. If the segments joining the points $A(a, b)$ and $B(c, d)$ subtends an angle θ at the origin, prove that : $\theta = \frac{ac + bd}{(a^2 + b^2)(c^2 + d^2)}$



[Watch Video Solution](#)

18. Show that the triangle, the coordinates of whose vertices are given by integers, can never be an equilateral triangle.



[Watch Video Solution](#)

19. In any triangle ABC, prove that $AB^2 + AC^2 = 2(AD^2 + BD^2)$, where D is the midpoint of BC.



[Watch Video Solution](#)

20. Let ABCD be a rectangle and P be any point in its plane. Show that $PA^2 + PC^2 = PB^2 + PD^2$ using coordinate geometry.



Watch Video Solution

21. Prove that the points $(0, 0)$, $(3, \frac{\pi}{2})$ and $(3, \frac{\pi}{6})$ are the vertices of an equilateral triangle.



Watch Video Solution

22. Find the coordinates of the point which divides the line segment joining the points $A(5, -2)$ and

$B(9, 6)$ in the ratio 3:1



[Watch Video Solution](#)

23. Find the lengths of the medians of a triangle whose vertices are $A(-1, 3)$, $B(1, -1)$ and $C(5, 1)$.



[Watch Video Solution](#)

24. In what ratio does the line $y - x + 2 = 0$ cut the line joining $(3, -1)$ and $(8, 9)$?



[Watch Video Solution](#)

25. The coordinates of three consecutive vertices of a parallelogram are $(1, 3)$, $(-1, 2)$ and $(2, 5)$. Then find the coordinates of the fourth vertex.



Watch Video Solution

26. In what ratio does the x-axis divide the line segment joining the points $(2, -3)$ and $(5, 6)$?



Watch Video Solution

27. If the coordinates of the mid-points of the sides of a triangle are $(1, 2)$, $(0, -1)$ and $(2, -1)$. Find the

coordinate of its vertices.



[Watch Video Solution](#)

28. Prove that the mid-point of the hypotenuse of a right triangle is equidistant from its vertices.



[Watch Video Solution](#)

29. The line-segment joining the mid-points of two sides of a triangle is parallel to the third side and Of it.



[Watch Video Solution](#)

30. Find the coordinates of a point which divides externally the line joining $(1, -3)$ and $(-3, 9)$ in the ratio $1 : 3$.



Watch Video Solution

31. The line segment joining $A(6, 3)$ to $B(-1, -4)$ is doubled in length by having its length added to each end, then the ordinates of new ends are



Watch Video Solution

32. Using section formula show that the points $(1,-1)$, $(2, 1)$ and $(4, 5)$ are collinear.

 [Watch Video Solution](#)

33. Find the ratio in which the point $(2, y)$ divides the line segment $(-4,3)$ and $(6,3)$. hence find the value of y

 [Watch Video Solution](#)

34. Find the harmonic conjugates of the point $R(5, 1)$ with respect to the points $P(2, 10)$ and $Q(6, -2)$

 [Watch Video Solution](#)

35. Two vertices of a triangle are $(-1, 4)$ and $(5, 2)$. If its centroid is $(0, -3)$, find the third vertex.

 [Watch Video Solution](#)

36. The vertices of a triangle are $(1, 2)$, $(h, -3)$ and $(-4, k)$. Find the value of $\sqrt{\{(h + k)^2 + (h + 3k)^2\}}$. If the centroid of the triangle be at point $(5, -1)$.

 [Watch Video Solution](#)

37. If $D(-2, 3)$, $E(4, -3)$ and $F(4, 5)$ are the mid-points of the sides BC , CA and AB of the triangle ABC , then find $\sqrt{(|AG|^2 + |BG|^2 - |CG|^2)}$ where, G is the centroid of $\triangle ABC$.

 [Watch Video Solution](#)

38. find the derivative of $\sin(4x-9)$

 [Watch Video Solution](#)

39. If G be the centroid of a triangle ABC , prove that, \hat{A}
 $AB^2 + BC^2 + CA^2 = 3(GA^2 + GB^2 + GC^2)$



Watch Video Solution

40. The vertices of a triangle are $(1, a)$, $(2, b)$ and $(c^2 - 3)$

Find the condition that the centroid may lie on the X-axis.



Watch Video Solution

41. The vertices of a triangle are $(1, a)$, $(2, b)$ and $(c^2, -3)$. (i) Prove that its centroid can not lie on the y-axis. (ii) Find the

condition that the centroid may lie on the x-axis for any value of $a, b, c \in \mathbb{R}$.

 [Watch Video Solution](#)

42. Find the coordinates of incentre of the triangle whose vertices are $(4, -2)$, $(-2, 4)$ and $(5, 5)$.

 [Watch Video Solution](#)

43. If $\left(\frac{3}{2}, 0\right)$, $\left(\frac{3}{2}, 6\right)$ and $(-1, 6)$ are mid-points of the sides of a triangle, then find

Incentre of the triangle

 [Watch Video Solution](#)

 [Watch Video Solution](#)

44. If $\left(\frac{3}{2}, 0\right)$, $\left(\frac{3}{2}, 6\right)$ and $(-1, 6)$ are mid-points of the sides of a triangle, then find

Centroid of the triangle

 [Watch Video Solution](#)

45. If a vertex of a triangle be $(1, 1)$ and the middle points of the sides through it be $(-2, 3)$ and $(5, 2)$, find the other vertices.

 [Watch Video Solution](#)

46. If G is the centroid and I the in-centre of the triangle, with vertices $A(-36, 7)$, $B(20, 7)$ and $C(0, -8)$, then, find the value of GI



Watch Video Solution

47. If the coordinates of the mid-points of the sides of a triangle are $(1, 1)$, $(2, -3)$ and $(3, 4)$, find the vertices of the triangle.



Watch Video Solution

48. In a ΔABC with vertices $A(1,2)$, $B(2,3)$ and $C(3, 1)$

and

$$\angle A = \angle B = \cos^{-1} \left(\frac{1}{\sqrt{10}} \right), \angle C = \cos^{-1} \left(\frac{4}{5} \right),$$

then find the circumcentre of ΔABC .



Watch Video Solution

49. Find the circumcentre of the triangle whose vertices are $(2, 2)$, $(4, 2)$ and $(0, 4)$.



Watch Video Solution

50. Find the orthocentre of $\triangle ABC$ if $A \equiv (0, 0)$, $B \equiv (3, 5)$ and $C \equiv (4, 7)$.

 [Watch Video Solution](#)

51. If a triangle has its orthocenter at $(1,1)$ and circumcentre $(\frac{3}{2}, \frac{3}{4})$ then centroid is:

 [Watch Video Solution](#)

52. The coordinates of A , B , C are $(6, 3)$, $(-3, 5)$ and $(4, -2)$ respectively and P is any point (x, y) .

Show that the ratio of the areas of triangles PBC

and ABC is $\left| \frac{x + y - 2}{7} \right|$.



[Watch Video Solution](#)

53. Find the area of the pentagon whose vertices are

$A(1, 1)$, $B(7, 21)$, $C(7, -3)$, $D(12, 2)$, and

$E(0, -3)$



[Watch Video Solution](#)

54. Prove that the points $(a, 0)$, $(0, b)$ and $(1, 1)$

are collinear if, $\frac{1}{a} + \frac{1}{b} = 1$



[Watch Video Solution](#)

 [Watch Video Solution](#)

55. Prove that the coordinates of the vertices of an equilateral triangle can not all be rational.

 [Watch Video Solution](#)

56. If the coordinates of two points A and B are $(3, 4)$ and $(5, -2)$ respectively. Find the coordinates of any point P , if $PA = PB$ and area of $PAB = 10$ sq. units.

 [Watch Video Solution](#)

57. Find the area of the triangle formed by the straight lines $7x - 2y + 10 = 0$, $7x + 2y - 10 = 0$ and $9x + y + 2 = 0$ (without sloving the vertices of the triangle).



[Watch Video Solution](#)

58. Find the locus of a point which moves such that its distance from the origin is three times its distance from x-axis.



[Watch Video Solution](#)

59. The locus of the moving point P such that $2PA = 3PB$, where A is (0,0) and B is (4,-3), is



[Watch Video Solution](#)

60. The sum of the squares of the distances of a moving point from two fixed points $(a, 0)$ and $(-a, 0)$ is equal to a constant quantity $2c^2$. Find the equation to its locus.



[Watch Video Solution](#)

61. A point moves so that the sum of its distances from $(ae, 0)$ and $(-ae, 0)$ is $2a$, prove that the equation to its locus is $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$, where $b^2 = a^2(1 - e^2)$.

 [Watch Video Solution](#)

62. Find the equation of the locus of a point which moves so that the difference of its distances from the points $(3, 0)$ and $(-3, 0)$ is 4 units.

 [Watch Video Solution](#)

63. The ends of the hypotenuse of a right angled triangle are $(6, 0)$ and $(0, 6)$. Find the locus of the third vertex.



Watch Video Solution

64. Differentiate x^4 with respect to x .



Watch Video Solution

65. Find the locus of a point whose coordinate are given by $x = t + t^2$, $y = 2t + 1$, where t is variable.



Watch Video Solution

66. A stick of length 10 units rests against the floor and a wall of a room. If the stick begins to slide on floor then the locus of its middle point is:

 [Watch Video Solution](#)

67. Find the locus of the point of intersection of lines $x \cos \alpha + y \sin \alpha = a$ and $x \sin \alpha - y \cos \alpha = b$ (α is a variable).

 [Watch Video Solution](#)

68. A variable line cuts X-axis at A, Y -axis at B, where

OA = a, OB = b (O as origin) such that $a^2 + b^2 = 1$.

Find the locus of circumcentre of ΔOAB



[Watch Video Solution](#)

69. Differentiate $\left(\frac{1}{5}\right)x^5$ with respect to x .



[Watch Video Solution](#)

70. Differentiate $2x^4$ with respect to x .



[Watch Video Solution](#)

71. Find the equation of the curve $2x^2 + y^2 - 3x + 5y - 8 = 0$ when the origin is transferred to the point $(-1, 2)$ without changing the direction of axes.



[Watch Video Solution](#)

72. The equation of curve referred to the new axes, axes retaining their directions, and origin $(4, 5)$ is $X^2 + Y^2 = 36$. Find the equation referred to the original axes.



[Watch Video Solution](#)

73. Shift the origin to a suitable point so that the equation $y^2 + 4y + 8x - 2 = 0$ will not contain a term in y and the constant term.



Watch Video Solution

74. At what point the origin be shifted, if the coordinates of a point $(-1, 8)$ become $(-7, 3)$?



Watch Video Solution

75. If the axes are turned through 45° , find the transformed form of the equation

$$3x^2 + 3y^2 + 2xy = 2.$$



[Watch Video Solution](#)

76. Differentiate $\log x$ with respect to x .



[Watch Video Solution](#)

77. Differentiate $\sin x$ with respect to x .



[Watch Video Solution](#)

78. If (x, y) and (X, Y) are the coordinates of the same point referred to two sets of rectangular axes with the same origin and if $ux + vy$, where u and v are independent of x and y , becomes $VX + UY$, show that $u^2 + v^2 = U^2 + V^2$.



[Watch Video Solution](#)

79. Differentiate $\sin x + x$ with respect to x .



[Watch Video Solution](#)

80. Given the equation $4x^2 + 2\sqrt{3}xy + 2y^2 = 1$.

Through what angle should the axes be rotated so that the term xy is removed from the transformed equation.



[Watch Video Solution](#)

81. Find λ if $(\lambda, \lambda + 1)$ is an interior point of $\triangle ABC$

where, $A \equiv (0, 3)$, $B \equiv (-2, 0)$ and $C \equiv (6, 1)$.



[Watch Video Solution](#)

82. Prove that the locus of the centroid of the triangle

whose vertices are

$(a \cos t, a \sin t)$, $(b \sin t, -b \cos t)$ and $(1, 0)$, where

t is a parameter, is circle.

A. $(3x - 1)^2 + (3y)^2 = a^2 - b^2$

B. $(3x - 1)^2 + (3y)^2 = a^2 + b^2$

C. $(3x + 1)^2 + (3y)^2 = a^2 + b^2$

D. $(3x + 1)^2 + 3y^2 = a^2 - b^2$

Answer: B



Watch Video Solution

83. Find the incentre of the triangle with vertices

$A(1, \sqrt{3})$, $B(0, 0)$ and $C(2, 0)$.

A. $\left(1, \frac{\sqrt{3}}{2}\right)$

B. $\left(\frac{2}{3}, \frac{1}{\sqrt{3}}\right)$

C. $\left(\frac{2}{3}, \frac{\sqrt{3}}{2}\right)$

D. $\left(1, \frac{1}{\sqrt{3}}\right)$

Answer: D



Watch Video Solution

84. The orthocentre of the triangle with vertices $(0, 0)$, $(3, 4)$, and $(4, 0)$ is

(a) $\left(3, \frac{5}{4}\right)$ (b) $(3, 12)$ $\left(3, \frac{3}{4}\right)$ (d) $(3, 9)$

A. $\left(3, \frac{5}{4}\right)$

B. $(3, 12)$

C. $\left(3, \frac{3}{4}\right)$

D. $(3, 9)$

Answer: C



Watch Video Solution

85. If x_1, x_2, x_3 as well as y_1, y_2, y_3 are in GP, with the same common ratio, then the points $(x_1, y_1), (x_2, y_2)$ and (x_3, y_3)

- A. lie on a straight line
- B. lie on an ellipse
- C. lie on a circle
- D. are vertices of a triangle

Answer: A



Watch Video Solution

86. Let A be the image of (2, -1) with respect to Y - axis

Without transforming the origin, coordinate axis are

turned at an angle 45° in the clockwise direction.

Then, the coordinates of A in the new system are

A. $\left(-\frac{1}{\sqrt{2}}, -\frac{3}{\sqrt{2}} \right)$

B. $\left(-\frac{3}{\sqrt{2}}, \frac{1}{\sqrt{2}} \right)$

C. $\left(\frac{1}{\sqrt{2}}, \frac{3}{\sqrt{2}} \right)$

D. $\left(\frac{3}{\sqrt{2}}, \frac{1}{\sqrt{2}} \right)$

Answer: A



Watch Video Solution

87. Let S_1, S_2, \dots be squares such that for each $n \geq 1$, the length of a side of S_n equals the length of a diagonal of S_{n+1} . If the length of a side of S_1 is 10cm , then the least value of n is the area of S_n less than 1 sq. cm?

A. 7

B. 8

C. 9

D. 10

Answer: B::C::D



Watch Video Solution

88. Differentiate $\tan x$ with respect to x .



Watch Video Solution

89. ABC is an isosceles triangle. If the coordinates of the base are B(1, 3) and C(-2, 7). The coordinates of vertex A can be

A. $(5/6, 6)$

B. $(-1/2, 5)$

C. $(1, 2)$

D. none of these

Answer: B::C::D



Watch Video Solution

90. Differentiate $\cos x$ with respect to x .



Watch Video Solution

91. Differentiate $\cos x + x$ with respect to x .



Watch Video Solution

92. If $A\left(\alpha, \frac{1}{\alpha}\right), B\left(\beta, \frac{1}{\beta}\right), C\left(\gamma, \frac{1}{\gamma}\right)$ be the vertices of a ΔABC , where α, β are the roots of $x^2 - 6ax + 2 = 0$, β, γ are the roots of $x^2 - 6bx + 3 = 0$ and γ, α are the roots of $x^2 - 6cx + 6 = 0$, a, b, c being positive.

The coordinates of centroid of ΔABC is

A. $\left(-\frac{1}{2}, -2\right)$

B. $\left(-\frac{1}{3}, -3\right)$

C. $\left(2, \frac{11}{18}\right)$

D. $\left(-\frac{1}{6}, -6\right)$

Answer: D



Watch Video Solution

93. Differentiate $\tan x + x$ with respect to x .



Watch Video Solution

94. Statement I : The area of the triangle formed by the points $A(100, 102)$, $B(102, 105)$, $C(104, 107)$ is same as the area formed by $A'(0, 0)$, $B'(2, 3)$, $C'(4, 5)$.

Statement II : The area of the triangle is constant with respect to translation.

A. Statement I is true, Statement II is true,

Statement II is a correct explanation for

Statement I.

B. Statement I is true, Statement II is true,

Statement II is not a correct explanation for

Statement I.

C. Statement I is true, Statement II is false.

D. Statement I is false, Statement II is true.

Answer: A



Watch Video Solution

95. Statement I : If centroid and circumcentre of a triangle are known its orthocentre can be found

Statement II : Centroid, orthocentre and circumcentre of a triangle are collinear.

A. Statement I is true, Statement II is true,

Statement II is a correct explanation for

Statement I.

B. Statement I is true, Statement II is true,

Statement II is not a correct explanation for

Statement I.

C. Statement I is true, Statement II is false.

D. Statement I is false, Statement II is true.

Answer: B



 [Watch Video Solution](#)

96. Differentiate $\cot x + x$ with respect to x .

 [Watch Video Solution](#)

97. Find the multiplication of 11×0

 [Watch Video Solution](#)

98. x coordinates of two points B and C are the roots of equation $x^2 + 4x + 3 = 0$ and their y coordinates are the roots of equation $x^2 - x - 6 = 0$. If x

coordinate of B is less than the x coordinate of C and y coordinate of B is greater than the y coordinate of C and coordinates of a third point A be $(3, -5)$, find the length of the bisector of the interior angle at A.



[Watch Video Solution](#)

99. Differentiate $\log x^2$ with respect to x .



[Watch Video Solution](#)

100. Differentiate 5^x with respect to x .



[Watch Video Solution](#)

101. Let S be a square of unit area. Consider any quadrilateral which has one vertex on each side of S . If a, b, c and d denote the lengths of sides of the quadrilateral, prove that $2 \leq a^2 + b^2 + c^2 + d^2 \leq 4$

 [Watch Video Solution](#)

102. The circumcentre of a triangle having vertices $A(a, a \tan \alpha), B(b, b \tan \beta), C(c, c \tan \gamma)$ is at origin, where $\alpha + \beta + \gamma = \pi$. Then the orthocentre lies on

 [Watch Video Solution](#)

Single Option Correct Type Questions

1. Differentiate $\cot x$ with respect to x .



Watch Video Solution

Exercise For Session 1

1. The polar coordinates of the point whose cartesian coordinates are $(-1, -1)$ is

A. $\left(\sqrt{2}, \frac{\pi}{4}\right)$

B. $\left(\sqrt{2}, \frac{3\pi}{4}\right)$

C. $\left(\sqrt{2}, -\frac{\pi}{4}\right)$

D. $\left(\sqrt{2}, -\frac{3\pi}{4}\right)$

Answer: D



Watch Video Solution

2. The cartesian coordinates of the point whose polar coordinates are $\left(13, \pi - \tan^{-1}\left(\frac{5}{12}\right)\right)$ is

A. (12, 5)

B. (-12, 5)

C. (-12, -5)

D. (12, -5)

Answer: B



Watch Video Solution

3. The transform equation of $r^2 \cos^2 \theta = a^2 \cos 2\theta$ to cartesian form is $(x^2 + y^2)x^2 = a^2 \lambda$, then value of λ is

A. $y^2 - x^2$

B. $x^2 - y^2$


C. xy

D. $x^2 y^2$

Answer: B

 Watch Video Solution

4. The coordinates of P' in the figure is



A. $\left(3, \frac{\pi}{3}\right)$

B. $\left(3, -\frac{\pi}{3}\right)$

C. $\left(-3, -\frac{\pi}{3}\right)$


D. $\left(-3, \frac{\pi}{3}\right)$

Answer: B



Watch Video Solution

5. The cartesian coordinates of the point Q in the figure is



- A. $(\sqrt{3}, 1)$
- B. $(-\sqrt{3}, 1)$
- C. $(-\sqrt{3}, -1)$
- D. $(\sqrt{3}, -1)$

Answer: B



Watch Video Solution

6. A point lies on X-axis at a distance 5 units from Y-axis. What are its coordinates ?



[Watch Video Solution](#)

7. A point lies on Y-axis at a distance 4 units from X-axis. What are its coordinates ?



[Watch Video Solution](#)

8. A point lies on negative direction of X-axis at a distance 6 units from Y-axis. What are its coordinates ?



[Watch Video Solution](#)

9. Transform the equation $y = x \tan \alpha$ to polar form.



[Watch Video Solution](#)

10. Transform the equation $r = 2a \cos \theta$ to cartesian form.



[Watch Video Solution](#)

Exercise For Session 2

1. If the distance between the points $(a, 2)$ and $(3, 4)$ be 8, then a equals to

A. $2 + 3\sqrt{3}$

B. $2 - 3\sqrt{15}$

C. $2 \pm 3\sqrt{15}$

D. $3 \pm 2\sqrt{15}$

Answer: D



Watch Video Solution

2. The three points $(-2, 2)$, $(8, -2)$ and $(-4, -3)$ are the vertices of

- A. an isosceles triangle
- B. an equilateral triangle
- C. a right angled triangle
- D. None of these

Answer: C

 [Watch Video Solution](#)

3. The distance between the points $\left(3, \frac{\pi}{4}\right)$ and $\left(7, \frac{5\pi}{4}\right)$

A. 8

B. 10

C. 12

D. 14

Answer: B



Watch Video Solution

4. Let $A(6, -1)$, $B(1, 3)$ and $C(x, 8)$ be three points such that $AB = BC$ then the value of x are

A. 3, 5

B. $-3, 5$

C. $3, -5$

D. $-3, -5$

Answer: B



Watch Video Solution

5. The points $(a + 1, 1)$, $(2a + 1, 3)$ and $(2a + 2, 2a)$ are collinear if

A. $a = -1, 2$

B. $a = \frac{1}{2}, 2$

C. $a = 2, 1$

$$D. a = -\frac{1}{2}, 2$$

Answer: D

 **Watch Video Solution**

6. Let $A = (3, 4)$ and B is a variable point on the lines $|x| = 6$. If $AB \leq 4$, then find the number of position of B with integral coordinates.

A. 5

B. 6

C. 10

D. 12

Answer: A



Watch Video Solution

7. The number of points on X-axis which are at a distance c units ($c < 3$) from $(2, 3)$ is

A. 1

B. 2

C. 0

D. 3

Answer: C



Watch Video Solution

8. The point on the axis of y which its equidistant from $(-1, 2)$ and $(3, 4)$, is

A. $(0, 3)$

B. $(0, 4)$

C. $(0, 5)$

D. $(0, -6)$

Answer: C



9. Find the distance between the points $(at_1^2, 2at_1)$ and $(at_2^2, 2at_2)$, where t_1 and t_2 are the roots of the equation $x^2 - 2\sqrt{3}x + 2 = 0$ and $a > 0$.



Watch Video Solution

10. If P and Q are two points whose coordinates are $(at^2, 2at)$ and $\left(\frac{a}{t^2}, \frac{2a}{t}\right)$ respectively and S is the point $(a, 0)$. Show that $\frac{1}{SP} + \frac{1}{sQ}$ is independent of t.



Watch Video Solution

11. Show that the points $(3, 4)$, $(8, -6)$ and $(13, 9)$ are the vertices of a right angled triangle.

 [Watch Video Solution](#)

12. Show that four points $(0, -1)$, $(6, 7)$, $(-2, 3)$ and $(8, 3)$ are the vertices of a rectangle. Also, find its area.

 [Watch Video Solution](#)

13. Find the circumcentre and circumradius of the triangle whose vertices are $(-2, 3)$, $(2, -1)$ and $(4,$

0).



[Watch Video Solution](#)

14. The vertices of a triangle are $A(1, 1)$, $B(4, 5)$ and $C(6, 13)$. Find $\cos A$.



[Watch Video Solution](#)

15. The opposite vertices of a square are $(2, 6)$ and $(0, -2)$. Find the coordinates of the other vertices.



[Watch Video Solution](#)

16. If the point (x, y) is equidistant from the points $(a + b, b - a)$ and $(a - b, a + b)$, prove that $bx = ay$.



[Watch Video Solution](#)

17. if a and b between 0 and 1 such that the points $(a, 1)$, $(1, b)$ and $(0, 0)$ form an equilateral triangle then the values of 'a' and 'b' respectively



[Watch Video Solution](#)

18. An equilateral triangle has two vertices at the points $(3, 4)$ and $(-2, 3)$, find the coordinates of the third vertex.



[Watch Video Solution](#)

19. If P be any point in the plane of square ABCD, prove that

$$PA^2 + PC^2 = PB^2 + PD^2$$



[Watch Video Solution](#)

1. The coordinates of the middle points of the sides of a triangle are $(4, 2)$, $(3, 3)$ and $(2, 2)$, then coordinates of centroid are

A. $(3, 7/3)$

B. $(3, 3)$

C. $(4, 3)$

D. $(3, 4)$

Answer: A



Watch Video Solution

2. The incentre of the triangle whose vertices are $(-36, 7)$, $(20, 7)$ and $(0, -8)$ is

A. $(0, -1)$

B. $(-1, 0)$

C. $(1, 1)$

D. $\left(\frac{1}{2}, 1\right)$

Answer: B



Watch Video Solution

3. If the orthocentre and centroid of a triangle are $(-3, 5)$ and $(3, 3)$ then its circumcentre is

A. $(6, 2)$

B. $(3, -1)$

C. $(-3, 5)$

D. $(-3, 1)$

Answer: A



Watch Video Solution

4. An equilateral triangle has each side to a. If the coordinates of its vertices are (x_1, y_1) , (x_2, y_2) and (x_3, y_3) then the square of the determinat

$$\begin{vmatrix} x_1 & y_1 & 1 \\ x_2 & y_2 & 1 \\ x_3 & y_3 & 1 \end{vmatrix}^2 \text{ equals}$$

A. $3a^4$

B. $\frac{3a^4}{2}$

C. $\frac{3}{4}a^4$

D. $\frac{3}{8}a^4$

Answer: C



Watch Video Solution

5. The vertices of a triangle are $A(0, 0)$, $B(0, 2)$ and $C(2, 0)$. The distance between circumcentre and orthocentre is

A. $\sqrt{2}$

B. $\frac{1}{\sqrt{2}}$

C. 2

D. $\frac{1}{2}$

Answer: A



Watch Video Solution

6. Area of the triangle with vertices (a, b) , (x_1, y_1) and (x_2, y_2) where a, x_1, x_2 are in G.P. with common ratio r and b, y_1, y_2 , are in G.P with common ratio s , is

A. (a) $ab(r - 1)(s - 1)(s - r)$

B. (b) $\frac{1}{2}ab(r + 1)(s + 1)(s - r)$

C. (c) $\frac{1}{2}ab(r - 1)(s - 1)(s - r)$

D. (d) $ab(r + 1)(s + 1)(r - s)$

Answer: C



Watch Video Solution

7. The points $(x + 1, 2)$, $(1, x + 2)$, $\left(\frac{1}{x + 1}, \frac{2}{x + 1}\right)$ are collinear, then x is equal to

A. -4

B. -8

C. 4

D. 8

Answer: A



Watch Video Solution

8. The vertices of a triangle are $(6, 0)$, $(0, 6)$ and $(6, 6)$.

The distance between its circumcentre and centroid is :

A. $2\sqrt{2}$

B. 2

C. $\sqrt{2}$

D. 1

Answer: C



Watch Video Solution

9. The centroid of the triangle with vertices $(1, \sqrt{3})$, $(0, 0)$ and $(2, 0)$ is

A. $\left(1, \frac{\sqrt{3}}{2}\right)$

B. $\left(\frac{2}{3}, \frac{1}{\sqrt{3}}\right)$

C. $\left(\frac{2}{3}, \frac{\sqrt{3}}{2}\right)$

D. $\left(1, \frac{1}{\sqrt{3}}\right)$

Answer: D



Watch Video Solution

10. The vertices of a triangle are $(0, 0)$, $(1,0)$ and $(0,1)$.

Then excentre opposite to $(0, 0)$ is

A. $\left(1 - \frac{1}{\sqrt{2}}, 1 + \frac{1}{\sqrt{2}}\right)$

B. $\left(1 + \frac{1}{\sqrt{2}}, 1 + \frac{1}{\sqrt{2}}\right)$

C. $\left(1 + \frac{1}{\sqrt{2}}, 1 - \frac{1}{\sqrt{2}}\right)$

D. $\left(1 - \frac{1}{\sqrt{2}}, 1 - \frac{1}{\sqrt{2}}\right)$

Answer: B



Watch Video Solution

11. If α, β, γ are the real roots of the equation $x^3 - 3px^2 + 3qx - 1 = 0$, then find the centroid of the triangle whose vertices are $\left(\alpha, \frac{1}{\alpha}\right)$, $\left(\beta, \frac{1}{\beta}\right)$ and $\left(\gamma, \frac{1}{\gamma}\right)$.



[Watch Video Solution](#)

12. If centroid of a triangle be $(1, 4)$ and the coordinates of its any two vertices are $(4, -8)$ and $(-9, 7)$, find the area of the triangle.



[Watch Video Solution](#)

13. Find the coordinates of the orthocentre of the triangle whose vertices are $(1, 2)$, $(2, 3)$ and $(4, 3)$.



Watch Video Solution

14. Show that the area of the triangle with vertices $(\lambda, \lambda - 2)$, $(\lambda + 3, \lambda)$ and $(\lambda + 2, \lambda + 2)$ is independent of λ .



Watch Video Solution

15. Prove that the points $(a, b + c)$, $(b, c + a)$ and $(c, a + b)$ are collinear.



[Watch Video Solution](#)

16. Prove that the points (a, b) , (c, d) and $(a-c, b-d)$ are collinear, if $ad = bc$.



[Watch Video Solution](#)

17. If the points (x_1, y_1) , (x_2, y_2) , and (x_3, y_3) are collinear show that

$$\frac{y_2 - y_3}{x_2 x_3} + \frac{y_3 - y_1}{x_3 x_1} + \frac{y_1 - y_2}{x_1 x_2} = 0$$



[Watch Video Solution](#)

18. The coordinates of points A,B,C and D are $(-3, 5)$, $(4, -2)$, $(x, 3x)$ and $(6, 3)$ respectively and Area of $\frac{\Delta ABC}{\Delta BCD} = \frac{2}{3}$, find x.



Watch Video Solution

19. Find the area of the hexagon whose consecutive vertices are $(5, 0)$, $(4, 2)$, $(1, 3)$, $(-2, 2)$, $(-3, -1)$ and $(0, -4)$



Watch Video Solution

1. The equation of the locus of points equidistant from $(-1, -1)$ and $(4, 2)$ is

A. $3x - 5y - 7 = 0$

B. $5x + 3y - 9 = 0$

C. $4x + 3y + 2 = 0$

D. $x - 3y + 5 = 0$

Answer: B



Watch Video Solution

2. The equation of the locus of a point which moves so that its distance from the point $(ak, 0)$ is k times its distance from the point $\left(\frac{a}{k}, 0\right)$ ($k \neq 1$) is

A. $x^2 - y^2 = a^2$

B. $2x^2 - y^2 = 2a^2$

C. $xy = a^2$

D. $x^2 + y^2 = a^2$

Answer: D



Watch Video Solution

3. If the coordinates of a variable point P be $\left(t + \frac{1}{t}, t - \frac{1}{t}\right)$, where t is the variable quantity, then the locus of P is

A. $xy = 8$

B. $2x^2 - y^2 = 8$

C. $x^2 - y^2 = 4$

D. $2x^2 + 3y^2 = 5$

Answer: C



Watch Video Solution

4. If the coordinates of a variable point be $(\cos \theta + \sin \theta, \sin \theta - \cos \theta)$, where θ is the parameter, then the locus of P is

A. $x^2 - y^2 = 4$

B. $x^2 + y^2 = 2$

C. $xy = 3$

D. $x^2 + 2y^2 = 3$

Answer: B



Watch Video Solution

5. If a point moves such that twice its distance from the axis of x exceeds its distance from the axis of y by 2, then its locus is

A. $x - 2y = 2$

B. $x + 2y = 2$

C. $2y - x = 2$

D. $2y - 3x = 5$

Answer: C



Watch Video Solution

6. The equation $4xy - 3x^2 = a^2$ become when the axes are turned through an angle $\tan^{-1} 2$ is

A. $x^2 + 4y^2 = a^2$

B. $x^2 - 4y^2 = a^2$

C. $4x^2 + y^2 = a^2$

D. $4x^2 - y^2 = a^2$

Answer: B



Watch Video Solution

7. Transform the equation

$x^2 - 3xy + 11x - 12y + 36 = 0$ to parallel axes

through the point $(-4, 1)$ becomes $ax^2 + bxy + 1 = 0$

then $b^2 - a =$

A. $\frac{1}{4}$

B. $\frac{1}{16}$

C. $\frac{1}{64}$

D. $\frac{1}{256}$

Answer: C



Watch Video Solution

8. Find the locus of a point equidistant from the point $(2,4)$ and the y -axis.



Watch Video Solution

9. Find the equation of the locus of the points twice as from $(-a, 0)$ as from $(a, 0)$.



Watch Video Solution

10. OA and OB are two perpendicular straight lines. A straight line AB is drawn in such a manner that $OA + OB = 8$. Find the locus of the mid point of AB .

 [Watch Video Solution](#)

11. The ends of a rod of length l move on two mutually perpendicular lines. Find the locus of the point on the rod which divides it in the ratio $1 : 2$.

 [Watch Video Solution](#)

12. The coordinates of three points O, A, B are $(0, 0)$, $(0, 4)$ and $(6, 0)$ respectively. A point P moves so that the area of $\triangle POA$ is always twice the area of $\triangle POB$. Find the equation to both parts of the locus of P .

 [Watch Video Solution](#)

13. What does the equation

$(a - b)(x^2 + y^2) - 2abx = 0$ become if the origin is shifted to the point $\left(\frac{ab}{a - b}, 0\right)$ without rotation?

 [Watch Video Solution](#)

14. The equation $x^2 + 2xy + 4 = 0$ is transformed to the parallel axes through the point $(6, \lambda)$. For what value of λ its new form passes through the new origin ?

 [Watch Video Solution](#)

15. Show that if the axes be turned through $7\frac{1}{2}^\circ$, the equation $\sqrt{3}x^2 + (\sqrt{3} - 1)xy - y^2 = 0$ become free of xy in its new form.



[Watch Video Solution](#)

16. Find the angle through which the axes may be turned so that the equation $Ax + By + C = 0$ may reduce to the form $x = \text{constant}$, and determine the value of this constant.



[Watch Video Solution](#)

17.

Transform

$$12x^2 + 7xy - 12y^2 - 17x - 31y - 7 = 0 \quad \text{to}$$

rectangular axes through the point $(1, -1)$ inclined at

an angle $\tan^{-1}\left(\frac{4}{3}\right)$ to the original axes.

[Watch Video Solution](#)

Exercise Single Option Correct Type Questions

1. Vertices of a variable triangle are $(3, 4)$, $(5 \cos \theta, 5 \sin \theta)$ and $(5 \sin \theta, -5 \cos \theta)$, where $\theta \in R$. Locus of its orthocentre is

A. $x^2 + y^2 + 6x + 8y - 25 = 0$

$$\text{B. } x^2 + y^2 - 6x + 8y - 25 = 0$$

$$\text{C. } x^2 + y^2 + 6x - 8y - 25 = 0$$

$$\text{D. } x^2 + y^2 - 6x - 8y - 25 = 0$$

Answer: D



Watch Video Solution

2. If a rod AB of length 2 units slides on coordinate axes in the first quadrant. An equilateral triangle ABC is completed with C on the side away from O. Then, locus of C is

$$\text{A. } x^2 + y^2 - xy + 1 = 0$$

B. $x^2 + y^2 - xy\sqrt{3} + 1 = 0$

C. $x^2 + y^2 + xy\sqrt{3} - 1 = 0$

D. $x^2 + y^2 - xy\sqrt{3} - 1 = 0$

Answer: D



Watch Video Solution

3. The sides of a triangle are $3x + 4y$, $4x + 3y$ and $5x + 5y$ units, where $x > 0$, $y > 0$. The triangle is

A. right angled

B. acute angled

C. obtuse angled

D. isosceles

Answer: C



Watch Video Solution

4. Let P and Q be the points on the line joining $A(-2, 5)$ and $B(3, 1)$ such that $AP = PQ = QB$.

Then, the mid-point of PQ is

A. $\left(\frac{1}{2}, 3\right)$

B. $\left(-\frac{1}{4}, 4\right)$

C. $(2, 3)$

D. $(-1, 4)$

Answer: A



Watch Video Solution

5. A triangle ABC right angled at A has points A and B as $(2, 3)$ and $(0, -1)$ respectively. If $BC = 5$ units, then the point C is

A. $(4, 2)$

B. $(-4, 2)$

C. $(-4, 4)$

D. $(4, -4)$

Answer: A



Watch Video Solution

6. The locus of a point P which divides the line joining $(1, 0)$ and $(2 \cos \theta, 2 \sin \theta)$ internally in the ratio $2 : 3$ for all θ is

- A. a straight line
- B. a circle
- C. a pair of straight lines
- D. a parabola

Answer: B



Watch Video Solution

7. The points with coordinates $(2a, 3a)$, $(3b, 2b)$ and (c, c) are collinear

- A. for no value of a, b, c
- B. for all values of a, b, c
- C. if $a, \frac{c}{5}, b$ are in HP
- D. if $a, \frac{2c}{5}, b$ are in HP

Answer: D



Watch Video Solution

8. The vertices of a triangle are $(0, 3)$, $(-3, 0)$ and $(3, 0)$.

The coordinates of its orthocentre are

A. $(0, -2)$

B. $(0, 2)$

C. $(0, 3)$

D. $(0, -3)$

Answer: C



Watch Video Solution

9. ABC is an equilateral triangle such that the vertices B and C lie on two parallel at a distance 6. If A lies between the parallel lines at a distance 4 from one of them then the length of a side of the equilateral triangle.

A. 8

B. $\sqrt{\frac{88}{3}}$

C. $\frac{4\sqrt{7}}{\sqrt{3}}$

D. None of these

Answer: C



Watch Video Solution

10. A, B, C are respectively the points (1,2), (4, 2), (4, 5).

If T_1, T_2 are the points of trisection of the line segment BC, the area of the Triangle AT_1T_2 is

A. 1

B. $\frac{3}{2}$

C. 2

D. $\frac{5}{2}$

Answer: B



Watch Video Solution

11. (i) The points $(-1, 0)$, $(4, -2)$ and $(\cos 2\theta, \sin 2\theta)$ are collinear

(ii) The points $(-1, 0)$, $(4, -2)$ and $\left(\frac{1 - \tan^2 \theta}{1 + \tan^2 \theta}, \frac{2 \tan \theta}{1 + \tan^2 \theta}\right)$ are collinear

A. both statements are equivalent

B. statement (i) has more solutions than statement (ii) for θ

C. statement (ii) has more solutions than statement (i) for θ

D. None of the above

Answer: B



12. If $\alpha_1, \alpha_2, \alpha_3, \beta_1, \beta_2, \beta_3$ are the values of n for which $\sum_{r=0}^{n-1} x^{2r}$ is divisible by $\sum_{r=0}^{n-1} x^r$, then the triangle having vertices $(\alpha_1, \beta_1), (\alpha_2, \beta_2)$ and (α_3, β_3) cannot be

- A. an isosceles triangle
- B. a right angled isosceles triangle
- C. a right angled triangle
- D. an equilateral triangle

Answer: D



13. A triangle ABC with vertices $A(-1, 0)$, $B\left(-2, \frac{3}{4}\right)$, and $C\left(-3, -\frac{7}{6}\right)$ has its orthocentre at H . Then, the orthocentre of triangle BCH will be

A. $(-30.6, 15.4)$

B. $(1, 3)$

C. $(-1, 2)$

D. None of these

Answer: D



Watch Video Solution

14.

If

$$\sum_{i=1}^4 (x_i^2 + y_i^2) \leq 2x_1x_3 + 2x_2x_4 + 2y_2y_3 + 2y_1y_4,$$

the points (x_1, y_1) , (x_2, y_2) , (x_3, y_3) , (x_4, y_4) are

- A. the vertices of a rectangle
- B. collinear
- C. the vertices of a trapezium
- D. None of these

Answer: A



Watch Video Solution

15. Without change of axes the origin is shifted to (h, k), then from the equation $x^2 + y^2 - 4x + 6y - 7 = 0$, the term containing linear powers are missing, then point (h, k) is

- A. (a) (3, 2)
- B. (b) (-3, 2)
- C. (c) (2, -3)
- D. (d) (-2, -3)

Answer: C



Watch Video Solution

Exercise More Than One Correct Option Type Questions

1. If $(-6, -4)$, $(3, 5)$, $(-2, 1)$ are the vertices of a parallelogram, then the remaining vertex can be $(0, -1)$ (b) $(7, 9)$ (c) $(-1, 0)$ (d) $(-11, -8)$

A. $(0, -1)$

B. $(-1, 0)$

C. $(-11, -8)$

D. $(7, 10)$

Answer: B::C::D



Watch Video Solution

2. If the point (x, y) is equidistant from the points $(a + b, b - a)$ and $(a - b, a + b)$, prove that $bx = ay$.

A. $ax = by$

B. $bx = ay$

C. $x^2 - y^2 = 2(ax + by)$

D. P can be (a, b)

Answer: B::D



Watch Video Solution

3. Evaluate $\int 7^x dx$



Watch Video Solution

4. Show that the following points are the vertices of a rectangle.

(i) A(-4, -1), B(-2, -4), C(4, 0) and D(2, 3)

(ii) A(2, -2), B(14, 10), C(11, 13) and D(-1, 1)

(iii) A(0, -4), B(6, 2), C(3, 5) and D(-3, -1)

A. parallelogram

B. rectangle

C. rhombus

D. square

Answer: A::B

 [Watch Video Solution](#)

5. The medians AD and BE of the triangle ABC with vertices A(0, b), B(0, 0) and C(a, 0) are mutually perpendicular if

A. (a) $b = a\sqrt{2}$

B. (b) $a = b\sqrt{2}$

C. (c) $b = -a\sqrt{2}$

D. (d) $a = -b\sqrt{2}$

Answer: B::D



Watch Video Solution

6. The points $A(x, y)$, $B(y, z)$ and $C(z, x)$ represents the vertices of a right angled triangle, if

A. (a) $x = y$

B. (b) $y = z$

C. (c) $z = x$

D. (d) $x = y = z$

Answer: A::B::C



Watch Video Solution

7. Let the base of a triangle lie along the line $x = a$ and be of length a . The area of this triangles is a^2 , if the vertex lies on the line

A. $x = -a$

B. $x = 0$

C. $x = \frac{a}{2}$

D. $x = 2a$

Answer: B::D



Exercise Passage Based Questions

1. ABC is a triangle right angled at A , $AB = 2AC$, $A = (1, 2)$, $B(-3, 1)$. The vertices of the triangles are in anticlockwise sense. $BCEF$ is a square with vertices in clockwise sense. Area of triangle ACF is:

A. $\frac{51}{8}$

B. $\frac{51}{4}$

C. $\frac{31}{5}$

D. $\frac{21}{4}$

Answer: B



Watch Video Solution

2. Evaluate $\int 4^x dx$



Watch Video Solution

3. Let $O(0, 0)$ and $B\left(1, \frac{1}{\sqrt{3}}\right)$ be the vertices of a triangle. Let R be the region consisting of all those points P inside ΔOAB satisfying $d(P, OA) \leq \min\{d(P, OB), d(P, AB)\}$, where d denotes the distance from the point P to the corresponding line. Let M be peak of region R .

The perimeter of region R is equal to

A. $\sqrt{3}$

B. $\frac{1}{\sqrt{3}}$

C. 3

D. $2 - \sqrt{3}$

Answer: D



Watch Video Solution

4. Let $O(0, 0)$, $A(2, 0)$, and $B\left(1, \frac{1}{\sqrt{3}}\right)$ be the vertices of a triangle. Let R be the region consisting of all those points P inside OAB which satisfy $d(P, OA) \leq \min [d(p, OB), d(P, AB)]$, where d

denotes the distance from the point to the corresponding line. Sketch the region R and find its area.

A. $2 - \sqrt{3}$

B. $2 + \sqrt{3}$

C. $4 + 3\sqrt{3}$

D. $2 + 4\sqrt{(2 - \sqrt{3})}$

Answer: D



Watch Video Solution

5. Differentiate 4^x with respect to x .



Watch Video Solution

Exercise Single Integer Answer Type Questions

1. If the area of the triangle formed by the points $(2a, b)$, $(a + b, 2b + a)$ and $(2b, 2a)$ be Δ_1 and the area of the triangle whose vertices are $(a + b, a - b)$, $(3b - a, b + 3a)$ and $(3a - b, 3b - a)$ be Δ_2 , then the value of Δ_2 / Δ_1 is



Watch Video Solution

2. The diameter of the nine point circle of the triangle with vertices $(3, 4)$, $(5 \cos \theta, 5 \sin \theta)$ and $(5 \sin \theta, -5 \cos \theta)$, where $\theta \in R$, is

 [Watch Video Solution](#)

3. Differentiate 7^x with respect to x .

 [Watch Video Solution](#)

4. If (x, y) is the incentre of the triangle formed by the points $(3, 4)$, $(4, 3)$ and $(1, 2)$, then the value of x^2 is

 [Watch Video Solution](#)

5. Let P and Q be points on the line joining A(-2, 5) and B(3, 1) such that $AP = PQ = QB$. If mid-point of PQ is (a, b), then the value of $\frac{b}{a}$ is



Watch Video Solution

Exercise 5

1. Differentiate 3^x with respect to x .



Watch Video Solution

2. Differentiate 6^x with respect to x .



Watch Video Solution

Exercise Statement I And Ii Type Questions

1. The vertices of a triangle are $A(1, 2)$, $B(-1, 3)$ and $C(3, 4)$. Let D, E, F divide BC, CA, AB respectively in the same ratio.

Statement I : The centroid of triangle DEF is $(1, 3)$.

Statement II : The triangle ABC and DEF have the same centroid.

- A. Statement I is true, Statement II is true,
Statement II is a correct explanation for
Statement I.
- B. Statement I is true, Statement II is true,
Statement II is not a correct explanation for
Statement I.
- C. Statement I is true, Statement II is false.
- D. Statement I is false, Statement II is true.

Answer: A



Watch Video Solution

2. Statement 1 : Let the vertices of a ABC be $A(-5, -2)$, $B(7, 6)$, and $C(5, -4)$. Then the coordinates of the circumcenter are $(1, 2)$

Statement 2 : In a right-angled triangle, the midpoint of the hypotenuse is the circumcenter of the triangle.



[Watch Video Solution](#)

3. Evaluate $\int 6^x dx$



[Watch Video Solution](#)

4. Transform the equation

$x^2 - 3xy + 11x - 12y + 36 = 0$ to parallel axes

through the point $(-4, 1)$ becomes $ax^2 + bxy + 1 = 0$

then $b^2 - a =$



Watch Video Solution

Exercise 7

1. If $A(x_1, y_1)$, $B(x_2, y_2)$ and $C(x_3, y_3)$ are the vertices of a $\triangle ABC$ and (x, y) be a point on the internal bisector of angle A, then prove that

$$b \begin{vmatrix} x & y & 1 \\ x_1 & y_1 & 1 \\ x_2 & y_2 & 1 \end{vmatrix} + c \begin{vmatrix} x & y & 1 \\ x_1 & y_1 & 1 \\ x_3 & y_3 & 1 \end{vmatrix} = 0$$

where, AC = b and AB = c.



[Watch Video Solution](#)

2. Differentiate 12^x with respect to x .



[Watch Video Solution](#)

3. Differentiate \tan^{-1} with respect to x



[Watch Video Solution](#)

4. Differentiate $\sin^{-1} x$ with respect to x



Watch Video Solution

Exercise Subjective Type Questions

1. Differentiate 2^x with respect to x .



Watch Video Solution

2. A line L intersects three sides BC , CA and AB of a triangle in P, Q, R respectively, show that

$$\frac{BP}{PC} \cdot \frac{CQ}{QA} \cdot \frac{AR}{RB} = -1$$



 [Watch Video Solution](#)

3. Differentiate 10^x with respect to x .

 [Watch Video Solution](#)

4. Differentiate 23^x with respect to x .

 [Watch Video Solution](#)

5. Differentiate $\cos^{-1} x$ with respect to x

 [Watch Video Solution](#)

Exercise Questions Asked In Previous 13 Years Exam

1. If a vertex of a triangle is $(1, 1)$ and the mid-points of two side through this vertex are $(-1, 2)$ and $(3, 2)$, then centroid of the triangle is

A. $\left(\frac{1}{3}, \frac{7}{3}\right)$

B. $\left(1, \frac{7}{3}\right)$

C. $\left(-\frac{1}{3}, \frac{7}{3}\right)$

D. $\left(-1, \frac{7}{3}\right)$

Answer: B



Watch Video Solution

2. Differentiate $\cot^{-1} x$ with respect to x .



Watch Video Solution

3. Let $A(h, k)$, $B(1, 1)$ and $C(2, 1)$ be the vertices of a right angled triangle with AC as its hypotenuse. If the area of the triangle is 1, then the set of values which k can take is given by (1) $\{1, 3\}$ (2) $\{0, 2\}$ (3) $\{-1, 3\}$
(4) $\{-3, -2\}$

A. $\{1, 3\}$

B. $\{0, 2\}$

C. $\{-1, 3\}$

D. $\{-3, -2\}$

Answer: C

 [Watch Video Solution](#)

4. Evaluate $\int 2^x dx$

 [Watch Video Solution](#)

5. The x-coordinate of the incentre of the triangle that has the coordinates of mid points of its sides as $(0, 1)$, $(1, 1)$ and $(1, 0)$ is

A. $2 + \sqrt{2}$

B. $2 - \sqrt{2}$

C. $1 + \sqrt{2}$

D. $1 - \sqrt{2}$

Answer: B



Watch Video Solution

6. Evaluate $\int 5^x dx$



Watch Video Solution

7. Let k be an integer such that the triangle with vertices $(k, -3k)$, $(5, k)$ and $(-k, 2)$ has area $28sq.$ units. Then the orthocentre of this triangle is at

the point : (1) $\left(1, -\frac{3}{4}\right)$ (2) $\left(2, \frac{1}{2}\right)$ (3) $\left(2, -\frac{1}{2}\right)$

(4) $\left(1, \frac{3}{4}\right)$

A. $\left(2, \frac{1}{2}\right)$

B. $\left(2, -\frac{1}{2}\right)$

C. $\left(1, \frac{3}{4}\right)$

D. $\left(1, -\frac{3}{4}\right)$

Answer: A



Watch Video Solution

