



## MATHS

### BOOKS - ARIHANT MATHS

#### VECTOR ALGEBRA

#### Example

1. Classify the following measures as scalars and vectors

(i) 20 m north-west

(ii) 10 newton

(iii) 30 km/h

(iv) 50m/s towards north

(v)  $10^{-19}$  coulomb



Watch Video Solution

## 2. Represent graphically

- (i) a displacement of 60 km,  $40^\circ$  east of north
- (ii) A displacement of 50 km south-east.




**Watch Video Solution**

## 3. In the following figure, which of the vectors are:

- (i) Collinear
- (ii) Equal
- (iii) Co-initial

(iv) collinear but not equal .



[Watch Video Solution](#)

4. Find a unit vector parallel to the vector  $-3\hat{i} + 4\hat{j}$ .



[Watch Video Solution](#)

5. Let  $a = 12\hat{i} + n\hat{j}$  and  $|a| = 13$ , find the value of  $n$ .



Watch Video Solution

6. Write two different vectors having same magnitude.



Watch Video Solution

7. If one side of a square be represented by the vectors  $3\hat{i} + 4\hat{j} + 5\hat{k}$ , then the area of the square is

A. 12

B. 13

C. 25

D. 50

Answer: D



Watch Video Solution

8. The direction cosines of the vector  $3\hat{i} - 4\hat{j} + 5\hat{k}$  are

A.  $\frac{3}{5}, \frac{-4}{5}, \frac{1}{5}$

B.  $\frac{3}{5\sqrt{2}}, \frac{-4}{5\sqrt{2}}, \frac{1}{\sqrt{2}}$

C.  $\frac{3}{\sqrt{2}}, \frac{-4}{\sqrt{2}}, \frac{1}{\sqrt{2}}$

D.  $\frac{3}{5\sqrt{2}}, \frac{4}{5\sqrt{2}}, \frac{1}{\sqrt{2}}$

Answer: B



Watch Video Solution

9. Show that the vector  $\vec{i} + \vec{j} + \vec{k}$  is equally inclined to the axes OX, OY and OZ.



Watch Video Solution

10. Let AB be a vector in two dimensional plane with the magnitude 4 units and making an angle of  $30^\circ$  with X-axis and lying in the first quadrant. Find the components of AB along the two axes of coordinates. Hence, represent AB in terms of unit vectors  $\hat{i}$  and  $\hat{j}$ .

 [Watch Video Solution](#)

11. Find the unit vector parallel to the resultant vector of  $2\hat{i} + 4\hat{j} - 5\hat{k}$  and  $\hat{i} + 2\hat{j} + 3\hat{k}$ .

 [Watch Video Solution](#)

12. If  $\vec{a}$ ,  $\vec{b}$ ,  $\vec{c}$  be the vectors represented by the sides of a triangle, taken in order, then prove that  $\vec{a} + \vec{b} + \vec{c} = \vec{0}$ .

 [Watch Video Solution](#)

13. If  $S$  is the mid-point of side  $QR$  of a  $\Delta PQR$ , then prove that  $PQ + PR = 2PS$ .

 [Watch Video Solution](#)

14. If  $ABCDEF$  is a regular hexagon and  $AB+AC+AD+AE+AF=\lambda AD$ , then  $\lambda$  is equal to

 [Watch Video Solution](#)

15. If  $A = (0, 1), B = (1, 0), C = (1, 2), D = (2, 1)$ , prove that  $\vec{AB} = \vec{CD}$ .

 [Watch Video Solution](#)

16. If the position vectors of  $A$  and  $B$  respectively are  $\hat{i} + 3\hat{j} - 7\hat{k}$  and  $5\hat{i} - 2\hat{j} + 4\hat{k}$ , then find  $AB$



Watch Video Solution

17. Vectors drawn from the origin  $O$  to the points  $A$ ,  $B$  and  $C$  are respectively  $\vec{a}$ ,  $\vec{b}$  and  $4\vec{a} - 3\vec{b}$ . Find  $\vec{AC}$  and  $\vec{BC}$ .



Watch Video Solution

18. Find the direction cosines of the vector joining the points  $A(1, 2, 3)$  and  $B(1, 2, 1)$ , directed from  $A$  to  $B$ .



Watch Video Solution

19. Let  $\alpha, \beta, \gamma$  be distinct real numbers. The points with position vectors  $\alpha\hat{i} + \beta\hat{j} + \gamma\hat{k}$ ,  $\beta\hat{i} + \gamma\hat{j} + \alpha\hat{k}$ ,  $\gamma\hat{i} + \alpha\hat{j} + \beta\hat{k}$

A. are collinear

B. form an equilateral triangle

C. form a scalene triangle



D. form a right angled triangle

**Answer:**



**Watch Video Solution**

20. If the position vectors of the vertices of a triangle be  $2\hat{i} + 4\hat{j} - \hat{k}$ ,  $4\hat{i} + 5\hat{j} + \hat{k}$  and  $3\hat{i} + 6\hat{j} - 3\hat{k}$ , then the triangle is

a. right angled

b. isosceles

c. equilateral

d. none of these

A. right angled

B. isosceles

C. equilateral

D. none of these

**Answer: A::B**



Watch Video Solution

21. The sides of a parallelogram are  $2\hat{i} + 4\hat{j} - 5\hat{k}$  and  $\hat{i} + 2\hat{j} + 3\hat{k}$ . The unit vector parallel to one of the diagonals is



Watch Video Solution

22. If  $\vec{a}$ ,  $\vec{b}$  are any two vectors, then give the geometrical interpretation of relation  $|\vec{a} + \vec{b}| = |\vec{a} - \vec{b}|$



Watch Video Solution

23. Can the magnitude of the resultant vector of the two given vectors is less than the magnitude of any of the given vectors?



Watch Video Solution

24. If  $\vec{a}$  is a non-zero vector of modulus  $a$  and  $m$  is a non-zero scalar, then  $m\vec{a}$  is a unit vector if

A.  $m = \pm 1$

B.  $m = |a|$

C.  $m = \frac{1}{|a|}$

D.  $m = \pm 2$

**Answer: C**



**Watch Video Solution**

25. For a non-zero vector  $a$ , the set of real number, satisfying

$|(5 - x)a| < |2a|$  consists of all  $x$  such that

A.  $0 < x < 3$

B.  $3 < x < 7$

C.  $-7 < x < -3$

D.  $-7 < x < 3$

**Answer: B**



[Watch Video Solution](#)

26. Find a vector of magnitude  $(5/2)$  units which is parallel to the vector  $3\hat{i} + 4\hat{j}$ .



[Watch Video Solution](#)

27. Find the power set of set  $A = \{1,2,3\}$



[Watch Video Solution](#)

28. Find the number of Element of power set of set A have  $2^n$  number of element .



[Watch Video Solution](#)

29. The position vectors of the vertices A, B and C of a triangle are  $\hat{i} - \hat{j} - 3\hat{k}$ ,  $2\hat{i} + \hat{j} - 2\hat{k}$  and  $-5\hat{i} + 2\hat{j} - 6\hat{k}$ , respectively. The length of the bisector AD of the  $\angle BAC$ , where D is on the segment BC, is

A.  $\frac{3}{4}\sqrt{3}$

B.  $\frac{1}{4}$

C.  $\frac{11}{2}$

D. None of these

**Answer: A**



**Watch Video Solution**

30. Which of the following is prime number

A. 144

B. 137

C. 125

D. 15

**Answer: B**



[Watch Video Solution](#)

**31.** The sum of the magnitudes of two forces acting at a point is 16 N. The resultant of these forces is perpendicular to the smaller force has a magnitude of 8 N. If the smaller force is magnitude  $x$ , then the value of  $x$  is

A. 13,5

B. 12,6

C. 10,6

D. 11,7

**Answer: A**



[Watch Video Solution](#)

32. The length of longer diagonal of the parallelogram constructed on  $5a + 2b$  and  $a - 3b$ . If it is given that  $|a| = 2\sqrt{2}$ ,  $|b| = 3$  and angle between  $a$  and  $b$  is  $\frac{\pi}{4}$  is

A. 15

B.  $\sqrt{113}$

C.  $\sqrt{593}$

D.  $\sqrt{369}$

Answer: C



Watch Video Solution

33. The vector  $\vec{c}$ , directed along the internal bisector of the angle between the vectors

$\vec{a} = 7\hat{i} - 4\hat{j} - 4\hat{k}$  and  $\vec{b} = -2\hat{i} - \hat{j} + 2\hat{k}$  with  $|\vec{c}| = 5\sqrt{6}$ , is

A. (a)  $\frac{5}{3}(\hat{i} - 7\hat{j} + 2\hat{k})$

B. (b)  $\frac{5}{3}(5\hat{i} + 5\hat{j} + 2\hat{k})$

C. (c)  $\frac{5}{3}(\hat{i} + 7\hat{j} + 2\hat{k})$

D. (d)  $\frac{5}{3}(-5\hat{i} + 5\hat{j} + 2\hat{k})$

**Answer: A**

 [Watch Video Solution](#)

**34.** Show that the vectors  $2\hat{i} - 3\hat{j} + 4\hat{k}$  and  $-4\hat{i} + 6\hat{j} - 8\hat{k}$  are collinear.

 [Watch Video Solution](#)

**35.** Prove that the points  $A(1, 2, 3)$ ,  $B(3, 4, 7)$ ,  $C(-3, -2, -5)$  are collinear and find the ratio in which B divides AC.

 [Watch Video Solution](#)



36. If the position vectors of A,B,C and D are

$2\hat{i} + \hat{j}$ ,  $\hat{i} - 3\hat{j}$ ,  $3\hat{i} + 2\hat{j}$  and  $\hat{i} + \lambda\hat{j}$  respectively and  $|AB||CD$ . Then  $\lambda$  will be

A.  $-8$

B.  $-6$

C.  $8$

D.  $6$

**Answer: B**



[Watch Video Solution](#)

37. The points with position vectors  $60\hat{i} + 3\hat{j}$ ,  $40\hat{i} - 8\hat{j}$ ,  $a\hat{i} - 52\hat{j}$  are collinear if a is :

a.  $-40$

b.  $40$

c.  $20$

d. none of these

A.  $-40$

B.  $40$

C.  $20$

D. none of these

**Answer: A**



[Watch Video Solution](#)

**38.** If  $a, b$  and  $c$  are three non-zero vectors such that no two of these are collinear. If the vector  $a+2b$  is collinear with  $c$  and  $b+3c$  is collinear with  $a$  ( $\lambda$  being some non-zero scalar), then  $a+2b+6c$  is equal to

A.  $0$

B.  $\lambda b$

C.  $\lambda c$

D.  $\lambda a$

**Answer: A**



**Watch Video Solution**

**39.** Check whether the given three vectors are coplanar or non-coplanar :

$$-2\hat{i} - 2\hat{j} + 4\hat{k}, \quad -2\hat{i} + 4\hat{j} - 2\hat{k}, \quad 4\hat{i} - 2\hat{j} - 2\hat{k}.$$



**Watch Video Solution**

**40.** If the vectors  $4\hat{i} + 11\hat{j} + m\hat{k}$ ,  $7\hat{i} + 2\hat{j} + 6\hat{k}$  and  $\hat{i} + 5\hat{j} + 4\hat{k}$  are coplanar, then  $m$  is equal to

a. 38

b. 0

c. 10

d. -10

A. 38

B. 0

C. 10

D.  $-10$

**Answer: C**



[Watch Video Solution](#)

41. If  $a, b$  and  $c$  are non-coplanar vectors, prove that  $3a-7b-4c$ ,  $3a-2b+c$  and  $a+b+2c$  are coplanar.



[Watch Video Solution](#)

42. The value of  $\lambda$  for which the four points  $2\hat{i} + 3\hat{j} - \hat{k}$ ,  $\hat{i} + 2\hat{j} + 3\hat{k}$ ,  $3\hat{i} + 4\hat{j} - 2\hat{k}$  and  $\hat{i} - \lambda\hat{j} + 6\hat{k}$  are coplanar.

a. 8

b. 0

c.  $-2$

d. 6

A. 8

B. 0

C.  $-2$

D. 6

**Answer: C**



[Watch Video Solution](#)

**43.** If  $A = \{0, 1, 2, 3, 5, 6\}$ ,  $B = \{1, 3, 5, 7, 9\}$  and  $C = \{0, 5, 10, 20, 40\}$ , find

A. 1)  $A \cup B$

B. 2)  $A \cup C$

C. 3)  $B \cup C$

D. 4)  $A \cap B$

**Answer:**



[Watch Video Solution](#)

44. Show that the vectors

$\hat{i} - 3\hat{j} + 2\hat{k}$ ,  $2\hat{i} - 4\hat{j} - \hat{k}$  and  $3\hat{i} + 2\hat{j} - \hat{k}$  and linearly independent.



Watch Video Solution

45. If  $\vec{a} = \hat{i} + \hat{j} + \hat{k}$ ,  $\vec{b} = 4\hat{i} + 3\hat{j} + 4\hat{k}$  and  $\vec{c} = \hat{i} + \alpha\hat{j} + \beta\hat{k}$

are linearly dependent vectors and  $|\vec{c}| = \sqrt{3}$  then:

A. (a)  $\alpha = 1, \beta = -1$

B. (b)  $\alpha = 1, \beta = \pm 1$

C. (c)  $\alpha = \pm 1, \beta = \pm 1$

D. (d)  $\alpha = \pm 1, \beta = 1$

Answer: D



Watch Video Solution

46. If  $|a| + |b| = |c|$  and  $a + b = c$ , then find angle between a and b.

A.  $\frac{\pi}{4}$

B.  $\frac{\pi}{2}$

C.  $\pi$

D. 0

Answer: C



Watch Video Solution

47. A unit vector  $\hat{a}$  makes an angle  $\frac{\pi}{4}$  with z-axis, if  $\hat{a} + \hat{i} + \hat{j}$  is a unit vector then  $\hat{a}$  is equal to

(A)  $\hat{i} + \hat{j} + \frac{\hat{k}}{2}$  (B)  $\frac{\hat{i}}{2} + \frac{\hat{j}}{2} - \frac{\hat{k}}{\sqrt{2}}$  (C)  $-\frac{\hat{i}}{2} - \frac{\hat{j}}{2} + \frac{\hat{k}}{\sqrt{2}}$  (D)

$\frac{\hat{i}}{2} - \frac{\hat{j}}{2} - \frac{\hat{k}}{\sqrt{2}}$

A. A.  $\frac{\hat{i}}{2} + \frac{\hat{j}}{2} + \frac{\hat{k}}{\sqrt{2}}$

B. B.  $\frac{\hat{i}}{2} + \frac{\hat{j}}{2} - \frac{\hat{k}}{\sqrt{2}}$

C. C.  $-\frac{\hat{i}}{2} - \frac{\hat{j}}{2} + \frac{\hat{k}}{\sqrt{2}}$

D. D. none of these

**Answer: C**



**Watch Video Solution**

**48.** If the resultant of two forces of magnitudes P and Q acting at a point at an angle of  $60^\circ$  is  $\sqrt{7}Q$ , then P/Q is

A. 1

B.  $\frac{3}{2}$

C. 2

D. 4

**Answer: C**



**Watch Video Solution**



49. The vector  $\vec{a}$  has the components  $2p$  and  $1$  w.r.t. a rectangular Cartesian system. This system is rotated through a certain angle about the origin in the counterclockwise sense. If, with respect to a new system,  $\vec{a}$  has components  $(p + 1)$  and  $1$ , then  $p$  is equal to

A.  $p=0$

B.  $p=1$  or  $p = -\frac{1}{3}$

C.  $p=-1$  or  $p = \frac{1}{3}$

D.  $p=1$  or  $p = -1$

**Answer: B**

 [Watch Video Solution](#)

50. ABC is an isosceles triangle right angled at A. forces of magnitude  $2\sqrt{2}$ ,  $5$  and  $6$  act along BC, CA and AB respectively. The magnitude of their resultant force is

A.  $4$

B. 5

C.  $11 + 2\sqrt{2}$

D. 30

**Answer: B**



[Watch Video Solution](#)

51. A line segment has length 63 and direction ratios are 3, -2, 6. The components of the line vector are

A. -27, 18, 54

B. 27, -18, 54

C. 27, -18, -54

D. -27, -18, -54

**Answer: B**



[Watch Video Solution](#)

52. If the vectors  $6\hat{i} - 2\hat{j} + 3\hat{k}$ ,  $2\hat{i} + 3\hat{j} - 6\hat{k}$  and  $3\hat{i} + 6\hat{j} - 2\hat{k}$  form a triangle, then it is

- A. right angled
- B. obtuse angled
- C. equilateral
- D. isosceles

**Answer: B**



[Watch Video Solution](#)

53. The position vectors of the points A, B, C are  $2\hat{i} + \hat{j} - \hat{k}$ ,  $3\hat{i} - 2\hat{j} + \hat{k}$  and  $\hat{i} + 4\hat{j} - 3\hat{k}$  respectively. These points

- A. form an isosceles triangle
- B. form a right angled triangle

C. are collinear

D. form a scalene triangle

**Answer: C**



[Watch Video Solution](#)

54. The position vector of a point C with respect to B is  $\hat{i} + \hat{j}$  and that of B with respect to A is  $\hat{i} - \hat{j}$ . The position vector of C with respect to A is

A.  $2\hat{i}$

B.  $2\hat{j}$

C.  $-2\hat{j}$

D.  $-2\hat{i}$

**Answer: A**



[Watch Video Solution](#)

55. Find the number of element of power set of set A have 4 element



Watch Video Solution

56. If  $\vec{a}$ ,  $\vec{b}$ ,  $\vec{c}$ ,  $\vec{d}$  are the position vector of point  $A, B, C$  and  $D$ , respectively referred to the same origin  $O$  such that no three of these point are collinear and  $\vec{a} + \vec{c} = \vec{b} + \vec{d}$ , then prove that quadrilateral  $ABCD$  is a parallelogram.

A. square

B. rhombus

C. rectangle

D. parallelogram

Answer: D



Watch Video Solution

57. P is a point on the side BC of  $\triangle ABC$  and Q is a point such that PQ is the resultant of AP, PB and PC. Then, ABQC is a

- A. square
- B. rectangle
- C. parallelogram
- D. trapezium

**Answer: C**



[Watch Video Solution](#)

58. Find the number of element of power set of set B have 3 element



[Watch Video Solution](#)

59.  $ABCD$  is a parallelogram whose diagonals meet at P. If O is a fixed point, then  $\vec{OA} + \vec{OB} + \vec{OC} + \vec{OD}$  equals :

A. (a)  $\overrightarrow{OP}$

B. (b)  $2\overrightarrow{OP}$

C. (c)  $3\overrightarrow{OP}$

D. (d)  $4\overrightarrow{OP}$

**Answer: D**

 [Watch Video Solution](#)

**60.** If C is the middle point of AB and P is any point outside AB, then

A.  $PA+PB=PC$

B.  $PA+PB=2PC$

C.  $PA+PB+PC=0$

D.  $PA+PB+2PC=0$

**Answer: B**

 [Watch Video Solution](#)

61. Which of the following is not prime number

A. 17

B. 19

C. 27

D. 29

**Answer: B**



[Watch Video Solution](#)

62. Five points given by A,B,C,D and E are in a plane. Three forces AC,AD and AE act at A and three forces CB,DB and EB act B. then, their resultant is

A. 2AC

B. 3AB



C. 3DB

D. 2BC

**Answer: B**



**Watch Video Solution**

63. In a regular hexagon  $ABCDEF$ ,  $\vec{AB} = a$ ,  $\vec{BC} = \vec{b}$  and  $\vec{CD} = c$  Then  $\vec{AE} =$



**Watch Video Solution**

64. If  $\vec{a} + \vec{b} + \vec{c} = \vec{0}$ ,  $|\vec{a}| = 3$ ,  $|\vec{b}| = 5$ ,  $|\vec{c}| = 7$ , then angle between  $\vec{a}$  and  $\vec{b}$  is : a.  $\frac{\pi}{2}$  b.  $\frac{\pi}{3}$  c.  $\frac{\pi}{4}$  d.  $\frac{\pi}{6}$

A.  $\frac{\pi}{2}$

B.  $\frac{\pi}{3}$

C.  $\frac{\pi}{4}$

D.  $\frac{\pi}{6}$

**Answer: B**



**Watch Video Solution**

65. If  $\vec{a}$  &  $\vec{b}$  are the position vectors of  $A$  &  $B$  respectively and  $C$  is a point on  $AB$  produced such that  $AC = 3AB$  then the position vector of  $C$  is:

A. (a)  $3\vec{a} - \vec{b}$

B. (b)  $3\vec{b} - \vec{a}$

C. (c)  $3\vec{a} - 2\vec{b}$

D. (d)  $3\vec{b} - 2\vec{a}$

**Answer: D**



**Watch Video Solution**

66. Let  $A$  and  $B$  be points with position vectors  $\vec{a}$  and  $\vec{b}$  with respect to origin  $O$ . If the point  $C$  on  $OA$  is such that  $2\vec{AC} = \vec{CO}$ ,  $\vec{CD}$  is parallel to  $\vec{OB}$  and  $|\vec{CD}| = 3|\vec{OB}|$  then  $\vec{AD}$  is (A)  $\vec{b} - \frac{\vec{a}}{9}$  (B)  $3\vec{b} - \frac{\vec{a}}{3}$  (C)  $\vec{b} - \frac{\vec{a}}{3}$  (D)  $\vec{b} + \frac{\vec{a}}{3}$

A.  $3b - \frac{a}{2}$

B.  $3b + \frac{a}{2}$

C.  $3b - \frac{a}{3}$

D.  $3b + \frac{a}{3}$

**Answer: C**

 [Watch Video Solution](#)

67. If the position vector of a point A is  $\vec{a} + 2\vec{b}$  and  $\vec{a}$  divides AB in the ratio 2 : 3, then the position vector of B, is

A.  $2a - b$

B.  $b - 2a$

C.  $a - 3b$

D.  $b$

**Answer:**

 [Watch Video Solution](#)

68. If D, E and F are respectively, the mid-points of AB, AC and BC in  $\triangle ABC$ , then  $BE + AF$  is equal to

A. DC

B.  $\frac{1}{2}BF$

C.  $2BF$

D.  $\frac{3}{2}BF$

**Answer: A**

 [Watch Video Solution](#)

69. In a quadrilateral  $PQRS$ ,  $\vec{PQ} = \vec{a}$ ,  $\vec{QR} = \vec{b}$ ,  $\vec{SP} = \vec{a} - \vec{b}$ ,  $M$  is the midpoint of  $\vec{QR}$  and  $X$  is a point on  $SM$  such that  $SX = \frac{4}{5}SM$ .

Prove that  $P$ ,  $X$  and  $R$  are collinear.

A.  $PX = \frac{1}{5}PR$

B.  $PX = \frac{3}{5}PR$

C.  $PX = \frac{2}{5}PR$

D. none of these

Answer: B



Watch Video Solution

70. Orthocenter of an equilateral triangle  $ABC$  is the origin  $O$ . If

$\vec{OA} = \vec{a}$ ,  $\vec{OB} = \vec{b}$ ,  $\vec{OC} = \vec{c}$ , then  $\vec{AB} + 2\vec{BC} + 3\vec{CA} =$

A.  $3\vec{c}$

B.  $3a$

C.  $0$

D.  $3b$

**Answer: B**



**Watch Video Solution**

71. If  $\vec{a}$ ,  $\vec{b}$  and  $\vec{c}$  are position vectors of A, B, and C respectively of  $\triangle ABC$  and if  $|\vec{a} - \vec{b}|$ ,  $|\vec{b} - \vec{c}| = 2$ ,  $|\vec{c} - \vec{a}| = 3$ , then the distance between the centroid and incenter of  $\triangle ABC$  is

A.  $1$

B.  $\frac{1}{2}$

C.  $\frac{1}{3}$

D.  $\frac{2}{3}$

**Answer: C**

[Watch Video Solution](#)

72. Let position vectors of point A, B and C of triangle ABC represents be  $\hat{i} + \hat{j} + 2\hat{k}$ ,  $\hat{i} + 2\hat{j} + \hat{k}$  and  $2\hat{i} + \hat{j} + \hat{k}$ . Let  $l_1$ ,  $l_2$  and  $l_3$  be the length of perpendicular drawn from the orthocenter 'O' on the sides AB, BC and CA, then  $(l_1 + l_2 + l_3)$  equals

A.  $\frac{2}{\sqrt{6}}$

B.  $\frac{3}{\sqrt{6}}$

C.  $\frac{\sqrt{6}}{2}$

D.  $\frac{\sqrt{6}}{3}$ .

**Answer: C**

[Watch Video Solution](#)

73. ABCDEF is a regular hexagon in the x-y plane with vertices in the anticlockwise direction. If  $\vec{AB} = 2\hat{i}$ , then  $\vec{CD}$  is

A.  $\hat{i} + 3\hat{j}$

B.  $\hat{i}9 + 2\hat{j}$

C.  $-\hat{i} + \sqrt{3}\hat{j}$

D. none of these

**Answer:**



**Watch Video Solution**

74. The vertices of a triangle are A(1,1,2), B (4,3,1) and C (2,3,5). The vector representing internal bisector of the angle A is

A.  $\hat{i} + \hat{j} + 2\hat{k}$

B.  $2\hat{i} - 2\hat{j} + \hat{k}$

C.  $2\hat{i} + 2\hat{j} + \hat{k}$

D. none of these

**Answer: C**



 Watch Video Solution

75. Let  $\vec{a} = (1, 1, -1)$ ,  $\vec{b} = (5, -3, -3)$  and  $\vec{c} = (3, -1, 2)$ . If  $\vec{r}$  is collinear with  $\vec{c}$  and has length  $\frac{|\vec{a} + \vec{b}|}{2}$ , then  $\vec{r}$  equals

A.  $\pm 3c$

B.  $\pm \frac{3}{2}c$

C.  $\pm c$

D.  $\pm \frac{2}{3}c$

Answer: C

 Watch Video Solution

76. In a trapezium ABCD the vector  $\vec{BC} = \lambda \vec{AD}$ . If  $\vec{p} = \vec{AC} + \vec{BD}$  is collinear with  $\vec{AD}$  such that  $\vec{p} = \mu \vec{AD}$ , then

A.  $\mu = \lambda + 1$

B.  $\lambda = \mu + 1$

C.  $\lambda + \mu = 1$

D.  $\mu = 2 + \lambda$

**Answer: A**



[Watch Video Solution](#)

77. If the position vectors of the points A,B and C be  $\hat{i} + \hat{j}$ ,  $\hat{i} - \hat{j}$  and  $a\hat{i} + b\hat{j} + c\hat{k}$  respectively, then the points A,B and C are collinear, if

A.  $a=b=c=1$

B.  $a=1$ ,  $b$  and  $c$  are arbitrary scalars

C.  $ab=c=0$

D.  $c=0$ ,  $a=1$  and  $b$  is arbitrary scalars

**Answer: D**



[Watch Video Solution](#)

78. Let  $a, b$  and  $c$  be distinct non-negative numbers and the vectors  $a\hat{i} + a\hat{j} + c\hat{k}, \hat{i} + \hat{k}, c\hat{i} + c\hat{j} + b\hat{k}$  lie in a plane, then the quadratic equation  $ax^2 + 2cx + b = 0$  has

- A. real and equal roots
- B. real and unequal roots
- C. unreal roots
- D. both roots real and positive

**Answer: A**



[Watch Video Solution](#)

79. Which one is an irrational number?

A. (a)  $\frac{22}{7}$

B. (b)  $\pi$

C. (c) 2

D. (d)  $\sqrt{36}$

**Answer: A**



**Watch Video Solution**

**80.** The points  $A(2 - x, 2, 2)$ ,  $B(2, 2 - y, 2)$ ,  $C(2, 2, 2 - z)$  and  $D(1, 1, 1)$  are coplanar, then locus of  $P(x, y, z)$  is

A.  $\frac{1}{x} + \frac{1}{y} + \frac{1}{z} = 1$

B.  $x + y + z = 1$

C.  $\frac{1}{1-x} + \frac{1}{1-y} + \frac{1}{1-z} = 1$

D. none of these

**Answer: A**



**Watch Video Solution**

81. Which one is an rational number?

A. (a)  $\sqrt{36}$

B. (b)  $\sqrt{2}$

C. (c)  $\sqrt{6}$

D. (d)  $\pi$

Answer: B



Watch Video Solution

82. If  $a_1$  and  $a_2$  are two values of  $a$  for which the unit vector

$a\vec{i} + b\vec{j} + \frac{1}{2}\vec{k}$  is linearly dependent with  $\vec{i} + 2\vec{j}$  and  $\vec{j} - 2\vec{k}$ ,

then  $\frac{1}{a_1} + \frac{1}{a_2}$  is equal to

A. (a) 1

B. (b)  $\frac{1}{8}$

C. (c)  $\frac{-16}{11}$

D. (d)  $\frac{-11}{16}$

Answer: C



Watch Video Solution

83. The smallest odd prime number is



Watch Video Solution

84. Every Whole number is Natural number?



Watch Video Solution

85. A, B, C and D are four points such that

$$\overrightarrow{AB} = m(2\hat{i} - 6\hat{j} + 2\hat{k}) \quad \overrightarrow{BC} = (\hat{i} - 2\hat{j}) \quad \text{and} \quad \overrightarrow{CD} = n(-6\hat{i} + 15\hat{j} - 3\hat{k})$$

. If CD intersects AB at some points E, then

A.  $m \geq \frac{1}{2}$

B.  $n \geq \frac{1}{3}$

C.  $m = n$

D.  $m < n$

**Answer: A:B**

 [Watch Video Solution](#)

**86.** Given three vectors  $a, b$  and  $c$  are non-zero and non-coplanar vectors.

Then which of the following are coplanar.

A.  $\frac{|a|}{|a| = 2|b|}a + \frac{|b|}{|a| + |b|}b$

B.  $\frac{|b|}{|a| + |b|}a + \frac{|a|}{|a| + |b|}b$

C.  $\frac{|a|}{|a| + |b|}a + \frac{|b|}{|a| + 2|b|}b$

D.  $\frac{|b|}{2|a| + |b|}a + \frac{|a|}{2|a| + |b|}b$

**Answer: B::D**

 [Watch Video Solution](#)

87. The vectors  $x\hat{i} + (x + 1)\hat{j} + (x + 2)\hat{k}$ ,  $(x + 3)\hat{i} + (x + 4)\hat{j} + (x + 5)\hat{k}$  and  $(x + 6)\hat{i}$  are coplanar if  $x$  is equal to

- A. a. 1
- B. b. -3
- C. c. 4
- D. d. 0

**Answer: A::B::C::D**

 [Watch Video Solution](#)

88. Given three vectors  $\vec{a}$ ,  $\vec{b}$  and  $\vec{c}$  are non-zero and non-coplanar vectors. Then which of the following are coplanar.



A.  $a + b, b + c, c + a$

B.  $a - b, b + c, c + a$

C.  $a + b, b - c, c + a$

D.  $a + b, b + c, c - a$

Answer: B::C::D



Watch Video Solution

89. In a four-dimensional space where unit vectors along the axes are  $\hat{i}, \hat{j}, \hat{k}$  and  $\hat{l}$ , and  $a_1, a_2, a_3, a_4$  are four non-zero vectors such that no vector can be expressed as a linear combination of other

$$(\lambda - 1)(a_1 - a_2) + \mu(a_2 + a_3) + \gamma(a_3 + a_4 - 2a_2) + a_3 + \delta a_4 = 0,$$

then

A. (a)  $\lambda = 1$

B. (b)  $\mu = -\frac{2}{3}$

C. (c)  $\gamma = \frac{2}{3}$

D. (d)  $\delta = \frac{1}{3}$

**Answer: A::B::D**



**Watch Video Solution**

90. Statement 1:  $|\vec{a}| = 3, |\vec{b}| = 4$  and  $|\vec{a} + \vec{b}| = 5$ , then  $|\vec{a} - \vec{b}| = 5$ . Statement 2:

The length of the diagonals of a rectangle is the same.

- A. (a) Statement-I and statement II are correct and Statement II is the correct explanation of statement I
- B. (b) Both statement I and statement II are correct but statement II is not the correct explanation of statement I
- C. (c) Statement I is correct but statement II is incorrect
- D. (d) Statement II is correct but statement I is incorrect

**Answer: A**



Watch Video Solution

91. Statement 1: If  $|\vec{a} + \vec{b}| = |\vec{a} - \vec{b}|$ , then  $\vec{a}$  and  $\vec{b}$  are perpendicular to each other. Statement 2: If the diagonal of a parallelogram are equal magnitude, then the parallelogram is a rectangle.

- A. Statement-I and statement II are correct and Statement III is the correct explanation of statement I
- B. Both statement I and statement II are correct but statement II is not the correct explanation of statement I
- C. Statement I is correct but statement II is incorrect
- D. Statement II is correct but statement I is incorrect

Answer: A



Watch Video Solution

92. Find the slope of line. The Equation of line is  $2x - 3y = 2$



Watch Video Solution

93. Find the slope of line . The Equation of line is  $2x - 5y = 4$



Watch Video Solution

94. Statement I: If  $a = 2\hat{i} + \hat{k}$ ,  $b = 3\hat{j} + 4\hat{k}$  and  $c = \lambda a + \mu b$  are coplanar, then  $c = 4a - b$ .

Statement II: A set vector  $a_1, a_2, a_3, \dots, a_n$  is said to be linearly independent, if every relation of the form

$l_1 a_1 + l_2 a_2 + l_3 a_3 + \dots + l_n a_n = 0$  implies that

$l_1 = l_2 = l_3 = \dots = l_n = 0$  (scalar).

A. Statement-I and statement II are correct and Statement II is the correct explanation of statement I

B. Both statement I and statement II are correct but statement II is not the correct explanation of statement I

C. Statement I is correct but statement II is incorrect

D. Statement II is correct but statement I is incorrect

**Answer: B**



[Watch Video Solution](#)

95. Find the Equation of line having slop 2 and point (2,3)



[Watch Video Solution](#)

96. Statement 1: Let  $\vec{a}$ ,  $\vec{b}$ ,  $\vec{c}$  and  $\vec{d}$  be the position vectors of four points  $A, B, C$  and  $D$  and  $3\vec{a} - 2\vec{b} + 5\vec{c} - 6\vec{d} = 0$ . Then points  $A, B, C,$  and  $D$  are coplanar. Statement 2: Three non-zero, linearly dependent coinitial vector  $(\vec{P}Q, \vec{P}R$  and  $\vec{P}S)$  are coplanar. Then  $\vec{P}Q = \lambda\vec{P}R + \mu\vec{P}S$ , where  $\lambda$  and  $\mu$  are scalars.

- A. Statement-II and statement II ar correct and Statement III is the correct explanation of statement I
- B. Both statement I and statement II are correct but statement II is not the correct explanation of statement I
- C. Statement I is correct but statement II is incorrect
- D. Statement II is correct but statement I is incorrect

**Answer: A**



**Watch Video Solution**

**97.** Given that  $p(3,2,-4)$  ,  $Q (5,4, -6)$  and  $R (9,8,-10)$  are collinear find the ratio in which  $Q$  divides  $PR$

- A. 1 : 2
- B. 1 : 3
- C. 3 : 1

D. 2:1

**Answer: C**



[Watch Video Solution](#)

98. Given that  $P(3,2,-4)$ ,  $Q(5,4,-6)$  and  $R(9,8,-10)$  are collinear find the ratio in which  $Q$  divides  $PR$

A. 1:2

B. 1:3

C. 3:1

D. 2:1

**Answer: B**



[Watch Video Solution](#)

99. ABCD is a parallelogram. L is a point on BC which divides BC in the ratio 1 : 2. AL intersects BD at P. M is a point on DC which divides DC in the ratio 1 : 2 and AM intersects BD in Q.

$PQ : DB$  is equal to

A.  $\frac{2}{3}$

B.  $\frac{1}{3}$

C.  $\frac{1}{2}$

D.  $\frac{3}{4}$

**Answer: B**



[Watch Video Solution](#)

100. Let A,B,C,D,E represent vertices of a regular pentagon ABCDE. Given the position vector of these vertices be  $a, a+b, b, \lambda a$  and  $\lambda b$  respectively.

Q. AD divides EC in the ratio



A.  $1 - \cos \frac{3\pi}{5} : \cos \frac{3\pi}{5}$

B.  $1 + 2\cos \frac{2\pi}{5} : \cos \frac{\pi}{5}$

C.  $1 + 2\cos \frac{\pi}{5} : 2\cos \frac{\pi}{5}$

D. none of these

**Answer: C**



**Watch Video Solution**

**101.** Let A,B,C,D,E represent vertices of a regular pentagon ABCDE. Given the position vector of these vertices be  $a, a+b, b, \lambda a$  and  $\lambda b$  respectively.

Q. AD divides EC in the ratio

A.  $\cos \frac{2\pi}{5} : 1$

B.  $\cos \frac{3\pi}{5} : 1$

C.  $1 : 2\cos \frac{\pi}{5}$

D.  $1 : 2$

**Answer: C**



**Watch Video Solution**

**102.** In a parallelogram  $OABC$ , vectors  $\vec{a}$ ,  $\vec{b}$ ,  $\vec{c}$  are respectively the positions of vectors of vertices  $A, B, C$  with reference to  $O$  as origin. A point  $E$  is taken on the side  $BC$  which divide the line  $2:1$  internally. Also the line segment  $AE$  intersect the line bisecting the angle  $O$  internally in point  $P$ . If  $CP$ , when extended meets  $AB$  in point  $F$ . Then The position vector of point  $P$ , is



**Watch Video Solution**

**103.** In a parallelogram  $OABC$  vectors  $a, b, c$  respectively, THE POSITION VECTORS OF VERTICES  $A, B, C$  with reference to  $O$  as origin. A point  $E$  is taken on the side  $BC$  which divides it in the ratio of  $2:1$  also, the line segment  $AE$  intersects the line bisecting the angle  $\angle AOC$  internally at

point P. if CP when extended meets AB in points F, then

Q. The position vector of point P is

A.  $\frac{2|a|}{||a| - 3|c| |}$

B.  $\frac{|a|}{||a| - 3|c| |}$

C.  $\frac{3|a|}{||a| - 3|c| |}$


D.  $\frac{3|c|}{3||c| - |a| |}$

**Answer: B**



**Watch Video Solution**

**104.** The direction ( $\theta$ ) of  $\vec{E}$  at point P due to uniformly charged finite rod will be



Watch Video Solution

105.  $P, Q$  have position vectors  $\vec{a}$  &  $\vec{b}$  relative to the origin 'O' &  $X, Y$  and  $P, Q$  internally and externally respectively in the ratio

2:1 Vector  $\vec{XY} = \frac{3}{2}(\vec{b} - \vec{a})$  b.  $\frac{4}{3}(\vec{a} - \vec{b})$  c.  $\frac{5}{6}(\vec{b} - \vec{a})$  d.  $\frac{4}{3}(\vec{b} - \vec{a})$

 [Watch Video Solution](#)

**106.**  $A(1, -1, -3), B(2, 1, -2)$  &  $C(-5, 2, -6)$  are the position vectors of the vertices of a triangle ABC. The length of the bisector of its internal angle at A is :

 [Watch Video Solution](#)

**107.** Let ABC be a triangle whose centroid is G, orthocentre is H and circumcentre is the origin 'O'. If D is any point in the plane of the triangle such that no three of O, A, C and D are collinear satisfying the relation.  $AD+BD+CH+3HG=\lambda HD$ , then what is the value of the scalar  $\lambda$ .

 [Watch Video Solution](#)

108. Let  $\vec{a}$ ,  $\vec{b}$  and  $\vec{c}$  be unit vectors such that  $\vec{a} + \vec{b} - \vec{c} = 0$ . If the area of triangle formed by vectors  $\vec{a}$  and  $\vec{b}$  is  $A$ , then what is the value of  $4A^2$ ?



Watch Video Solution

109. The values of  $x$  for which the angle between the vectors  $\vec{a} = x\hat{i} - 3\hat{j} - \hat{k}$  and  $\vec{b} = 2x\hat{i} + x\hat{j} - \hat{k}$  is acute, and the angle, between the vector  $\vec{b}$  and the axis of ordinates is obtuse, are



Watch Video Solution

110. If the points  $a(\cos \alpha + \hat{i} \sin \alpha)$ ,  $b(\cos \beta + \hat{i} \sin \beta)$  and  $c(\cos \gamma + \hat{i} \sin \gamma)$  are collinear, then the value of  $|z|$  is \_\_\_\_\_ where  $z = bc \sin(\beta - \gamma) + ca \sin(\gamma - \alpha) + ab \sin(\alpha + \beta) + 3\hat{i}$



Watch Video Solution

111. A particle, in equilibrium, is subjected to four forces  $\vec{F}_1, \vec{F}_2, \vec{F}_3$  and  $\vec{F}_4$ ,

$$\vec{F}_1 = -10\hat{k}, \vec{F}_2 = u\left(\frac{4}{13}\hat{i} - \frac{12}{13}\hat{j} + \frac{3}{13}\hat{k}\right), \vec{F}_3 = v\left(-\frac{4}{13}\hat{i} - \frac{12}{13}\hat{j} + \frac{3}{13}\hat{k}\right)$$

then find the values of  $u, v$  and  $w$

 [Watch Video Solution](#)

112. Find the all the values of lamda such that  $(x, y, z) \neq (0, 0, 0)$  and

$$x(\hat{i} + \hat{j} + 3\hat{k}) + y(3\hat{i} - 3\hat{j} + \hat{k}) + z(-4\hat{i} + 5\hat{j}) = \lambda(x\hat{i} + y\hat{j} + z\hat{k})$$

 [Watch Video Solution](#)

113. If  $G$  is the centroid of  $\triangle ABC$  and  $G'$  is the centroid of

$$\triangle A'B'C' \text{ then } \vec{AA'} + \vec{BB'} + \vec{CC'} =$$

 [Watch Video Solution](#)

114. If D,E and F are the mid-points of the sides BC,CA and AB, respectively of a  $\Delta ABC$  and O is any point, show that

(i)  $AD+BE+CF=0$

 [Watch Video Solution](#)

115. If  $\vec{A}$  and  $\vec{B}$  are two vectors and  $k$  any scalar quantity greater than zero, then prove that  $\left| \vec{A} + \vec{B} \right|^2 \leq (1+k) \left| \vec{A} \right|^2 + \left( 1 + \frac{1}{k} \right) \left| \vec{B} \right|^2$ .

 [View Text Solution](#)

116. If O is the circumcentre, G is the centroid and O' the orthocenter of  $\Delta ABC$  prove that

(i)  $SA+SB+SC=3SG$ , where S is any point in the plane of  $\Delta ABC$ .

(ii)  $OA+OB+OC=OO'$

Where, AP is diameter of the circumcircle.

 [Watch Video Solution](#)



117. If  $A = \{2, 4, 6, 8, 10\}$  and  $B = \{3, 4, 6, 7, 9\}$  then  $A - B = ?$



Watch Video Solution

118. Statement -1 : If a transversal cuts the sides OL, OM and diagonal ON of a parallelogram at A, B, C respectively, then

$$\frac{OL}{OA} + \frac{OM}{OB} = \frac{ON}{OC}$$

Statement -2 : Three points with position vectors  $\vec{a}$ ,  $\vec{b}$ ,  $\vec{c}$  are collinear

iff there exist scalars  $x$ ,  $y$ ,  $z$  not all zero such that

$$x\vec{a} + y\vec{b} + z\vec{c} = \vec{0}, \text{ where } x + y + z = 0.$$



Watch Video Solution

119. If D, E and F are three points on the sides BC, CA and AB, respectively, of a triangle ABC such that the lines AD, BE and CF are concurrent, then show that

$$\frac{BD}{CD} \cdot \frac{CE}{AE} \cdot \frac{AF}{BF} = 1$$



Watch Video Solution

120.

Let

$$\vec{A}(t) = f_1(t)\hat{i} + f_2(t)\hat{j} \text{ and } \vec{B}(t) = g_1(t)\hat{i} + g_2(t)\hat{j}, t \in [0, 1], f_1, f_2, g_1, g_2$$

are continuous functions. If  $\vec{A}(t)$  and  $\vec{B}(t)$  are non-zero vectors for all

$$t \text{ and } \vec{A}(0) = 2\hat{i} + 3\hat{j}, \vec{A}(1) = 6\hat{i} + 2\hat{j}, \vec{B}(0) = 3\hat{i} + 2\hat{j} \text{ and } \vec{B}(1) = 2\hat{i} + 3\hat{j}$$

Then, show that  $\vec{A}(t)$  and  $\vec{B}(t)$  are parallel for some  $t$ .



Watch Video Solution

121. Prove that if  $\cos \alpha \neq 1$ ,  $\cos \beta \neq 1$  and  $\cos \gamma \neq 1$ , then the vectors

$$a = \hat{i} \cos \alpha + \hat{j} + \hat{k}, b = \hat{i} + \hat{j} \cos \beta + \hat{k} \text{ and } c = \hat{i} + \hat{j} + \hat{k} \cos \gamma \text{ can}$$

never be coplanar.



Watch Video Solution

122. If the vectors  $x\hat{i} + \hat{j} + \hat{k}$ ,  $\hat{i} + y\hat{j} + \hat{k}$  and  $\hat{i} + \hat{j} + z\hat{k}$  are coplanar

where,  $x \neq 1$ ,  $y \neq 1$  and  $z \neq 1$ , then prove that

$$\frac{1}{1-x} + \frac{1}{1-y} + \frac{1}{1-z} = 1$$

 [Watch Video Solution](#)

**123.** If  $\vec{a}$ ,  $\vec{b}$  and  $\vec{c}$  are any three non-coplanar vectors, then prove that points

$$l_1 \vec{a} + m_1 \vec{b} + n_1 \vec{c}, l_2 \vec{a} + m_2 \vec{b} + n_2 \vec{c}, l_3 \vec{a} + m_3 \vec{b} + n_3 \vec{c}, l_4 \vec{a} + m_4 \vec{b} + n_4 \vec{c}$$

are coplanar if 
$$\begin{vmatrix} l_1 & l_2 & l_3 & l_4 \\ m_1 & m_2 & m_3 & m_4 \\ n_1 & n_2 & n_3 & n_4 \\ 1 & 1 & 1 & 1 \end{vmatrix} = 0$$

 [Watch Video Solution](#)

**124.** Let  $r_1, r_2, r_3, \dots, r_n$  be the position vectors of points  $P_1, P_2, P_3, \dots, P_n$  relative to an origin O. show that if then a similar equation will also hold good with respect to any other origin O'. If  $a_1 + a_2 + a_3 + \dots + a_n = 0$ .

 [Watch Video Solution](#)

125. For what value of  $\lambda$ , the vectors  $(\lambda - 2)\vec{a} + \vec{b}$  and  $(4\lambda - 2)\vec{a} + 3\vec{b}$  are collinear

 [Watch Video Solution](#)

### Exercise For Session 1

1. Classify the following measure as scalar and vector:  $10Kg$

 [Watch Video Solution](#)

2. Represent the following graphically: A displacement of  $70km, 40^\circ$  north of west.

 [Watch Video Solution](#)


3. In the following figure, which of the vectors are:

(i) Collinear

(ii) Equal

(iii) Co-initial

(iv) collinear but not equal .



Watch Video Solution

4. Answer the following as true or false.

(i)  $\vec{a}$  and  $-\vec{a}$  are collinear.

(ii) Two collinear vectors are always equal in magnitude.

(iii) Two vectors having same magnitude are collinear.

(iv) Two collinear vectors having the same magnitude



[Watch Video Solution](#)

5. Find the perimeter of a triangle with sides

$3\hat{i} + 4\hat{j} + 5\hat{k}$ ,  $4\hat{i} - 3\hat{j} - 5\hat{k}$  and  $7\hat{i} + \hat{j}$ .



[Watch Video Solution](#)

6. Find the angle of vector  $\vec{a} = 6\hat{i} + 2\hat{j} - 3\hat{k}$  with  $x$ -axis.



[Watch Video Solution](#)

7. Write the direction ratios of the vector  $r = \hat{i} - \hat{j} + 2\hat{k}$  and hence calculate its direction cosines.

 [Watch Video Solution](#)

## Exercise For Session 2

1. If  $a = 2\hat{i} - \hat{j} + 2\hat{k}$  and  $b = -\hat{i} + \hat{j} - \hat{k}$ , then find  $a+b$ . Also, find a unit vector along  $a+b$ .

 [Watch Video Solution](#)

2. Find a unit vector in the direction of the resultant of the vectors  $(\hat{i} + 2\hat{j} + 3\hat{k})$ ,  $(-\hat{i} + 2\hat{j} + \hat{k})$  and  $(3\hat{i} + \hat{j})$ .

 [Watch Video Solution](#)

3. Find the direction cosines of the resultant of the vectors  $(\hat{i} + \hat{j} + \hat{k})$ ,  $(-\hat{i} + \hat{j} + \hat{k})$ ,  $(\hat{i} - \hat{j} + \hat{k})$  and  $(\hat{i} + \hat{j} - \hat{k})$ .

 [Watch Video Solution](#)

4. In a regular hexagon ABCDEF,  $AB=a, BC=b$  and  $CD=c$ . Then,  $\overrightarrow{AE}$  is equal to

 [Watch Video Solution](#)

5. Prove that  $3\overrightarrow{OD} + \overrightarrow{DA} + \overrightarrow{DB} + \overrightarrow{DC}$  is equal to  $\overrightarrow{OA} + \overrightarrow{OB} + \overrightarrow{OC}$ .

 [Watch Video Solution](#)

6. In a regular hexagon ABCDEF,  $\overline{AB} + \overline{AC} + \overline{AD} + \overline{AE} + \overline{AF} = k\overline{AD}$  then k is equal to

 [Watch Video Solution](#)



7.  $ABCDE$  is a pentagon. Prove that the resultant of forces

$\vec{AB}, \vec{AE}, \vec{BC}, \vec{DC}, \vec{ED}$  and  $\vec{AC}$  is  $3\vec{AC}$ .



[Watch Video Solution](#)

8. find the area of square whose side is 25 cm.



[Watch Video Solution](#)

9. If  $P(-1, 2)$  and  $Q(3, -7)$  are two points, express the vector  $PQ$  in terms of unit vectors  $\hat{i}$  and  $\hat{j}$  also, find distance between point  $P$  and  $Q$ .

What is the unit vector in the direction of  $PQ$ ?



[Watch Video Solution](#)

10. If  $\vec{OP} = 2\hat{i} + 3\hat{j} - \hat{k}$  and  $\vec{OQ} = 3\hat{i} - 4\hat{j} + 2\hat{k}$  find the modulus and direction cosines of  $\vec{PQ}$ .





Watch Video Solution

11. Show that the points :  
 $A(2\hat{i} - \hat{j} + \hat{k})$ ,  $B(\hat{i} - 3\hat{j} - \hat{k})$ ,  $C(3\hat{i} - 4\hat{j} - 4\hat{k})$  are the vertices of a right-angled triangle.



Watch Video Solution

12. If  $a = 2\hat{i} + 2\hat{j} - \hat{k}$  and  $|x\vec{a}| = 1$ , then find x.



Watch Video Solution

13. If  $p = 7\hat{i} - 2\hat{j} + 3\hat{k}$  and  $q = 3\hat{i} + \hat{j} + 5\hat{k}$ , then find the magnitude of  $p-2q$ .



Watch Video Solution

14. Find a vector in the direction of  $5\hat{i} - \hat{j} + 2\hat{k}$ , which has magnitude 8 units.

 [Watch Video Solution](#)

15. If  $a = \hat{i} + 2\hat{j} + 2\hat{k}$  and  $b = 3\hat{i} + 6\hat{j} + 2\hat{k}$ , then find a vector in the direction of  $a$  and having magnitude as  $|b|$ .

 [Watch Video Solution](#)

16. Find the position vector of a point  $R$  which divides the line joining the points  $P(\hat{i} + 2\hat{j} - \hat{k})$  and  $Q(\hat{i} + 2\hat{j} + 2\hat{k})$  internally in the ratio 2:1.

 [Watch Video Solution](#)

17. If the position vector of one end of the line segment  $AB$  be  $2\hat{i} + 3\hat{j} - \hat{k}$  and the position vector of its middle point be  $3(\hat{i} + \hat{j} + \hat{k})$ ,

then find the position vector of the other end.



[Watch Video Solution](#)

### Exercise For Session 3

1. Show that the points  $A(1,3,2)$ ,  $B(-2,0,1)$  and  $C(4,6,3)$  are collinear.



[Watch Video Solution](#)

2. If the position vectors of the points  $A, B$  and  $C$  be  $a, b$  and  $3a-2b$  respectively, then prove that the points  $A, B$  and  $C$  are collinear.



[Watch Video Solution](#)

3. The position vectors of four points  $P, Q, R$  and  $S$  are  $2a+4c, 5a+3\sqrt{3}b+4c, -2\sqrt{3}b+c$  and  $2a+c$  respectively, prove that  $PQ$  is parallel to  $RS$ .



Watch Video Solution

4. If three points A,B and C have position vectors  $(1,x,3)$ ,  $(3,4,7)$  and  $(y,-2,-5)$ , respectively and if they are collinear, then find  $(x,y)$ .



Watch Video Solution

5. Show that the three points with position vectors  $-2\hat{i} + 3\hat{j} + 5\hat{k}$ ,  $\hat{i} + 2\hat{j} + 3\hat{k}$  and  $7\hat{i} - \hat{k}$  are collinear.



Watch Video Solution

6.  $a$  and  $b$  are non-collinear vectors. If  $c = (x - 2)a + b$  and  $d = (2x + 1)a - b$  are collinear vectors, then find the value of  $x$ .



Watch Video Solution

7. Let  $a, b, c$  are three vectors of which every pair is non-collinear, if the vectors  $a+b$  and  $b+c$  are collinear with  $c$  and  $a$  respectively, then find  $a+b+c$ .

 [Watch Video Solution](#)

8. Show that the vectors  $\hat{i} - \hat{j} - \hat{k}$ ,  $2\hat{i} + 3\hat{j} + \hat{k}$  and  $7\hat{i} + 3\hat{j} - 4\hat{k}$  are coplanar.

 [Watch Video Solution](#)

9. If the vectors  $2\hat{i} - \hat{j} + \hat{k}$ ,  $\hat{i} + 2\hat{j} - 3\hat{k}$  and  $3\hat{i} + a\hat{j} + 5\hat{k}$  are coplanar, then prove that  $a = -4$ .

 [Watch Video Solution](#)

10. Show that the vectors  $a - 2b + 4c$ ,  $-2a + 3b - 6c$  and  $-b + 2c$  are coplanar vector, where  $a, b, c$  are non-coplanar vectors.



Watch Video Solution

11. If  $\vec{a}$ ,  $\vec{b}$  and  $\vec{c}$  are non-coplanar vectors, prove that the four points  $2\vec{a} + 3\vec{b} - \vec{c}$ ,  $\vec{a} - 2\vec{b} + 3\vec{c}$ ,  $3\vec{a} + 4\vec{b} - 2\vec{c}$  and  $\vec{a} - 6\vec{b} + 6\vec{c}$  are coplanar.



Watch Video Solution

### Exercise Single Option Correct Type Questions

1. If  $a = 3\hat{i} - 2\hat{j} + \hat{k}$ ,  $b = 2\hat{i} - 4\hat{j} - 3\hat{k}$  and  $c = -\hat{i} + 2\hat{j} + 2\hat{k}$ , then  $a+b+c$  is

A.  $3\hat{i} - 4\hat{j}$

B.  $3\hat{i} + 4\hat{j}$

C.  $4\hat{i} - 4\hat{j}$

D.  $4\hat{i} + 4\hat{j}$

**Answer: C**



**Watch Video Solution**

2. What should be added in vector  $a = 3\hat{i} + 4\hat{j} - 2\hat{k}$  to get its resultant a unit vector  $\hat{i}$ ?

A.  $-2\hat{i} - 4\hat{j} + 2\hat{k}$

B.  $-2\hat{i} + 4\hat{j} - 2\hat{k}$

C.  $2\hat{i} + 4\hat{j} - 2\hat{k}$

D. none of these

**Answer: A**



**Watch Video Solution**

3. If  $a = 2\hat{i} + 2\hat{j} - 8\hat{k}$  and  $b = \hat{i} + 3\hat{j} - 4\hat{k}$ , then the magnitude of  $a+b$  is equal to



A. 13

B.  $\frac{13}{5}$

C.  $\frac{3}{13}$

D.  $\frac{4}{13}$

**Answer: A**

 [Watch Video Solution](#)

4. If  $a = 2\hat{i} + 5\hat{j}$  and  $b = 2\hat{i} - \hat{j}$ , then the unit vector along  $a+b$  will be

A.  $\frac{\hat{i} - \hat{j}}{\sqrt{2}}$

B.  $\hat{i} + \hat{j}$

C.  $\sqrt{2}(\hat{i} + \hat{j})$

D.  $\frac{\hat{i} + \hat{j}}{\sqrt{2}}$

**Answer: D**

 [Watch Video Solution](#)

5. Find the unit vector parallel to the resultant vector of  $2\hat{i} + 4\hat{j} - 5\hat{k}$  and  $\hat{i} + 2\hat{j} + 3\hat{k}$ .

A.  $\frac{1}{7}(3\hat{i} + \hat{j} + \hat{k})$

B.  $\frac{\hat{i} + \hat{j} + \hat{k}}{\sqrt{3}}$

C.  $\frac{\hat{i} + \hat{j} + 2\hat{k}}{\sqrt{6}}$

D.  $\frac{1}{\sqrt{69}}(-\hat{i} - \hat{j} + 8\hat{k})$

**Answer: A**



**Watch Video Solution**

6. If  $a = \hat{i} + 2\hat{j} + 3\hat{k}$ ,  $b = -\hat{i} + 2\hat{j} + \hat{k}$  and  $c = 3\hat{i} + \hat{j}$ , then the unit vector along its resultant is

A.  $3\hat{i} + 5\hat{j} + 4\hat{k}$

B.  $\frac{3\hat{i} + 5\hat{j} + 4\hat{k}}{50}$

C.  $\frac{3\hat{i} + 5\hat{j} + 4\hat{k}}{5\sqrt{2}}$

D. none of these

**Answer: C**



**Watch Video Solution**

7. If  $a = (2, 5)$  and  $b = (1, 4)$ , then vector parallel to  $(a+b)$  is

A. (3,5)

B. (1,1)

C. (1,3)

D. (8,5)

**Answer: C**



**Watch Video Solution**

8. In the  $\triangle ABC$ ,  $AB = a$ ,  $AC = c$  and  $BC = b$ , then

A.  $a+b+c=0$

B.  $a+b-c=0$

C.  $a-b+c=0$

D.  $-a + b + c = 0$

**Answer: B**



[Watch Video Solution](#)

9. If O is origin and the position vector of A is  $4\hat{i} + 5\hat{j}$ , then unit vector parallel to OA is

A.  $\frac{4}{\sqrt{41}}\hat{i}$

B.  $\frac{5}{\sqrt{41}}\hat{i}$

C.  $\frac{1}{\sqrt{41}}(4\hat{i} + 5\hat{j})$

D.  $\frac{1}{\sqrt{41}}(4\hat{i} - 5\hat{j})$

**Answer: C**



**Watch Video Solution**

10. The position vectors of the points A, B and C are  $\hat{i} + 2\hat{j} - \hat{k}$ ,  $\hat{i} + \hat{j} + \hat{k}$  and  $2\hat{i} + 3\hat{j} + 2\hat{k}$ , respectively. If A is chosen as the origin, then the position vectors of B and C are

A.  $\hat{i} + 2\hat{k}$ ,  $\hat{i} + \hat{j} + 3\hat{k}$

B.  $\hat{j} + 2\hat{k}$ ,  $\hat{i} + \hat{j} + 3\hat{k}$

C.  $-\hat{j} + 2\hat{k}$ ,  $\hat{i} - \hat{j} + 3\hat{k}$

D.  $-\hat{j} + 2\hat{k}$ ,  $\hat{i} + \hat{j} + 3\hat{k}$

**Answer: D**



**Watch Video Solution**

11. The position vectors of  $P$  and  $Q$  are  $5\hat{i} + 4\hat{j} + a\hat{k}$  and  $-\hat{i} + 2\hat{j} - 2\hat{k}$ , respectively. If the distance between them is 7, then find the value of  $a$ .

A.  $-5, 1$

B.  $5, 1$

C.  $0, 5$

D.  $1, 0$

**Answer: A**



[Watch Video Solution](#)

12. If position vector of points  $A, B$  and  $C$  are respectively  $\hat{i}, \hat{j}$ , and  $\hat{k}$  and  $AB = CX$ , then position vector of point  $X$  is

A.  $-\hat{i} + \hat{j} + \hat{k}$

B.  $\hat{i} - \hat{j} + \hat{k}$

C.  $\hat{i} + \hat{j} - \hat{k}$

D.  $\hat{i} + \hat{j} + \hat{k}$

**Answer: A**



**Watch Video Solution**

13. The position vectors of A and B are  $2\hat{i} - 9\hat{j} - 4\hat{k}$  and  $6\hat{i} - 3\hat{j} + 8\hat{k}$  respectively, then the magnitude of AB is

A. 11

B. 12

C. 13

D. 14

**Answer: D**



**Watch Video Solution**

14. If the position vectors of P and Q are  $(\hat{i} + 3\hat{j} - 7\hat{k})$  and  $(5\hat{i} - 2\hat{j} + 4\hat{k})$ , then  $|PQ|$  is

A.  $\sqrt{158}$

B.  $\sqrt{160}$

C.  $\sqrt{161}$

D.  $\sqrt{162}$

**Answer: D**



[Watch Video Solution](#)

15. If the position vectors of P and Q are  $\hat{i} + 2\hat{j} - 7\hat{k}$  and  $5\hat{i} - 2\hat{j} + 4\hat{k}$  respectively, the cosine of the angle between PQ and Z-axis is

A.  $\frac{4}{\sqrt{162}}$

B.  $\frac{11}{\sqrt{162}}$

C.  $\frac{5}{\sqrt{162}}$



D.  $\frac{-5}{\sqrt{162}}$

**Answer: B**



**Watch Video Solution**

16. If the position vectors of A and B are  $\hat{i} + 3\hat{j} - 7\hat{k}$  and  $5\hat{i} - 2\hat{j} + 4\hat{k}$ , then the direction cosine of AB along Y-axis is

A.  $\frac{4}{\sqrt{162}}$

B.  $-\frac{5}{\sqrt{162}}$

C.  $-5$

D. 11

**Answer: B**



**Watch Video Solution**

17. The direction cosines of vector  $a = 3\hat{i} + 4\hat{j} + 5\hat{k}$  in the direction of positive axis of X, is

A. A.  $\pm \frac{3}{\sqrt{50}}$

B. B.  $\frac{4}{\sqrt{50}}$

C. C.  $\frac{3}{\sqrt{50}}$

D. D.  $-\frac{4}{\sqrt{50}}$

**Answer: C**



**Watch Video Solution**

18. The direction cosines of the vector  $3\hat{i} - 4\hat{j} + 5\hat{k}$  are

A. A.  $\frac{3}{5}, -\frac{4}{5}, \frac{1}{5}$

B. B.  $\frac{3}{5\sqrt{2}}, \frac{-4}{5\sqrt{2}}, \frac{1}{\sqrt{2}}$

C. C.  $\frac{3}{\sqrt{2}}, \frac{-4}{\sqrt{2}}, \frac{1}{\sqrt{2}}$

D. D.  $\frac{3}{5\sqrt{2}}, \frac{4}{5\sqrt{2}}, \frac{1}{\sqrt{2}}$

**Answer: B**



[Watch Video Solution](#)

19. The point having position vectors

$2\hat{i} + 3\hat{j} + 4\hat{k}$ ,  $3\hat{i} + 4\hat{j} + 2\hat{k}$  and  $4\hat{i} + 2\hat{j} + 3\hat{k}$  are the vertices of

A. A right angled triangle

B. B. isosceles triangle

C. C. equilateral triangle

D. D. collinear

**Answer: C**



[Watch Video Solution](#)

20. If the position vectors of the vertices A,B and C of a  $\triangle ABC$  are

$7\hat{j} + 10\hat{k}$ ,  $-\hat{i} + 6\hat{j} + 6\hat{k}$  and  $-4\hat{i} + 9\hat{j} + 6\hat{k}$ , respectively, the

triangle is

- A. A. equilateral
- B. B. isosceles
- C. C. scalene
- D. D. right angled and isosceles also

**Answer: D**



**Watch Video Solution**

21. If  $a, b$  and  $c$  are the position vectors of the vertices  $A, B$  and  $C$  of the  $\triangle ABC$ , then the centroid of  $\triangle ABC$  is

A. A.  $\frac{a + b + c}{3}$

B. B.  $\frac{1}{2} \left( a + \frac{b + c}{2} \right)$

C. C.  $a + \frac{b + c}{2}$

D. D.  $\frac{a + b + c}{2}$

**Answer: A**



**Watch Video Solution**

**22.** If  $a$  and  $b$  are position vector of two points A,B and C divides AB in ratio 2:1, then position vector of C is

A.  $\frac{a + 2b}{3}$

B.  $\frac{2a + b}{3}$

C.  $\frac{a + 2}{3}$

D.  $\frac{a + b}{2}$

**Answer: A**



**Watch Video Solution**

**23.** Find the position vector of the point which divides the join of the points  $\left(2\vec{a} - 3\vec{b}\right)$  and  $\left(3\vec{a} - 2\vec{b}\right)$  (i) internally and (ii) externally in

the ratio 2:3 .



Watch Video Solution

24. If O is origin and C is the mid - point of A (2, -1) and B ( -4, 3) . Then value of OC is

A.  $\hat{i} + \hat{j}$

B.  $\hat{i} - \hat{j}$

C.  $-\hat{i} + \hat{j}$

D.  $-\hat{i} - \hat{j}$

Answer: C



Watch Video Solution

25. If the position vectors of the points A and B are  $\hat{i} + 3\hat{j} - \hat{k}$  and  $3\hat{i} - \hat{j} - 3\hat{k}$ , then what will be the position vector of the

mid-point of AB

A.  $\hat{i} + 2\hat{j} - \hat{k}$

B.  $2\hat{i} + \hat{j} - 2\hat{k}$

C.  $2\hat{i} + \hat{j} - \hat{k}$

D.  $\hat{i} + \hat{j} - 2\hat{k}$

**Answer: B**



**Watch Video Solution**

**26.** The position vectors of A and B are  $\hat{i} - \hat{j} + 2\hat{k}$  and  $3\hat{i} - \hat{j} + 3\hat{k}$ . The position vector of the middle points of the line AB is

A.  $\frac{1}{2}\hat{i} - \frac{1}{2}\hat{j} + \hat{k}$

B.  $2\hat{i} - \hat{j} + \frac{5}{2}\hat{k}$

C.  $\frac{3}{2}\hat{i} - \frac{1}{2}\hat{j} + \frac{3}{2}\hat{k}$

D. none of these

**Answer: B**



**Watch Video Solution**

27. If the vector  $\vec{b}$  is collinear with the vector  $\vec{a} (2\sqrt{2}, -1, 4)$  and  $|\vec{b}| = 10$ , then

A.  $a \pm b = 0$

B.  $a \pm 2b = 0$

C.  $2a \pm b = 0$

D. none of these

**Answer: C**



**Watch Video Solution**

28. If  $\vec{a}, \vec{b}$  are the position vectors of the points  $(1, -1), (-2, m)$ , find the value of  $m$  for which  $\vec{a}$  and  $\vec{b}$  are collinear.



A. 4

B. 3

C. 2

D. 0

**Answer: C**



**Watch Video Solution**

**29.** The points with position vectors  $10\hat{i} + 3\hat{j}$ ,  $12\hat{i} - 5\hat{j}$  and  $a\hat{i} + 11\hat{j}$  are collinear, if  $a$  is equal to

A.  $-8$

B. 4

C. 8

D. 12

**Answer: C**

 [Watch Video Solution](#)

30. The vectors  $\hat{i} + 2\hat{j} + 3\hat{k}$ ,  $\lambda\hat{i} + 4\hat{j} + 7\hat{k}$ ,  $-3\hat{i} - 2\hat{j} - 5\hat{k}$  are collinear, of  $\lambda$  is equal to

(A) 3

(B) 4

(C) 5

(D) 6

A. 3

B. 4

C. 5

D. 6

**Answer: A**

 [Watch Video Solution](#)

31. If the points  $a + b$ ,  $a - b$  and  $a + kb$  be collinear, then  $k$  is equal to

A. A. 0

B. B. 2

C. C.  $-2$

D. D. any real number

**Answer: D**



[Watch Video Solution](#)

32. If the position vectors of A, B, C and D are  $2\hat{i} + \hat{j}$ ,  $\hat{i} - 3\hat{j}$ ,  $3\hat{i} + 2\hat{j}$  and  $\hat{i} + \lambda\hat{j}$ , respectively and  $AB \parallel CD$ , then  $\lambda$  will be

A.  $-8$

B.  $-6$

C. 8

D. 6

**Answer: B**



[Watch Video Solution](#)

33. If the vectors  $3\hat{i} + 2\hat{j} - \hat{k}$  and  $6\hat{i} - 4x\hat{j} + y\hat{k}$  are parallel, then the value of x and y will be

A.  $-1, -2$

B.  $1, -2$

C.  $-1, 2$

D.  $1, 2$

**Answer: A**



[Watch Video Solution](#)

34. If  $a$  and  $b$  are two non collinear vectors; then every vector  $r$  coplanar with  $a$  and  $b$  can be expressed in one and only one way as a linear combination:  $xa+yb=0$ .

- A. (a) $x=0$ , but  $y$  is not necessarily zero
- B. (b) $y=0$ , but  $x$  is not necessarily zero
- C. (c) $x=0,y=0$
- D. (d)none of these

**Answer: C**



[Watch Video Solution](#)

35. Four non-zero vectors will always be

- A. linearly dependent
- B. linearly independent
- C. either (a) or (b)

D. none of these

**Answer: A**



[Watch Video Solution](#)

**36.** The vectors  $a, b$  and  $a+b$  are

A. collinear

B. coplanar

C. non-coplanar

D. none of these

**Answer: B**



[Watch Video Solution](#)

37. Find the all the values of lamda such that  $(x, y, z) \neq (0, 0, 0)$  and  $x(\hat{i} + \hat{j} + 3\hat{k}) + y(3\hat{i} - 3\hat{j} + \hat{k}) + z(-4\hat{i} + 5\hat{j}) = \lambda(x\hat{i} + y\hat{j} + z\hat{k})$

A. A.  $-2, 0$

B. B  $0, -2$

C. C.  $-1, 0$

D. D.  $0, -1$

**Answer: D**



[Watch Video Solution](#)

38. The number of integral values of  $p$  for which  $(p + 1)\hat{i} - 3\hat{j} + p\hat{k}, p\hat{i} + (p + 1)\hat{j} - 3\hat{k}$  and  $-3\hat{i} + p\hat{j} + (p + 1)\hat{k}$  are linearly dependent vectors is  $q$

A. 0

B. 1

C. 2

D. 3

**Answer: B**



[Watch Video Solution](#)

39. If the vectors  $AB = 3\hat{i} + 4\hat{k}$  and  $AC = 5\hat{i} - 2\hat{j} + 4\hat{k}$  are the sides of a  $\triangle ABC$ , then the length of the median through A is

A.  $\sqrt{18}$

B.  $\sqrt{72}$

C.  $\sqrt{33}$

D.  $\sqrt{288}$


**Answer: C**



[Watch Video Solution](#)



40. In the figure, a vectors  $x$  satisfies the equation  $x-w=v$ . then,  $x$  is equal to



A.  $2a + b + c$

B.  $a + 2b + c$

C.  $a + b + 2c$

D.  $a + b + c$

**Answer: B**

 [Watch Video Solution](#)

41. Vectors  $\vec{a} = \hat{i} + 2\hat{j} + 3\hat{k}$ ,  $\vec{b} = 2\hat{i} - \hat{j} + \hat{k}$  and  $\vec{c} = 3\hat{i} + \hat{j} + 4\hat{k}$  are so placed that the end point of one vector is the starting point of the next vector. Then the vectors are

- A. not coplanar
- B. coplanar but cannot form a triangle
- C. coplanar and form a triangle
- D. coplanar and can form a right angled triangle.

**Answer: B**



[Watch Video Solution](#)

42. If  $OP=8$  and  $OP$  makes angles  $45^\circ$  and  $60^\circ$  with  $OX$ -axis and  $OY$ -axis respectively, then  $OP$  is equal to

A.  $8(\sqrt{2}\hat{i} + \hat{j} \pm \hat{k})$

B.  $4(\sqrt{2}\hat{i} + \hat{j} \pm \hat{k})$

C.  $\frac{1}{4}(\sqrt{2}\hat{i} + \hat{j} \pm \hat{k})$

D.  $\frac{1}{8}(\sqrt{2}\hat{i} + \hat{j} \pm \hat{k})$

**Answer: B**



**Watch Video Solution**

**43.** Let  $a, b$  and  $c$  be three unit vectors such that  $3a + 4b + 5c = 0$ . Then which of the following statements is true?

A.  $a$  is parallel to  $b$

B.  $a$  is perpendicular to  $b$

C.  $a$  is neither parallel nor perpendicular to  $b$

D. none of these

**Answer: D**



**Watch Video Solution**

44. if A,B,C,D and E are five coplanar points, then

$\vec{DA} + \vec{DB} + \vec{DC} + \vec{AE} + \vec{BE} + \vec{CE}$  is equal to :

A. DE

B. 3DE

C. 2DE

D. 4ED

**Answer: B**



[Watch Video Solution](#)

45. If the vectors  $\vec{a}$  and  $\vec{b}$  are linearly independent satisfying

$(\sqrt{3}\tan\theta + 1)\vec{a} + (\sqrt{3}\sec\theta - 2)\vec{b} = 0$ , then the most general values

of  $\theta$  are

A.  $n\pi - \frac{\pi}{6}, n \in Z$

$$B. 2n\pi \pm \frac{11\pi}{6}n \in Z$$

$$C. n\pi \pm \frac{\pi}{6}, n \in Z$$

$$D. 2n\pi + \frac{11\pi}{6}, n \in Z$$

**Answer: D**



**Watch Video Solution**

**46.** Find the slope of the normal having point (3,2) and (4,1)



**Watch Video Solution**

**47.** A line passes through the points whose position vectors are

$\hat{i} + \hat{j} - 2\hat{k}$  and  $\hat{i} - 3\hat{j} + \hat{k}$ . The position vector of a point on it at unit

distance from the first point is (A)  $\frac{1}{5}(5\hat{i} + \hat{j} - 7\hat{k})$  (B)

$\frac{1}{5}(5\hat{i} + 9\hat{j} - 13\hat{k})$  (C)  $(\hat{i} - 4\hat{j} + 3\hat{k})$  (D)  $\frac{1}{5}(\hat{i} - 4\hat{j} + 3\hat{k})$

A. A.  $\frac{1}{5}(5\hat{i} + \hat{j} - 7\hat{k})$

B.  $\frac{1}{5}(4\hat{i} + 9\hat{j} - 15\hat{k})$

C.  $(\hat{i} - 4\hat{j} + 3\hat{k})$

D.  $\frac{1}{5}(\hat{i} - 4\hat{j} + 3\hat{k})$

**Answer: A**



**Watch Video Solution**

**48.** Find the slop of line . The Equation of line is  $2x - 3y = 2$



**Watch Video Solution**

**49.** If P and Q are the middle points of the sides BC and CD of the parallelogram ABCD, then AP+AQ is equal to

A. AC

B.  $\frac{1}{2}AC$

C.  $\frac{2}{3}AC$

D.  $\frac{3}{2}AC$

**Answer: D**



**Watch Video Solution**

50. If the figure formed by the four points  $\hat{i} + \hat{j} - \hat{k}$ ,  $2\hat{i} + 3\hat{j}$ ,  $3\hat{i} + 5\hat{j} - 2\hat{k}$  and  $\hat{k} - \hat{j}$  is

A. rectangle

B. parallelogram

C. trapezium

D. none of these

**Answer: C**



**Watch Video Solution**

51. A and B are two points. The position vector of A is  $6b-2a$ . A point P divides the line AB in the ratio 1:2. If  $a-b$  is the position vector of P, then the position vector of B is given by

A.  $7a-15b$

B.  $7a+15b$

C.  $15a-7b$

D.  $15a+7b$

**Answer: A**



[Watch Video Solution](#)

52. If three points A, B and C are collinear, whose position vectors are  $\hat{i} - 2\hat{j} - 8\hat{k}$ ,  $5\hat{i} - 2\hat{k}$  and  $11\hat{i} + 3\hat{j} + 7\hat{k}$  respectively, then the ratio in which B divides AC is

A.  $1:2$



B. B. 2:3

C. C. 2:1

D. D. 1:1

**Answer: B**



[Watch Video Solution](#)

53. If in a triangle  $AB=a, AC=b$  and  $D, E$  are the mid-points of  $AB$  and  $AC$  respectively, then  $DE$  is equal to

A.  $\frac{a}{4} - \frac{b}{4}$

B.  $\frac{a}{2} - \frac{b}{2}$

C.  $\frac{b}{4} - \frac{a}{4}$

D.  $\frac{b}{2} - \frac{a}{2}$

**Answer: D**



[Watch Video Solution](#)

54. The two adjacent sides of a parallelogram are  $2\hat{i} + 4\hat{j} - 5\hat{k}$  and  $\hat{i} + 2\hat{j} + 3\hat{k}$ . Find the unit vectors along the diagonals of the parallelogram.

A.  $\frac{1}{\sqrt{69}}(\hat{i} + 2\hat{j} - 8\hat{k})$

B.  $\frac{1}{69}(\hat{i} + 2\hat{j} - 8\hat{k})$

C.  $\frac{1}{\sqrt{69}}(-\hat{i} - 2\hat{j} + 8\hat{k})$

D.  $\frac{1}{69}(-\hat{i} - 2\hat{j} + 8\hat{k})$

**Answer: C**



**Watch Video Solution**

55. If A, B and C are the vertices of a triangle with position vectors  $\vec{a}$ ,  $\vec{b}$  and  $\vec{c}$  respectively and G is the centroid of  $\triangle ABC$ , then  $\vec{GA} + \vec{GB} + \vec{GC}$  is equal to

A. 0

B.  $A + B + C$

C.  $\frac{a + b + c}{3}$

D.  $\frac{a + b - c}{3}$

**Answer: A**



**Watch Video Solution**

56. If ABCDEF is a regular hexagon then  $\overrightarrow{AD} + \overrightarrow{EB} + \overrightarrow{FC}$  equals :

A. 0

B. 2AB

C. 3AB

D. 4AB

**Answer: D**



**Watch Video Solution**

57. Find the discriminant of the quadratic equation  $x^2 + 3x + 2$

 [Watch Video Solution](#)

58. In a regular hexagon ABCDEF, prove that  $AB+AC+AD+AE+AF=3AD$ .

 [Watch Video Solution](#)

59. Find the dot product of the vector  $2\hat{i} + 3\hat{j} + 1\hat{k}$  and  $1\hat{i} + 2\hat{j} + 3\hat{k}$

 [Watch Video Solution](#)

60. If a and b are two non-zero and non-collinear vectors then  $a+b$  and  $a-b$  are

A. linearly dependent vectors

B. linearly independent vectors

C. linearly dependent and independent vectors

D. none of these

**Answer: B**



[Watch Video Solution](#)

61. If  $|\vec{a} + \vec{b}| < |\vec{a} - \vec{b}|$ , then the angle between  $\vec{a}$  and  $\vec{b}$  can lie in the interval

A.  $(\pi/2, \pi/2)$

B.  $(0, \pi)$

C.  $(\pi/2, 3\pi/2)$

D.  $(0, 2\pi)$

**Answer: C**



[Watch Video Solution](#)

62. The magnitudes of mutually perpendicular forces  $a, b$  and  $c$  are 2, 10 and 11 respectively. Then the magnitude of its resultant is

- A. 12
- B. 15
- C. 9
- D. none of these

**Answer: B**



[Watch Video Solution](#)

63. If  $\hat{i} - 3\hat{j} + 5\hat{k}$  bisects the angle between  $\hat{a}$  and  $-\hat{i} + 2\hat{j} + 2\hat{k}$ , where  $\hat{a}$  is a unit vector, then

A.  $a = \frac{1}{105} (41\hat{i} + 88\hat{j} - 40\hat{k})$

B.  $a = \frac{1}{105} (41\hat{i} + 88\hat{j} + 40\hat{k})$

$$C. a = \frac{1}{105} (-41\hat{i} + 88\hat{j} - 40\hat{k})$$

$$D. a = \frac{1}{105} (41\hat{i} - 88\hat{j} - 40\hat{k})$$

**Answer: D**



**Watch Video Solution**

**64.** Find the discriminant of the quadratic equation  $x^2 + 5x + 4 = 0$  and also find the nature of its roots.



**Watch Video Solution**

**65.** Given three vectors

$$\vec{a} = 6\hat{i} - 3\hat{j}, \vec{b} = 2\hat{i} - 6\hat{j} \text{ and } \vec{c} = -2\hat{i} + 21\hat{j} \quad \text{such that}$$

$\vec{a} = \vec{a} + \vec{b} + \vec{c}$  Then the resolution of the vector  $\vec{a}$  into components with respect to  $\vec{a}$  and  $\vec{b}$  is given by a.  $3\vec{a} - 2\vec{b}$  b.  $3\vec{b} - 2\vec{a}$  c.  $2\vec{a} - 3\vec{b}$  d.  $\vec{a} - 2\vec{b}$

**A. 3a-2b**

B.  $3b-2a$

C.  $2a-3b$

D.  $a-2b$

**Answer: C**



**Watch Video Solution**

66. 'I' is the incentre of triangle  $ABC$  whose corresponding sides are  $a, b, c$ , respectively.  $a\vec{IA} + b\vec{IB} + c\vec{IC}$  is always equal to a.  $\vec{0}$  b.

$(a + b + c)\vec{BC}$  c.  $(\vec{a} + \vec{b} + \vec{c})\vec{AC}$  d.  $(a + b + c)\vec{AB}$

A. 0

B.  $(a+b+c)BC$

C.  $(a+b+c)AC$

D.  $(a+b+c)AB$

**Answer: A**



 [Watch Video Solution](#)

67. If  $\vec{x}$  and  $\vec{y}$  are two non-collinear vectors and ABC is a triangle with side lengths a, b and c satisfying  $(20a - 15b)\vec{x} + (15b - 12c)\vec{y} + (12c - 20a)(\vec{x} \times \vec{y}) = \vec{0}$ , then triangle ABC is

- A. an acute angled triangle
- B. an obtuse angled triangle
- C. a right angled triangle
- D. a scalane triangle

**Answer: C**

 [Watch Video Solution](#)

68. If  $\vec{x}$  and  $\vec{y}$  are two non-collinear vectors and a, b, and c represent the sides of a  $\triangle ABC$  satisfying

$(a - b)\vec{x} + (b - c)\vec{y} + (c - a)(\vec{x} \times \vec{y}) = 0$ , then  $ABC$  is (where  $\vec{x} \times \vec{y}$  is perpendicular to the plane of  $x$  and  $y$ )

- A. an acute angled triangle
- B. an obtuse angled triangle
- C. a right angled triangle
- D. a scalene triangle

**Answer: A**



[Watch Video Solution](#)

69. If the resultant of two forces is equal in magnitude to one of the components and perpendicular to it direction, find the other components using the vector method.

- A.  $P\sqrt{2}$
- B.  $P$
- C.  $P\sqrt{3}$

D. none of these

**Answer: A**



**Watch Video Solution**

70. If  $\vec{b}$  is a vector whose initial point divides the join of  $5\hat{i}$  and  $5\hat{j}$  in the ratio  $k:1$  and whose terminal point is the origin and  $|\vec{b}| \leq \sqrt{37}$ , then,  $k$  lies in the interval

a.  $[-6, -1/6]$

b.  $(-\infty, -6] \cup [-1/6, \infty)$

c.  $[0, 6]$

d. none of these



**Watch Video Solution**

71. find the term independent of  $x$  in the expansion of  $\left(2x - \frac{1}{x}\right)^2$  ?



**Watch Video Solution**

72. If  $\vec{a}$  and  $\vec{b}$  are two unit vectors and  $\theta$  is the angle between them, then the unit vector along the angular bisector of  $\vec{a}$  and  $\vec{b}$  will be given by

A.  $\frac{a - b}{2 \cos(\theta/2)}$

B.  $\frac{a + b}{2 \cos(\theta/2)}$

C.  $\frac{a - b}{\cos(\theta/2)}$

D. none of these

**Answer: B**



**Watch Video Solution**

73. A, B, C and D have position vectors  $\vec{a}$ ,  $\vec{b}$ ,  $\vec{c}$  and  $\vec{d}$ , respectively, such that  $\vec{a} - \vec{b} = 2(\vec{d} - \vec{c})$ . Then

A. AB and CD bisect each other

B. BD and AC bisect each other

C. AB and CD trisect each other

D. BD and AC trisect each other

**Answer: D**



[Watch Video Solution](#)

74. if  $\alpha$  and  $\beta$  are the root of the quadratic polynomial

$f(x) = x^2 - 5x + 6$ , find the value of  $(\alpha^2\beta + \beta^2\alpha)$

A. 20

B. 30

C. 50

D. none of these

**Answer: B**



[Watch Video Solution](#)

75. If  $a + b + c = \alpha d$ ,  $b + c + d = \beta a$  and  $a, b, c$  are non-coplanar, then the sum of  $a + b + c + d =$

- A. 0
- B.  $\alpha a$
- C.  $\beta b$
- D.  $(\alpha + \beta)c$

**Answer: A**

 [Watch Video Solution](#)

76. The position vectors of the points P and Q with respect to the origin O are  $\vec{a} = \hat{i} + 3\hat{j} - 2\hat{k}$  and  $\vec{b} = 3\hat{i} - \hat{j} - 2\hat{k}$ , respectively. If M is a point on PQ, such that OM is the bisector of POQ, then  $\vec{OM}$  is

- A.  $2(\hat{i} - \hat{j} + \hat{k})$

B.  $2\hat{i} + \hat{j} - 2\hat{k}$

C.  $2(-\hat{i} + \hat{j} - \hat{k})$

D.  $2(\hat{i} + \hat{j} + \hat{k})$

**Answer: B**



**Watch Video Solution**

77.  $ABCD$  is a quadrilateral.  $E$  is the point of intersection of the line joining the midpoints of the opposite sides. If  $O$  is any point and  $\vec{OA} + \vec{OB} + \vec{OC} + \vec{OD} = x\vec{OE}$ , then  $x$  is equal to a. 3 b. 9 c. 7 d. 4

A. 3

B. 9

C. 7

D. 4

**Answer: D**

[Watch Video Solution](#)

78. In the  $\triangle OAB$ , M is the midpoint of AB, C is a point on OM, such that  $2OC = CM$ . X is a point on the side OB such that  $OX = 2XB$ . The line XC is produced to meet OA in Y. Then  $\frac{OY}{YA} =$

A.  $\frac{1}{3}$

B.  $\frac{2}{7}$

C.  $\frac{3}{2}$

D.  $\frac{2}{5}$

**Answer: B**

[Watch Video Solution](#)

79. Evaluate  $\int \left( \frac{\ln x}{x} \right) dx$

A.  $\ln x + c$



B.  $\frac{1}{2}\ln^2 x + c$

C.  $\ln^2 x + c$

D. none of these

**Answer: A**



**Watch Video Solution**

**80.** Find the value of  $\lambda$  so that the points P, Q, R and S on the sides OA, OB, OC and AB, respectively, of a regular tetrahedron OABC are coplanar.

It is given that  $\frac{OP}{OA} = \frac{1}{3}$ ,  $\frac{OQ}{OB} = \frac{1}{2}$ ,  $\frac{OR}{OC} = \frac{1}{3}$  and  $\frac{OS}{AB} = \lambda$ .

A.  $\lambda = \frac{1}{2}$

B.  $\lambda = -1$

C.  $\lambda = 0$

D. fo no value of  $\lambda$

**Answer: B**

[Watch Video Solution](#)

81. OABCDE is a regular hexagon of side 2 units in the XY-plane in the first quadrant. O being the origin and OA taken along the x-axis. A point P is taken on a line parallel to the z-axis through the centre of the hexagon at a distance of 3 units from O in the positive Z direction. Then find vector  $\overrightarrow{AP}$ .

A.  $-\hat{i} + 3\hat{j} + \sqrt{5}\hat{k}$

B.  $\hat{i} - \sqrt{3}\hat{j} + 5\hat{k}$

C.  $-\hat{i} + \sqrt{3}\hat{j} + \sqrt{5}\hat{k}$

D.  $\hat{i} + \sqrt{3}\hat{j} + \sqrt{5}\hat{k}$

**Answer: C**

[Watch Video Solution](#)

1. Find  $\frac{dy}{dx}$  if  $y = \frac{1}{2} - x^4$

 [Watch Video Solution](#)

### Exercise More Than One Correct Option Type Questions

1. If the vectors  $\hat{i} - \hat{j}$ ,  $\hat{j} + \hat{k}$  and  $\vec{a}$  form a triangle then  $\vec{a}$  may be

A.  $-\hat{i} - \hat{k}$

B.  $\hat{i} - 2\hat{j} - \hat{k}$

C.  $2\hat{j} + \hat{j} + \hat{k}$

D.  $\hat{i} + \hat{k}$

**Answer: A::B::D**

 [Watch Video Solution](#)

2. If the resultant of three forces  $\vec{F}_1 = p\hat{i} + 3\hat{j} - \hat{k}$ ,  $\vec{F}_2 = 6\hat{i} - \hat{k}$  and  $\vec{F}_3 = -5\hat{i} + \hat{j} + 2\hat{k}$  acting on a particle has a magnitude equal to 5 units, then the value of  $p$  is

- A. -6
- B. -4
- C. 2
- D. 4

**Answer: B::C**



[Watch Video Solution](#)

3. Let ABC be a triangle, the position vectors of whose vertices are  $7\hat{j} + 10\hat{k}$ ,  $-\hat{i} + 6\hat{j} + 6\hat{k}$  and  $-4\hat{i} + 9\hat{j} + 6\hat{k}$ . Then  $\Delta ABC$  is

- A. isosceles
- B. equilateral

C. right angled

D. none of these

**Answer: A::C**



**Watch Video Solution**

4. The sides of a parallelogram are  $2\hat{i} + 4\hat{j} - 5\hat{k}$  and  $\hat{i} + 2\hat{j} + 3\hat{k}$ . The unit vector parallel to one of the diagonals is

A.  $\frac{1}{7} (3\hat{i} + 6\hat{j} - 2\hat{k})$

B.  $\frac{1}{7} (3\hat{i} - 6\hat{j} - 2\hat{k})$

C.  $\frac{1}{\sqrt{69}} (\hat{i} + 2\hat{j} + 8\hat{k})$

D.  $\frac{1}{\sqrt{69}} (-\hat{i} - 2\hat{j} + 8\hat{k})$

**Answer: A::D**



**Watch Video Solution**

5. If  $A(-4, 0, 3)$  and  $B(14, 2, -5)$ , then which one of the following points lie on the bisector of the angle between  $\vec{OA}$  and  $\vec{OB}$  (O is the origin of reference) ? (A) (2,2,4) (B) (2,11,5) (C) (2,11,5) (D) (1,1,2)

A. (2,2,4)

B. (2,11,5)

C. (-3,-3,-6)

D. (1,1,2)

**Answer: A::C::D**



**Watch Video Solution**

6. If points  $\hat{i} + \hat{j}$ ,  $\hat{i} - \hat{j}$  and  $p\hat{i} + q\hat{j} + r\hat{k}$  are collinear, then

A.  $p=1$

B.  $r=0$

C.  $q \in R$

D.  $q \neq 1$

**Answer: A::B::D**



**Watch Video Solution**

7. If  $\vec{a}$ ,  $\vec{b}$  and  $\vec{c}$  are non-coplanar vectors and  $\lambda$  is a real number, then the vectors  $\vec{a} + 2\vec{b} + 3\vec{c}$ ,  $\lambda\vec{b} + \mu\vec{c}$  and  $(2\lambda - 1)\vec{c}$  are coplanar when

A.  $\mu \in R$

B.  $\lambda = \frac{1}{2}$

C.  $\lambda = 0$

D. no value of  $\lambda$

**Answer: A::B::C::D**



**Watch Video Solution**

## Exercise Statement I And II Type Questions

1. Statement 1 : In  $\Delta ABC$ ,  $\vec{AB} + \vec{BC} + \vec{CA} = 0$

Statement 2 : If  $\vec{OA} = \vec{a}$ ,  $\vec{OB} = \vec{b}$ , then  $\vec{AB} = \vec{a} + \vec{b}$

- A. Both Statement I and Statement II are correct and statement II is the correct explanation of statement I
- B. Both statement I and statement II are correct but statement II is not the correct explanation of statement I
- C. Statement I is correct but statement II is incorrect
- D. Statement II is correct but statement I is incorrect

**Answer: C**



**Watch Video Solution**

2. Statement I:  $a = \hat{i} + p\hat{j} + 2\hat{k}$  and  $b = 2\hat{i} + 3\hat{j} + q\hat{k}$  are parallel vectors, iff  $p = \frac{3}{2}$  and  $q = 4$ .



Statement II:  $a = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$  and  $b = b_1\hat{i} + b_2\hat{j} + b_3\hat{k}$  are parallel

$$\frac{a_1}{b_1} = \frac{a_2}{b_2} = \frac{a_3}{b_3}.$$

- A. Both Statement I and Statement II are correct and statement II is the correct explanation of statement I
- B. Both statement I and statement II are correct but statement II is not the correct explanation of statement I
- C. Statement I is correct but statement II is incorrect
- D. Statement II is correct but statement I is incorrect

**Answer: A**



[Watch Video Solution](#)

3. Statement 1: if three points  $P, Q$  and  $R$  have position vectors  $\vec{a}, \vec{b}$ , and  $\vec{c}$ , respectively, and  $2\vec{a} + 3\vec{b} - 5\vec{c} = 0$ , then the points  $P, Q$ , and  $R$  must be collinear. Statement 2: If for three points  $A, B$ , and  $C$ ,  $\vec{AB} = \lambda\vec{AC}$ , then points  $A, B$ , and  $C$  must be collinear.

- A. Both Statement I and Statement II are correct and statement II is the correct explanation of statement I
- B. Both statement I and statement II are correct but statement II is not the correct explanation of statement I
- C. Statement I is correct but statement II is incorrect
- D. Statement II is correct but statement I is incorrect


**Answer: A**



[Watch Video Solution](#)

## Exercise Passage Based Questions

1. Let OABCD be a pentagon in which the sides OA and CB are parallel and the sides OD and AB are parallel. Also,  $OA:CB=2:1$  and  $OD:AB=1:3$ .



Q. The ratio  $\frac{AX}{XD}$  is

A.  $3/4$

B.  $1/3$

C.  $2/5$

D.  $1/2$

Answer: C



Watch Video Solution

2. If  $x = \cos(2t)$  and  $y = \sin^2 t$  then what is  $\frac{d^2y}{dx^2}$



[Watch Video Solution](#)

3. If ABCDEF is regular hexagon, then  $AD+EB+FC$  is

A. (a)  $2AB$

B. (b)  $3AB$

C. (c)  $4AB$

D. (d) none of these

**Answer: C**



[Watch Video Solution](#)

4. Consider the regular hexagon ABCDEF with centre at O (origin).

Q. Five forces  $AB, AC, AD, AE, AF$  act at the vertex A of a regular hexagon ABCDEF. Then, their resultant is

A.  $3AO$

B. 2AO

C. 4AO

D. 6AO

**Answer: D**



[Watch Video Solution](#)

5. Three points A, B, and C have position vectors  $-2\vec{a} + 3\vec{b} + 5\vec{c}$ ,  $\vec{a} + 2\vec{b} + 3\vec{c}$  and  $7\vec{a} - \vec{c}$  with reference to an origin O. Answer the following questions?

Which of the following is true?

A.  $AC=2AB$

B.  $AC=-3AB$

C.  $AC=3AB$

D. none of these

**Answer: C**



**Watch Video Solution**

6. Three points A, B and C have position vectors  $-2a + 3b + 5c$ ,  $a + 2b + 3c$  and  $7a - c$  with reference to an origin O. answer the following questions.

Q. Which of the following is true?

A.  $2OA - 3OB + OC = 0$

B.  $2OA + 7OB + 9OC = 0$

C.  $OA + OB + OC = 0$

D. none of these

**Answer: A**



**Watch Video Solution**

7. Three points A, B, and C have position vectors  $-2\vec{a} + 3\vec{b} + 5\vec{c}$ ,  $\vec{a} + 2\vec{b} + 3\vec{c}$  and  $7\vec{a} - \vec{c}$  with reference to an origin O. Answer the following questions?

B divided AC in ratio

A. 2:1

B. 2:3

C. 2: -3

D. 1:2

**Answer: D**



[Watch Video Solution](#)

8. If two vectors OA and OB are there, then their resultant  $OA+OB$  can be found by completing the parallelogram OACB and  $OC=OA+OB$ . Also, if  $|OA|=|OB|$ , then the resultant will bisect the angle between them.

Q. A vector C directed along internal bisector of angle between vectors

$A = 7\hat{i} - 4\hat{j} - 4\hat{k}$  and  $B = -2\hat{i} - \hat{j} + 2\hat{k}$  with  $|C| = 5\sqrt{6}$  is

a.  $\frac{5}{3}(\hat{i} - \hat{j} + \hat{k})$

b.  $\frac{5}{3}(\hat{i} - 7\hat{j} + 2\hat{k})$

c.  $\frac{5}{3}(5\hat{i} + 5\hat{j} + 2\hat{k})$

d.  $\frac{5}{3}(-5\hat{i} + 5\hat{j} + 3\hat{k})$

A.  $\frac{5}{3}(\hat{i} - \hat{j} + \hat{k})$

B.  $\frac{5}{3}(\hat{i} - 7\hat{j} + 2\hat{k})$

C.  $\frac{5}{3}(5\hat{i} + 5\hat{j} + 2\hat{k})$

D.  $\frac{5}{3}(-5\hat{i} + 5\hat{j} + 3\hat{k})$

**Answer: B**



[Watch Video Solution](#)

9. Find  $\frac{dy}{dx}$  if  $2x - 3y = \sin x$



[Watch Video Solution](#)



10. Solve  $\int \frac{2x}{1+x^2} dx$

 Watch Video Solution

11. If  $z = 10$ , find the value of  $z^3 - 3(z - 10)$ .

 Watch Video Solution

### Exercise Matching Type Questions

1. a and b form the consecutive sides of a regular hexagon ABCDEF.

Column I	Column II
a. If $\mathbf{CD} = x\mathbf{a} + y\mathbf{b}$ , then	p. $x = -2$
b. If $\mathbf{CE} = x\mathbf{a} + y\mathbf{b}$ , then	q. $x = -1$
c. If $\mathbf{AE} = x\mathbf{a} + y\mathbf{b}$ , then	r. $y = 1$
d. If $\mathbf{AD} = -x\mathbf{b}$ , then	s. $y = 2$

 Watch Video Solution

## Exercise Single Integer Answer Type Questions

1. If the resultant of three forces  $F_1 = p\hat{i} + 3\hat{j} - \hat{k}$ ,  $F_2 = 6\hat{i} - \hat{k}$  and  $F_3 = -5\hat{i} + \hat{j} + 2\hat{k}$  acting on a particle has a magnitude equal to 5 units, then the value of  $p$  is

 [Watch Video Solution](#)

2. If ABCD is parallelogram,  $AB = 2\hat{i} + 4\hat{j} - 5\hat{k}$  and  $AD = \hat{i} + 2\hat{j} + 3\hat{k}$ , then the unit vectors in the direction of BD is

 [Watch Video Solution](#)

3. If vectors  $\vec{a} = \hat{i} + 2\hat{j} - \hat{k}$ ,  $\vec{b} = 2\hat{i} - \hat{j} + \hat{k}$  and  $\vec{c} = \lambda\hat{i} + \hat{j} + 2\hat{k}$  are coplanar, then find the value of  $(\lambda - 4)$ .

 [Watch Video Solution](#)

4. If determinant of  $A = 5$  and  $A$  is a square matrix of order 3 then find the determinant of  $\text{adj}(A)$

 [Watch Video Solution](#)

5. Let  $p$  be the position vector of orthocentre and  $g$  is the position vector of the centroid of  $\Delta ABC$ , where circumcentre is the origin. If  $p = kg$ , then the value of  $k$  is

 [Watch Video Solution](#)

6. In a  $\Delta ABC$ , a line is drawn passing through centroid dividing  $AB$  internally in ratio 2:1 and  $AC$  in  $\lambda : 1$  (internally). The value of  $\lambda$  is

 [Watch Video Solution](#)

7. The vector  $\vec{a}$  has the components  $2p$  and  $1$  w.r.t. a rectangular Cartesian system. This system is rotated through a certain angle about the origin in the counterclockwise sense. If, with respect to a new system,  $\vec{a}$  has components  $(p + 1)$  and  $1$ , then  $p$  is equal to

 [Watch Video Solution](#)

### Exercise Subjective Type Questions

1. A vector  $a$  has components  $a_1, a_2, a_3$  in a right handed rectangular cartesian coordinate system  $OXYZ$  the coordinate axis is rotated about  $z$  axis through an angle  $\frac{\pi}{2}$ . The components of  $a$  in the new system

 [Watch Video Solution](#)

2. Find the magnitude and direction of  $r_1 - r_2$  when  $|r_1| = 5$  and points North-East while  $|r_2| = 5$  but points North-West.

 [Watch Video Solution](#)

 Watch Video Solution

3. Let  $OACB$  be a parallelogram with  $O$  at the origin and  $OC$  a diagonal. Let  $D$  be the midpoint of  $OA$ . using vector methods prove that  $BD$  and  $CO$  intersect in the same ratio. Determine this ratio.

 Watch Video Solution

4.  $\triangle ABC$  is a triangle with the point  $P$  on side  $BC$  such that  $3BP=2PC$ , the point  $Q$  is on the line  $CA$  such that  $4CQ=QA$ . Find the ratio in which the line joining the common point  $R$  of  $AP$  and  $BQ$  and the point  $S$  divides  $AB$ .

 Watch Video Solution

5. In  $\triangle ABC$  internal angle bisector  $AI, BI$  and  $CI$  are produced to meet opposite sides in  $A', B', C'$  respectively. Prove that the maximum value of  $\frac{AI \times BI \times CI}{AA' \times BB' \times CC'}$  is  $\frac{8}{27}$

 Watch Video Solution

6. IF  $a_1, a_2, a_3, \dots, a_{10}$  be in AP and  $h_1, h_2, h_3, \dots, h_{10}$  be in HP. If  $a_1 = h_1 = 2$  and  $a_{10} = h_{10} = 3$ , then find value of  $a_4 h_7$ .



[Watch Video Solution](#)

7. Let OABCD be a pentagon in which the sides OA and CB are parallel and the sides OD and AB are parallel as shown in figure. Also,  $OA:CB=2:1$  and  $OD:AB=1:3$ . if the diagonals OC and AD meet at x, find  $OX:XC$ .



[Watch Video Solution](#)

8. P and Q have position vectors  $a$  and  $b$  relative to the origin O and X,Y divide PQ internally and externally respectively in the ratio 2:1, vector XY is  $\lambda a + \mu b$ , then the value of  $|\lambda + \mu|$  is



[Watch Video Solution](#)

## Exercise Questions Asked In Previous 13 Years Exam

1. If vectors  $\overrightarrow{AB} = -3\hat{i} + 4\hat{k}$  and  $\overrightarrow{AC} = 5\hat{i} - 2\hat{j} + 4\hat{k}$  are the sides of a  $\Delta ABC$ , then the length of the median through A is

A.  $\sqrt{18}$

B.  $\sqrt{72}$

C.  $\sqrt{33}$

D.  $\sqrt{45}$

**Answer: C**



[Watch Video Solution](#)

2. Let  $a, b$  and  $c$  be three non-zero vectors which are pairwise non-collinear. If  $a+3b$  is collinear with  $c$  and  $b+2c$  is collinear with  $a$ , then  $a+3b+6c$  is

A.  $a+c$

B. a

C. c

D. 0

**Answer: D**



[Watch Video Solution](#)

3. The non-zero vectors  $a, b$  and  $c$  are related by  $a=8b$  and  $c=-7b$  angle between  $a$  and  $c$  is

A.  $\pi$

B. 0

C.  $\frac{\pi}{4}$

D.  $\frac{\pi}{2}$

**Answer: A**



[Watch Video Solution](#)



4. If C is the middle point of AB and P is any point outside AB, then

A.  $PA+PB+PC=0$

B.  $PA+PB+2PC=0$

C.  $PA+PB=PC$

D.  $PA+PB=2PC$

**Answer: D**



[Watch Video Solution](#)

5. Let  $a, b$  and  $c$  be three non-zero vectors such that no two of these are collinear. If the vector  $a+2b$  is collinear with  $c$  and  $b+3c$  is collinear with  $a$  ( $\lambda$  being some non-zero scalar), then  $a + 2b + 6c$  is equal to

A.  $\lambda a$

B.  $\lambda b$

C.  $\lambda c$

D. 0

**Answer: D**



[Watch Video Solution](#)

6. If  $\vec{a}$ ,  $\vec{b}$  and  $\vec{c}$  are non-coplanar vectors and  $\lambda$  is a real number, then the vectors  $\vec{a} + 2\vec{b} + 3\vec{c}$ ,  $\lambda\vec{b} + \mu\vec{c}$  and  $(2\lambda - 1)\vec{c}$  are coplanar when

A. all value of  $\lambda$

B. all except one value of  $\lambda$

C. all except two value of  $\lambda$

D. no value of  $\lambda$

**Answer: C**



[Watch Video Solution](#)

7. Area of a rectangle having vertices A, B, C and D with position vectors :

$$-\hat{i} + \left(\frac{1}{2}\right)\hat{j} + 4\hat{k}, \hat{i} + \left(\frac{1}{2}\right)\hat{j} + 4\hat{k}, \hat{i} - \left(\frac{1}{2}\right)\hat{j} + 4\hat{k} \quad \text{and}$$

$$-\hat{i} - \left(\frac{1}{2}\right)\hat{j} + 4\hat{k}, \text{ respectively is:}$$

A. square

B. rhombus

C. rectangle

D. none of these

**Answer: D**



[Watch Video Solution](#)

8. If a, b, and c are all different and if

$$\begin{vmatrix} a & a^2 & 1 + a^3 \\ b & b^2 & 1 + b^3 \\ c & c^2 & 1 + c^3 \end{vmatrix} = 0 \text{ Prove that } abc = -1.$$

A. 2

B.  $-1$

C.  $1$

D.  $0$

**Answer: B**



**Watch Video Solution**

9. The vector  $\hat{i} + x\hat{j} + 3\hat{k}$  is rotated through an angle  $\theta$  and doubled in magnitude, then it becomes  $4\hat{i} + (4x - 2)\hat{j} + 2\hat{k}$ . Then value of  $x$  are  $-\frac{2}{3}$  (b)  $\frac{1}{3}$  (c)  $\frac{2}{3}$  (d)  $2$

A.  $\left\{ -\frac{2}{3}, 2 \right\}$

B.  $\left( \frac{1}{3}, 2 \right)$

C.  $\left\{ \frac{2}{3}, 0 \right\}$

D.  $\{2, 7\}$

**Answer: A**



**Watch Video Solution**