



MATHS

BOOKS - CENGAGE PUBLICATION

MONOTONICITY AND MAXIMA MINIMA OF FUNCTIONS

Illustration

1. Prove that the function $f(x) = (\log)_e(x^2 + 1) - e^{-x} + 1$ is strictly increasing $\forall x \in R$.



Watch Video Solution

2. Find the least value of k for which the function $x^2 + kx + 1$ is an increasing function in the interval $(1,2)$



Watch Video Solution

3. Find the range of values of a if $f(x) = 2e^x - ae^{-x} + (2a + 1)x - 3$ is monotonically increasing for all values of x .



[Watch Video Solution](#)

4. If $f(x)$ and $g(x)$ are differentiable and increasing functions then which of the following functions always increases?

A. $f(x)+g(x)$

B. $f(x)*g(x)$

C. $f(x)-g(x)$

D. $f(x)/g(x)$

Answer: A



[View Text Solution](#)

5.


Let

$$g(x) = (f(x))^3 - 3(f(x))^2 + 4f(x) + 5x + 3 \sin x + 4 \cos x \quad \forall x \in \mathbb{R}.$$

Then prove that g is increasing whenever f is increasing.


[Watch Video Solution](#)

6. For each of the following graphs, comment whether $f(x)$ is increasing or decreasing or neither increasing nor decreasing at $x = a$.


[View Text Solution](#)

7. Find the complete set of values of α for which the function

$$f(x) = \begin{cases} x + 1, & x < 1 \\ \alpha, & x = 1 \\ x^2 - x + 3, & x > 1 \end{cases}$$

is strictly increasing for all x .

$$x^2 - x + 3, \quad x > 1$$



[View Text Solution](#)

8. Prove that function $f(x) = \begin{cases} -2x^3 + 3x^2 - 6x + 5, & x < 0 \\ -x^2 - x + 1, & x \geq 0 \end{cases}$ is decreasing for all x .



[View Text Solution](#)

9. If $f \circ g \circ h(x)$ is an increasing function, then which of the following is not possible? $f(x), g(x),$ and $h(x)$ are increasing $f(x)$ and $g(x)$ are decreasing and $h(x)$ is increasing $f(x), g(x),$ and $h(x)$ are decreasing



[Watch Video Solution](#)

10. Let $f(x)$ and $g(x)$ be two continuous functions defined from \overrightarrow{RR} , such that $f(x_1) > f(x_2)$ and $g(x_1) < g(x_2)$ for all $x_1 > x_2$. Then what is the solution set of $f(g(\alpha^2 - 2\alpha)) > f(g(3\alpha - 4))$

 [Watch Video Solution](#)

11. Let $f: [0, \infty) \rightarrow [0, \infty)$ and $g: [0, \infty) \rightarrow [0, \infty)$ be non-increasing and non-decreasing functions, respectively, and $h(x) = g(f(x))$. If f and g are differentiable for all points in their respective domains and $h(0) = 0$, then show $h(x)$ is always, identically zero.

 [Watch Video Solution](#)

12. Find the values of p if $f(x) = \cos x - 2px$ is invertible.

 [View Text Solution](#)

13. Find the critical points(s) and stationary points (s) of the function

$$f(x) = (x - 2)^{2/3}(2x + 1)$$



[View Text Solution](#)

14. Separate the intervals of monotonicity of the following function:

(i) $f(x) = 3x^4 - 8x^3 - 6x^2 + 24x + 7$

(ii) $F(x) = -\sin^3 x + 3\sin^2 x + 5, x \in (-\pi/2, \pi/2)$

(iii) $f(x) = (2^x - 1)(2^x - 2)^2$



[Watch Video Solution](#)

15. अंतराल ज्ञात कीजिए जिन पर

$$f(x) = \frac{4 \sin x - 2x - x \cos x}{2 + \cos x}$$

से प्रदत्त फलन f(i) निरंतर वर्धमान (ii) निरंतर ह्रासमान है |



[View Text Solution](#)

16. Find the interval of monotonicity of the function $f(x) = |x - 1|x^2$.



Watch Video Solution

17. Find the intervals of decrease and increase for the function

$$f(x) = \cos\left(\frac{\pi}{x}\right)$$



Watch Video Solution

18. Let $g(x) = f(x) + f(1 - x)$ and $f''(x) > 0 \forall x \in (0, 1)$. Find the intervals of increase and decrease of $g(x)$.



Watch Video Solution

19. Find the range of the function $f(x) = x \sin x - \frac{1}{2} \sin^2 x$ for $x \in \left(0, \frac{\pi}{2}\right)$



View Text Solution

20. Find the range of $f(x) = \frac{1}{\pi} \sin^{-1} x + \tan^{-1} x + \frac{x+1}{x^2+2x+5}$

 [Watch Video Solution](#)

21. Find the number of roots of the equation

$$\log_e(1+x) - \frac{\tan^{-1} x}{1+x} = 0$$

 [View Text Solution](#)

22. Find the number of roots of the function

$$f(x) = \frac{1}{(x+1)^3} - 3x + \sin x$$

 [View Text Solution](#)

23. Prove that $\log_e(1+x) < xf$ or $x > 0$

 [Watch Video Solution](#)

24. Let f and g be differentiable on \mathbb{R} and suppose $f(0) = g(0)$ and $f'(x) \leq g'(x)$ for all $x \geq 0$. Then show that $f(x) \leq g(x)$ for all $x \geq 0$.

 [Watch Video Solution](#)

25. Show that $1 + \xi n \left(x + \sqrt{x^2 + 1} \right) \geq \sqrt{1 + x^2}$ for all $x \geq 0$.

 [Watch Video Solution](#)

26. If $a, b > 0$ and

 [Watch Video Solution](#)

27. Prove that $|\cos \alpha - \cos \beta| \leq |\alpha - \beta|$

 [Watch Video Solution](#)

28. If $P(1) = 0$ and $\frac{dP(x)}{dx} > P(x)$ for all $x > 1$. Prove that $P(x) > 0$ for all $x > 1$

 [Watch Video Solution](#)

29. For $0 < x < \frac{\pi}{2}$, prove that $x > \sin x$

 [View Text Solution](#)

30. Prove that $e^x \geq 1 + x$ and hence $e^x + \sqrt{1 + e^{2x}} \geq (1 + x) + \sqrt{2 + 2x + x^2} \forall x \in \mathbb{R}$

 [Watch Video Solution](#)

31. Separate the intervals of concavity of the following functions

(i) $f(x) = \sin^{-1} x$, (ii) $f(x) = x + \sin x$

 [View Text Solution](#)

32. If graph of the function $f(x) = 3x^4 + 2x^3 + ax^2 - x + 2$ is concave upward for all real x , then find values of a ,

 [View Text Solution](#)

33. Prove that for any two numbers x_1 and x_2

$$\frac{2e^{x_1} + e^{x_2}}{3} > e^{\frac{2x_1 + x_2}{3}}$$

 [Watch Video Solution](#)

34. If $A+B+C=\pi$ Prove that in triangle ABC, $\sin A + \sin B + C \leq \frac{3\sqrt{3}}{2}$

 [View Text Solution](#)

35. Prove that $a_1^m + a_2^m + \dots + a_n^m \frac{1}{n} < \frac{a_1 + a_2 + \dots + a_n}{(n)^m}$

If $0 < m < 1$ and $a_i > 0$ for all i .

 [View Text Solution](#)

36. Find the points of inflection for $f(x) = \sin x$ $f(x) = 3x^4 - 4x^3$

$$f(x) = x^{\frac{1}{3}}$$

 [Watch Video Solution](#)

37. Find the coordinates of the point of inflection of the curve $f(x) = e^{-(x^2)}$

 [View Text Solution](#)

38. The function $y = \frac{ax + b}{x - 1}(x - 4)$ has turning point at $P(2, -1)$. Then find the values of a and b .

 [View Text Solution](#)

39. How many critical points function $f(x)=(x-1)|x-3|-4x+12$ has? How many of these are points of local extrema?

 [View Text Solution](#)

40. Discuss the extremum of the function $f(x)=(x-1)(x-2)(x-3)$. How many critical points $f(x)$ has?

 [View Text Solution](#)

41. Find the points of extrema of the function $f(x)=2 \sec x+3 \operatorname{cosec} x$

 [View Text Solution](#)


42. The function $f(x) = (x^2 - 4)^n (x^2 - x + 1)$, $n \in N$, assumes a local minimum value at $x = 2$. Then find the possible values of n

 [View Text Solution](#)

43. Test $f(x)=\{x\}$ for the existence of a local maximum and minimum at $x=1$, where $\{ \cdot \}$ represents the fractional part function.

[View Text Solution](#)

44. Consider the following graphs of the functions. Check each for the extrema at $x=a$



[View Text Solution](#)

45. Let $f(x) = \begin{cases} x^3 + x^2 + 10x, & x < 0 \\ -3 \sin x, & x \geq 0 \end{cases}$. Investigate $x=0$ for local

maxima/minima.



[View Text Solution](#)

46. Let $f(x) = \frac{a}{x} + x^2$. If it has a maximum at $x = -3$, then find the value of a .



[Watch Video Solution](#)

47. Discuss the extremum of $f(x) = \sin x(1 + \cos x)$, $x \in \left(0, \frac{\pi}{2}\right)$



[Watch Video Solution](#)

48. Find the points at which the function f given by $f(x) = (x - 2)^4(x + 1)^3$ has local maxima and local minima and points of inflexion.



[Watch Video Solution](#)

49. Discuss the extremum of $f(x) = 40(3x^4 + 8x^3 - 18x^2 + 60)$.



[View Text Solution](#)

50. Let $f(x) = \sin^3 x + \lambda \sin^2 x$, $-\frac{\pi}{2} < x < \frac{\pi}{2}$ Find the intervals in which λ should lie in order that $f(x)$ has exactly one minimum and exactly one maximum.



[View Text Solution](#)

51. Let $f(x) = \sin^3 x + \lambda \sin^2 x$, $-\frac{\pi}{2} < x < \frac{\pi}{2}$ Find the intervals in which λ should lie in order that $f(x)$ has exactly one minimum and exactly one maximum.



[View Text Solution](#)

52. Discuss the extrema of $f(x) = \frac{x}{1 + x \tan x}$, $x \in \left(0, \frac{\pi}{2}\right)$

 [Watch Video Solution](#)

53. Determine which is bigger, $\frac{1}{(\pi)^{\frac{1}{e}}}$ or $\frac{1}{(e)^{\frac{1}{\pi}}}$?

 [View Text Solution](#)

54. Discuss the extrema of the following functions

(i) $f(x) = |x|$, (ii) $f(x) = e^{-|x|}$, (iii) $f(x) = x^{2/3}$

 [View Text Solution](#)

55. If $f(x) = \begin{cases} x^2, & x \leq 0 \\ x^3, & x > 0 \end{cases}$ Investigate the functions at x for maxima/minima

 [View Text Solution](#)

56. Discuss the extremum of $f(x) = 2x + 3x^{\frac{2}{3}}$



Watch Video Solution

57. $f(x) = \begin{cases} \frac{\cos(\pi x)}{2}, & x > 0 \\ x + a, & x \leq 0 \end{cases}$ Find the values of a if $x = 0$ is a point of maxima.



Watch Video Solution

58. about to only mathematics



View Text Solution

59. $f(x) = |ax - b| + c|x| \forall x \in (-\infty, \infty)$, where $a > 0, b > 0, c > 0$. Find the condition if $f(x)$ attains the minimum value only at one point.



Watch Video Solution

60. Let $f(x) = 2x^3 - 9x^2 + 12x + 6$. Discuss the global maxima and minima of $f(x) \in [0, 2]$ and $(1, 3)$ and, hence, find the range of $f(x)$ for corresponding intervals.

 [Watch Video Solution](#)

61. Find the absolute maximum and absolute minimum values of $f(x) = 3x^4 - 8x^3 + 12x^2 - 48x + 25$ in $[0, 3]$

 [Watch Video Solution](#)

62. Find the range of the function $f(x) = 4 \cos^3 x - 8 \cos(2)x + 1$.

 [View Text Solution](#)

63. Find the range of the function $f(x) = 2\sqrt{x-2} + \sqrt{4-x}$

 [Watch Video Solution](#)

64. A function $y = f(x)$ is represented parametrically as following

$$x = \phi(t) = t^5 - 20t + 7$$

$$y = \psi(t) = 4t^3 - 3t^2 - 18t + 3$$

where t in $[-2, 2]$

Find the intervals of monotonicity and also find the points of extrema. Also find the range of function.



[View Text Solution](#)

65. Find how many roots of the equations $x^4 + 2x^2 - 8x + 3 = 0$.



[Watch Video Solution](#)

66. Draw the graph of $y = \frac{x^2}{\sqrt{x+1}}$



[Watch Video Solution](#)

67. Draw the graph of $y = xe^x$. Find the range of the function. Also find the point of inflection.

 [Watch Video Solution](#)

68. Minimum integral value of k for which the equation $e^x = kx^2$ has exactly three real distinct solution,

 [Watch Video Solution](#)

69. Draw the graph of $f(x) = \log_e(\sqrt{1-x^2})$. Find the range of the function. Also find the values of k if $f(x)$ has two distinct real roots.

 [View Text Solution](#)

70. Draw the graph of $f(x) = \frac{x^2 - 5x + 6}{x^2 - x}$

 [View Text Solution](#)

71. Find the value of a if $x^3 - 3x + a = 0$ has three distinct real roots.

 [Watch Video Solution](#)

72. Prove that there exist exactly two non-similar isosceles triangles ABC such that $\tan A + \tan B + \tan C = 100$.

 [Watch Video Solution](#)

73. If t is a real number satisfying the equation $2t^3 - 9t^2 + 30 - a = 0$, then find the values of the parameter a for which the equation $x + \frac{1}{x} = t$ gives six real and distinct values of x .

 [Watch Video Solution](#)

74. The tangent to the parabola $y = x^2$ has been drawn so that the abscissa x_0 of the point of tangency belongs to the interval $[1,2]$. Find x_0

for which the triangle bounded by the tangent, the axis of ordinates, and the straight line $y = x^2$ has the greatest area.

 [Watch Video Solution](#)

75. Find the point (α, β) on the ellipse $4x^2 + 3y^2 = 12$, in the first quadrant, so that the area enclosed by the lines $y = x$, $y = \beta$, $x = \alpha$, and the x-axis is maximum.

 [Watch Video Solution](#)

76. LL' is the latus rectum of the parabola $y^2 = 4ax$ and PP' is a double ordinate drawn between the vertex and the latus rectum. Show that the area of the trapezium $PP'LL'$ is maximum when the distance PP' from the vertex is $a/9$.

 [Watch Video Solution](#)

77. Find the points on the curve $5x^2 - 8xy + 5y^2 = 4$ whose distance from the origin is maximum or minimum.

 [Watch Video Solution](#)

78. Rectangles are inscribed inside a semi-circle of radius r . Find the rectangle with maximum area.

 [Watch Video Solution](#)

79. A running track of 440 ft is to be laid out enclosing a football field, the shape of which is a rectangle with a semi-circle at each end. If the area of the rectangular portion is to be maximum, then find the length of its sides.

 [Watch Video Solution](#)

80. If the sum of the lengths of the hypotenuse and another side of a right-angled triangle is given, show that the area of the triangle is maximum when the angle between these sides is $\frac{\pi}{3}$.



[View Text Solution](#)

81. The tangent to the parabola $y = x^2$ has been drawn so that the abscissa x_0 of the point of tangency belongs to the interval $[1,2]$. Find x_0 for which the triangle bounded by the tangent, the axis of ordinates, and the straight line $y = x^2$ has the greatest area.



[Watch Video Solution](#)

82. Find the point (α, β) on the ellipse $4x^2 + 3y^2 = 12$, in the first quadrant, so that the area enclosed by the lines $y = x, y = \beta, x = \alpha$, and the x-axis is maximum.



[Watch Video Solution](#)

83. LL' is the latus rectum of the parabola $y^2 = 4ax$ and PP' is a double ordinate drawn between the vertex and the latus rectum. Show that the area of the trapezium $PP'LL'$ is maximum when the distance PP' from the vertex is $a/9$.



[Watch Video Solution](#)

84. Find the points on the curve $5x^2 - 8xy + 5y^2 = 4$ whose distance from the origin is maximum or minimum.



[Watch Video Solution](#)

85. A sheet of area $40m^2$ is used to make an open tank with square base. Find the dimensions of the base such that the volume of this tank is maximum.



[Watch Video Solution](#)

86. A sheet of area $40m^2$ is used to make an open tank with square base. Find the dimensions of the base such that the volume of this tank is maximum.

 [Watch Video Solution](#)

87. The lateral edge of a regular hexagonal pyramid is $1cm$. If the volume is maximum, then find its height.

 [View Text Solution](#)

88. Show that the height of the cylinder of maximum volume that can be inscribed in a sphere of radius R is $\frac{2R}{\sqrt{3}}$.

 [Watch Video Solution](#)

89. Show that height of the cylinder of greatest volume which can be inscribed in a right circular cone of height h and semi vertical angle is

one-third that of the cone and the greatest volume of cylinder is

$$\frac{4}{27}\pi h^3 \tan^2 \alpha.$$

 [Watch Video Solution](#)

Solved Examples

1. Find the possible values of a such that $f(x) = e^{2x} - (a + 1)e^x + 2x$ is monotonically increasing for $x \in \mathbb{R}$.

 [Watch Video Solution](#)

2. Of

$$f(x) = \frac{2}{\sqrt{3}} \tan^{-1} \left(\frac{2x + 1}{\sqrt{3}} \right) - \log(x^2 + x + 1) (\lambda^2 - 5\lambda + 3)x + 10$$

is a decreasing function for all $x \in \mathbb{R}$, find the permissible values of λ .

 [Watch Video Solution](#)

3. Let $f(x) = x^3 + ax^2 + bx + 5 \sin^2 x$ be an increasing function on the set R . Then find the condition on a and b .

 [Watch Video Solution](#)

4. Prove that $\left[\lim_{x \rightarrow 0} \frac{\sin x \cdot \tan x}{x^2} \right] = 1$, where $[.]$ represents greatest integer function.

 [View Text Solution](#)

5. Using the relation $2(1 - \cos x) < x^2$, $x = 0$ or prove that $\sin(\tan x) \geq x$, $\forall x \in \left[0, \frac{\pi}{4}\right]$

 [Watch Video Solution](#)

6. If $f'(x) > 0$, $f(x) > 0 \forall x \in (0, 1)$ and $f(0) = 0$, $f(1) = 1$, then prove that $f(x)f^{-1}(x) < x^2 \forall x \in (0, 1)$

 [View Text Solution](#)

7. Discuss the monotonicity of $Q(x)$, where $Q(x) = 2f\left(\frac{x^2}{2}\right) + f(6 - x^2) \forall x \in R$ It is given that $f^x > 0 \forall x \in R$. Find also the point of maxima and minima of $Q(x)$.

 [View Text Solution](#)

8. Prove that $\left(\tan^{-1}\left(\frac{1}{e}\right)\right)^2 + \frac{2e}{\sqrt{e^2 + 1}} < \left(\tan^{-1} e\right)^2 + \frac{2}{\sqrt{e^2 + 1}}$

 [Watch Video Solution](#)

9. Prove that $\sin^2 \theta \theta \sin(\sin \theta)$ for $\theta > 0$

 [View Text Solution](#)

10. Let $f(x) = 1 + 4x - x^2, \forall x \in R$

$$g(x) = \max \{f(t), x \leq t \leq (x + 1), 0 \leq x < 3\} = \min \{(x + 3), 3 \leq x < 5\}$$

Verify continuity of $g(x)$, for all $x \in [0, 5]$



[Watch Video Solution](#)

11. Show that $5x \leq 8 \cos x - 2 \cos 2x \leq 6x$ for $x \leq x \leq \frac{\pi}{3}$



[View Text Solution](#)

12. Let $f(x), x \geq 0$, be a non-negative continuous function. If

$$f'(x) \cos x \leq f(x) \sin x \forall x \geq 0, \text{ then find } f\left(\frac{5\pi}{3}\right)$$



[Watch Video Solution](#)

13. If $ax^2 + \frac{b}{x} \geq c$ for all positive x where $a > 0$ and $b > 0$, show that

$$27ab^2 \geq 4c^3.$$

 [Watch Video Solution](#)

14. Prove that for $x \in \left[0, \frac{\pi}{2}\right]$, $\sin x + 2x \geq \frac{3x(x+1)}{\pi}$.

 [View Text Solution](#)

15. Consider a curve $C: y = \cos^{-1}(2x - 1)$ and a straight line $L: 2px - 4y + 2\pi - p = 0$. Statement 1: The set of values of p for which the line L intersects the curve at three distinct points is $[-2\pi, -4]$. Statement 2: The line L is always passing through point of inflection of the curve C .

 [Watch Video Solution](#)

16. From a fixed point A on the circumference of a circle of radius r , the perpendicular AY falls on the tangent at P . Find the maximum area of triangle APY .

 [Watch Video Solution](#)

17. P and Q are two points on a circle of centre C and radius α . The angle PCQ being 2θ , find the value of $\sin\theta$ when the radius of the circle inscribed in the triangle CPQ is maximum.

 [Watch Video Solution](#)

18. The lower corner of a leaf in a book is folded over so as to reach the inner edge of the page. Show that the fraction of the width folded over when the area of the folded part is minimum is:

 [Watch Video Solution](#)

19. Let $A(p^2, -p)$, $B(q^2, q)$, $C(r^2, -r)$ be the vertices of triangle ABC . A parallelogram $AFDE$ is drawn with D, E , and F on the line segments BC , CA and AB , respectively. Using calculus, show that the maximum area of such a parallelogram is $\frac{1}{2}(p+q)(q+r)(p-r)$.



[Watch Video Solution](#)

20. A window of perimeter P (including the base of the arch) is in the form of a rectangle surrounded by a semi-circle. The semi-circular portion is fitted with the colored glass while the rectangular part is fitted with the clear glass that transmits three times as much light per square meter as the colored glass does. What is the ratio for the sides of the rectangle so that the window transmits the maximum light?

[Watch Video Solution](#)

Concept Application Exercise 6.1

1. Prove that the following functions are strictly increasing:

$$f(x) = \log(1 + x) - \frac{2x}{2 + x} \text{ for } x > -1$$

[Watch Video Solution](#)

2. Separate the intervals of monotonicity for the following function:

(a) $f(x) = -2x^3 - 9x^2 - 12x + 1$

(b) $f(x) = x^2 e^{-x}$

(c) $f(x) = \sin x + \cos x, x \in (0, 2\pi)$

(d) $f(x) = 3 \cos^4 x + \cos x, x \in (0, 2\pi)$

(e) $f(x) = (\log_e x)^2 + (\log_e x)$

 [View Text Solution](#)

3. Discuss monotonicity of $f(x) = \frac{x}{\sin x}$ and $g(x) = x/(\tan x)$, where

 [Watch Video Solution](#)

4. A function $y = f(x)$ is given by $x = \frac{1}{1+t^2}$ and $y = \frac{1}{t(1+t^2)}$ for all $t > 0$ then f is

 [Watch Video Solution](#)

5. Find the value of a for which the function $(a + 2)x^3 - 3ax^2 + 9ax - 1$ decreases monotonically for all real x .

 [Watch Video Solution](#)

6. Find the value of a in order that $f(x) = \sqrt{3}\sin x - \cos x - 2ax + b$ decreases for all real values of x .

 [Watch Video Solution](#)

7. Discuss the monotonicity of function $f(x) = 2\log|x - 1| - x^2 + 2x + 3$.

 [Watch Video Solution](#)

8. If $f(x) = \sin x + \log_e|\sec x + \tan x| - 2xf$ or $x \in \left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$ then check the monotonicity of $f(x)$

 [View Text Solution](#)

9. Find the interval of the monotonicity of the function $f(x) = \log_e \left(\frac{\log_e x}{x} \right)$

 [View Text Solution](#)

10. Let $g(x) = f(\log x) + f(2 - \log x)$ and $f''(x) < 0 \forall x \in (0, 3)$. Then find the interval in which $g(x)$ increases.

 [Watch Video Solution](#)

Concept Application Exercise 6.2

1. Find the range of $f(x) = \tan^{-1} x - \frac{1}{2} \log_e x \in \left(\frac{1}{\sqrt{3}}, \sqrt{3} \right)$

 [View Text Solution](#)

2. Find the range of $f(x) = \frac{\sin x}{x} + \frac{x}{\tan x} \in \left(0, \frac{\pi}{2}\right)$



[View Text Solution](#)

3. Column I, Column II Range of $f(x) = \sin^{-1} x + \cos^{-1} x + \cot^{-1} \xi s$, p.

$\left[0, \frac{\pi}{2}\right] \cup \left[\frac{\pi}{2}, \pi\right]$ Range of $f(x) = \cot^{-1} x + \tan^{-1} x + \cos ec^{-1} \xi s$, q.

$\left[\frac{\pi}{2}, \frac{3\pi}{2}\right]$ Range of $f(x) = \cot^{-1} x + \tan^{-1} x + \cos^{-1} \xi s$, r. $[0, \pi]$

Range of $f(x) = \sec^{-1} x + \cos ec^{-1} x + \sin^{-1} \xi s$, s. $\left[\frac{3\pi}{4}, \frac{5\pi}{4}\right]$



[Watch Video Solution](#)

4. Find the number of solution of the equation

$$x^3 + 2x + \cos x + \tan x = 0 \in \left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$$



[View Text Solution](#)

5. Show that $\frac{x}{(1+x)} < 1n(1+x)$ for $x > 0$

 [Watch Video Solution](#)

6. For $0 < x \leq \frac{\pi}{2}$, show that $x - \frac{x^3}{6} < \sin(x) < x$.

 [Watch Video Solution](#)

7. Show that $\tan^{-1} x > \frac{x}{1 + \frac{x^2}{3}}$ if $x \in (0, \infty)$.

 [Watch Video Solution](#)

8. Show that $2\sin x + \tan x \geq 3x$, where $0 \leq x < \pi/2$

 [Watch Video Solution](#)

9. Prove that $f(x) = \frac{\sin x}{x}$ is monotonically decreasing in $\left[0, \frac{\pi}{2}\right]$. Hence, prove that $\sin(2x) < 2\sin x$.

 [Watch Video Solution](#)

10. For θ



[View Text Solution](#)

Concept Application Exercise 6.3

1. Show that graph of the function $y = \log_e(x - 2) - \frac{1}{x}$ always concave downwards.



[View Text Solution](#)

2. Separate the interval of concavity of $y = x \log_e x - \frac{x^2}{2} + \frac{1}{2}$



[View Text Solution](#)

3. consider $f(x) = \cos 2x + 2x\lambda^2 + (2\lambda + 1)(\lambda - 1)x^2$, $\lambda \in \mathbb{R}$

if $\alpha \neq \beta$ and $\frac{f(\alpha + \beta)}{2} < \frac{f(\alpha) + f(\beta)}{2}$ for α and β then find the values of λ

 [View Text Solution](#)

4. Find the values of x where function $f(x) = (\sin x + \cos x)(e^x)$ in $(0, 2\pi)$ has point of inflection

 [View Text Solution](#)

5. Prove that $a_1^m + a_2^m + \dots + \frac{a_n^m}{n} > \left(\frac{a_1 + a_2 + \dots + a_n}{n} \right)^m$, if $m < 0$ or $m > 1$ and $a_i > 0 \forall i$

 [View Text Solution](#)

1. Find the least value of $\sec A + \sec B + \sec C$ in an acute angled triangle.

 [Watch Video Solution](#)

2. Find the critical (stationary) points of the function $f(x) = \frac{x^5}{20} - \frac{x^4}{12}$

+5Name these points .Also find the point of inflection

 [View Text Solution](#)

3. The curve $f(x) = \frac{x^2 + ax + b}{x - 10}$ has a stationary point at $(4, 1)$. Find the values of a and b . Also, show that $f(x)$ has point of maxima at this point.

 [Watch Video Solution](#)

4. If the function $f(x) = ax^2 + bx^3$ has maximum value at $x=2$ such that $f(2) = 1$, then find the values of a and b

 [View Text Solution](#)

5. Discuss the extremum of $f(x) = \frac{1}{3} \left(x + \frac{1}{x} \right)$

 [Watch Video Solution](#)

6. Discuss the extremum of $f(x) = 1 + 2 \sin x + 3 \cos^2 x$, $x \leq x \leq \frac{2\pi}{3}$

 [Watch Video Solution](#)

7. Discuss the extremum of $f(x) = \sin x + \frac{1}{2} \sin 2x + \frac{1}{3} \sin 3x$, $0 \leq x \leq \pi$.

 [Watch Video Solution](#)

8. Let $f(x) = -\sin^3 x + 3\sin^2 x + 5\cos x$ on $\left[0, \frac{\pi}{2}\right]$. Find the local maximum and local minimum of $f(x)$.

 [Watch Video Solution](#)

9. Discuss the extremum of $f(\theta) = \sin^p \theta \cos^q \theta$, $p, q > 0$, $0 < \theta < \frac{\pi}{2}$

 [View Text Solution](#)

10. Find the maximum and minimum values of the function $y = (\log)_e (3x^4 - 2x^3 - 6x^2 + 6x + 1) \forall x \in (0, 2)$ Given that $(3x^4 - 2x^3 - 6x^2 + 6x + 1) > 0 \forall x \in (0, 2)$

 [View Text Solution](#)

11. Discuss the extremum of $f(x) = x(x^2 - 4)^{-\frac{1}{3}}$

 [Watch Video Solution](#)

12. Discuss the maxima and minima of the function $f(x) = x^{\frac{2}{3}} - x^{\frac{4}{3}}$.

Draw the graph of $y = f(x)$ and find the range of $f(x)$.

 [Watch Video Solution](#)

13. Discuss the extremum of $f(x) = |x^2 - 2|$, $-1 \leq x$

 [View Text Solution](#)

14. Discuss the extremum of

$$f(x) = \begin{cases} 1 + \sin x, & x < 0 \\ x^2 - x + 1, & x \geq 0 \end{cases} \text{ at } x = 0$$

 [Watch Video Solution](#)

15. Find the minimum value of $|x| + \left|x + \frac{1}{2}\right| + |x - 3| + \left|x - \frac{5}{2}\right|$.

 [View Text Solution](#)

16. Let $f(x)$ be defined as $f(x) = \tan^{-1} \alpha - 5x^2, 0$

 [View Text Solution](#)

17.

Let

$$f(x) = \begin{cases} x^3 - x^2 + 10x - 5, & x \leq 1, \\ -2x + (\log)_2(b^2 - 2), & x > 1 \end{cases}$$

Find the values of b for which $f(x)$ has the greatest value at $x = 1$.

 [Watch Video Solution](#)

Concept Application Exercise 6.5

1. Draw the graph of $f(x) = \frac{x^2 - x + 1}{x^2 + x + 1}$.

 [View Text Solution](#)

2. Discuss the number of roots of the equation $e(k - x \log x) = 1$ for different value of k .

 [Watch Video Solution](#)

3. Draw the graph of $y = (x + 1)^{2/3} + (x - 1)^{2/3}$

 [Watch Video Solution](#)

4. Draw the graph of $f(x) = \log_e(1 - \log_e x)$. Find the point of inflection

 [View Text Solution](#)

5. Draw and graph of $f(x) = \frac{4 \log_e x}{x^2}$. Also find the range.

 [Watch Video Solution](#)

Concept Application Exercise 6.6

1. A private telephone company serving a small community makes a profit of Rs. 12.00 per subscriber, if it has 725 subscribers. It decides to reduce the rate by a fixed sum for each subscriber over 725, thereby reducing the profit by 1 paise per subscriber. Thus, there will be profit of Rs. 11.99 on each of the 726 subscribers, Rs. 11.98 on each of the 727 subscribers, etc. What is the number of subscribers which will give the company the maximum profit?

 [View Text Solution](#)

2. एक वृत्त और एक वर्ग के परिमाण का योग k है, जहां k एक अचर है सिद्ध कीजिए कि उनके क्षेत्रफल का योग निम्नतम है जब वर्ग की भुजा वृत्त की त्रिज्या की दुगुनी है।

 [View Text Solution](#)

3. A figure is bounded by the curves $y = x^2 + 1$, $y = 0$, $x = 0$, and $x = 1$. At what point (a, b) should a tangent be drawn to curve $y = x^2 + 1$ for it to cut off a trapezium of greatest area from the figure?

 [Watch Video Solution](#)

4. Find the minimum length of radius vector of the curve $\frac{a^2}{x^2} + \frac{b^2}{y^2} = 1$

 [Watch Video Solution](#)

5. Find the point at which the slope of the tangent of the function $f(x) = e^x \cos x$ attains minima, when $x \in [0, 2\pi]$.

 [Watch Video Solution](#)

6. An electric light is placed directly over the centre of a circular plot of lawn 100 m in diameter. Assuming that the intensity of light varies directly as the sine of the angle at which it strikes an illuminated surface and inversely as the square of its distance from its surface, how should the light be hung in order that the intensity may be as great as possible at the circumference of the plot?

 [Watch Video Solution](#)

7. A swimmer S is in the sea at a distance d km from the closest point A on a straight shore. The house of the swimmer is on the shore at a distance L km from A . He can swim at a speed of u km/hr and walk at a speed of v km/hr ($v > u$). At what point on the shore should he land so that he reaches his house in the shortest possible time?

 [Watch Video Solution](#)

8. Let (h, k) be a fixed point, where $h > 0, k > 0$. A straight line passing through this point cuts the positive direction of the coordinate axes at the point P and Q . Find the minimum area of triangle OPQ , O being the origin.

 [Watch Video Solution](#)

9. A point P is given on the circumference of a circle of radius r . Chord QR is parallel to the tangent at P . Determine the maximum possible area of the triangle PQR .

 [Watch Video Solution](#)

Concept Application Exercise 6.7

1. A metal box with a square base and vertical sides is to contain 1024cm^3 of water, the material for the top and bottom costs Rs. 5per cm^2 and the material for the costs Rs. 2.50per cm^2 . Find the least cost of the box.



[View Text Solution](#)

2. 100cm^2 आयतन वाले डिब्बे सभी बंद बेलनकार (लंब वृतीय) डिब्बों में से न्यूनतम पृष्ठ क्षेत्रफल वाले डिब्बे की विमाएँ ज्ञात कीजिए



[View Text Solution](#)

3. The lateral edge of a regular rectangular pyramid is $acmlong$. The lateral edge makes an angle α with the plane of the base. Find the value of α for which the volume of the pyramid is greatest.



[View Text Solution](#)

4. Show that the cone of the greatest volume which can be inscribed in a given sphere has an altitude equal to $2/3$ of the diameter of the sphere.



[Watch Video Solution](#)

5. A regular square based pyramid is inscribed in a sphere of given radius R so that all vertices of the pyramid belong to the sphere. Find the greatest value of the volume of the pyramid.



[View Text Solution](#)

Exercise

1. A function is matched below against an interval where it is supposed to be increasing. Which of the following parts is incorrectly matched?

Interval, Function $[2, \infty)$, $2x^3 - 3x^2 - 12x + 6$ $(-\infty, \infty)$,
 $x^3 = 3x^2 + 3x + 3$ $(-\infty - 4)$, $x^3 + 6x^2 + 6$ $\left(-\infty, \frac{1}{3}\right)$,
 $3x^2 - 2x + 1$

- A. $[2, \infty)$, $2x^3 - 3x^2 - 12x + 6$
- B. $(-\infty, \infty)$, $x^3 - 3x^2 - 3x + 3$
- C. $(-\infty, -4)$, $x^3 - 6x^2 + 6$
- D. $\left[-\infty, \frac{1}{3}\right]$, $3x^2 - 2x + 1$

Answer: 4



Watch Video Solution

2. On which of the following intervals is the function $x^{100} + \sin x - 1$ decreasing? (a) $(0, \frac{\pi}{2})$ (b) $(0, 1)$ (c) $(\frac{\pi}{2}, \pi)$ (d) none of these

A. $(0, \pi/2)$

B. $(0,1)$

C. $(\pi/2, \pi)$

D. None of these

Answer: 4



Watch Video Solution

3. The function $f(x) = \tan^{-1}(\sin x + \cos x)$ is an increasing function in (a) $(-\frac{\pi}{2}, \frac{\pi}{4})$ (b) $(0, \frac{\pi}{2})$ (c) $(-\frac{\pi}{2}, \frac{\pi}{2})$ (d) $(\frac{\pi}{4}, \frac{\pi}{2})$

A. $(0, \pi/2)$

B. $(0, \frac{\pi}{2})$

C. $(-\frac{\pi}{2}, \frac{\pi}{2})$

D. $(\frac{\pi}{4}, \frac{\pi}{2})$

Answer: 1



Watch Video Solution

4. Assertion Consider the following statements in S and RS : Both $\sin x$ and $\cos x$ are decreasing function in the interval $(\frac{\pi}{2}, \pi)$ *Reason* If a differentiable function decreases in an interval (a, b) , then its derivative also decrease in (a, b) . Which of the following is true? (a) Both S and R are wrong. (b) Both S and R are correct, but R is not the correct explanation of S . (c) S is correct and R is the correct explanation for S . (d) S is correct and R is wrong.

A. Both S and R are wrong

B. Both S and R are correct , but R is not the correct explanation ofS.

C. S is correct and R is the correct explanation for S.

D. S is correct and R is wrong.

Answer: 4



[Watch Video Solution](#)

5. The length of the longest interval in which the function $3 \sin x - 4 \sin^3 x$ is increasing is $\frac{\pi}{3}$ (b) $\frac{\pi}{2}$ (c) $\frac{3\pi}{2}$ (d) π

A. $\frac{\pi}{3}$

B. $\frac{\pi}{2}$

C. $\frac{\pi}{2}$

D. (π)

Answer: 1



[Watch Video Solution](#)

6. The function x^x decreases in the interval (a) $(0, e)$ (b) $(0, 1)$ (c) $\left(0, \frac{1}{e}\right)$

(d) none of these

A. $(0, e)$

B. $(0, 1)$

C. $\left(0, \frac{1}{e}\right)$

D. none of these

Answer: 3



[View Text Solution](#)

7. Let $f(x) = x\sqrt{4ax - x^2}$, ($a > 0$). Then $f(x)$ is (a). increasing in $(0, 3a)$
decreasing in $(3a, 4a)$ (b). increasing in $(a, 4a)$ decreasing in $(5a, \infty)$ (c).
increasing in $(0, 4a)$ (d). none of these

A. increasing in $(0, 3a)$ decreasing in $(3a, 4a)$

B. increasing in $(a, 4a)$, decreasing in $(5a, \infty)$

C. increasing in $(0, 4a)$

D. none of these

Answer: 1



Watch Video Solution

8. Function $f(x) = |x| - |x - 1|$ is monotonically increasing when (a) $x < 0$ (b) $x > 1$ (c) $x < 1$ (d) $0 < x < 1$

A. $x < 0$

B. $x > 0$

C. $x < 0$

D. $0 < x < 1$

Answer: 4



Watch Video Solution

9. If $f'(x) = |x| - \{x\}$, where $\{x\}$ denotes the fractional part of x , then $f(x)$ is decreasing in (a) $\left(-\frac{1}{2}, 0\right)$ (b) $\left(-\frac{1}{2}, 2\right)$ $\left(-\frac{1}{2}, 2\right)$ (d) $\left(\frac{1}{2}, \infty\right)$
- A. $\left(-\frac{1}{2}, 0\right)$
 B. $\left(-\frac{1}{2}, 2\right)$
 C. $\left(-\frac{1}{2}, 2\right)$
 D. $\left(\frac{1}{2}, \infty\right)$

Answer: 1



Watch Video Solution

10. The length of the largest continuous interval in which the function $f(x) = 4x - \tan 2x$ is monotonic is (a) $\frac{\pi}{2}$ (b) $\frac{\pi}{4}$ (c) $\frac{\pi}{8}$ (d) $\frac{\pi}{16}$

A. $\pi/2$

B. $\pi/4$

C. $\pi/8$

D. $\pi/16$

Answer: 2



Watch Video Solution

11. $f(x) = (x - 2)|x - 3|$ is monotonically increasing in $\left(-\infty, \frac{5}{2}\right) \cup (3, \infty)$ (b) $\left(\frac{5}{2}, \infty\right)$ (c) $(2, \infty)$ (d) $(-\infty, 3)$

A. $(-\infty, 5/2) \cup (3, \infty)$

B. $5/2, \infty)$

C. $(2, \infty)$

D. $(-\infty, 3)$

Answer: 1



Watch Video Solution

12. $f(x) = (x - 8)^4(x - 9)^5$, $0 \leq x \leq 10$, monotonically decreases in

(a) $\left(\frac{76}{9}, 10\right]$ (b) $\left(8, \frac{76}{9}\right)$ (c) $(0, 8)$ (d) $\left(\frac{76}{9}, 10\right)$

A. $\left(\frac{76}{9}, 10\right)$

B. $\left(8, \frac{76}{9}\right)$

C. $[0, 8)$

D. $\left(\frac{76}{9}, 10\right)$

Answer: 2



Watch Video Solution

13. For all $x \in (0, 1)$ (a) $e^x < 1 + x$ (b) $(\log)_e(1 + x) < x$ (c) $\sin x > x$

(d) $(\log)_e x > x$

A. $e^x < 1 + x$

B. $\log_e(1 + x) < x$

C. $\sin x > x$

D. $\log_e x > x$

Answer: 2

 [Watch Video Solution](#)

14. If $f(x) = xe^{x(x-1)}$, then a) $f(x)$ is increasing on $\left[-\frac{1}{2}, 1\right]$ b) decreasing on \mathbb{R} c) increasing on \mathbb{R} (d) decreasing on $\left[-\frac{1}{2}, 1\right]$

A. increasing on $\left[-\frac{1}{2}, 1\right]$

B. decreasing on \mathbb{R}

C. increasing on \mathbb{R}

D. decreasing on $\left[-\frac{1}{2}, 1\right]$

Answer: 1

 [Watch Video Solution](#)

15. If $f(x) = kx^3 - 9x^2 + 9x + 3$ monotonically increasing in R , then

A. $k < 3$

B. $k \leq 2$

C. $k \geq 3$

D. none of these

Answer: 3



Watch Video Solution

16. If the function $f(x) = \frac{K \sin x + 2 \cos x}{\sin x + \cos x}$ is strictly increasing for all

values of x , then $K < 1$ (b) $K > 1$ $K < 2$ (d) $K > 2$

A. $k < 1$

B. $k \leq 2$

C. $k \geq 3$

D. none of these

Answer: 4



Watch Video Solution

17. Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be a function such that $f(x) = ax + 3 \sin x + 4 \cos x$.

Then $f(x)$ is invertible if $a \in (-5, 5)$ (b) $a \in (-\infty, 5)$

$a \in (-5, +\infty)$ (d) none of these

A. $k < 1$

B. $k > 1$

C. $k < 2$

D. $k > 2$

Answer: 4



Watch Video Solution

18. Which of the following statement is always true? If $f(x)$ is increasing, the $f^{-1}(x)$ is decreasing. If $f(x)$ is increasing, then $\frac{1}{f(x)}$ is also increasing. If f and g are positive functions and f is increasing and g is decreasing, then $\frac{f}{g}$ is a decreasing function. If f and g are positive functions and f is decreasing and g is increasing, the $\frac{f}{g}$ is a decreasing function.

A. If $f(x)$ is increasing then $f^{-1}(x)$ is also decreasing

B. If $f(x)$ is increasing then $1/f(x)$ is also increasing

C. If f and g are positive functions and f is increasing and g is decreasing then f/g is decreasing function

D. If f and g are positive functions and f is decreasing and g is increasing then f/g is decreasing function

Answer: 4



Watch Video Solution

19. Let $f: \overrightarrow{RR}$ be a differentiable function for all values of x and has the property that $f(x)$ and $f'(x)$ has opposite signs for all value of x . Then, $f(x)$ is an increasing function $f(x)$ is an decreasing function $f^2(x)$ is an decreasing function $|f(x)|$ is an increasing function

- A. $f(x)$ is an iincreasing function
- B. $f(x)$ is a decreasing function
- C. $f^2(x)$ is a decreasing function
- D. $|f(x)|$ is an increasing function

Answer: 3

 [Watch Video Solution](#)

20. Let $f: R \rightarrow R$ be a differentiable function $\forall x \in R$. If the tangent drawn to the curve at any point $x \in (a, b)$ always lies below the curve, then

$$f'(x) < 0, f''(x) < 0 \forall x \in (a, b) \qquad f'(x) > 0, f''(x) > 0 \forall x \in (a, b)$$

$$f'(x) > 0, f''(x) > 0 \forall x \in (a, b) \text{ none of these}$$

A. $f'(x) > 0, f''(x) < 0 \forall x \in (a, b)$

B. $f'(x) < 0, f''(x) < 0 \forall x \in (a, b)$

C. $f'(x) > 0, f''(x) > 0 \forall x \in (a, b)$

D.

Answer: 3

 [View Text Solution](#)

21. Let $f(x)$ be a function such that $f'(x) = (\log)_{\frac{1}{3}} [(\log)_3(\sin x + a)]$. If

$f(x)$ is decreasing for all real values of x , then $a \in (1, 4)$ (b) $a \in (4, \infty)$

$a \in (2, 3)$ (d) $a \in (2, \infty)$

A. $a \in (1, 4)$

B. $a \in (4, \infty)$

C. $a \in (2, 3)$

D. $a \in (2, \infty)$

Answer: 2



Watch Video Solution

22. If $f(x) = x^3 + 4x^2 + \lambda x + 1$ is a monotonically decreasing function of x in the largest possible interval $\left(-2, -\frac{2}{3}\right)$. Then $\lambda = 4$ (b) $\lambda = 2$
 $\lambda = -1$ (d) λ has no real value

A. $\lambda = 4$

B. $\lambda = 2$

C. $\lambda = -1$

D. λ has no real value

Answer: 1



Watch Video Solution

23. $f(x) = |x \log_e x|$ monotonically decreases in $\left(0, \frac{1}{e}\right)$ (b) $\left(\frac{1}{e}, 1\right)$
(1, ∞) (d) $\left(\frac{1}{e}, \infty\right)$

A. $(0, 1/e)$

B. $(1/e, 1)$

C. $(1, \infty)$

D. $(1/e, \infty)$

Answer: 2



[View Text Solution](#)

24. The set of value(s) of a for which the function $f(x) = \frac{ax^3}{3} + (a+2)x^2 + (a-1)x + 2$ possesses a negative point of inflection is (a) $(-\infty, -2) \cup (0, \infty)$ (b) $\left\{-\frac{4}{5}\right\}$ (c) $(-2, 0)$ (d) empty set

A. $(-\infty, -2) \cup (0, \infty)$

B. $\{ - 4/5 \}$

C. $(- 2, 0)$

D. empty set

Answer: 1



Watch Video Solution

25. The maximum value of the function $f(x) = \frac{(1+x)^{0.6}}{1+x^{0.6}}$ in the interval $[0, 1]$ is $2^{0.4}$ (b) $2^{-0.4}$ 1 (d) $2^{0.6}$

A. $2^{0.4}$

B. $2^{-0.4}$

C. 1

D. $2^{0.6}$

Answer: 3



Watch Video Solution

26. Suppose that f is a polynomial of degree 3 and that $f'' \neq 0$ at any of the stationary point. Then f has exactly one stationary point f must have no stationary point f must have exactly two stationary points f has either zero or two stationary points.

- A. f has exactly one stationary point
- B. f must have no stationary point
- C. f must have exactly two stationary points
- D. f has either zero or two stationary points

Answer: 4



[Watch Video Solution](#)

27. A function $g(x)$ is defined as $g(x) = \frac{1}{4}f(2x^2 - 1) + \frac{1}{2}f(1 - x^2)$ and $f(x)$ is an increasing

function. Then $g(x)$ is increasing in the interval. (a) $(-1, 1)$ (b)

$\left(-\sqrt{\frac{2}{3}}, 0\right) \cup \left(\sqrt{\frac{2}{3}}, \infty\right)$ (c) $\left(-\sqrt{\frac{2}{3}}, \sqrt{\frac{2}{3}}\right)$ (d) none of these

A. $(-1, 1)$

B. $-\left(\frac{\sqrt{2}}{3}, 0\right) \cup \left(\frac{\sqrt{2}}{3}, \infty\right)$

C. $-\frac{\sqrt{2}}{3}, \frac{\sqrt{2}}{3}$

D. none of these

Answer: 2



Watch Video Solution

28. If $\varphi(x)$ is a polynomial function and

$\varphi'(x) > \varphi(x) \forall x \geq 1$ and $\varphi(1) = 0$, then $\varphi(x) \geq 0 \forall x \geq 1$ $\varphi(x)$

A. $\phi(x) \geq 0 \forall x \geq 1$

B. $\phi(x) < 0 \forall x \geq 1$

C. $\phi(x) = 0 \forall x \geq 1$

D. none of these

Answer: 1



Watch Video Solution

29.

If

$f''(x) > \forall x \in R, f(3) = 0$ and $g(x) = f(\tan^2 x - 2 \tan x + 4)$ $0 < x < \pi$

,then $g(x)$ is increasing in

A. $\left(0, \frac{\pi}{4}\right)$

B. $\left(\frac{\pi}{6}, \frac{\pi}{3}\right)$

C. $\left(0, \frac{\pi}{3}\right)$

D. $\left(\frac{\pi}{4}, \frac{\pi}{2}\right)$

Answer: 4



Watch Video Solution

30. If $f(x) = x + \sin x$, $g(x) = e^{-x}$, $u = \sqrt{c+1} - \sqrt{c}$, $v = \sqrt{c} - \sqrt{c-1}$, ($c > 1$), then $\text{fog}(u) \text{gof}(v)$ (d) $\text{fog}(u)$

A. $\text{fog}(u) < \text{fog}(v)$

B. $\text{gof}(u) < \text{gof}(v)$

C. $\text{gof}(u) > \text{gof}(v)$

D. $\text{fog}(u) < \text{fog}(v)$

Answer: 4

 [Watch Video Solution](#)

31. The number of solutions of the equation $x^3 + 2x^2 + 6x + 2 \cos x = 0$ where $x \in [0, 2\pi]$ is (a) one (b) two (c) three (d) zero

A. one

B. two

C. three

D. zero

Answer: 4



[Watch Video Solution](#)

32. Let $f(x) = \cos \pi x + 10x + 3x^2 + x^3$, $-2 \leq x \leq 3$. The absolute minimum value of $f(x)$ is 0 (b) -15 (c) $3 - 2\pi$ none of these

A. 0

B. -15

C. $3 - 2\pi$

D. none of these

Answer: 2



[Watch Video Solution](#)

33. The global maximum value of $f(x) = (\log)_{10}(4x^3 - 12x^2 + 11x - 3), x \in [2, 3]$, is $-\frac{3}{2}(\log)_{10}3$ (b) $1 + (\log)_{10}3$ (c) $(\log)_{10}3$ (d) $\frac{3}{2}(\log)_{10}3$

A. $-\frac{3}{2}\log_{10}^3$

B. $1 + \log 10^3$

C. $\log 10^3$

D. $\frac{3}{2}\log 10^3$

Answer: 2



Watch Video Solution

34. $f: \mathbb{R} \rightarrow \mathbb{R}$, $f(x)$ is differentiable such that $f'(x) = k(x^5 + x), k \neq 0$. Then $f(x)$ is always increasing (b) decreasing either increasing or decreasing non-monotonic

A. increasing

B. decreasing on $[-1/2, \infty]$

C. either increasing or decreasing

D. non monotonic

Answer: 3



Watch Video Solution

35. The value of a for which the function $f(x) = a \sin x + \left(\frac{1}{3}\right) \sin 3x$ has an extremum at $x = \frac{\pi}{3}$ is (a) 1 (b) -1 (c) 0 (d) 2

A. 1

B. -1

C. 0

D. 2

Answer: 5



Watch Video Solution

36. If $f(x) = a \log|x| + bx^2 + x$ has extreme values at $x = -1$ and at $x = 2$, then find a and b .

A. $a=2, b=-1$

B. $a=2, b=-1/2$

C. $a=-2, b=1/2$

D. none of these

Answer: 2



[Watch Video Solution](#)

37. If a function $f(x)$ has $f'(a) = 0$ and $f''(a) = 0$, then (a) $x = a$ is a maximum for $f(x)$ (b) $x = a$ is a minimum for $f(x)$ (c) can not conclude anything about its maxima and minima (d) $f(x)$ is necessarily a constant function.

- A. $x=a$ is a maximum for $f(x)$
- B. $x=a$ is a minimum for $f(x)$
- C. it is difficult to say (a) and (b)
- D. $f(x)$ is necessarily a constant function

Answer: 3

 [Watch Video Solution](#)

38. The function $f(x) = \sin^4 x + \cos^4 x$ increasing if $x \in (0, \frac{\pi}{2})$

- A. It is monotonic increasing $\forall x$ in \mathbb{R} .
- B. $f(x)$ fails to exist for three distinct real values of x
- C. $f(x)$ changes its sign twice as x varies from $-\infty \rightarrow \infty$
- D. The function attains its extreme values at x_1 and x_2 such that

$$x_1 x_2 > 0$$

Answer: 3



Watch Video Solution

39. The function $f(x) = \sin^4 x + \cos^4 x$ increasing if $x \in$

- A. $0 < x < \pi/8$
- B. $\pi/40 < x < 3\pi/8$
- C. $3\pi/8 < x < 5\pi/8$
- D. $5\pi/8 < x < 3\pi/4$

Answer: 2



Watch Video Solution

40. If $f(x) = x^5 - 5x^4 + 5x^3 - 10$ has local maximum and minimum at $x = p$ and $x = q$, respectively, then $(p, q) =$ (a) $(0, 1)$ (b) $(1, 3)$ (c) $(1, 0)$
(d) none of these

A. $(0, 1)$

B. (1,3)

C. (1,0)

D. none of these

Answer: 2



Watch Video Solution

41. Let $P(x) = a_0 + a_1x^2 + a_2x^4 + \dots + a_nx^{2n}$ be a polynomial in a real variable x with $0 < a_0 < a_1 < a_2 < \dots < a_n$. The function $P(x)$ has a.

neither a maximum nor a minimum b. only one maximum c. only one minimum d. only one maximum and only one minimum e. none of these

A. neither a maximum nor a minimum

B. only one maximum

C. only one minimum

D. only one maximum and only one minimum

Answer: 3



[Watch Video Solution](#)

42. Let $f(x) = \begin{cases} |x| & \text{for } 0 < |x| \leq 2 \\ 1 & \text{for } x = 0 \end{cases}$ Then at $x=0$, $f(x)$ has

- (a) a local maximum
- (b) no local maximum
- (c) a local minimum
- (d) no extremum

- A. a local maximum
- B. no local maximum
- C. a local minimum
- D. no extremum

Answer: 1



[Watch Video Solution](#)

43. If $f(x) = x^3 + bx^2 + cx + d$ and $\Delta < 0$

- A. $f(x)$ is strictly increasing function
- B. $f(x)$ has local maxima
- C. $f(x)$ is a strictly decreasing function
- D. $f(x)$ is bounded

Answer: 1



[View Text Solution](#)

44. If $f(x) = \begin{cases} \sin(x^2 - 3x), & x \leq 0 \\ 6x + 5x^2, & x > 0 \end{cases}$ then at $x=0$, $f(x)$ is?

- A. $f(x)$ has a local minima
- B. $f(x)$ has a local maxima
- C. $f(x)$ has point of inflection
- D. none of these

Answer: 1



[Watch Video Solution](#)

45. The greatest value of $f(x) = \cos(xe^{[x]} + 7x^2 - 3x)$, $x \in [-1, \infty]$, is (where $[.]$ represents the greatest integer function). – 1 (b) 1 (c) 0 (d) none of these

A. -1

B. 1

C. 0

D. none of these

Answer: 2



[Watch Video Solution](#)

46. The function $f(x) = (4\sin^2 x - 1)^n (x^2 - x + 1)$, $n \in \mathbb{N}$, has a local minimum at $x = \frac{\pi}{6}$. Then n is any even number n is an odd number n is odd prime number n is any natural number

- A. n is any even integer
- B. n is an odd integer
- C. n is odd prime number
- D. n is any natural number

Answer: 1



[Watch Video Solution](#)

47. The true set of real values of x for which the function $f(x) = x \ln x - x + 1$ is positive is

- A. $(1, \infty)$
- B. $(1/e, \infty)$

C. $[e, \infty)$

D. $(0, 1) \cup (1, \infty)$

Answer: 1



[View Text Solution](#)

48. All possible value of $f(x) = (x + 1)^{\frac{1}{3}} - (x - 1)^{\frac{1}{3}}$ on $[0,1]$ is 1 (b) 2 (c)

3 (d) $\frac{1}{3}$

A. 1

B. 2

C. 3

D. $\frac{1}{3}$

Answer: 2



[Watch Video Solution](#)

49. The function $f(x) = \frac{\ln(\pi + x)}{\ln(e + x)}$ is

- A. increasing in $(0, \infty)$
- B. decreasing oin $(0, \infty)$
- C. increasing in $(0, \pi/e)$, *decrea* sin $g \in (\pi/e, \infty)$
- D. decreasing in $(0, \pi/e)$ increasing in $(\pi/e, \infty)$

Answer: 2



Watch Video Solution

50. If the function $f(x) = 2x^3 - 9ax^2 + 12a^2x + 1$, where $a > 0$, attains its maximum and minimum at p and q , respectively, such that $p^2 = q$, then a equal to (a) 1 (b) 2 (c) $\frac{1}{2}$ (d) 3

A. 1

B. 2

C. $\frac{1}{2}$

D. 3

Answer: 2



[Watch Video Solution](#)

51. Let $f(x) = \begin{cases} x + 2, & -1 \leq x < 0 \\ 1, & x = 0 \\ \frac{x}{2}, & 0 < x \leq 1 \end{cases}$

A. a point of minima

B. a point of maxima

C. both points of minima and maxima

D. neither a point of minima nor that of maxima

Answer: 4



[Watch Video Solution](#)

52. If $f(x) = \begin{cases} \sin^{-1}(\sin x), & x > 0 \\ \frac{\pi}{2}, & x = 0, \end{cases}$ then $\cos^{-1}(\cos x), x < 0$

- A. $x=0$ is a point of maxima
- B. $x=0$ is a point of minima
- C. $x=0$ is a point of intersection
- D. none of these

Answer: 1



Watch Video Solution

53. A function f is defined by $f(x) = |x|^m |x - 1|^n \forall x \in \mathbb{R}$. The local maximum value of the function is $(m, n \in \mathbb{N}), 1 < m < n$ (b) $m \wedge n$

$\frac{m^m n^n}{(m+n)^{m+n}}$ (d) $\frac{(mn)^{mn}}{(m+n)^{m+n}}$

- A. 1
- B. $m^n n^m$

C. $\frac{m^m n^n}{(m+n)^{m+n}}$

D. $\frac{mn^{mn}}{m+n^{m+n}}$

Answer: 3



Watch Video Solution

54. Let the function $f(x)$ be defined as follows: $f(x) = x^3 + x^2 - 10x$, $-1 \leq x < 0$
 $\cos x$, $0 \leq x < 2\pi$ $1 + \sin x$, $2\pi \leq x \leq \pi$, then which of the following statement(s)
 is/are correct

- A. a local minimum at $x = \pi/2$
- B. a global maximum at $x = \pi/2$
- C. a absolute maximum at $x = -1$
- D. a absolute maximum at $x = \pi$

Answer: 3



Watch Video Solution

55. Consider the function $f: (-\infty, \infty) \rightarrow (-\infty, \infty)$ defined by $f(x) = \frac{x^2 + a}{x^2 + a}$, $a > 0$, which of the following is not true? maximum value of f is not attained even though f is bounded. $f(x)$ is increasing on $(0, \infty)$ and has minimum at $x = 0$ $f(x)$ is decreasing on $(-\infty, 0)$ and has minimum at $x = 0$. $f(x)$ is increasing on $(-\infty, \infty)$ and has neither a local maximum nor a local minimum at $x = 0$.

A. Maximum value of f is not attained even though f is bounded

B. $f(x)$ is increasing on $(0, \infty)$ and has minimum at $x=0$

C. $f(x)$ is decreasing on $(-\infty, 0)$ and has minimum at $x=0$

D. $f(x)$ is increasing on $(-\infty, \infty)$ and has neither a local maximum nor a local minimum at $x=0$

Answer: 4



Watch Video Solution

56. $F(x) = 4 \tan x - \tan^2 x + \tan^3 x, x \neq n\pi + \frac{\pi}{2}$

- A. is monotonically increasing
- B. is monotonically decreasing
- C. has a point of maxima
- D. has a point of minima

Answer: 1



[View Text Solution](#)

57. Let $h(x) = x^{\frac{m}{n}}$ for $x \in R$, where m and n are odd numbers where $0 < m < n$. Then $y = h(x)$ has

- a. no local extremums
- b. one local maximum
- c. one local minimum
- d. none of these

- A. no local extremums
- B. one local maximum
- C. one local minimum

D. none of these

Answer: 1



Watch Video Solution

58. The greatest value of the function $f(x) = \frac{\sin 2x}{\sin\left(x + \frac{\pi}{4}\right)}$ on the interval $\left(0, \frac{\pi}{2}\right)$ is

A. $\frac{1}{\sqrt{2}}$

B. $\sqrt{2}$

C. 1

D. $-\sqrt{2}$

Answer: 3



Watch Video Solution

59. The minimum value of $e^{(2x^2 - 2x + 1) \sin^2 x}$ is a. e (b) $\frac{1}{e}$ (c) 1 (d) 0

A. e

B. $1/e$

C. 1

D. 0

Answer: 3



[Watch Video Solution](#)

60. The maximum value of $x^4 e^{-x^2}$ is (a) e^2 (b) e^{-2} (c) $12e^{-2}$ (d) $4e^{-2}$

A. e^2

B. e^{-2}

C. $12e^{-2}$

D. $4e^{-2}$

Answer: 4



Watch Video Solution

61. If $a^2x^4 + b^2y^4 = c^6$, then the maximum value of xy is $\frac{c^2}{\sqrt{ab}}$ (b) $\frac{c^3}{ab}$
 $\frac{c^3}{\sqrt{2ab}}$ (d) $\frac{c^3}{2ab}$

A. $\frac{c^2}{\sqrt{ab}}$

B. $\frac{c^3}{\sqrt{ab}}$

C. $\frac{c^3}{\sqrt{2ab}}$

D. $\frac{c^3}{\sqrt{2ab}}$

Answer: 3



Watch Video Solution

62. Least natural number a for which

$$x + ax^{-2} > 2, \forall x \in (0, \infty) \text{ is}$$

A. 1

B. 2

C. 5

D. none of these

Answer: 2



Watch Video Solution

63. $f(x) = \begin{cases} 4x - x^3 + \ln(a^2 - 3a + 3), & 0 \leq x < 3 \\ 3x - 18, & x \geq 3 \end{cases}$

complete set of values of a such that $f(x)$ as a local minima at $x = 3$ is

[- 1, 2] (- ∞, 1) ∪ (2, ∞) [1, 2] (d) (- ∞, - 1) ∪ (2, ∞)

A. [- 1, 2]

B. (- ∞, 1) ∪ (2, ∞)

C. [1, 2]

D. (- ∞, - 1) ∪ (2, ∞)

Answer: 2



Watch Video Solution

64.

$f(x) = \begin{cases} 2 - |x^2 + 5x + 6|, & x \neq 2a^2 + 1, \\ x = -2 \end{cases}$ Then the range of a is

so that $f(x)$ has maxima at $x = -2$, is (a) $|a| \geq 1$ (b) $|a| < 1$ (c) $a > 1$ (d)

$a < 1$

A. $|a| \geq 1$

B. $|a| \leq 1$

C. $a > 1$

D. $a < 1$

Answer: 4



Watch Video Solution

65. A differentiable function $f(x)$ has a relative minimum at $x = 0$. Then the function $f = f(x) + ax + b$ has a relative minimum at $x = 0$ for

(a) all a and all b (b) all b if $a = 0$ (c) all $b > 0$ (d) all $a > 0$

A. all a and all b

B. all b if $a = 0$

C. all $b > 0$

D. all $a > 0$

Answer: 2



Watch Video Solution

66. if $f(x) = 4x^3 - x^2 - 2x + 1$ and $g(x) = \{ \min f(t) : 0 \leq t \leq x; 0 \leq x \leq 1, 3 - x : 1 \}$ then $g\left(\frac{1}{4}\right) + g\left(\frac{3}{4}\right) + g\left(\frac{5}{4}\right)$ is equal to

A. $7/4$

B. $9/4$

C. $13/4$

D. $5/2$

Answer: 4



Watch Video Solution

67. If $f: \vec{R} \rightarrow \vec{R}$ and $g: \vec{R} \rightarrow \vec{R}$ are two functions such that $f(x) + f''(x) = -xg(x)f'(x)$ and $g(x) > 0 \forall x \in \vec{R}$. Then the function $f^2(x) + f'(x)^2$ has a maxima at $x = 0$ a minima at $x = 0$ a point of inflexion at $x = 0$ none of these

A. a maxima at $x = 0$

B. a minima at $x = 0$

C. a point of inflexion at $x = 0$

D. a point of inflexion at $x = 0$

Answer: 1



Watch Video Solution

68. If $A > 0, B > 0, A + B = \pi/3$, and maximum value of $\tan A \tan B$ is M then the value of $1/M$ is _____.

A. $\frac{1}{\sqrt{3}}$

B. $\frac{1}{3}$

C. 3

D. $\sqrt{3}$

Answer: 2



Watch Video Solution

69. If $f(x) = \frac{t + 3x - x^2}{x - 4}$, where t is a parameter that has minimum and maximum, then the range of values of t is (a) $(0, 4)$ (b) $(0, \infty)$ (c)

$(-\infty, 4)$ (d) $(4, \infty)$

A. $(0, 4)$

B. $(0, \infty)$

C. $-\infty, 4)$

D. $(4, \infty)$

Answer: 3



Watch Video Solution

70. The least value of a for which the equation $\frac{4}{\sin x} + \frac{1}{1 - \sin x} = a$ has at least one solution in the interval $\left(0, \frac{\pi}{2}\right)$ 9 (b) 4 (c) 8 (d) 1

A. 9

B. 4

C. 8

D. 1

Answer: 3



Watch Video Solution

71. If $f(x) = -x^3 - 3x^2 - 2x + a$, $a \in R$ then the real values of x satisfying $f(x^2 + 1) > f(2x^2 + 2x + 3)$ will be

A. $(-\infty, \infty)$

B. $(0, \infty)$

C. $(-\infty, 0)$

D. ϕ

Answer: 1



View Text Solution

72. If the equation $4x^3 + 5x + k = 0$ ($k \in R$) has a negative real root then (a) $k = 0$ (b) $-\infty < k < 0$ (c) $0 < k < \infty$ (d) $-\infty < k < \infty$

A. $k=0$

B. $-\infty < k < 0$

C. $0 < k < \infty$

D. $-\infty < k < \infty$

Answer: 3

 [View Text Solution](#)

73. Tangent is drawn to ellipse $\frac{x^2}{27} + y^2 = 1$ at $(3\sqrt{3}\cos\theta, \sin\theta)$ [where $\theta \in (0, \frac{\pi}{2})$] Then the value of θ such that sum of intercepts on axes made by this tangent is minimum is (a) $\frac{\pi}{3}$ (b) $\frac{\pi}{6}$ (c) $\frac{\pi}{8}$ (d) $\frac{\pi}{4}$

A. $\frac{\pi}{3}$

B. $\frac{\pi}{6}$

C. $\frac{\pi}{8}$

D. $\frac{\pi}{4}$

Answer: 2



Watch Video Solution

74. The largest term in the sequence $a_n = \frac{n^2}{n^3 + 200}$ is given by $\frac{529}{49}$ (b)
 $\frac{8}{89}$ $\frac{49}{543}$ (d) none of these

A. $\frac{529}{49}$

B. $\frac{8}{89}$

C. $\frac{49}{543}$

D. none of these

Answer: 3



Watch Video Solution

75. A factory D is to be connected by a road with a straight railway line on which a town A is situated. The distance DB of the factory to the

railway line is $5\sqrt{3}km$. Length AB of the railway line is 20km. Freight charges on the road are twice the charges on the railway. The point P (A P

- A. $BP=5$ km
- B. $AP=5$ km
- C. $BP=7.5$ km
- D. none of these

Answer: 1

 [Watch Video Solution](#)

76. The volume of the greatest cylinder which can be inscribed in a cone of height 30 cm and semi-vertical angle 30° is $4000\frac{\pi}{\sqrt{3}}$ (b) $400\frac{\pi}{3}cm^3$

$4000\frac{\pi}{\sqrt{3}cm^3}$ (d) none of these

- A. $4000\pi / 3cm^3$
- B. $400\pi / 3cm^3$

C. $4000\pi / \sqrt{3}cm^3$

D. $4000\pi / 3cm^3$ none of these

Answer: 1



Watch Video Solution

77. A rectangle of the greatest area is inscribed in a trapezium $ABCD$, one of whose non-parallel sides AB is perpendicular to the base, so that one of the rectangles lies on the larger base of the trapezium. The base of trapezium are 6cm and 10 cm and AB is 8 cm long. Then the maximum area of the rectangle is $24cm^2$ (b) $48cm^2$ $36cm^2$ (d) none of these

A. $24cm^2$

B. $48cm^2$

C. $36cm^2$

D. none of these

Answer: 2



Watch Video Solution

78. A bell tent consists of a conical portion above a cylindrical portion near the ground. For a given volume and a circular base of a given radius, the amount of the canvas used is a minimum when the semi-vertical angle of the cone is $\frac{\cos^{-1} 2}{3}$ (b) $\frac{\sin^{-1} 2}{3}$ $\frac{\cos^{-1} 1}{3}$ (d) none of these

A. $\cos^{-1} 2/3$

B. $\sin^{-1} 2/3$

C. $\cos^{-1} 1/3$

D. none of these

Answer: 1



Watch Video Solution

79. A rectangle is inscribed in an equilateral triangle of side length $2a$ units. The maximum area of this rectangle can be (a) $\sqrt{3}a^2$ (b) $\frac{\sqrt{3}a^2}{4}$ a^2

(d) $\frac{\sqrt{3}a^2}{2}$

A. $\sqrt{3}a^2$

B. $\frac{\sqrt{3}a^2}{4}$

C. a^2

D. $\frac{\sqrt{3}a^2}{2}$

Answer: 4



Watch Video Solution

80. Tangents are drawn to $x^2 + y^2 = 16$ from the point $P(0, h)$. These tangents meet the x -axis at A and B . If the area of triangle PAB is minimum, then $h = 12\sqrt{2}$ (b) $h = 6\sqrt{2}$ (c) $h = 8\sqrt{2}$ (d) $h = 4\sqrt{2}$

A. $h=12\sqrt{2}$

B. $h=6\sqrt{2}$

C. $h=8\sqrt{2}$

D. $h=4\sqrt{2}$

Answer: 4



Watch Video Solution

81. The largest area of the trapezium inscribed in a semi-circle of radius

R , if the lower base is on the diameter, is (a) $\frac{3\sqrt{3}}{4}R^2$ (b) $\frac{\sqrt{3}}{2}R^2$

$\frac{3\sqrt{3}}{8}R^2$ (d) R^2

A. $\left(\frac{3\sqrt{3}}{4}R^2\right)$

B. $\left(\frac{\sqrt{3}}{2}R^2\right)$

C. $\left(\frac{3\sqrt{3}}{8}R^2\right)$

D. R^2

Answer: 1



[Watch Video Solution](#)

82. In the formula $\angle A + \angle B + \angle C = 180^\circ$, if $\angle A = 90^\circ$ and $\angle B = 55^\circ$, then $\angle C =$ _____ (a) $\frac{\pi}{4}$ (b) $\frac{\pi}{6}$ (c) $\frac{\pi}{3}$ (d) none of these

A. $\pi/4$

B. $\pi/6$

C. $\pi/3$

D. none of these

Answer: 3



[View Text Solution](#)

83. Two runner A and B start at the origin and run along positive x axis ,with B running three times as fast as A. An obsever, standeing one unit above the origin , keeps A and B in view.Then the maximum angle theta of sight between the observer's view of A and B is

A. $\pi / 8$

B. $\pi / 6$

C. $\pi / 3$

D. $\pi / 4$

Answer: 2



[View Text Solution](#)

84. The fuel charges for running a train are proportional to the square of the speed generated in km/h, and the cost is Rs. 48 at 16 km/h. If the fixed charges amount to Rs. 300/h, the most economical speed is 60 km/h (b) 40 km/h 48 km/h (d) 36 km/h

A. $60 \text{ km} / \text{h}$

B. $40 \text{ km} / \text{h}$

C. $48 \text{ km} / \text{h}$

D. $36 \text{ km} / \text{h}$

Answer: 2



[Watch Video Solution](#)

85. A cylindrical gas container is closed at the top and open at the bottom. If the iron plate of the top is $\frac{5}{4}$ times as thick as the plate forming the cylindrical sides, the ratio of the radius to the height of the cylinder using minimum material for the same capacity is 3:4 (b) 5:6 (c) 4:5 (d) none of these

A. 3:4

B. 5:6

C. 4:5

D. none of these

Answer: 3



[Watch Video Solution](#)

86. about to only mathematics

A. $4\sqrt{3r}$

B. $2\sqrt{3r}$

C. $6\sqrt{3r}$

D. $8\sqrt{3r}$

Answer: 3



[View Text Solution](#)

87. A given right cone has volume p , and the largest right circular cylinder that can be inscribed in the cone has volume q . Then $p:q$ is (a) 9:4 (b) 8:3 (c) 7:2 (d) none of these

A. 9:4

B. 8:3

C. 7:2

D. none of these

Answer: 1



[Watch Video Solution](#)

88. Find the cosine of the angle at the vertex of an isosceles triangle having the greatest area for the given constant length e of the median drawn to its lateral side.

A. 0.4

B. 0.5

C. 0.6

D. 0.8

Answer: 4



[View Text Solution](#)

89. A box, constructed from a rectangular metal sheet, is 21 cm by 16cm by cutting equal squares of sides x from the corners of the sheet and then turning up the projected portions. The value of x os that volume of the box is maximum is 1 (b) 2 (c) 3 (d) 4

A. 1

B. 2

C.

D. 3

Answer: 3



[Watch Video Solution](#)

90. The vertices of a triangle are $(0,0)$, $(x, \cos x)$, and $(\sin^3 x, 0)$, where $x \in (0, \frac{\pi}{2})$

A. $3 \frac{\sqrt{3}}{32}$

B. $\frac{\sqrt{3}}{32}$

C. $\frac{4}{32}$

D. $6\frac{\sqrt{3}}{32}$

Answer: 1



[Watch Video Solution](#)

91. The maximum area of the rectangle whose sides pass through the vertices of a given rectangle of sides a and b is $2(ab)$ (b) $\frac{1}{2}(a + b)^2$
 $\frac{1}{2}(a^2 + b^2)$ (d) *none of these*

A. $2(ab)$

B. $\frac{1}{2}(a + b)^2$

C. $\frac{1}{2}(a + b)^2$

D. none of these

Answer: 2



[Watch Video Solution](#)

92. The base of prism is equilateral triangle. The distance from the centre of one base to one of the vertices of the other base is l . Then altitude of the prism for which the volume is greatest is (a) $\frac{l}{2}$ (b) $\frac{l}{\sqrt{3}}$ (c) $\frac{l}{3}$ (d) $\frac{l}{4}$

A. $\frac{l}{2}$

B. $\frac{l}{\sqrt{3}}$

C. $\frac{l}{3}$

D. $\frac{l}{4}$

Answer: 2



Watch Video Solution

Multiple correct answers type

1. Let $f(x) = \begin{cases} x^2 + 3x, & -1 \leq x < 0 \\ -\sin x, & 0 \leq x < \pi/2 \\ -1 - \cos x, & \frac{\pi}{2} \leq x \leq \pi \end{cases}$. Draw the graph of the

function and find the following

(a) Range of the function

(b) Point of inflection

(c) Point of local minima

A. $f(x)$ has global minimum value -2

B. global maximum value occurs at $x=0$

C. global maximum value occurs at $x=\pi$

D. $x = \pi/2$ is point of local minima

Answer: 1,2,3,4



Watch Video Solution

2. Let $f(x) = x^4 - 4x^3 + 6x^2 - 4x + 1$. Then, (a) f increase on $[1, \infty]$

(b) f decreases on $[1, \infty]$ (c) f has a minimum at $x = 1$ (d) f has neither

maximum nor minimum

- A. f increases on $[1, \infty]$
- B. f decreases on $[1, \infty]$
- C. f has a minimum at $x=1$
- D. f has neither maximum nor minimum

Answer: 1,3



Watch Video Solution

3. Let $f(x) = 2x - \sin x$ and $g(x) = 3\sqrt{x}$. Then

- A. range of $g \circ f$ is \mathbb{R}
- B. $g \circ f$ is one-one
- C. both f and g are one-one
- D. both f and g are onto

Answer: 1,2,3,4



Watch Video Solution

4. about to only mathematics

A. increases in $[0, \infty)$

B. idecreases in $[0, \infty)$

C. neither increases nor decreases in $[0, \infty)$

D. increases in $(-\infty, \infty)$

Answer: 1,4



View Text Solution

5. Let $f(x) = |x^2 - 3x - 4|$, $-1 \leq x \leq 4$ Then $f(x)$ is monotonically increasing in $\left[-1, \frac{3}{2}\right]$ $f(x)$ monotonically decreasing in $\left(\frac{3}{2}, 4\right)$ the maximum value of $f(x)$ is $\frac{25}{4}$ the minimum value of $f(x)$ is 0

A. $f(x)$ is monotonically increasing in $[-1, 3/2]$

B. $f(x)$ is monotonically decreasing in $(3/2, 4]$

C. the maximum value of $f(x)$ is $\frac{25}{4}$

D. the minimum value of $f(x)$ is 0

Answer: 1,2,3,4



Watch Video Solution

6. If $f(x) = \int_0^x \frac{\sin t}{t} dt, x > 0$, then

A. $f(x)$ has a local maxima at $x = n\pi (n = 2k, k \in I^+)$

B. $f(x)$ has a local minimum at $x = n\pi (n = 2k, k \in I^+)$

C. $f(x)$ has neither maxima nor minima at $x = n\pi (n \in I^+)$

D. $f(x)$ has local maxima at $x = n\pi (n = 2k - 1, k \in I^+)$

Answer: 2,4



Watch Video Solution

7. The values of parameter a for which the point of minimum of the function $f(x) = 1 + a^2x - x^3$ satisfies the inequality $\frac{x^2 + x + 2}{x^2 + 5x + 6} < 0$ are (a) $(2\sqrt{3}, 3\sqrt{3})$ (b) $(-3\sqrt{3}, -2\sqrt{3})$ (c) $(-2\sqrt{3}, 3\sqrt{3})$ (d) $(-2\sqrt{2}, 2\sqrt{3})$

A. $2\sqrt{3}, 3\sqrt{3}$

B. $-3\sqrt{3}, -2\sqrt{3}$

C. $-2\sqrt{3}, 3\sqrt{3}$

D. $-3\sqrt{2}, 2\sqrt{3}$

Answer: 1,2

 [Watch Video Solution](#)

8. Let $f(x) = ax^2 - b|x|$, where a and b are constants. Then at $x = 0$, $f(x)$ has a maxima whenever $a > 0, b > 0$ a maxima whenever $a > 0, b < 0$ minima whenever $a < 0, b < 0$ neither a maxima nor a minima whenever $a < 0, b > 0$

A. a maxima whenever $a > 0, b > 0$

B. a maxima whenever $a > 0, b < 0$

C. minima whenever $a > 0, b < 0$

D. neither a maxima nor a minima whenever $a > 0, b < 0$

Answer: 1,3



Watch Video Solution

9. The function $y = \frac{2x - 1}{x - 2}, (x \neq 2)$ is its own inverse decrease at all values of x in the domain has a graph entirely above the $x - a$ is unbounded

A. is its own inverse

B. decreases at all values of x in the domain

C. has a graph entirely above the x axis

D. is unbounded

Answer: 1,2,4



Watch Video Solution

10. Let $f(x) = ax^2 + bx + c$ and $f(-1) < 1$, $f(1) > -1$, $f(3) < -4$ and $a \neq 0$, then

- A. (a) $a > 0$
- B. (b) $a < 0$
- C. (c) Sign of a can't be determined
- D. (a) none of the above

Answer: 2,3



Watch Video Solution

11. If $f(x) = x^3 - x^2 + 100x + 2002$, then $f(1000) > f(1001)$
 $f\left(\frac{1}{2000}\right) > f\left(\frac{1}{2001}\right)$ $f(x-1) > f(x-2)$ $f(2x-3) > f(2x)$

A. $f(1000) < f(1001)$

B. $f\left(\frac{1}{2000}\right) > f\left(\frac{1}{2001}\right)$

C. $f(x - 1) > f(x - 2)$

D. $f(2x - 3) > f(2x)$

Answer: 2,3



Watch Video Solution

12. If $f'(x) = g(x)(x - a)^2$, where $g(a) \neq 0$, and g is continuous at $x = a$, then (a) f is increasing in the neighbourhood of a if $g(a) > 0$ (b) f is increasing in the neighbourhood of a if $g(a) < 0$ (c) f is decreasing in the neighbourhood of a if $g(a) > 0$ (d) f is decreasing in the neighbourhood of a if $g(a) < 0$

A. f is increasing in the neighborhood of a if $g(x) > 0$

B. f is increasing in the neighborhood of a if $g(x) < 0$

C. f is decreasing in the neighborhood of a if $g(x) > 0$

D. f is decreasing in the neighborhood of a if $g(x) < 0$

Answer: 1,4



Watch Video Solution

13. The value of a for which the function $f(x) = (4a - 3)(x + \log 5) + 2(a - 7)\cot\left(\frac{x}{2}\right)\sin^2\left(\frac{x}{2}\right)$ does not possess critical points is (a) $\left(-\infty, -\frac{4}{3}\right)$ (b) $(-\infty, -1)$ (c) $[1, \infty)$ (d) $(2, \infty)$

A. $(-\infty, -4/3)$

B. $(-\infty, -1)$

C. $[1, \infty)$

D. $(2, \infty)$

Answer: 1,4



Watch Video Solution

14. Let $f(x) = (x^2 - 1)^{n+1} \cdot (x^2 + x + 1)$. Then $f(x)$ has local extremum at $x = 1$, when n is (A) $n = 2$ (B) $n = 4$ (C) $n = 3$ (D) $n = 5$

A. a maxima at $x=1$ if n is odd

B. a maxima at $x=1$ if n is even

C. a minima $x=1$ if n is even

D. a minima at $x=2$ if n is even

Answer: 1,3,4



[View Text Solution](#)

15. Let $f(x) = \sin x + ax + b$. Then which of the following is/are true?

(a) $f(x) = 0$ has only one real root which is positive if $a > 1, b < 0$. (b)

$f(x) = 0$ has only one real root which is negative if $a > 1, b < 0$. (c)

$f(x) = 0$ has only one real root which is negative if $a > 1, b > 0$. (d)

none of these

A. $f(x) = 0$ has only one real root which is positive if $a > 1, b < 0$

B. $f(x) = 0$ has only one real root which is negative if $a > 1, b > 0$

C. $f(x) = 0$ has only one real root which is negative if $a < -1, b < 0$

D. None of these

Answer: 1,2,3



Watch Video Solution

16. The function $\frac{\sin(x+a)}{\sin(x+b)}$ has no maxima or minima if

$b - a = n\pi, n \in I$ $b - a = (2n + 1)\pi, n \in I$ $b - a = 2n\pi, n \in I$ (d)

none of these

A. $b - a = n\pi, n \in I$

B. $b - a = (2n + 1)\pi, n \in I$

C. $b - a = 2n\pi, n \in I$

D. none of these

Answer: 1,2,3



Watch Video Solution

17. Consider $f(x) = ax^4 + cx^2 + dx + e$ has no point of inflection. Then which of the following is/are possible? (a) $a > 0, c < 0$ (b) $a < 0, c > 0$ (c) $a, c < 0$ (d) $a, c > 0$

A. $a > 0, c < 0$

B. $a < 0, c > 0$

C. $a, c < 0$

D. $a, c > 0$

Answer: 3,4



Watch Video Solution

18. Let $f(x) = \begin{cases} \frac{(x-1)(6x-1)}{2x-1}, & \text{if } x \neq \frac{1}{2} \\ 0, & \text{if } x = \frac{1}{2} \end{cases}$ Then at $x = \frac{1}{2}$, which of the following is/are not true? f has a local maxima f has a local minima f has an inflection point. f has a removable discontinuity.

- A. f has a local maxima
- B. f has a local minima
- C. f has an inflection point
- D. f has a removable discontinuity

Answer: 1,2,4

 [Watch Video Solution](#)

19. In which of the following graphs is $x = c$ the point of inflection?

figure (b) figure (c) figure (d) figure

A. 

B. 

C. 

D. 

Answer: 1,2,4



View Text Solution

20. Let $f(x)$ be an increasing function defined on $(0, \infty)$. If $f(2a^2 + a + 1) > f(3a^2 - 4a + 1)$, then the possible integers in the range of a is/are (a) 1 (b) 2 (c) 3 (d) 4

A. 1

B. 2

C. 3

D. 4

Answer: 2,3,4



Watch Video Solution

21. Let $f(x) = (x - 1)^4(x - 2)^n$, $n \in \mathbb{N}$. Then $f(x)$ has (a) a maximum at $x = 1$ if n is odd (b) a maximum at $x = 1$ if n is even (c) a minimum at $x = 1$ if n is even (d) a minima at $x = 2$ if n is even

A. local maximum , if n is odd

B. local minimum, if n is odd

C. local maximum if n is even

D. local minimum if n is even

Answer: 1,4



Watch Video Solution

22. For the cubic function $f(x) = 2x^3 + 9x^2 + 12x + 1$, which one of the following statement/statements hold good? 1. $f(x)$ is non-monotonic. 2. $f(x)$ increases in $(-\infty, -2) \cup (-1, \infty)$ and decreases

in $(-2, -1)$ 3. $f: \overrightarrow{RR}$ is bijective. 4. Inflection point occurs at $x = -\frac{3}{2}$.

A. $f(x)$ is non monotonic

B. $f(x)$ increases in $(-\infty, -2) \cup (-1, \infty)$ and decreases in $(-2, -1)$

C. $f: \mathbb{R} \rightarrow \mathbb{R}$ is objective

D. O inflection point occurs at $x = -3/2$

Answer: 1,2,4



Watch Video Solution

23. Let $f(x) = a_5x^5 + a_4x^4 + a_3x^3 + a_2x^2 + a_1x$, where a_i 's are real and $f(x) = 0$ has a positive root α_0 . Then $f'(x) = 0$ has a positive root α_1 such that $0 < \alpha_1 < \alpha_0$

A. $f(x)=0$ has a root α_1 such that $0 < \alpha_1 < \alpha_0$

B. $f(x) = 0$ has at least two real roots

C. $f(x) = 0$ has at least one real root

D. none of these

Answer: 1,2,3



Watch Video Solution

24. If $f(x)$ and $g(x)$ are two positive and increasing functions, then which of the following is not always true? (a) $[f(x)]^{g(x)}$ is always increasing (b) $[f(x)]^{g(x)}$ is decreasing, when $f(x) < 1$ (c) $[f(x)]^{g(x)}$ is increasing, then $f(x) > 1$. (d) If $f(x) > 1$, then $[f(x)]^{g(x)}$ is increasing.

- A. $[f(x)]^{g(x)}$ is always increasing
- B. $[f(x)]^{g(x)}$ is decreasing when $f(x) < 1$
- C. If $[f(x)]^{g(x)}$ is increasing then $f(x) > 1$
- D. If $f(x) > 1$, then $[f(x)]^{g(x)}$ is increasing

Answer: 1,2,3



Watch Video Solution

25. An extremum of the function

$$f(x) = \frac{2-x}{\pi} \cos \pi(x+3) + \frac{1}{\pi^2} \sin \pi(x+3) \text{ where } x \in (0, 4) \text{ occurs}$$

at (a) $x = 1$ (b) $x = 2$ (c) $x = 3$ (d) $x = \pi$

A. $x=1$

B. $x=2$

C. $x=3$

D. $x=\pi$

Answer: 1,3



[Watch Video Solution](#)

26. For the function $f(x) = x^4(12(\log)_e x - 7)$, the point (1,7) is the point of inflection. $x = e^{\frac{1}{3}}$ is the point of minima the graph is concave downwards in (0,1) the graph is concave upwards in $(1, \infty)$

A. the point (1,7) is the point of minima

B. $x=e^{1/3}$ is the point of minima

C. the graph is concave downwards in $(0,1)$

D. the graph is concave upwards in $(1, \infty)$

Answer: 1,2,3,4



Watch Video Solution

27. Let $f(x) = \log(2x - x^2) + \frac{\sin(\pi x)}{2}$. Then which of the following is/are true? Graph of f is symmetrical about the line $x = 1$ Maximum value of f is 1. Absolute minimum value of f does not exist. none of these

A. Graph of f is symmetrical about the line $x=1$

B. maximum value of f is 1

C. absolute minimum value of f does not exist

D. none of these

Answer: 1,2,3

 [Watch Video Solution](#)

28. Which of the following hold(s) good for the function

$$f(x) = 2x - 3x^{\frac{2}{3}} \quad \forall x \in \mathbb{R}$$

- A. $f(x)$ has two points of extremum
- B. $f(x)$ is convave upward $\forall x \in \mathbb{R}$
- C. $f(x)$ is non differentiable function
- D. $f(x)$ is continuous function

Answer: 1,2,3,4

 [Watch Video Solution](#)

29. For the function $f(x) = \frac{e^x}{1 + e^x}$, which of the following hold good?

f is monotonic in its entire domain. Maximum of f is not attained even though f is bounded f has a point of inflection. f has one asymptote.

- A. f is monotonic in its entire domain
- B. maximum of f is not attained even though
- C. f is bounded
- D. f has a point of inflection

Answer: 1,2,3

 [Watch Video Solution](#)

- 30.** Which of the following is true about point of extremum $x = a$ of function $y = f(x)$? At $x = a$, function $y = f(x)$ may be discontinuous. At $x = a$, function $y = f(x)$ may be continuous but non-differentiable. At $x = a$, function $y = f(x)$ may have point of inflection. none of these
- A. At $x = a$, function $y = f(x)$ may be discontinuous
 - B. At $x = a$ function $y = f(x)$ may be continuous but non differentiable
 - C. At $x = a$ function $y = f(x)$ may have point of inflection
 - D. none of these

Answer: 1,2,3



Watch Video Solution

31. Which of the following function has point of extremum at $x = 0$?

$$f(x) = e^{-|x|} \quad f(x) = \sin|x| \quad f(x) = \begin{cases} x^2 + 4x + 3, & x < 0 \\ -x, & x \geq 0 \end{cases}$$

$f(x) = \begin{cases} |x|, & x < 0 \\ \{x\}, & 0 \leq x < 1 \end{cases}$ (where $\{x\}$ represents fractional part function).

A. $f(x) = e^{-|x|}$

B. $f(x) = \sin |x|$

C. $f(x) = \begin{cases} x^2 + 4x + 3 & x < 0 \\ -x & x \geq 0 \end{cases}$

D. $f(x) = \begin{cases} x^2 + 4x + 3 & x < 0 \\ -x & x \geq 0 \end{cases}$

Answer: 1,2,4



Watch Video Solution

32. Which of the following function/functions has/have point of inflection? $f(x) = x^{\frac{6}{7}}$ (b) $f(x) = x^6$ $f(x) = \cos x + 2x$ (d) $f(x) = x|x|$

A. $f(x)=x^{6/7}$

B. $f(x) = x^6$

C. $f(x)=\cos x+2x$

D. $f(x)=x|x|$

Answer: 3,4



Watch Video Solution

33. The function $f(x) = x^2 + \frac{\lambda}{x}$ has a minimum at $x = 2$ if $\lambda = 16$
maximum at $x = 2$ if $\lambda = 16$ maximum for no real value of λ point of
inflection at $x = 1$ if $\lambda = -1$

A. minimum at $x=2$ if $\lambda =16$

B. maximum at $x =2$ if $\lambda =16$

C. maximum for no real value of λ

D. point of inflection at $x=1$ if $\lambda=-1$

Answer: 1,3,4



Watch Video Solution

34. The function $f(x) = x^{\frac{1}{3}}(x - 1)$ has two inflection points has one point of extremum is non-differentiable has range $\left[-3 \times 2^{-\frac{8}{3}}, \infty \right)$

A. has two inflection points

B. has one point of extremum

C. is non differentiable

D. has range $\left[-3 \times 2^{-8/3}, \infty \right)$

Answer: 1,2,3,4



Watch Video Solution

35. Let f be the function $f(x) = \cos x - \left(1 - \frac{x^2}{2}\right)$. Then (a) $f(x)$ is an increasing function in $(0, \infty)$ (b) $f(x)$ is a decreasing function in $(-\infty, \infty)$ (c) $f(x)$ is an increasing function in $(-\infty, \infty)$ (d) $f(x)$ is a decreasing function in $(-\infty, 0)$

- A. $f(x)$ is an increasing function in $(0, \infty)$
- B. $f(x)$ is a decreasing function in $(-\infty, \infty)$
- C. $f(x)$ is an increasing function in $(-\infty, \infty)$
- D. $f(x)$ is a decreasing function in $(-\infty, 0)$

Answer: 1,4

 [Watch Video Solution](#)

36. Consider the function $f(x) = x \cos x - \sin x$. Then identify the statement which is correct. f is neither odd nor even. f is monotonic decreasing at $x = 0$ f has a maxima at $x = \pi$ f has a minima at $x = -\pi$

A. f is odd

B. f is monotonic decreasing at $x=0$

C. f has point of inflection at $x=0$

D. f has a maxima at $x= \pi$

Answer: 1,2,3



Watch Video Solution

37. If $f(x) = \frac{x^2}{2 - 2 \cos x}$; $g(x) = \frac{x^2}{6x - 6 \sin x}$ where $0 < x < 1$, then

(A) both ' f ' and ' g ' are increasing functions

A. f is increasing function

B. g is increasing function

C. f is decreasing function

D. g is decreasing function

Answer: 1,4



[View Text Solution](#)

38. Find the greatest value of $f(x) = \frac{1}{2ax - x^2 - 5a^2}$ in $[-3, 5]$ depending upon the parameter a .

A. $f(5)$ if $a = 1$

B. $f(-3)$ if $a = 1$

C. $f(5)$ if $a < 1$

D. $f(-3)$ if $a > 1$

Answer: 1,2,3,4



[Watch Video Solution](#)

39. For any acute angled $\triangle ABC$, $\frac{\sin A}{A} + \frac{\sin B}{B} + \frac{\sin C}{C}$ can

A. 1

B. 2

C. 3

D. 4

Answer: 1,2



[View Text Solution](#)

40. Let $f(x)$ be a non negative continuous and bounded function for all $x \geq 0$. If $(\cos x)f(x) < (\sin x - \cos x)f(x) \forall x \geq 0$, then which of the following is/are correct?

A. $f(6) + f(5) > 0$

B. $x^2 - 3x + 2 + f(7) = 0$ has 2 distinct solution

C. $f(5)f(7)-f(5)=0$

D. $\lim_{x \rightarrow 6} \frac{f(x) - \sin(\pi x)}{x - 6} = 1$

Answer: 2,3



[View Text Solution](#)

41. A rectangular sheet of fixed perimeter with sides having their lengths in the ratio 8:15 is converted into an open rectangular box by folding after removing squares of equal area from all four corners. If the total area of removed squares is 100, the resulting box has maximum volume. Then the length of the sides of the rectangular sheet are 24 (b) 32 (c) 45 (d) 60

A. 24

B. 32

C. 45

D. 60

Answer: 1,3



Watch Video Solution

42. The function $f(x) = 2|x| + |x + 2| = ||x|2| - 2|x||$ has a local minimum or a local maximum at $x = -2$ (b) $-\frac{2}{3}$ (c) 2 (d) $\frac{2}{3}$

A. -2

B. $-\frac{2}{3}$

C. 2

D. $\frac{2}{3}$

Answer: 1,2



[View Text Solution](#)

43. about to only mathematics

A. $(f(c))^2 + 3f(c) = (g(c))^2 + 3g(c)$ for some $c \in [0,1]$

B. $(f(c))^2 + f(c) = (g(c))^2 + 3g(c)$ for some $c \in [0,1]$

C. $(f(c))^2 + 3f(c) = (g(c))^2 + g(c)$ for some $c \in [0,1]$

D. $(f(c))^2 = (g(c))^2$ for some $c \in [0,1]$

Answer: 1,4



View Text Solution

44. Let $a \in R$ and let $f: R \rightarrow R$ be given by $f(x) = x^5 - 5x + a$, then (a) $f(x)$ has three real roots if $a > 4$ (b) $f(x)$ has only one real root if $a > 4$ (c) $f(x)$ has three real roots if $a < -4$ (d) $f(x)$ has three real roots if $-4 < a < 4$

A. $f(x)$ has three real roots if $a > 4$

B. $f(x)$ has only one real root if $a > 4$

C. $f(x)$ has three real roots if $a < -4$

D. $f(x)$ has three real roots if $-4 < a < 4$

Answer: 2,4



Watch Video Solution

45. Let $f: \mathbb{R} \rightarrow (0, \infty)$ and $g: \mathbb{R} \rightarrow \mathbb{R}$ be twice differentiable functions such that f'' and g'' are continuous functions of \mathbb{R} suppose $f'(2) = g(2) = 0$, $f''(2) \neq 0$ and $g'(2) \neq 0$. If

$$\lim_{x \rightarrow 2} \frac{f(x)g(x)}{f'(x)g'(x)} = 1, \text{ then}$$

- A. f has a local minimum at $x=2$
- B. f has a local maximum at $x=2$
- C. $f''(2) > f(x)$
- D. $\forall x, f'(x) - f''(x) = 0$ for at least one x in \mathbb{R}

Answer: 1,4

 [View Text Solution](#)

46. If $f: \mathbb{R} \rightarrow \mathbb{R}$ is a differentiable function such that $f(x) > 2f(x)$ for all $x \in \mathbb{R}$ and $f(0) = 1$, then

- A. $f(x) > e^{2x} \in (0, \infty)$

B. $f(x)$ is decreasing in $(0, \infty)$

C. $f(x)$ is increasing in $(0, \infty)$

D. $f'(x) < e^{2x}$ in $(0, \infty)$

Answer: 1,3



Watch Video Solution

47. If $f(x) = \begin{vmatrix} \cos(2x) & \cos(2x) & \sin(2x) \\ -\cos x & \cos x & -\sin x \\ \sin x & \sin x & \cos x \end{vmatrix}$, then

A. $f'(x) = 0$ at exactly three points in $(-\pi, \pi)$

B. $f(x)$ attains its maximum at $x=0$

C. $f(x)$ attains its minimum at $x=0$

D. $f'(x) = 0$ at more than three points in $(-\pi, \pi)$

Answer: 2,4



View Text Solution

48. Let $f: (0, \pi) \rightarrow \mathbb{R}$ be a twice differentiable function such that

$$\lim_{t \rightarrow x} \frac{f(x)\sin t - f(t)\sin x}{t - x} = \sin^2 x \quad \text{for all } x \in (0, \pi). \quad \text{If}$$
$$f\left(\frac{\pi}{6}\right) = \left(-\frac{\pi}{12}\right) \text{ then which of the following statement (s) is (are)}$$

TRUE?

A. $f\left(x\frac{\pi}{4}\right) = \frac{\pi}{4\sqrt{2}}$

B. $f(x) < \frac{x^4}{6} - x^2 f$ or $\text{all } x \in (0, \pi)$

C. There exist $\alpha \in (0, \pi)$ such that $f'(\alpha) = 0$

D. $f''\left(\frac{\pi}{2}\right) + f\left(\frac{\pi}{2}\right) = 0$

Answer: 2,3,4



View Text Solution

Linked comprehension type

1. $f(x) = \sin^{-1} x + x^2 - 3x + \frac{x^3}{3}, x \in [0, 1]$

A. $f(x)$ has a point of maxima

B. $f(x)$ has a point of minima

C. $f(x)$ is increasing

D. $f(x)$ is decreasing

Answer: 2

 [View Text Solution](#)

2. which of the following is true for $x \in [0, 1]$?

A. $\sin^{-1} x + x^2 - x \frac{9 - x^2}{3} \leq 0$

B. $\sin^{-1} x + x^2 - x \frac{9 - x^2}{3} \geq 0$

C. $\sin^{-1} x + x^2 - x \frac{9 - x^2}{3} \leq 0$

D. $\sin^{-1} x + x^2 - x \frac{9 - x^2}{3} \geq 0$

Answer: 1

 [View Text Solution](#)

3. Let $f'(\sin x) < 0$ and $f''(\sin x) > 0 \forall x \in \left(0, \frac{\pi}{2}\right)$ and $g(x)$

$$= f(\sin x) + f(\cos x)$$

which of the following is true?

- A. g' is increasing
- B. g' is decreasing
- C. g' has a point of minima
- D. g' has a point of maxima

Answer: 1



[View Text Solution](#)

4. Let $f'(\sin x) < 0$ and $f''(\sin x) > 0 \forall x \in \left(0, \frac{\pi}{2}\right)$ and $g(x)$

$$= f(\sin x) + f(\cos x)$$

which of the following is true?

A. $g(x)$ is decreasing in $\left(\frac{\pi}{4}, \frac{\pi}{2}\right)$

B. $g(x)$ increasing in $\left(0, \frac{\pi}{4}\right)$

C. $g(x)$ is nonotonically increasing in $\left(0, \frac{\pi}{2}\right)$

D. none of these

Answer: 4

 [View Text Solution](#)

5. Let $f'(\sin x) < 0$ and $f''(\sin x) > 0 \forall x \in \left(0, \frac{\pi}{2}\right)$ and $g(x)$

$=f(\sin x)+f(\cos x)$

which of the following is true?

A. $f(x)$ is discounting function for all ordered pairs (a,b)

B. $f(x)$ is contionuous for finite number of ordered pairs $(,ab)$

C. $f(x)$ can be differentiable

D. $f(x)$ is continous for infinite ordered pairs (a,b)

Answer: 4



[View Text Solution](#)

6. Let $f'(\sin x) < 0$ and $f''(\sin x) > 0 \forall x \in \left(0, \frac{\pi}{2}\right)$ and $g(x) = f(\sin x) + f(\cos x)$

If $x = 3$ is the only point of minima in its neighborhood and $x = 4$ is neither a point of maxima nor a point minima, then which of the following can be true?

A. $a > 0, b < 0$

B. $a < 0, b < 0$

C. $a > 0, b \in \mathbb{R}$

D. none of these

Answer: 1



[View Text Solution](#)

7. Let $f'(\sin x) < 0$ and $f''(\sin x) > 0 \forall x \in \left(0, \frac{\pi}{2}\right)$ and $g(x) = f(\sin x) + f(\cos x)$

If $x = 3$ is the only point of minima in its neighborhood and $x = 4$ is neither a point of maxima nor a point minima, then which of the following can be true?

A. $a < 0, b > 0$

B. $a > 0, b < 0$

C. $a < 0, b < 0$

D. not possible

Answer: 4



[View Text Solution](#)

8. Let $f'(\sin x) < 0$ and $f''(\sin x) > 0 \forall x \in \left(0, \frac{\pi}{2}\right)$ and $g(x) = f(\sin x) + f(\cos x)$

If $x = 3$ is the only point of minima in its neighborhood and $x = 4$ is neither

a point of maxima nor a point minima, then which of the following can be true?

A. $a < 0, b > 0$

B. $a > 0, b < 0$

C. $a > 0, b > 0$

D. not possible

Answer: 3



[View Text Solution](#)

9. If $\phi(x)$ is a differentiable real valued function satisfying $\phi(x) + 2\phi' \leq 1$,

then it can be adjucted as

$$e^{2x}\phi(x) + 2e^{2x}\phi'(x) \leq e^{2x} \text{ or } \frac{d}{dx} \left(e^{2x}\phi(x) - \frac{e^{2x}}{2} \right) \leq \text{ or } \frac{d}{dx} e^{2x} \left(\phi(x) - \frac{1}{2} \right) \leq 0$$

Here e^{2x} is called integrating factor which helps in creating single differential coefficient as shown above. Answer the following question:

If $p(1)=0$ and $dP \frac{x}{dx} < P(x)$ for all $x \geq 1$ then

A. $P(x) > 0 \forall x > 1$

B. $P(x)$ is a constant function

C. $P(x) < 0 \forall x > 1$

D. none of these

Answer: 1

 [View Text Solution](#)

10. If $H(x_0) = 0$ for some $x = x_0$ and $\frac{d}{dx}H(x) > 2cxH(x)$ for all $x \geq x_0$, where $c > 0$, then prove that $H(x)$ cannot be zero for any $x > x_0$.

A. $H(x) = 0$ has root for $x > x_0$

B. $H(x) = 0$ has no root for $x > x_0$

C. $H(x)$ is a constant function

D. none of these

Answer: 2



[View Text Solution](#)

11. Let $h(x) = f(x) - a(f(x))^3$ for every real number x

$h(x)$ increase as $f(x)$ increases for all real values of x if

A. $a \in (0, 3)$

B. $a \in (-2, 2)$

C. $[3, \infty)$

D. none of these

Answer: 1



[Watch Video Solution](#)

12. Let $h(x) = f(x) - a(f(x))^3$ for every real number x

$h(x)$ increase as $f(x)$ increases for all real values of x if

A. $a \in (0, 3)$

B. $a \in (-2, 2)$

C. $[3, \infty)$

D. none of these

Answer: 4



[View Text Solution](#)

13. If $f(x)$ is strictly increasing function then $h(x)$ is non monotonic function given

A. $a \in (0, 3)$

B. $a \in (-2, 2)$

C. $(3, \infty)$

D. $a \in (-\infty, 0) \cup (3, \infty)$

Answer: 4

14. Let $f(x) = x^3 - 9x^2 + 24x + c = 0$ have three real and distinct roots α, β and λ .

(i) Find the possible values of c .

(ii) If $[\alpha] + [\beta] + [\lambda] = 8$, then find the values of c , where $[\cdot]$ represents the greatest integer function.

(ii) If $[\alpha] + [\beta] + [\lambda] = 7$, then find the values of c , where $[\cdot]$ represents the greatest integer function.s

A. (-20 , -16)

B. (-20,-18)

C. (-18,-16)

D. none of these

Answer: 1

15. Let $f(x) = x^3 - 9x^2 + 24x + c = 0$ have three real and distinct roots α , β and λ .

(i) Find the possible values of c .

(ii) If $[\alpha] + [\beta] + [\lambda] = 8$, then find the values of c , where $[\cdot]$ represents the greatest integer function.

(ii) If $[\alpha] + [\beta] + [\lambda] = 7$, then find the values of c , where $[\cdot]$ represents the greatest integer function.s

A. (-20, -16)

B. (-20,-18)

C. (-18,-16)

D. none of these

Answer: 3



Watch Video Solution

16. Let $f(x) = x^3 - 9x^2 + 24x + c = 0$ have three real and distinct roots α , β and λ .

(i) Find the possible values of c .

(ii) If $[\alpha] + [\beta] + [\lambda] = 8$, then find the values of c , where $[\cdot]$ represents the greatest integer function.

(ii) If $[\alpha] + [\beta] + [\lambda] = 7$, then find the values of c , where $[\cdot]$ represents the greatest integer function.s

A. (-20,-16)

B. (-20,-18)

C. (-18,-16)

D. none of these

Answer: 2



Watch Video Solution

17. consider the graph of $y=g(x)=f'(x)$ given that $f(c) = 0$, where $y=f(x)$ is a polynomial function



The graph of $y=f(x)$ will intersect the x axis

- A. twice
- B. once
- C. never
- D. none of these

Answer: 2



[View Text Solution](#)

18. Consider the graph of $y = g(x) = f'(x)$ given that $f(c) = 0$, where $y = f(x)$ is a polynomial function



The equation $f(x) = 0, a \leq x \leq b$, has

- A. four real roots
- B. no real roots
- C. two distinct real roots
- D. at least three repeated roots

Answer: 4

 [View Text Solution](#)

19. consider the graph of $y=g(x)=f'(x)$ given that $f(c) = 0$, where $y=f(x)$ is a polynomial function



The graph of $y = f(x)$, $a \leq x \leq b$ has

- A. two points of inflection
- B. one point of inflection
- C. no point of inflection
- D. none of these

Answer: 2



[View Text Solution](#)

20. consider the graph of $y=g(x)=f'(x)$ given that $f(c) = 0$, where $y=f(x)$ is a polynomial function



The function $y = f(x)$, $a \leq x \leq b$, has

- A. exactly one local maxima
- B. one local minima and one maxima
- C. exactly one local minima
- D. none of these

Answer: 4



[View Text Solution](#)

21. consider the graph of $y=g(x)=f'(x)$ given that $f(c) = 0$, where $y=f(x)$ is a polynomial function



The equation $f''(x)=0$

- A. has no real root
- B. has at least one real root
- C. has at least two distinct roots
- D. none of these

Answer: 2



[View Text Solution](#)

22. Let $f(x) = 4x^2 - 4ax + a^2 - 2a + 2$ and the global minimum value of $f(x)$ for x in $[0,2]$ is equal to 3

The number of values of a for which the global minimum value equal to 3 for x in $[0,2]$ occurs at the endpoint of interval $[0,2]$ is

A. 1

B. 2

C. 3

D. 0

Answer: 2



[View Text Solution](#)

23. Let $f(x) = 4x^2 - 4ax + a^2 - 2a + 2$ and the global minimum value of $f(x)$ for x in $[0,2]$ is equal to 3

The number of values of a for which the global minimum value equal to 3 for x in $[0,2]$ occurs at the endpoint of interval $[0,2]$ is

A. 1

B. 2

C. 3

D. 0

Answer: 4



[View Text Solution](#)

24. Let $f(x) = 4x^2 - 4ax + a^2 - 2a + 2$ and the global minimum value of $f(x)$ for x in $[0,2]$ is equal to 3

The number of values of a for which the global minimum value equal to 3 for x in $[0,2]$ occurs at the endpoint of interval $[0,2]$ is

A. $a \leq 0$ or $a \geq 4$

B. $0 \leq a \leq 4$

C. $a > 0$

D. none of these

Answer: 1



[View Text Solution](#)

25. Let $f(x) = x^3 - 3(7 - a)x^2 - 3(9 - a^2)x + 2$

The values of parameter a if $f(x)$ has a negative point of local minimum are

A. π

B. $(-3, 3)$

C. $\left(-\infty, \frac{58}{14}\right)$

D. none of these

Answer: 1



Watch Video Solution

26. Let $f(x) = x^3 - 3(7 - a)x^2 - 3(9 - a^2)x + 2$

The values of parameter a if $f(x)$ has a positive point of local maxima are

A. π

B. $-\infty, -3 \cup \left(3, \frac{29}{7}\right)$

C. $-\infty, \frac{58}{14}$

D. none of these

Answer: 2



Watch Video Solution

27. Let $f(x) = x^3 - 3(7 - a)x^2 - 3(9 - a^2)x + 2$

The values of parameter a if $f(x)$ has points of extrema which are opposite in sign are

A. π

B. $(-3, 3)$

C. $\left(-\infty, \frac{58}{14}\right)$

D. none of these

Answer: 2



View Text Solution

28. consider the function $f(x) = 1 \left(1 + \frac{1}{x} \right)^x$

The domain of $f(x)$ is

- A. $(-1, 0) \cup (0, \infty)$
- B. $\mathbb{R} - \{0\}$
- C. $(-\infty, -1) \cup (0, \infty) \cup (0, \infty)$
- D. $(0, \infty)$

Answer: 3



[View Text Solution](#)

29. consider the function $f(x) = 1 \left(1 + \frac{1}{x} \right)^x$

The domain of $f(x)$ is

- A.
- B.

C.

D.



[View Text Solution](#)

30. consider the function $f(x) = \left(1 + \left(\frac{1}{x}\right)\right)^x$

The range of the function $f(x)$ is

A. $(0, \infty)$

B. $(-\infty, e)$

C. $1, \infty)$

D. $(1, e) \cup (e, \infty)$

Answer: 4



[View Text Solution](#)

31. consider the function $f(x) = x + \cos x$

which of the following is not true about $y = f(x)$?

- A. It is an increasing function
- B. It is a monotonic function
- C. It has infinite points of inflection s
- D. None of these

Answer: 4



[View Text Solution](#)

32. consider the function $f(x) = x + \cos x - a$

values of a which $f(x) = 0$ has exactly one positive root are

- A. (0,1)
- B. $(-\infty, 1)$
- C. (-1,1)

D. $(1, \infty)$

Answer: 4



[View Text Solution](#)

33. consider the function $f(x) = x + \cos x - a$

values of a for which $f(x) = 0$ has exactly one negative root are

A. $(0, 1)$

B. $(-\infty, 1)$

C. $(-1, 1)$

D. $(1, \infty)$

Answer: 2



[View Text Solution](#)

34. consider the function $f(X) = 3x^4 + 4x^3 - 12x^2$

$Y = f(X)$ increase in the interval

A. $(-1, 0) \cup (2, \infty)$

B. $(-\infty, 0) \cup (1, 2)$

C. $(-2, 0) \cup (1, \infty)$

D. none of these

Answer: 3



[View Text Solution](#)

35. consider the function $f(X) = 3x^4 + 4x^3 - 12x^2$

The range of the function $y = f(x)$ is

A. $(-\infty, \infty)$

B. $[-32, \infty)$

C. $[0, \infty)$

D. none of these

Answer: 2



[View Text Solution](#)

36. consider the function $f(x) = 3x^4 + 4x^3 - 12x^2$

The range of values of a for which $f(x) = a$ has no real

A. $(4, \infty)$

B. $(10, \infty)$

C. $(20, \infty)$

D. none of these

Answer: 4



[View Text Solution](#)

37. consider the function $f: R \rightarrow R, f(x) = \frac{x^2 - 6x + 4}{x^2 + 2x + 4}$

$f(x)$ is

- A. unbounded function
- B. one one function
- C. onto function
- D. none of these

Answer: 4



[Watch Video Solution](#)

38. consider the function $f: R \rightarrow R, f(x) = \frac{x^2 - 6x + 4}{x^2 + 2x + 4}$

which of the following is not true about $f(x)$?

- A. $f(x)$ has two points of extremum
- B. $f(x)$ has only one asymptote
- C. $f(x)$ is differentiable for all x in R

D. none of these

Answer: 4



[View Text Solution](#)

39. consider the function $f: R \rightarrow R, f(x) = \frac{x^2 - 6x + 4}{x^2 + 2x + 4}$

Range of $f(x)$ is

A. $\left(-\infty, -\frac{2}{3}\right] \cup [2, 0)$

B. $\left[\frac{-1}{3}, 5\right]$

C. $(-\infty, 2) \cup \left[\frac{7}{3}, \infty\right)$

D. $(20, \infty)$

Answer: 2



[Watch Video Solution](#)

40. Consider a polynomial $y = P(x)$ of the least degree passing through $A(-1,1)$ and whose graph has two points of inflection $B(1,2)$ and C with abscissa 0 at which the curve is inclined to the positive axis of abscissa at an angle of $\sec^{-1} \sqrt{2}$

The value of $P(2)$ is

A. -1

B. $-\frac{3}{2}$

C. $\frac{5}{2}$

D. $\frac{7}{2}$

Answer: 3



[View Text Solution](#)

41. Consider a polynomial $y = P(x)$ of the least degree passing through $A(-1,1)$ and whose graph has two points of inflection $B(1,2)$ and C with abscissa 0 at which the curve is inclined to the positive axis of abscissa at

an angle of $\sec^{-1} \sqrt{2}$

The value of $P(0)$ is

A. 1

B. 0

C. $\frac{3}{4}$

D. $\frac{1}{2}$

Answer: 4



[View Text Solution](#)

42. Consider a polynomial $y = P(x)$ of the least degree passing through $A(-1,1)$ and whose graph has two points of inflection $B(1,2)$ and C with abscissa 0 at which the curve is inclined to the positive axis of abscissa at an angle of $\sec^{-1} \sqrt{2}$

The equation $P(x) = 0$ has

A. three distinct real roots

B. one real root

C. three real roots such that one root is repeated

D. none of these

Answer: 3



[View Text Solution](#)

43. Let $f(x)$ be real valued continuous function on \mathbb{R} defined as $f(x) = x^2 e^{-|x|}$

The values of k for which the equation $x^2 e^{-|x|} = k$ has four real roots are

A. $0 < k < e$

B. $0 < k < \frac{8}{e^2}$

C. $0 < k < \frac{4}{e^2}$

D. none of these

Answer: 3



[View Text Solution](#)

44. Let $f(x)$ be real valued continuous function on \mathbb{R} defined as $f(x) = x^2 e^{-|x|}$

Number of points of inflection for $y = f(x)$ is (a) 1 (b) 2 (c) 3 (d) 4

A. $y = f(x)$ has two points of maxima

B. $y = f(x)$ has only one asymptote

C. $f(x) = 0$ has three real roots

D. none of these

Answer: 4



[View Text Solution](#)

45. Let $f(x)$ be real valued continuous function on \mathbb{R} defined as $f(x) = x^2 e^{-|x|}$

Number of points of inflection for $y = f(x)$ is (a) 1 (b) 2 (c) 3 (d) 4

A. 1

B. 2

C. 3

D. 4

Answer: 4



[View Text Solution](#)

46. $P(x)$ be a polynomial of degree 3 satisfying $P(-1) = 10$, $P(1) = -6$ and $p(x)$ has maxima at $x = -1$ and $p(x)$ has minima at $x=1$ then The value of $P(2)$ is
(a) -15 (b) -16 (c) -17 (d) -22 (c) -17 (d) -22

A. -15

B. -16

C. -17

D. -22

Answer: 3



[View Text Solution](#)

47. $P(x)$ be a polynomial of degree 3 satisfying $P(-1) = 10$, $P(1) = -6$ and $p(x)$ has maxima at $x = -1$ and $p(x)$ has minima at $x=1$ then The value of $P(1)$ is

A. -12

B. -10

C. 15

D. 21

Answer: 1



[Watch Video Solution](#)

48. The graph of $y = g(x) = f(x)$ is as shown in the following figure analyse this graph and answer the following question



The graph of $y=f(x)$ for $a < x < b$ has


- A. no point of extremum
- B. one point of extrema
- C. two points extrema
- D. can't say anything

Answer: 3



[View Text Solution](#)

49. The graph of $y = g(x) = f(x)$ is as shown in the following figure analyse this graph and answer the following question



Number of points of inflection on the graph of $y = f(x)$ for $a < x < b$ has

- A. 0
- B. 1
- C. 2
- D. can't say anything

Answer: 3



[View Text Solution](#)

50. The graph of $y = g(x) = f(x)$ is as shown in the following figure analyse this graph and answer the following question



Which of the following is not true about the graph of $y = f(x)$, $a < x < b$

- A. always increasing
- B. discontinuous at one point
- C. first increases then decreases
- D. none of these

Answer: 3



[View Text Solution](#)

Matrix Match Type

1. consider function $f(x) = x^4 - 14x^2 + 24x - 3$. Now match the following lists:



 **View Text Solution**

2. Match the following lists:

List I	List II
<p>a. If $f(x)$ is an integrable function for $x \in \left[\frac{\pi}{6}, \frac{\pi}{3} \right]$ and</p> $I_1 = \int_{\pi/6}^{\pi/3} \sec^2 \theta f(2 \sin 2\theta) d\theta, \text{ and}$ $I_2 = \int_{\pi/6}^{\pi/3} \operatorname{cosec}^2 \theta f(2 \sin 2\theta) d\theta, \text{ then } I_1/I_2 =$	<p>p. 3</p>
<p>b. If $f(x+1) = f(3+x) \forall x$, and the value of $\int_a^{a+b} f(x) dx$ is independent of a, then the value of b can be</p>	<p>q. 1</p>
<p>c. The value of $2 \int_1^4 \frac{\tan^{-1}[x^2]}{\tan^{-1}[x^2] + \tan^{-1}[25 + x^2 - 10x]} dx$ (where $[.]$ denotes the greatest integer function) is</p>	<p>r. 2</p>
<p>d. If $I = \int_0^2 \sqrt{x + \sqrt{x + \sqrt{x + \dots \infty}}} dx$ (where $x > 0$), then $[I]$ is equal to (where $[.]$ denotes the greatest integer function)</p>	<p>s. 4</p>



[View Text Solution](#)

3. Let $f(x) = (x - 1)^m(2 - x)^n$, $mn \in N$ and $m, n < 2$. Then match the following lists:



[View Text Solution](#)

4. The function $f(x) = \sqrt{ax^3 + bx^2 + cx + d}$ has its non-zero local minimum and local maximum values at $x = -2$ and $x = 2$, respectively. If a is a root of $x^2 - x - 6 = 0$, then find a, b, c and d .



[Watch Video Solution](#)

5. Match the following lists:

List I	List II
<p>a. If $f(x)$ is an integrable function for $x \in \left[\frac{\pi}{6}, \frac{\pi}{3} \right]$ and</p> $I_1 = \int_{\pi/6}^{\pi/3} \sec^2 \theta f(2 \sin 2\theta) d\theta, \text{ and}$ $I_2 = \int_{\pi/6}^{\pi/3} \operatorname{cosec}^2 \theta f(2 \sin 2\theta) d\theta, \text{ then } I_1/I_2 =$	<p>p. 3</p>
<p>b. If $f(x+1) = f(3+x) \forall x$, and the value of $\int_a^{a+b} f(x) dx$ is independent of a, then the value of b can be</p>	<p>q. 1</p>
<p>c. The value of $2 \int_1^4 \frac{\tan^{-1}[x^2]}{\tan^{-1}[x^2] + \tan^{-1}[25 + x^2 - 10x]} dx$ (where $[.]$ denotes the greatest integer function) is</p>	<p>r. 2</p>
<p>d. If $I = \int_0^2 \sqrt{x + \sqrt{x + \sqrt{x + \dots \infty}}} dx$ (where $x > 0$), then $[I]$ is equal to (where $[.]$ denotes the greatest integer function)</p>	<p>s. 4</p>

 [View Text Solution](#)

6. Match the following lists:

List I	List II
<p>a. If $f(x)$ is an integrable function for $x \in \left[\frac{\pi}{6}, \frac{\pi}{3} \right]$ and</p> $I_1 = \int_{\pi/6}^{\pi/3} \sec^2 \theta f(2 \sin 2\theta) d\theta, \text{ and}$ $I_2 = \int_{\pi/6}^{\pi/3} \operatorname{cosec}^2 \theta f(2 \sin 2\theta) d\theta, \text{ then } I_1/I_2 =$	<p>p. 3</p>
<p>b. If $f(x+1) = f(3+x) \forall x$, and the value of $\int_a^{a+b} f(x) dx$ is independent of a, then the value of b can be</p>	<p>q. 1</p>
<p>c. The value of $2 \int_1^4 \frac{\tan^{-1}[x^2]}{\tan^{-1}[x^2] + \tan^{-1}[25 + x^2 - 10x]} dx$ (where $[\cdot]$ denotes the greatest integer function) is</p>	<p>r. 2</p>
<p>d. If $I = \int_0^2 \sqrt{x + \sqrt{x + \sqrt{x + \dots \infty}}} dx$ (where $x > 0$), then $[I]$ is equal to (where $[\cdot]$ denotes the greatest integer function)</p>	<p>s. 4</p>



View Text Solution

7. Match the following lists:

List I	List II
<p>a. If $f(x)$ is an integrable function for $x \in \left[\frac{\pi}{6}, \frac{\pi}{3} \right]$ and</p> $I_1 = \int_{\pi/6}^{\pi/3} \sec^2 \theta f(2 \sin 2\theta) d\theta, \text{ and}$ $I_2 = \int_{\pi/6}^{\pi/3} \operatorname{cosec}^2 \theta f(2 \sin 2\theta) d\theta, \text{ then } I_1/I_2 =$	<p>p. 3</p>
<p>b. If $f(x+1) = f(3+x) \forall x$, and the value of $\int_a^{a+b} f(x) dx$ is independent of a, then the value of b can be</p>	<p>q. 1</p>
<p>c. The value of $2 \int_1^4 \frac{\tan^{-1}[x^2]}{\tan^{-1}[x^2] + \tan^{-1}[25 + x^2 - 10x]} dx$ (where $[\cdot]$ denotes the greatest integer function) is</p>	<p>r. 2</p>
<p>d. If $I = \int_0^2 \sqrt{x + \sqrt{x + \sqrt{x + \dots \infty}}} dx$ (where $x > 0$), then $[I]$ is equal to (where $[\cdot]$ denotes the greatest integer function)</p>	<p>s. 4</p>



View Text Solution

8. Match the following lists:

List I	List II
<p>a. If $f(x)$ is an integrable function for $x \in \left[\frac{\pi}{6}, \frac{\pi}{3} \right]$ and</p> $I_1 = \int_{\pi/6}^{\pi/3} \sec^2 \theta f(2 \sin 2\theta) d\theta, \text{ and}$ $I_2 = \int_{\pi/6}^{\pi/3} \operatorname{cosec}^2 \theta f(2 \sin 2\theta) d\theta, \text{ then } I_1/I_2 =$	<p>p. 3</p>
<p>b. If $f(x+1) = f(3+x) \forall x$, and the value of $\int_a^{a+b} f(x) dx$ is independent of a, then the value of b can be</p>	<p>q. 1</p>
<p>c. The value of $2 \int_1^4 \frac{\tan^{-1}[x^2]}{\tan^{-1}[x^2] + \tan^{-1}[25 + x^2 - 10x]} dx$ (where $[\cdot]$ denotes the greatest integer function) is</p>	<p>r. 2</p>
<p>d. If $I = \int_0^2 \sqrt{x + \sqrt{x + \sqrt{x + \dots \infty}}} dx$ (where $x > 0$), then $[I]$ is equal to (where $[\cdot]$ denotes the greatest integer function)</p>	<p>s. 4</p>



View Text Solution

9. $f(x)$ is polynomial function of degree 6, which satisfies $(\lim)_{x \rightarrow 0} \left(1 + \frac{f(x)}{x^3}\right)^{\frac{1}{x}} = e^2$ and has local maximum at $x = 1$ and local minimum at $x = 0$ and $x = 2$. then $5f(3)$ is equal to

 [Watch Video Solution](#)

10. Match the statement /expressions in List I with the open intervals in

List II



 [View Text Solution](#)

11. Let $f(x) = x + \log_e x - x \log_e x, x \in (0, \infty)$

List I contains information about zero of $f(x)$, $f'(x)$ and $f''(x)$

List II contains information about the limiting behaviour of $f(x)$, $f'(x)$ and $f''(x)$ at infinity

List III contains information about increasing /decreasing nature of $f(x)$

and $f'(x)$



which of the following options is the only CORRECT combination?

A. (iv)(i)(S)

B. (I)(ii)(R)

C. (III)(iv)(P)

D. (II)(ii)(S)

Answer: 4



[View Text Solution](#)

Numerical Value Type

1. If α is an integer satisfying $|\alpha| \leq 4 - |[x]|$, where x is a real number for which $2x \tan^{-1} x$ is greater than or equal to $\ln(1 + x^2)$, then the number of maximum possible values of a (where $[.]$ represents the greatest integer function) is_____



[Watch Video Solution](#)

2. From a given solid cone of height H , another inverted cone is carved whose height is h such that its volume is maximum. Then the ratio $\frac{H}{h}$ is



[View Text Solution](#)

3. Let $f(x) = \begin{cases} |x^3 + x^2 + 3x + \sin x| \left(3 + \sin\left(\frac{1}{x}\right)\right) & x \neq 0 \\ 0 & x = 0 \end{cases}$ then

the number of point where $f(x)$ attains its minimum value is ____



[Watch Video Solution](#)

4. Let $f(x)$ be a cubic polynomial which has local maximum at $x = -1$ and $f(x)$ has a local minimum at $x = 1$. If $f(-1) = 10$ and $f(3) = -22$, then one fourth of the distance between its two horizontal tangents is _____



[Watch Video Solution](#)

5. Consider $P(x)$ to be a polynomial of degree 5 having extremum at $x = -1, 1$, and $(\lim)_{x \rightarrow 0} \left(\frac{p(x)}{x^3} - 2 \right) = 4$. Then the value of $[P(1)]$ is (where $[.]$ represents greatest integer function)___

 [Watch Video Solution](#)

6. If m is the minimum value of $f(x, y) = x^2 - 4x + y^2 + 6y$ when x and y are subjected to the restrictions $0 \leq x \leq 1$ and $0 \leq y \leq 1$, then the value of $|m|$ is_____

 [Watch Video Solution](#)

7. For a cubic function $y = f(x)$, $f^x = 4x$ at each point (x, y) on it and it crosses the x -axis at $(-2, 0)$ at an angle of 45° with positive direction of the x -axis. Then the value of $\left| \frac{f(1)}{5} \right|$ is_____

 [Watch Video Solution](#)

8. Number of integral values of b for which the equation $\frac{x^3}{3} - x = b$ has three distinct solutions is ___

 [Watch Video Solution](#)

9. Let $f(x) = \begin{cases} x + 2, & x < -1 \\ x^2, & -1 \leq x < 1 \\ (x - 2)^2, & x \geq 1 \end{cases}$ Then number of times $f'(x)$ changes its sign in $(-\infty, \infty)$ is ___

 [Watch Video Solution](#)

10. The number of nonzero integral values of a for which the function $f(x) = x^4 + ax^3 + \frac{3x^2}{2} + 1$ is concave upward along the entire real line is _____

 [Watch Video Solution](#)

11. Let $f(x) = \begin{cases} x^{\frac{3}{5}}, & \text{if } x \leq 1 \\ 1 - (x - 2)^3, & \text{if } x > 1 \end{cases}$. Then the number of critical points on the graph of the function is ___

 [Watch Video Solution](#)

12. A right triangle is drawn in a semicircle of radius $\frac{1}{2}$ with one of its legs along the diameter. If the maximum area of the triangle is M , then the value of $32\sqrt{3}Mc$ is _____

 [Watch Video Solution](#)

13. A rectangle with one side lying along the x-axis is to be inscribed in the closed region of the xy plane bounded by the lines $y = 0$, $y = 3x$, and $y = 30 - 2x$. If M is the largest area of such a rectangle, then the value of $\frac{2M}{27}$ is _____

 [Watch Video Solution](#)

14. The least integral value of x where $f(x) = (\log)_{\frac{1}{2}}(x^2 - 2x - 3)$ is monotonically decreasing is _____



Watch Video Solution

15. The least area of a circle circumscribing any right triangle of area $\frac{9}{\pi}$ is _____



Watch Video Solution

16. Let $f(x) = \begin{cases} |x^2 - 3x| + a, & 0 \leq x < \frac{3}{2} \\ -2x + 3, & x \geq \frac{3}{2} \end{cases}$ If $f(x)$ has a local maxima at $x = \frac{3}{2}$, then greatest value of $|4a|$ is _____



Watch Video Solution

17. Let $f(x) = 30 - 2x - x^3$, the number of the positive integral values of x which does satisfy $f(f(f(x))) > f(f(-x))$ is _____.



View Text Solution

18. Let $f(x) = \begin{cases} x(x-1)(x-2), & (0 \leq x < n) \\ \sin(\pi x), & (n \leq x \leq 2n) \end{cases}$

least value of n for which $f(x)$ has more points of minima than maxima in $[0, 2n]$ is ____.



[View Text Solution](#)

19. Number of critical point of the function $f(x) = x + \sqrt{|x|}$ is _____.



[View Text Solution](#)

20. consider $f(x) = \frac{1}{1+|x|} + \frac{1}{1+|x-1|}$ Let x_1 and x_2 be point where $f(x)$ attains local minimum and global maximum respectively .If $k = f(x_1) + f(x_2)$ then $6k-9=$ _____.



[View Text Solution](#)

21. Let f be a function defined on R (the set of all real numbers) such that $f'(x) = 2010(x - 2009)(x - 2010)^2(x - 2011)^3(x - 2012)^4$, for all $x \in R$. If g is a function defined on R with values in the interval $(0, \infty)$ such that $f(x) = \ln(g(x))$, for all $x \in R$, then the number of point is R at which g has a local maximum is ___



Watch Video Solution

22. Let $f: R \rightarrow R$ be defined as $f(x) = |x| + |x^2 - 1|$. The total number of points at which f attains either a local maximum or a local minimum is _____



Watch Video Solution

23. Let $p(x)$ be a real polynomial of least degree which has a local maximum at $x = 1$ and a local minimum at $x = 3$. If $p(1) = 6$ and $p(3) = 2$, then $p'(0)$ is _____



Watch Video Solution

24. A cylindrical container is to be made from certain solid material with the following constraints: It has a fixed inner volume of Vm^3 , has a 2 mm thick solid wall and is open at the top. The bottom of the container is a solid circular disc of thickness 2mm and is of radius equal to the outer radius of the container. If the volume the material used to make the container is minimum when the inner radius of the container is $10mm$. then the value of $\frac{V}{250\pi}$ is

 [Watch Video Solution](#)

Single Correct Answer Type

1. Given $P(x) = x^4 + ax^3 + bx^2 + cx + d$ such that $x = 0$ is the only real root of $P'(x) = 0$. If $P(-1) < P(1)$, then in the interval $[-1, 1]$

A. (a) $P(-1)$ is the minimum and $P(1)$ is the maximum of P

B. (b) $P(-1)$ is not minimum but $P(1)$ is the maximum of P

C. (c) $P(-1)$ is not minimum and $P(1)$ is not the maximum of P

D. (d) neither $P(-1)$ is the minimum nor $P(1)$ is the maximum of P

Answer: 2



[View Text Solution](#)

2. Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be defined by $f(x) = \begin{cases} k - 2x, & \text{if } x \leq -1 \\ -2x + 3, & \text{if } x > -1 \end{cases}$. If f has a local minimum at $x = -1$, then a possible value of k is (1) 0 (2) $-\frac{1}{2}$ (3) -1 (4) 1

A. -1

B. 1

C. 0

D. $\frac{1}{2}$

Answer: 1



[View Text Solution](#)

3. Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be a continuous function defined by $f(x) = \frac{1}{e^x + 2e^{-x}}$

. Statement-1: $f(c) = \frac{1}{3}$, for some $c \in \mathbb{R}$. Statement-2:

$0 < f(x) \leq \frac{1}{2\sqrt{2}}$, for all $x \in \mathbb{R}$. (1) Statement-1 is true, Statement-2 is

true; Statement-2 is not the correct explanation for Statement-1 (2)

Statement-1 is true, Statement-2 is false (3) Statement-1 is false,

Statement-2 is true (4) Statement-1 is true, Statement-2 is true;

Statement-2 is the correct explanation for Statement-1

A. statement 1 is false statement 2 is true

B. statement 1 is true , statement 2 is true statement 2 is a correct explanation for statement 1

C. statement 1 is true statement 2 is true statement 2 is not a correct explanation for statement 2

D. statement 1 is true , statement 2 is false

Answer: 2

4. Let $a, b \in \mathbb{R}$ be such that the function f given by $f(x) = \ln|x| + bx^2 + ax, x \neq 0$ has extreme values at $x = 1$ and $x = 2$. Statement 1: f has local maximum at $x = 1$ and at $x = 2$.

Statement 2: $a = \frac{1}{2}$ and $b = \frac{-1}{4}$ (1) Statement 1 is false, statement 2 is

true (2) Statement 1 is true, statement 2 is true; statement 2 is a correct explanation for statement 1 (3) Statement 1 is true, statement 2 is true;

statement 2 is not a correct explanation for statement 1 (4) Statement 1 is true, statement 2 is false

A. statement 1 is false statement 2 is true

B. statement 1 is true statement 2 is true, statement 2 is a correct explanation for statement 1

C. statement 1 is true statement 2 is true, statement 2 is not a correct explanation for statement 1

D. statement 1 is true statement 2 is false

Answer: 2



[View Text Solution](#)

5. find the values k for which the quadratic equation $2x^2 + Kx + 3 = 0$ has two real equal roots

- A. lies between 1 and 2
- B. lies between 2 and 3
- C. lies between -1 and 0
- D. does not exist

Answer: 4



[View Text Solution](#)

6. If $x = -1$ and $x = 2$ are extreme points of $f(x) = \alpha \log|x| + \beta x^2 + x$, then

A. $\alpha = -6, \beta = \frac{1}{3}$

B. $\alpha = -6, \beta = -\frac{1}{2}$

C. $\alpha = 2, \beta = -\frac{1}{2}$

D. $\alpha = 2, \beta = \frac{1}{2}$

Answer: 3



Watch Video Solution

7. Let $f(x)$ be a polynomial of degree four having extreme values at $x = 1$

and $x = 2$. If $\lim_{x \rightarrow 0} \left[1 + \frac{f(x)}{x^2} \right] = 3$ then $f(2)$ is equal to

A. -8

B. -4

C. 0

D. 4

Answer: 3



[Watch Video Solution](#)

8. A wire of the length 2 units is cut into two parts which are bent respectively to form a square of side= x units and a circle of radius = r units. If the sum of the areas of the square and the circle so formed is minimum, then

A. $(4 - \pi)x = \pi r$

B. $x=2r$

C. $2x=r$

D. $2x = (\pi + 4)r$

Answer: 2



[Watch Video Solution](#)

9. Twenty metres of wire is available for fencing off a flower-bed in the form of a circular sector. Then the maximum area (in sqm) of the flower-

bed is: (1) 25 (2) 30 (3) 12.5 (4) 10

A. 30

B. 12.5

C. 10

D. 25

Answer: 4



[View Text Solution](#)

10. Let $f(x) = x^2 + \left(\frac{1}{x^2}\right)$ and $g(x) = x - \frac{1}{x} \quad \xi n R - \{-1, 0, 1\}$. If $h(x) = \left(\frac{f(x)}{g(x)}\right)$ then the local minimum value of $h(x)$ is: (1) 3 (2) -3 (3) $-2\sqrt{2}$ (4) $2\sqrt{2}$

A. $2\sqrt{2}$

B. 3

C. -3

D. $-2\sqrt{2}$

Answer: 1



[View Text Solution](#)

11. The function $f: [0, 3] \xrightarrow{1, 29}$, defined by $f(x) = 2x^3 - 15x^2 + 36x + 1$, is one-one and onto onto but not one-one one-one but not onto neither one-one nor onto

- A. one-one and onto
- B. onto but not one -one
- C. one-one but not onto
- D. neither one-one nor on to

Answer: 2



[Watch Video Solution](#)

12. The number of points in $(-\infty, \infty)$, for which $x^2 - x \sin x - \cos x = 0$, is 6 (b) 4 (c) 2 (d) 0

A. 6

B. 4

C. 2

D. 0

Answer: 3

 [Watch Video Solution](#)

13. If $f: R \rightarrow R$ is a twice differentiable function such that $f''(x) > 0$ for all $x \in R$, and $f\left(\frac{1}{2}\right) = \frac{1}{2}$ and $f(1) = 1$, then

A. (a) $0 < f'(1) \leq \frac{1}{2}$

B. (b) $f'(1) \leq 0$

C. (c) $f'(1) > 1$

$$D. (d) \frac{1}{2} < f'(1) \leq 1$$

Answer: 3



[View Text Solution](#)

Linked comprehension Type

1. Let $f: [0, 1] \rightarrow \mathbb{R}$ be a function. Suppose the function f is twice differentiable, $f(0) = f(1) = 0$ and satisfies $f''(x) - 2f'(x) + f(x) \geq e^x, x \in [0, 1]$ Which of the following is true for $0 < x < 1$?

A. $0 < f(x) < \infty$

B. $-\frac{1}{2} < f(x) < \frac{1}{2}$

C. $-\frac{1}{4} < f(x) < 1$

D. $-\infty < f(x) < 0$

Answer: 4



[View Text Solution](#)

2. If the function $e^{-x} f(x)$ assumes its minimum in the interval $[0,1]$ at $x = 1/4$ which of the following is true ?

A. $f'(x) < f(x), 1/4 < x < 3/4$

B. $f'(x) > f(x), 0 < x < 1/4$

C. $f'(x) < f(x), 0 < x < 1/4$

D. $f'(x) < f(x), 3/4 < x < 1$

Answer: 3



[View Text Solution](#)