


CHEMISTRY

BOOKS - NARENDRA AWASTHI

CHEMICAL KINETIC & NUCLEAR CHEMISTRY

Exercise

1. The differential rate law equation for the elementary reaction



A. $-\frac{d[A]}{dt} = \frac{d[B]}{dt} = \frac{d[C]}{dt} = k[A][B]^2$

B. $-\frac{d[A]}{dt} = \frac{1}{2} \frac{d[b]}{dt} = \frac{1}{3} \frac{d[C]}{dt} = k[A]^2[B]$

C. $-\frac{d[A]}{dt} = \frac{1}{2} \frac{d[b]}{dt} = \frac{1}{3} \frac{d[C]}{dt} = k[A][B]^2$

D. None of these

Answer: C







Watch Video Solution

2. The rate of a reaction is expressed in different ways as follows:

$$+\frac{1}{2} \frac{d[C]}{dt} = -\frac{1}{3} \frac{d[D]}{dt} = +\frac{1}{4} \frac{d[A]}{dt} = -\frac{d[B]}{dt}$$

the reaction is



Answer: B



Watch Video Solution

3. In the reaction, $A + 2B \rightarrow 6C + 2D$, if the initial rate $-\frac{d[A]}{dt}$ at $t=0$ is $2.6 \times 10^{-2} M \text{ sec}^{-1}$, what will be the value of $-\frac{d[B]}{dt}$ at $t=0$?

A. $8.5 \times 10^{-2} M \text{ sec}^{-1}$

B. $2.5 \times 10^{-2} M \text{ sec}^{-1}$

C. $5.2 \times 10^{-2} M \text{ sec}^{-1}$

D. $7.5 \times 10^{-2} M \text{ sec}^{-1}$

Answer: C

 Watch Video Solution

4. For the reaction $2A \rightarrow B + 3C$, if $-\frac{d[A]}{dt} = k_1[A]^2$, $-\frac{d[B]}{dt} = k_2[A]^2$, $-\frac{d[C]}{dt} = k_3[A]^2$ the correct reaction between k_1 , k_2 and k_3 is :

A. $k_1 = k_2 = k_3$

B. $2k_1 = k_2 = 3k_3$

C. $4k_1 = k_2 = 3k_3$

D. $\frac{k_1}{2} = k_2 = \frac{k_3}{3}$

Answer: D

 [Watch Video Solution](#)

5. The rate constant of n^{th} order has units :

A. $\text{litre}^{1-n} \text{mol}^{1-n} \text{sec}^{-1}$

B. $\text{Mol}^{1-n} \text{litre}^{1-n} \text{sec}$

C. $\text{Mol}^{1-n^2} \text{litre}^{n^2} \text{sec}^{-1}$

D. $\text{Mole}^{1-n} \text{litre}^{n-1} \text{sec}^{-1}$

Answer: D

 [Watch Video Solution](#)

6. Which of the following statement is incorrect?

A. Unit of rate of disappearance is $M s^{-1}$

B. Unit of rate of reaction is $M s^{-1}$

C. Unit of rate constant k depends upon order

D. Unit of k for first order reaction is $M s^{-1}$

Answer: D

 [Watch Video Solution](#)

7. Which of the following relation is correct for k_f and k_b in an equilibrium process that contains equal moles of reactants and products.

A. $k_f = k_b$

B. $k_f > k_b$


C. $k_f < k_b$

D. we cannot predict

Answer: D

 [Watch Video Solution](#)

8. Listed in the table are forward and reverse rate constants for the reaction



Temperature(K)	$k_f(M^{-1}s^{-1})$	$k_b(M^{-1}s^{-1})$
----------------	---------------------	---------------------

1400	0.29	1.1×10^{-6}
------	------	----------------------

1500	1.3	1.4×10^{-5}
------	-----	----------------------

Select the correct statement:

a) Reaction is exothermic and value of equilibrium constant(K_{eq}) at 1400

K is 3.79×10^{-6}

b) Reaction is endothermic and value of k_{eq} 1400 K is 2.63×10^5

c) Reaction is exothermic and value of k_{eq} 1400 K is 2.63×10^5

d) Reaction is endothermic and value of k_{eq} 1500 K is 9.28×10^4

A. Reaction is exothermic and value of equilibrium constant(K_{eq}) at

1400 K is 3.79×10^{-6}

B. Reaction is endothermic and value of k_{eq} 1400 K is 2.63×10^5

C. Reaction is exothermic and value of k_{eq} 1400 K is 2.63×10^5

D. Reaction is endothermic and value of k_{eq} 1500 K is 9.28×10^4

Answer: C



[Watch Video Solution](#)

9. The rate constant depends on

A. temperature

B. pressure

C. extent of reaction

D. Initial concentraton of the reactant

Answer: A



[Watch Video Solution](#)

10. In the following reaction, how is the rate of appearance of the Product related to the rate of disappearance of the

underlined reactant ?



A. $-\frac{d[\text{BrO}_3^-]}{dt} = \frac{d[\text{Br}_2]}{dt}$

B. $-\frac{1}{3} \frac{d[\text{BrO}_3^-]}{dt} = \frac{d[\text{Br}_2]}{dt}$

C. $-\frac{d[\text{BrO}_3^-]}{dt} = \frac{1}{3} \frac{d[\text{Br}_2]}{dt}$



D. None of these

Answer: C





Watch Video Solution


11. Consider a reaction $A(\text{g}) \xrightarrow{k=0.1\text{M}^{-1}\text{min}^{-1}} 2B(\text{g})$. If initial concentration of A is 0.5 M then select correct graph.




a) b)




c) d)




A.



B.



C.



D.

Answer: C

[▶ Watch Video Solution](#)

12. Which of the following statement is incorrect?

- A. A second order reaction must be a bimolecular elementary reaction
- B. A bimolecular elementary reaction must be a second order reacton
- C. Zero order reaction must be a complex reaction
- D. First order reaction may be complex or elementary reaction

Answer: A



[Watch Video Solution](#)

13. The molecularity of a complex reaction given below is :



A. 1

B. 2

C. 3


D. no meaning

Answer: D




[Watch Video Solution](#)


14. Decomposition of $NH_4NO_2(aq)$ into $N_2(g)$ and $2H_2O(l)$ is first order reaction.




A.



B.



C.



D.





Answer: D





Watch Video Solution

15. Decomposition of $HI(g)$ on Gold surface is zero order reaction.

Initially, few moles of H_2 are present in container then which of the following graph is correct ?









Answer: B

 [Watch Video Solution](#)

16. Consider the plots for the types of reaction



These plots respectively correspond to the reaction orders :

a) 0,2,1

b) 0,1,2

c) 1,1,2

d) 1,0,2

A. 0,2,1

B. 0,1,2

C. 1,1,2


D. 1,0,2

Answer: D




[Watch Video Solution](#)


17. If decomposition reaction $A(g) \rightarrow B(g)$ follows first order kinetics then the graph of rate of formation (R) of B against time t will be :




A.



B.



C.







D.

Answer: C



Watch Video Solution

18. Consider the plots for the types of reaction



These plots respectively correspond to the reaction orders :

- A. 0,1,2
- B. 1,2,0
- C. 1,0,2
- D. None of these

Answer: C



Watch Video Solution

19. For a zero order reaction, the plot of concentration, vs time is linear with


- A. +ve slope and zero intercept
- B. -ve slope and zero intercept
- C. +ve slope and non-zero intercept
- D. -ve slope and non-zero intercept

Answer: D



Watch Video Solution

20. What will be the order of reaction for a chemical change having $\log t_{\frac{1}{2}}$ vs $\log a$? (where a = initial concentration of reactant, $t_{\frac{1}{2}}$ = half life)




- A. zero order
- B. First order
- C. Second order
- D. None of these

Answer: A



Watch Video Solution

21. A graph between $\log t_{\frac{1}{2}}$ and $\log a$ (abscissa), a being the initial concentration of A in the reaction For reaction $A \rightarrow \text{Product}$, the rate law is :



A. $-\frac{d[A]}{dt} = K$

B. $-\frac{d[A]}{dt} = K[A]$

C. $-\frac{d[A]}{dt} = K[A]^2$


D. $-\frac{d[A]}{dt} = K[A]^3$

Answer: C




Watch Video Solution


22. For the reaction $A \rightarrow B$, for which graph between half life ($t_{1/2}$) and initial concentration (a) of the reactant is given below




Hence graph between $-\frac{d[A]}{dt}$ and time will be:




A.



B.



C.



D.

Answer: C



Watch Video Solution

23. For the ideal gaseous reaction, the rate is generally expressed in terms of $\frac{dP}{dt}$ instead of $\frac{dC}{dt}$ or $\frac{dn}{dt}$ (where $C = \frac{n}{V}$ is concentration and n the no. of moles). What is the relation among these three expressions if T and V are constant?

$$A. \frac{dC}{dt} = \frac{dn}{dt} = \frac{dP}{dt}$$


$$B. \frac{dC}{dt} = \frac{1}{V} \frac{dn}{dt} = \frac{1}{Rt} \left(\frac{dP}{dt} \right)$$

$$C. RT \frac{dC}{dt} = \frac{dn}{dt} = \frac{dP}{dt}$$

D. None of these

Answer: B

 [Watch Video Solution](#)



Initial $[A_2]$	Initial $[B_2]$	R. O. R. (r) $M s^{-1}$
0.2	0.2	0.04
0.1	0.4	0.04
0.2	0.4	0.08

Order of reaction with respect to A_2 and B_2 are respectively :

A. a=1,b=1

B. a=2,b=0

C. a=2,b=1

D. None

Answer: A

 [Watch Video Solution](#)

25. For a reaction initial rate is given as : $R_0 = k[A_0]^2[B_0]$. By what factor, the initial rate of reaction will increase if initial concentration is taken 1.5 times and B is tripled?

A. 4.5

B. 2.25

C. 6.75

D. None of these

Answer: C

 [Watch Video Solution](#)

26. For $A_{(s)} + B_{(s)} \rightarrow C_{(s)}$, $\text{rate} = k[A]^{1/2}[B]^2$, if initial concentration of A and B are increased by factors 4 and 2 respectively, then the initial rate is changed by the factor:

A. 4

B. 6

C. 8

D. None of these

Answer: C



[Watch Video Solution](#)

27. Reaction $A \rightarrow B$ follows second order kinetics. Doubling the concentration of A will increase the rate of formation of B by a factor of :

A. $1/4$

B. $1/2$

C. 2


D. 4

Answer: D



Watch Video Solution

28. The reaction of A_2 and B_2 follows the equation



The following data were observed

$[A_2]_0$	$[B_2]_0$	Initial rate of appearance of $AB(g)$ (in $M s^{-1}$)
0.10	0.10	2.5×10^{-4}
0.20	0.10	5×10^{-4}
0.20	0.20	10×10^{-4}

The value of rate constant for the above reaction is :

- a) 2.5×10^{-4}
- b) 2.5×10^{-2}
- c) 1.25×10^{-2}
- d) None of these

A. 2.5×10^{-4}

B. 2.5×10^{-2}

C. 1.25×10^{-2}

D. None of these

Answer: C

 [Watch Video Solution](#)

29. The unit of rate constant of elementary reaction depends upon the

A. temperature of the reaction

B. concentration of reactants

C. activation energy of the reaction

D. Molecularity of the reaction

Answer: D

 [Watch Video Solution](#)

30. Select the rate law that corresponds to the datashown for the

reaction $A + B \rightarrow C$

<i>Exp.</i>	$[A]$	$[B]$	<i>Rate</i>
1	0.012	0.035	0.10
2	0.024	0.070	0.80
3	0.024	0.035	0.10
4	0.012	0.070	0.80

a) $\text{Rate} = k[B]^3$

b) $\text{Rate} = k[B]^4$

c) $\text{Rate} = k[A][B]^3$

d) $\text{Rate} = k[A]^2[B]^2$

A. $\text{Rate} = k[B]^3$

B. $\text{Rate} = k[B]^4$

C. $\text{Rate} = k[A][B]^3$

D. $\text{Rate} = k[A]^2[B]^2$

Answer: A



Watch Video Solution

31. An elementary reaction A and B is second order reaction. Which of the following rate equation must be correct?

A. $r = k[A]^2[B]^0$

B. $r = k[A]^{3/2}[B]^{1/2}$

C. $r = k[A]^0[B]^2$

D. $r = k[A][B]$

Answer: D



Watch Video Solution

32. If a is the initial concentration of the reactant, the half life period of the reaction of n^{th} order is inversely proportional to :

A. a^{n-1}

B. a^n

C. a^{1-n}

D. a^{n+1}

Answer: A



Watch Video Solution

33. Which of the following expressions is correct for zero order and first order respectively [Where a is initial concentration]?

A. $t_{1/2}\alpha a, t_{1/2}\alpha \frac{1}{a}$

B. $t_{1/2}\alpha a, t_{1/2}\alpha a^0$

C. $t_{1/2}\alpha a^0, t_{1/2}\alpha a$

D. $t_{1/2}\alpha a, t_{1/2}\alpha \frac{1}{a^2}$

Answer: B



Watch Video Solution

34. Unit of rate constant for zero order reaction is



[Watch Video Solution](#)

35. The unit of rate of reaction and rate of rate constant are same for a :

- a) zero order reaction
- b) first order reaction
- c) Second order reaction
- d) third order reaction


- A. zero order reaction
- B. first order reaction
- C. Second order reaction
- D. third order reaction

Answer: A



[Watch Video Solution](#)

36.



. What type of reaction is this?

- a) unimolecular elementary
- b) Pseudo first order
- c) Zero order reaction must be a complex reaction
- d) Second order

A. unimolecular elementary

B. Pseudo first order

C. Zero order reaction must be a complex reaction

D. Second order

Answer: B



Watch Video Solution

37. When ethyl acetate was hydrolysed in presence of 0.1 M HCl, the rate constant was found to be $5.4 \times 10^{-5} s^{-1}$. But in presence of 0.1 M H_2SO_4 the rate constant was found to be $6.25 \times 10^{-5} s^{-1}$. Thus it may be concluded that:

- a) H_2SO_4 furnishes more H^+ than HCl
- b) H_2SO_4 furnishes less H^+ than HCl
- c) both have the same strength
- d) will depend on concentration of ethyl acetate

A. H_2SO_4 furnishes more H^+ than HCl

B. H_2SO_4 furnishes less H^+ than HCl

C. both have the same strength

D. will depend on concentration of ethyl acetate

Answer: A



Watch Video Solution

38. For an elementary reaction $2A + B \rightarrow A_2B$ if the volume of vessel is quickly reduced to half of its original volume then rate of reaction will :

- a) remain unchanged
- b) increase four times
- c) increase eight times
- d) decrease eight times

A. remain unchanged

B. increase four times

C. increase eight times

D. decrease eight times

Answer: C



[Watch Video Solution](#)

39. In the reaction $A \rightarrow B + C$, rate constant is 0.001 Ms^{-1} . If we start with 1 M of A then conc. Of A and B after 10 minutes are respectively :

a) 0.5 M, 0.5 M

b) 0.6 M, 0.4 M

c) 0.4 M, 0.6 M

d) none of these

A. 0.5 M, 0.5 M

B. 0.6 M, 0.4 M

C. 0.4 M, 0.6 M

D. none of these

Answer: C



[Watch Video Solution](#)

40. For a reaction $A \xrightarrow{k_r = 0.6 M^{-1} \text{ min}^{-1}} 2B$

starting with 1 M of 'A' only, concentration of B (in M) after 100 sec. and 200 sec. is respectively?

a) 2 and 4

b) 1 and 2

c) 2 and 3

d) None of these

A. 2 and 4

B. 1 and 2

C. 2 and 3

D. None of these

Answer: D



[Watch Video Solution](#)

41. For the zero order reaction $A \rightarrow B + C$, initial concentration of A is 0.1 M. If $[A]=0.08$ M after 10 minutes, then its half-life and completion time are respectively :

a) 10 min, 20 min

b) 2×10^{-3} min , 4×10^{-3} min

c) 25 min, 50 min

d) 250 min, 500 min

A. 10 min, 20 min

B. 2×10^{-3} min , 4×10^{-3} min

C. 25 min, 50 min

D. 250 min, 500 min

Answer: C



Watch Video Solution

42. For an elementary reaction , $X(g) \rightarrow Y(g) + Z(g)$

the half life period is 10 min. In what period of time would the concentration of X be reduced to 10% of original concentration?

A. 20 min

B. 33 min

C. 15 min

D. 25 min

Answer: B



[Watch Video Solution](#)

43. In the presence of acid, the initial concentration, of cane sugar was reduced from $0.2M$ to 0.1 in $5hr$ and to $0.05M$ in $10hr$. The reaction must be of:

- a) Zero order
- b) First order
- c) Second order
- d) Third order

- A. Zero order
- B. First order
- C. Second order
- D. Third order

Answer: B





Watch Video Solution

44. A first order reaction is 75% completed in 100 minutes. How long time will it take for its 87.5% completion?

A. 125 min

B. 150 min

C. 175 min

D. 200 min

Answer: B



Watch Video Solution

45. The rate constant for a first order reaction whose half life is 480 sec, is :

A. $1.44 \times 10^{-3} \text{ sec}^{-1}$

B. $1.44 \times \text{sec}^{-1}$

C. $0.72 \times 10^{-3} \text{sec}^{-1}$

D. $2.88 \times 10^{-3} \text{sec}^{-1}$

Answer: A

 [Watch Video Solution](#)

46. Rate constant $k=2.303 \text{ min}^{-1}$ for a particular reaction. The initial concentration of the reactant is 1 mol /litre then rate of reaction after 1 minute is:

A. 2.303 M min^{-1}

B. $0.2303 \text{ M min}^{-1}$

C. 0.1 M min^{-1}

D. None of these

Answer: B



Watch Video Solution

47. For the reaction $3A(g) \xrightarrow{k} B(g) + C(g)$ k is $10^{-4} L/mol \cdot \text{min}$.

If $[A] = 0.5M$ then the value of $-\frac{d[A]}{dt}$ (in ms^{-1} is:

A. 7.5×10^{-5}

B. 3×10^{-4}

C. 2.5×10^{-5}

D. None of these

Answer: D



Watch Video Solution

48. 99 % at a first order reaction was completed in 32 min . When will 99.9 % of the reaction complete.

A. 50 min

B. 46 min

C. 48 min

D. 49 min

Answer: C



Watch Video Solution

49. Which of the following represent the expression for $\frac{3}{4}$ th life of first order reaction:

a) $\frac{k}{2.303} \log 4/3$

b) $\frac{2.303}{k} \log 3/4$

c) $\frac{2.303}{k} \log 4$

d) $\frac{2.303}{k} \log 3$

A. $\frac{k}{2.303} \log 4/3$

B. $\frac{2.303}{k} \log 3/4$

C. $\frac{2.303}{k} \log 4$


D. $\frac{2.303}{k} \log 3$

Answer: C



Watch Video Solution

50. Consider following two competing first order reactions,



if 50% of the reaction of P was completed when 96% of Q was complete, then the ratio (k_2/k_1) will be :

a) 4.06

b) 0.215

c) 1.1

d) 4.65

A. 4.06

B. 0.215

C. 1.1


D. 4.65

Answer: D





Watch Video Solution

51. For the reaction



(ii) $B \xrightarrow{k_{II}} Q$, following observation is made.



Calculate $\frac{k_I}{k_{II}}$, where k_I and k_{II} are rate constants for the respective reaction.

a) 2.303

b) 1

c) 0.36

d) 0.693

A. 2.303

B. 1

C. 0.36

D. 0.693

Answer: D

 [Watch Video Solution](#)

52. For the homogenous gaseous reaction $A \rightarrow 3B$, if pressure after time t was P_t and after completion of reaction, pressure was P_∞ then select correct relation

a) $k = \frac{1}{t} \ln \left(\frac{P_\infty}{3(P_\infty - P_t)} \right)$

b) $k = \frac{1}{t} \ln \left(\frac{2P_\infty}{(P_\infty - P_T)} \right)$

c) $k = \frac{1}{t} \ln \left(\frac{3P_\infty}{2P_\infty - P_t} \right)$

d) $k = \frac{1}{t} \ln \left(\frac{P_\infty}{3(P_\infty - P_T)} \right)$

A. $k = \frac{1}{t} \ln \left(\frac{P_\infty}{3(P_\infty - P_t)} \right)$

B. $k = \frac{1}{t} \ln \left(\frac{2P_\infty}{(P_\infty - P_T)} \right)$

$$C. k = \frac{1}{t} \ln \left(\frac{3P_{\infty}}{2P_{\infty} - P_t} \right)$$

$$D. k = \frac{1}{t} \ln \left(\frac{P_{\infty}}{3(P_{\infty} - P_T)} \right)$$

Answer: D



Watch Video Solution

53. The half-life of first order decomposition of NH_4NO_3 is 2.10 hr at 288 K temperature

$NH_4NO_3(aq) \rightarrow N_2O(g) + 2H_2O(l)$. If 6.2 of NH_4NO_3 is allowed to decompose, the required for NH_4NO_3 to decompose 90 % is :

A. 6.978 hr

B. 0.319

C. 0.319 hr

D. None of these

Answer: A



Watch Video Solution

54. For a first order homogenous gaseous reaction, $A \rightarrow 2B + C$

then initial pressure was P_i while total pressure after time 't' was P_t . The

right expression for the rate constants k in terms of P_i , P_t and t is :

a) $k = \frac{2.303}{t} \log \left(\frac{2P_i}{3P_i - P_t} \right)$

b) $k = \frac{2.303}{t} \log \left(\frac{2P_i}{2P_t - P_i} \right)$

c) $k = \frac{2.303}{t} \log \left(\frac{P_i}{P_i - P_t} \right)$

d) None of these

A. $k = \frac{2.303}{t} \log \left(\frac{2P_i}{3P_i - P_t} \right)$

B. $k = \frac{2.303}{t} \log \left(\frac{2P_i}{2P_t - P_i} \right)$

C. $k = \frac{2.303}{t} \log \left(\frac{P_i}{P_i - P_t} \right)$

D. None of these

Answer: A



Watch Video Solution

55. The decomposition of azo methane, at certain temperature according to the equation $(CH_3)_2N_2 \rightarrow C_2H_6 + N_2$ is a first order reaction.

After 40 minutes from the start, the total pressure developed is found to be 350 mm Hg in place of initial pressure 200 mm Hg of azo methane. The value of rate constant k is :

A. $2.88 \times 10^{-4} \text{ sec}^{-1}$

B. $1.25 \times 10^{-4} \text{ sec}^{-1}$

C. $5.77 \times 10^{-4} \text{ sec}^{-1}$

D. None of these

Answer: C



Watch Video Solution

56. The hydrolysis of sucrose was studied with the help of calorimeter and following data were

Collected time (min.) : 0 70 ∞
observed rotation (degrees) : 44 16.5 - 11

the time taken when reaction mixture will be optically inactive ? (Given : $\ln 2=0.7, \ln 3=1.1, \ln 5=1.6$)

- A. 16 min
- B. 69.47 min
- C. 160 min
- D. None of these

Answer: C

 [Watch Video Solution](#)

57. For a particular reaction with initial conc. Of the reactants as a_1 and a_2 , the half-life periods are t_1 and t_2 respectively. The order of the reaction (n) is given by :

- A. $n = 1 + \frac{\log(t_2/t_1)}{\log(a_2/a_1)}$
- B. $n = \frac{\log(t_2/t_1)}{\log(a_2/a_1)}$

C. $n = 1 + \frac{\log(t_1/t_2)}{\log(a_2/a_1)}$

D. None of these

Answer: C



Watch Video Solution

58. The value of $\frac{t_{0.875}}{t_{0.50}}$ for n^{th} order reaction is

A. $2^{(2n-2)}$

B. $2^{(2n-2) - 1}$

C. $\frac{8^{n-1} - 1}{2^{n-1} - 1}$

D. None of these

Answer: C



Watch Video Solution

59. $A \rightarrow B$ first order reaction A is optically active and B is optically inactive. A series of experiment were conducted on a solution of A

Time	0	60 min	∞
optical rotation	82°	77°	2°

assume some impurity is present calculate the optical rotation for 5 hours.)

(Given in $1.066=0.064$, $e^{0.16}=1.17$)

- A. 60
- B. 30
- C. 20
- D. 120

Answer: A



[Watch Video Solution](#)

60. At 300 K the half-life of a sample of a gaseous compound initially at 1 atm is 100 sec. When the pressure is 0.5 atm the half-life is 50 sec. The order of reaction is :

A. 0

B. 1

C. 2

D. 3

Answer: A



Watch Video Solution

61. For an endothermic reaction, where ΔH represents the enthalpy of reaction in kJmol^{-1} , the minimum value for the energy of activation will be

A. less than ΔH

B. more than ΔH


C. equal to ΔH

D. Zero

Answer: B

 [Watch Video Solution](#)

62. The activation energy of the reaction, $A + B \rightarrow C + D + 38 \text{ kcal}$ is 20 kcal. What would be the activation energy of the following reaction.



A. 20 kcal

B. -20 kcal

C. 18 kcal

D. 58 kcal

Answer: D

 [Watch Video Solution](#)

63. When the activation energies of the forward and backward reactions are equal, then :

A. $\Delta U=0, \Delta S=0$

B. $\Delta U=0, \Delta G=0$

C. $\Delta S=0, \Delta G=0$

D. Only $\Delta U=0$

Answer: D




[Watch Video Solution](#)


64. For an exothermic chemical process occurring in two steps as follows




The progress of reaction can be best described by :




A.



B.



C.







D.

Answer: C



Watch Video Solution

65. Select the correct diagram for an endothermic reaction that proceeds through two steps with the second steps is rate determining :



Answer: D



[Watch Video Solution](#)

66. $\frac{k_{34^\circ}}{k_{35^\circ}} < 1$, then :

- a) Rate increase with the rise in temperature
- b) rate decrease with rise in temperature
- c) rate does not change with rise in temperature
- d) None of the above

A. Rate increase with the rise in temperature

B. rate decrease with rise in temperature

C. rate does not change with rise in temperature

D. None of the above

Answer: A



[Watch Video Solution](#)

67. The plot of $\ln k$ v/s $1/T$ is linear with slope of :

A. $-E_a / R$

B. E_a / R)

C. $E_a / 2.303R$

D. $-E_a / 2.303R$

Answer: A



[Watch Video Solution](#)

68. Rate constant for a chemical reaction taking place at 500K is expressed as $K = A \cdot e^{-1000}$ The activation energy of the reaction is :

A. 100 cal/mol

B. 1000 kcal/mol

C. 10^4 kcal/mol

D. 10^6 kcal/mol

Answer: B



Watch Video Solution

69. For a complex reaction $A \xrightarrow{k} \text{products}$

$$E_{a1} = 180\text{kJ/mole}, E_{a2} = 80\text{kJ/mol}, E_{a3} = 50\text{kJ/mol}$$

Overall rate constant k is related to individual rate constant by the

equation $k = \left(\frac{k_1 k_2}{k_3} \right)^{2/3}$. Activation energy (kJ/mol) for the overall

reaction is :

a) 100

b) 43.44

c) 150

d) 140

A. 100

B. 43.44

C. 150

D. 140

Answer: D



Watch Video Solution

70. For reaction $A \rightarrow B$, the rate constant $K_1 = A_1 \left(e^{-E_{a1}/RT} \right)$ and the reaction $X \rightarrow Y$, the rate constant $K_2 = A_2 \left(e^{-E_{a2}/RT} \right)$. If $A_1 = 10^9$, $A_2 = 10^{10}$ and $E_{a1} = 1200$ cal/mol and $E_{a2} = 1800$ cal/mol, then the temperature at which $K_1 = K_2$ is : (Given, $R = 2$ cal/K-mol)

- a) $300K$
- b) $300 \times 2.303K$
- c) $\frac{300}{2.303}K$
- d) None of these

A. $300K$

B. $300 \times 2.303K$

C. $\frac{300}{2.303}K$

D. None of these

Answer: C

 [Watch Video Solution](#)

71. The activation energies of the forward and backward reactions in the case of a chemical reaction are 30.5 and 45.5 KJ/mol , respectively . The reaction is:

- A. exothermic
- B. endothermic
- C. neither exothermic
- D. independent of temperature

Answer: A

 [Watch Video Solution](#)

72. A reaction rate constant is given by : $K = 1.2 \times 10^{14} e^{\frac{-25000}{RT}} \text{ sec}^{-1}$. It

means :

- A. log K versus log T will give a straight line with a slope as 25000
- B. log K versus log T will give a straight line with a slope as -25000
- C. log k versus T will give a straight line with a slope as -25000
- D. log K versus 1/T will give a straight line

Answer: D



[Watch Video Solution](#)




73. The temperature coefficient of a reaction is:


- A. the rate constant
- B. the rate constant at a fixed temperature
- C. the ratio of rate constant at two temperature
- D. the ratio of rate constant differing by 10° C preferably k_{308}
 k_{298}

Answer: D

 Watch Video Solution

74. Which graph shows zero activation energy ?






Answer: C

[▶ Watch Video Solution](#)

75. A homogenous catalytic reaction takes place through the three alternative plots A, B, and C shown in the given figure. Which one of the following indicates the relative ease with which the reaction cant take

place?



A. $A > B > C$

B. $C > B > A$

C. $A > C > B$

D. $A=B=C$

Answer: B



76. The rate of a reaction gets doubled when temperature increase by 27°C to 37° . By what factor will it change for the temperature range 17°C to 27°C

- a) 1.81
- b) 1.71
- c) 2.1
- d) 2.41

A. 1.81

B. 1.71

C. 2.1

D. 2.41

Answer: C



77. Which of the following explains the increase of the reaction rate by catalyst?

- A. Catalyst decreases the rate of backward reaction so that the rate of forward reaction increases
- B. Catalyst provides extra energy to reacting molecules so that they may reduce effective collisions
- C. Catalyst provides an alternative path of lower activation energy to the reactants.
- D. Catalyst increases the number of collisions between the reacting molecules.

Answer: C



[Watch Video Solution](#)

78. Collision theory is satisfactory for:

- A. First order reactions
- B. Zero order reactions
- C. Bimolecular elementary reactions
- D. Any order reactions

Answer: C

 [Watch Video Solution](#)

79. For the first order reaction $A \rightarrow B + C$, carried out at 27°C . If $3.8 \times 10^{-16} \%$ of the reactant molecules exists in the activated state, the E_a (activation energy) of the reaction is:

- a) 12 KJ/mol
- b) 831.4 KJ/mol
- c) 100 KJ/mol
- d) 88.57 KJ/mol

A. 12 KJ/mol

B. 831.4 KJ/mol

C. 100 KJ/mol



D. 88.57 KJ/mol

Answer: C



Watch Video Solution

80. A following mechanism has been proposed for a reaction



The rate law expression for the reaction by RDS method is:

A. $r = k[A]^2[B]$

B. $r = k[A][B]$

C. $r = k[A]^2$



D. $r = k[A][C]$

Answer: B



[Watch Video Solution](#)

81. A hypothetical reaction $A_2 + B_2 \rightarrow 2AB$ follows the mechanism as given below:



The order of the overall reaction is

A. 2

B. 1

C. $\frac{3}{2}$

D. 0

Answer: C



[Watch Video Solution](#)

82. Chemical reaction occurs as a result of collisions between reacting molecules. Therefore, the reaction rate is given by

- A. total number of collisions occurring in a unit volume per second
- B. fraction of molecules which passes energy less than the threshold energy
- C. total number of effective collisions which have enough activation energy
- D. none of the above

Answer: C



Watch Video Solution

83. Radioactivity is affected by:

- A. temperature

B. pressure

C. electric and magnetic field

D. none of these

Answer: D



Watch Video Solution

84. The radiation from naturally occurring radioactive substance as seen after deflection by a magnetic field in one direction are :

a) α -rays

b) β -rays

c) both α and β rays

d) either α or β -rays

A. α -rays

B. β -rays

C. both α and β rays

D. either α or β -rays

Answer: D



[Watch Video Solution](#)

85. During α -decay:

- A. $\frac{n}{p}$ ratio decreases
- B. $\frac{n}{p}$ ratio increases
- C. $\frac{n}{p}$ ratio remains emission constant
- D. $\frac{n}{p}$ ratio may increase or decrease

Answer: B



[Watch Video Solution](#)

86. Atoms ${}_{7}X^A$, ${}_{8}Y^B$ and ${}_{9}Z^{17}$ are such that ${}_{8}Y$ is an isobar of ${}_{7}X$ and atom ${}_{9}Z^{17}$ is isotone of ${}_{8}Y$. Mass no. of X and no of neutrons in Y are respectively :

A. 8,8

B. 17,7

C. 9,8

D. 16,8

Answer: D



[Watch Video Solution](#)

87. ${}_{90}Th^{234}$ disintegrates to give ${}_{82}Pb^{206}$ as the final product. How many alpha and beta particles are emitted during this process ?

A. 6

B. 7

C. 8

D. 13

Answer: D



Watch Video Solution

88. An isotone of ${}^{76}_{32}\text{Ge}$ is-

(a) ${}^{77}_{32}\text{Ge}$

(b) ${}^{77}_{33}\text{As}$

(c) ${}^{77}_{34}\text{Se}$

(d) ${}^{78}_{34}\text{Se}$

A. – $(32)^{77}\text{Ge}$

B. – $(33)^{77}\text{As}$

C. – $(34)^{77}\text{Se}$

D. – $(36)^{77}\text{Se}$

Answer: B



Watch Video Solution

89. Isodiaphers are atoms having:

A. n/p same

B. p/n same

C. $(n-p)$ same

D. $(n-p)$ different

Answer: C



Watch Video Solution

90. The number of neutrons accompanying the formation of ${}_{54}\text{Xe}^{139}$ and ${}_{38}\text{Sr}^{94}$ from the absorption of a slow neutron by ${}_{92}\text{U}^{235}$, followed by nuclear fission is

A. 0

B. 1

C. 2





D. 3

Answer: D






Watch Video Solution

91. Complete the following nuclear equation by supplying the symbol for the other product of the fission :



d) none of these





D. none of these

Answer: B



Watch Video Solution

92. $X \rightarrow {}_{82}^{206}\text{Pb} + {}_2^4\text{He}$. In this reaction predict the position of group of X:

a) II B

b) IV B

c) VI A

d) VI B

A. II B

B. IV B

C. VI A

D. VI B

Answer: C



Watch Video Solution

93. ${}_{90}Th$ is a member of III group . After losing α -particle it forms a new element belonging to :

- a) I group
- b) II group
- c) III group
- d) IV group

A. I group

B. II group

C. III group

D. IV group

Answer: B



Watch Video Solution

94. Alpha decay of ${}_{92}^{238}U$ forms ${}_{90}^{234}Th$. What kind of decay from ${}_{90}^{234}Th$ produces ${}_{84}^{234}Ac$?

- a) α
- b) β
- c) β^+ (positron)
- d) γ -emission

A. α

B. β

C. β^+ (positron)

D. γ -emission

Answer: C



Watch Video Solution

95. A radioactive sample had an initial activity of 56 dpm . After 69.3 minutes, it was found to have an activity of 28 dpm . Find the number of

atoms in a sample having an activity of 100 dpm.

- A. 693
- B. 100
- C. 1000
- D. 10000

Answer: D



[Watch Video Solution](#)

96. A radioactive sample has initial activity of 28 dpm 30 minutes later its activity 14 dpm . How many atoms of nuclide were present initially?

- A. 2800
- B. 1212
- C. 528
- D. 2802

Answer: B



[Watch Video Solution](#)

97. The value of decay constant of Co^{60} is $2.5 \times 10^{-7} \text{ min}^{-1}$. The activity of 2.0 g of the sample is nearly :

A. 5×10^5 dpm

B. 2.5×10^{10} dpm

C. 5×10^{15} dpm

D. 10^{10} dpm

Answer: C



[Watch Video Solution](#)

98. Half-life ($t_{1/2}$) for a radioactive decay is 6930 sec. The time required to fall the rate of decay to $\left(\frac{1}{100}\right)^{th}$ of its initial value is:

- A. 69.3 sec
- B. 20000 sec
- C. 46060 sec
- D. None of these

Answer: C

 [Watch Video Solution](#)

99. A sample of radioactive substance is found 90% of its initial amount after one day . What % of the original sample can be found after 3 days?

- A. 81
- B. 72.9
- C. 25
- D. 65.61

Answer: B

 [Watch Video Solution](#)

100. If time t is required for a radioactive substance to become one third of its initial amount, what fraction would be left after $0.5 t$?

A. $\frac{1}{2}$

B. $\frac{1}{\sqrt{3}}$

C. $\frac{1}{3}$

D. $\sqrt{\frac{2}{3}}$

Answer: B

 [Watch Video Solution](#)

101. The present activity of the hair of Egypt mummy is $1.75 \text{ dpm } t_{1/2}$ of ${}^{14}_6\text{C}$ is 5770 year and disintegration rate of fresh sample of C^{14} is 14 dpm . Find out age of mummy.

a) 23080 year

b) 138480 year

c) 11998.3 year

d) 17315 year

A. 23080 year

B. 138480 year

C. 11998.3 year

D. 17315 year

Answer: D



Watch Video Solution

102. A 0.50g sample of rock was found to have 2.5×10^{-6} mol of ${}_{19}^{40}\text{K}$ ($t_{1/2} = 1.3 \times 10^9$ yr) and 7.5×10^{-6} mol of ${}_{20}^{40}\text{Ca}$. How old is the rock?

A. 6.5×10^8 yr

B. 1.3×10^9 yr

C. 2.6×10^9 yr

D. 5.2×10^9 yr

Answer: C



Watch Video Solution

103. Indium -112 is radioactive and has a very short half-life ($t_{1/2} = 14$ min). Its decay constant and average life are respectively:

a) 0.0495 min^{-1} , 9.7 min

b) 0.495 min^{-1} , 20.2 min

c) 9.7 min^{-1} , 20.2 min

d) 0.0495 min^{-1} , 20.2 min

A. 0.0495 min^{-1} , 9.7 min

B. 0.495 min^{-1} , 20.2 min

C. 9.7 min^{-1} , 20.2 min

D. 0.0495 min^{-1} , 20.2 min

Answer: D



[Watch Video Solution](#)

104. The half-life of a radioactive element is 100 minutes . The time interval between the stage to 50% and 87.5% decay will be:

- a) 100 min
- b) 50 min
- c) 200 min
- d) 25 min

- A. 100 min
- B. 50 min
- C. 200 min
- D. 25 min

Answer: C



[Watch Video Solution](#)

105. The half-life of Tc^{99} is 6.0 hr. The total residual activity in a patient 30 hr after receiving an injection containing Tc^{99} must be more than $0.01 \mu Ci$. What is the maximum activity (in μCi) that the sample injected can have?

- a) 0.16
- b) 0.32
- c) 0.64
- d) 0.08

A. 0.16

B. 0.32

C. 0.64

D. 0.08

Answer: B



Watch Video Solution

106. A pure radio-chemical preparation was observed to disintegrate at the rate of 2140 counts/minutes at 12.35 P.M. . At 3.55 P.M. of the same day, the disintegration rate of the sample was only 535 count/minutes.

What is the half-life of the material?

- a) 50 min
- b) 100 min
- c) 200 min
- d) None of these

A. 50 min

B. 100 min

C. 200 min

D. None of these

Answer: B



Watch Video Solution

107. A radioactive substance decay 25% in 10 minutes . If at start there are 4×10^{20} atoms present. After how much time will the number of atoms be reduced to 10^{20} atoms? (given $\ln 3=1.098$)

- a) 10.98 min
- b) 21.97 min
- c) 48.19 min
- d) None of these

A. 10.98 min

B. 21.97 min

C. 48.19 min

D. None of these

Answer: C



Watch Video Solution

108. Time taken of decay for a nuclear reaction is given by $t = 4t_{1/2}$. The relation between the mean life (T) and time of decay (t) is given by :

a) $2 T \ln 2$

b) $4 T \ln 2$

c) $2T^4 \ln 2$

d) $\frac{1}{T^2} \ln 2$

A. $2 T \ln 2$

B. $4 T \ln 2$

C. $2T^4 \ln 2$

D. $\frac{1}{T^2} \ln 2$

Answer: B



Watch Video Solution

109. Two radioactive nuclides A and B have half-lives 50 min and 10 min respectively . A fresh sample contains the nuclides of B to be eight times

that of A. How much time should elapse so that the number of nuclides of A becomes double of B ?

- a) 30
- b) 40
- c) 50
- d) 100

A. 30

B. 40

C. 50

D. 100

Answer: C



[Watch Video Solution](#)

110. A radioactive nuclide is produced at a constant rate of α per second . It's decay constant is λ . If N_0 be the no. of nuclei at time $t=0$, then max. no. nuclei possible are :

a) N_0

b) α / λ

c) $N_0 + \frac{\alpha}{\lambda}$

d) $\frac{\lambda}{\sigma} + N_0 s$

A. N_0

B. α / λ

C. $N_0 + \frac{\alpha}{\lambda}$

D. $\frac{\lambda}{\sigma} + N_0 s$

Answer: B



Watch Video Solution

111. An analysis of the rock shows that the relative number of Sr^{87} and Rb^{87} ($t_{1/2} = 4.7 \times 10^{10}$ year) atoms is 0.05 . What is the age of the rock?

Assume all the Sr^{87} have been formed from Rb^{87} only

a) 7.26×10^9 year

b) 1.43×10^9 year

c) 3.28×10^9 year

d) 4.32×10^8 year

A. 7.26×10^9 year

B. 1.43×10^9 year

C. 3.28×10^9 year

D. 4.32×10^8 year

Answer: C



Watch Video Solution

112. A radioactive substance (parent) decays to its daughter element . The age of radioactive substance (t) is related to the daughter (d)/parent (p) ratio by the equation :

a) $t = \frac{1}{\lambda} \ln\left(1 + \frac{p}{d}\right)$

b) $t = \frac{1}{\lambda} \ln\left(1 + \frac{d}{p}\right)$

c) $t = \frac{1}{\lambda} \ln\left(\frac{d}{p}\right)$

d) $t = \frac{1}{\lambda} \ln\left(\frac{p}{d}\right)$

$$\text{A. } t = \frac{1}{\lambda} \ln\left(1 + \frac{p}{d}\right)$$

$$\text{B. } t = \frac{1}{\lambda} \ln\left(1 + \frac{d}{p}\right)$$

$$\text{C. } t = \frac{1}{\lambda} \ln\left(\frac{d}{p}\right)$$

$$\text{D. } t = \frac{1}{\lambda} \ln\left(\frac{p}{d}\right)$$

Answer: B

 [Watch Video Solution](#)

113. The reaction $\text{A(g)} + 2\text{B(g)} \rightarrow \text{C(g)}$ is an elementary reaction. In an experiment involving this reaction, the initial pressures of A and B are $P_A = 0.40$ atm and $P_B = 1.0$ atm respectively. When $P_C = 0.3$ atm, the rate of the reaction relative to the initial rate is:-

A. $\frac{1}{12}$

B. $\frac{1}{50}$

C. $\frac{1}{25}$

D. none of these

Answer: C



Watch Video Solution

114. Which of the following is incorrect statement ?

- a) Stoichiometry of a reaction tells about the order of the elementary reaction.
- b) For a zero order reaction. Rate and the rate constant are identical .
- c) A zero order reaction is controlled by factors other than concentration of reactants
- d) A zero order reaction is an elementary reaction

A. Stoichiometry of a reaction tells about the order of the elementary reaction.

B. For a zero order reaction. Rate and the rate constant are identical .

C. A zero order reaction is controlled by factors other than concentration of reactants

D. A zero order reaction is an elementary reaction

Answer: D



Watch Video Solution

115. Two first order reaction have half-lives in the ratio 8:1 Calculate the ratio of time intervals $t_1 : t_2$ and t_1 and t_2 are the time period for $\left(\frac{1}{4}\right)^{th}$ and $\left(\frac{3}{4}\right)^{th}$ completion.

- a) 1:0.301
 - b) 0.125:0.602
 - c) 1:0.602
 - d) none of these
-
- A. 1:0.301
 - B. 0.125:0.602
 - C. 1:0.602
 - D. none of these

Answer: C

116. Reaction $A + B \rightarrow C + D$ follows rate law $r = k[A]^{1/2}[B]^{1/2}$

Starting with 1 M of A and B each. What is the time taken for concentration of A to become 0.1M?

(Given $k = 2.303 \times 10^{-2} \text{ sec}^{-1}$)

- a) 10 sec
- b) 100 sec
- c) 1000 sec
- d) 434 sec

- A. 10 sec
- B. 100 sec
- C. 1000 sec
- D. 434 sec

Answer: B

117. For a first order homogenous gaseous reaction, $A \rightarrow 2B + C$

then initial pressure was P_i while total pressure after time 't' was P_t . The

right expression for the rate constants k in terms of P_i , P_t and t is :

a) $k = \frac{2.303}{t} \log\left(\frac{2P_i}{3P_i - P_t}\right)$

b) $k = \frac{2.303}{t} \log\left(\frac{2P_i}{2P_t - P_i}\right)$

c) $k = \frac{2.303}{t} \log\left(\frac{P_i}{P_i - P_t}\right)$

d) None of these

A. $k = \frac{2.303}{t} \log\left(\frac{P_\infty}{P_\infty - P_t}\right)$

B. $k = \frac{2.303}{t} \log\left(\frac{2P_\infty}{P_\infty - P_t}\right)$

C. $k = \frac{2.303}{t} \log\left(\frac{2P_\infty}{3(P_\infty - P_t)}\right)$

D. none of these

Answer: C



Watch Video Solution

118. A hydrogenation reaction is carried out at $500K$. If the same reaction is carried out in the presence of a catalyst at the same rate, the temperature required is $400K$. Calculate the activation energy of the reaction if the catalyst lowers the activation barrier by $20kJmol^{-1}$.

- a) 100 kJ/mol
- b) 80 kJ/mol
- c) 60 kJ/mol
- d) none of these

A. 100 kJ/mol

B. 80 kJ/mol

C. 60 kJ/mol


D. none of these

Answer: B





Watch Video Solution

119. The following mechanism has been proposed for the exothermic catalyzed complex reaction:




If k_1 is much smaller than k_2 , the most suitable qualitative plot of potential energy (PE) versus reaction coordinates for the above reaction is






B.



C.



D.

Answer: B



Watch Video Solution

120. A radioactive isotope X with half-life of 693×10^9 years decay to Y which is stable. A sample of rock from of the moon was found to contain both the elements X Y in the mole ratio 1: 7 . What is the age of the rock

?

a) 2.079×10^{12} years

b) 1.94×10^{10} years

c) 1.33×10^9 years

d) 10^{10} years

A. 2.079×10^{12} years

B. 1.94×10^{10} years

C. 1.33×10^9 years

D. 10^{10} years

Answer: A



[Watch Video Solution](#)

121. The ratio of activities of two radio nuclides X and Y in a mixture at time $t = 0$ was found to be 4:1. After two hours, the ratio activities become 1:1. If the $t_{1/2}$ of radio nuclide X is 20 min then $t_{1/2}$ [in minutes] of radio nuclide Y is ,

- a) 10
- b) 20
- c) 30
- d) 40

A. 10

B. 20

C. 30


D. 40

Answer: C



Watch Video Solution

122. Two consecutive irreversible first order reactions can be represented by



The rate equation for A is readily integrated to obtain

$$[A]_t = [A]_0 \cdot e^{-k_1 t}, \text{ and } [B]_t = \frac{k_1 [A]_0}{k_2 - k_1} \left[e^{-k_1(t)} - e^{-k_2(t)} \right]$$

At what time will B be present in maximum concentration ?

- a) $\frac{K_1}{K_2 - K_1}$
b) $\frac{1}{k_1 - k_2} \ln \frac{k_1}{k_2}$
c) $\frac{1}{k_2 - k_1} \ln \frac{k_1}{k_2}$
d) none of these

A. $\frac{K_1}{K_2 - K_1}$

B. $\frac{1}{k_1 - k_2} \ln \frac{k_1}{k_2}$

C. $\frac{1}{k_2 - k_1} \ln \frac{k_1}{k_2}$

D. none of these


Answer: B



[Watch Video Solution](#)

123. Two consecutive irreversible first order reactions can be represented






by




The rate equation for A is readily integrated to obtain

$$[A]_t = [A]_0 \cdot e^{-k_1 t}, \text{ and } [B]_t = \frac{k_1 [A]_0}{k_2 - k_1} \left[e^{-k_1(t)} - e^{-k_2(t)} \right]$$

When $k_1 = 1s^{-1}$ and $k_2 = 500s^{-1}$, select most appropriate graph






D.

Answer: B

 [Watch Video Solution](#)

124. Two consecutive irreversible first order reactions can be represented by



The rate equation for A is readily integrated to obtain

$$[A]_t = [A]_0 \cdot e^{-k_1(t)}, \text{ and } [B]_t = \frac{k_1[A]_0}{k_2 - k_1} \left[e^{-k_1(t)} - e^{-k_2(t)} \right]$$

Select the correct statement for given reaction:

- A decreases linearly
- B rise to a max. and then constant
- B rises to a max and the falls
- The slowest rate of increases of C occurring where B is max

- A. A decreases linearly
- B. B rise to a max. and then constant
- C. B rises to a max and the falls
- D. The slowest rate of increases of C occurring where B is max

Answer: C

 [Watch Video Solution](#)

125. Arrhenius studies the effect of temperature on the rate of a reaction and postulated that rate constant varies with temperature exponentially as $k = Ae^{E_a/RT}$. This method is generally used for finding the activation energy of a reaction. Keeping temperature constant, the effect of catalyst on the activation energy has also been studied.

The pre-exponential factor in the Arrhenius equation of a first order reaction has the unit :

- a) $\text{molL}^{-1}\text{s}^{-1}$
- b) $\text{Lmol}^{-1}\text{s}^{-1}$

c) s^{-1}

d) dimensionless

A. $\text{mol L}^{-1} \text{s}^{-1}$

B. $\text{L mol}^{-1} \text{s}^{-1}$

C. s^{-1}

D. dimensionless

Answer: C



[Watch Video Solution](#)

126. Arrhenius studies the effect of temperature on the rate of a reaction and postulated that rate constant varies with temperature exponentially as $k = Ae^{-E_a/RT}$. This method is generally used for finding the activation energy of a reaction. Keeping temperature constant, the effect of catalyst on the activation energy has also been studied.

If x is the fraction of molecules having energy greater than E_a it will be given by :

a) $x = -\frac{E_a}{RT}$

b) $\ln x = -\frac{E_a}{RT}$

c) $x = e^{E_a/RT}$

d) Any of these

A. $x = -\frac{E_a}{RT}$

B. $\ln x = -\frac{E_a}{RT}$

C. $x = e^{E_a/RT}$

D. Any of these

Answer: B



Watch Video Solution

127. Arrhenius studies the effect of temperature on the rate of a reaction and postulated that rate constant varies with temperature exponentially as $k = Ae^{E_a/RT}$. This method is generally used for finding the activation energy of a reaction. Keeping temperature constant, the effect of catalyst on the activation energy has also been studied.

If the rate of reaction doubles for $10^\circ C$ rise of temperature from 290K to 300K, the activation energy of the reaction will be approximately:

a) 40 Kcal mol^{-1}

b) 12 Kcal mol^{-1}

c) 60 Kcal mol^{-1}

d) 70 Kcal mol^{-1}

A. 40 Kcal mol^{-1}

B. 12 Kcal mol^{-1}

C. 60 Kcal mol^{-1}

D. 70 Kcal mol^{-1}

Answer: B




Watch Video Solution

128. An important parameter of a photochemical reaction is the quantum efficiency or quantum yield (ϕ) which is defined as

$$\phi = \frac{\text{moles of the substance reaction}}{\text{moles of photons absorbed}}$$

Absorption of UV radiation decompose acetone according to the reaction



The quantum yield of a reaction at $\lambda = 330nm$ is 0.4 . A sample of acetone absorbs monochromatic radiation at $\lambda 330mn$ at the rate of $7.2 \times 10^{-3} Js^{-1}$ (given : $N_A = 6 \times 10^{23}$, $h = 6.6 \times 10^{-34}$ in S.I unit).

The rate of formation of $CO(mol/s)$ is :

- a) 2×10^{-8}
- b) 8×10^{-8}
- c) 8×10^{-9}
- d) none of these

A. 2×10^{-8}

B. 8×10^{-8}

C. 8×10^{-9}

D. none of these


Answer: C

 [Watch Video Solution](#)

129. An important parameter of a photochemical reaction is the quantum efficiency or quantum yield (ϕ) which is defined as

$$\phi = \frac{\text{moles of the substance reaction}}{\text{moles of photons absorbed}}$$

Absorption of UV radiation decompose acetone according to the reaction



The quantum yield of a reaction at $\lambda = 330nm$ is 0.4 . A sample of acetone absorbs monochromatic radiation at $\lambda 330mn$ at the rate of $7.2 \times 10^{-3} Js^{-1}$ (given : $N_A = 6 \times 10^{23}$, $h = 6.6 \times 10^{-34}$ in S.I unit).

The rate of formation of $CO(mol/s)$ is :

- a) 2×10^{-8}
- b) 8×10^{-8}
- c) 8×10^{-9}
- d) none of these

A. 2×10^{-8}

B. 1.6×10^{-9}

C. 16×10^{-9}

D. 8×10^{-9}

Answer: C



Watch Video Solution

130. Radioactive disintegration is a first order reaction and its rate depends only upon the nature of nucleus and does not depend upon external factors like temperature and pressure. The rate of radioactive disintegration (Activity) is represented as

$-\frac{dN}{dt} = \lambda N$ Where $\lambda =$ decay constant, $N =$ number of nuclei at time t ,

$N_0 =$ initial no. of nuclei. The above equation after integration can be represented as

$$\lambda = \frac{2.303}{t} \log\left(\frac{N_0}{N}\right)$$

Half-life period of U is 2.5×10^5 years. In how much time will the amount of U^{237} remaining be only 25 % of the original amount ?

a) 2.5×10^5 year

b) 1.25×10^5 years

c) 5×10^5 years

d) none of these

A. 2.5×10^5 year

B. 1.25×10^5 years

C. 5×10^5 years

D. none of these

Answer: C

 [Watch Video Solution](#)

131. Radioactive disintegration is a first order reaction and its rate depends only upon the nature of nucleus and does not depend upon external factors like temperature and pressure. The rate of radioactive disintegration (Activity) is represented as

$$-\frac{dN}{dt} = \lambda N \text{ Where } \lambda = \text{decay constant, } N = \text{number of nuclei at time } t,$$

N_0 = initial no. of nuclei. The above equation after integration can be represented as

$$\lambda = \frac{2.303}{t} \log \left(\frac{N_0}{N} \right)$$

Calculate the half-life period of a radioactive element which remains only

1/16 of its original amount in 4740 years:

- a) 1185 years
- b) 2370 years
- c) 52.5 years
- d) none of these

A. 1185 years

B. 2370 years

C. 52.5 years

D. none of these

Answer: A



[Watch Video Solution](#)

132. Radioactive disintegration is a first order reaction and its rate depends only upon the nature of nucleus and does not depend upon external factors like temperature and pressure. The rate of radioactive disintegration (Activity) is represented as

$-\frac{dN}{dt} = \lambda N$ Where $\lambda =$ decay constant, $N =$ number of nuclei at time t ,

$N_0 =$ initial no. of nuclei. The above equation after integration can be represented as

$$\lambda = \frac{2.303}{t} \log\left(\frac{N_0}{N}\right)$$

What is the activity in Ci (curie) of 1.0mole plutonium -239 ?

($t_{1/2} = 24000$ years)

- a) 1.49 Ci
- b) 14.9 Ci
- c) 5.513×10^{11} Ci
- d) None of these

A. 1.49 Ci

B. 14.9 Ci

C. 5.513×10^{11} Ci

D. None of these

Answer: B



Watch Video Solution

133. Size of nucleus was obtained by the equation $r = R_0 A^{1/3}$, Where r is the radius of nucleus of mass no A . and R_0 is a constant whose value is equal to 1.5×10^{-15} metre.

(Given : 1 amu = 1.66×10^{-24} g)

What is the density of a nucleus of mass number A ?

a) $\frac{4}{3}\pi(1.5 \times 10^{-15})^3 A$

b) $1.17 \times 10^{17} \text{ kg/cm}^3$

c) $1.17 \times 10^{17} \text{ kg/m}^3$

d) none of these

A. $\frac{4}{3}\pi(1.5 \times 10^{-15})^3 A$

B. $1.17 \times 10^{17} \text{ kg/cm}^3$

C. $1.17 \times 10^{-17} \text{ kg/m}^3$

D. none of these

Answer: B



Watch Video Solution

134. Size of nucleus was obtained by the equation $r = R_0 A^{1/3}$, Where r is the radius of nucleus of mass no A . and R_0 is a constant whose value is equal to 1.5×10^{-15} metre.

(Given : $1 \text{ amu} = 1.66 \times 10^{-24} \text{ g}$)

Nucleus radius of ${}_6\text{C}^{12}$ is 3×10^{-15} metre. What is density ratio of $d_c/d_{\text{H}_2\text{O}}$?

- a) 1.76×10^{17}
- b) 1.76×10^{14}
- c) 17.6×10^7
- d) 17.6×10^{17}

A. 1.76×10^{17}

B. 1.76×10^{14}

C. 17.6×10^7

D. 17.6×10^{17}

Answer: B



Watch Video Solution

135. Select the correct statement(s):

- A. Rate constant are never negative
- B. Partial orders are never negative
- C. Molecularity and order of reaction both are equal for elementary reaction
- D. Order of reaction may be change with change in practical conditions (temp. and pressure)

Answer: A::C::D

 [Watch Video Solution](#)

136. Select the correct statement(s):

- A. the rate of reaction decreases with decreases in temperature
- B. The rate of reaction is uniform in zero order reaction

- C. The rate of reaction depends upon the surface area of the solid reactants
- D. Average and instantaneous rate of reaction defined for macro and micro-scopic time interval respectively

Answer: A::B::C

 [Watch Video Solution](#)

137. Select the correct statement(s) :

A. The rate law of the elementary reaction , $2A \rightarrow B + C$, must be

$$r = k[A]^2$$

B. The rate law for the complex reaction $A + B \rightarrow C$, might not be

$$r = k[A][B]$$

C. If the partial orders differ from the stoichiometric coefficients in the balanced reaction, the reaction must be complex

D. If the partial orders are equal to corresponding coefficients in the balanced reaction, the reaction must be elementary

Answer: A::B::C

 [Watch Video Solution](#)

138. Select the correct statement(s) :

A. Every substance that appears in the rate law of reaction must be a reactant or product in that reaction

B. If we know the rate law of a reaction, we can deduce its mechanism

C. If the reaction has rate $r = k[A][B]^{3/2}$ then reaction may be elementary

D. A zero order reaction must be a complex reaction

Answer: D

 [Watch Video Solution](#)

139. Select the correct statement(s) :

a) In the reaction ${}_{92}^{235}\text{U} + {}_0^1\text{n} \rightarrow {}_{56}^{140}\text{Ba} + {}_{36}^{94}\text{Kr} + 2{}_0^1\text{n} + x$, x is

b) In the reaction ${}_{11}^{23}\text{Na} + z \rightarrow {}_{12}^{23}\text{Mg} + {}_0^1\text{n}$, the bombarding particle z is deuteron

c) Very large amount of energy is produced during nuclear fusion and nuclear fission

d) In a fission reaction, a loss in mass occurs releasing a vast amount of energy

A. In the reaction ${}_{92}^{235}\text{U} + {}_0^1\text{n} \rightarrow {}_{56}^{140}\text{Ba} + {}_{36}^{94}\text{Kr} + 2{}_0^1\text{n} + x$, x is

B. In the reaction ${}_{11}^{23}\text{Na} + z \rightarrow {}_{12}^{23}\text{Mg} + {}_0^1\text{n}$, the bombarding particle z is deuteron

C. Very large amount of energy is produced during nuclear fusion and nuclear fission


D. In a fission reaction, a loss in mass occurs releasing a vast amount of energy

Answer: A::C::D



Watch Video Solution

140. In the decay process:



- a) A and B are isodiaphers
- b) A and C are isotones
- c) A and C are isotopes
- d) B , C and D are isobars

A. A and B are isobars

B. A and D are isotopes

C. B , C and D are isobars

D. A and C are isotones

Answer: B::C



Watch Video Solution

141. Match the following columns

Column-I

- (A) Unit of k is always equals to
- (B) Unit of k in zero order
- (C) Unit of k in first order
- (D) Unit of k in second order

Column-II

- (P) 1/time
- (Q) M/time
- (R) $\text{Time}^{-1} M^{-1}$
- (S) Unit of A (pre-exponential factor)

 [Watch Video Solution](#)

142. Match the following columns

Column-I

- (A) Molecularity of a reaction
- (B) Order of reaction
- (C) The dissociation of H_2O_2 (aq) is
- (D) $\text{H}_2(\text{g}) + \text{Cl}_2(\text{g}) \xrightarrow{h\nu} 2\text{HCl}$ is

Column-II

- (P) 0, 1 Possible
- (Q) 1, 2 Possible
- (R) First order reaction
- (S) Zero order reaction

 [Watch Video Solution](#)

143. Match the following columns

Column-I (Curve)

- (A) C vs t (abscissa) for zero order
(B) $\log C$ vs t (abscissa) for first order
(C) $\left(\frac{-dc}{dt}\right)$ vs C for zero order
(D) $\ln\left(\frac{-dc}{dt}\right)$ vs $\ln C$ for first order

Column-II (Slope)

- (P) unity
(Q) zero
(R) $-k$
(S) $-\frac{k}{2.303}$

 [Watch Video Solution](#)

Column-I

- (A) Isotones
(B) Isobars
(C) Isotopes
(D) Isodiaphers

Column-II

- (P) ${}^{234}_{91}\text{Pa}$ and ${}^{234}_{90}\text{Th}$
(Q) ${}^{12}_6\text{C}$ and ${}^{14}_6\text{C}$
(R) ${}^{39}_{19}\text{K}$ and ${}^{19}_9\text{F}$
(S) ${}^{39}_{18}\text{Ar}$ and ${}^{40}_{19}\text{K}$

144.

 [Watch Video Solution](#)

145. Each question contains STATEMENTS-1 (Assertion) and STATEMENT -2 (Reason).

Examine the statements carefully and mark the correct answer according to the instructions given below:

STATEMENT-1: Molecularity has no meaning for a complex reactions

STATEMENT-2: Molecularity defined only for RDS

- a) If both the statements are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1
- b) If both the statements are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1
- c) If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- d) If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

A. If both the statements are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1

B. If both the statements are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1

C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE

D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

Answer: C



[Watch Video Solution](#)

146. Each question contains STATEMENTS-1 (Assertion) and STATEMENT -2 (Reason).

Examine the statements carefully and mark the correct answer according to the instructions given below:

STATEMENT-1: An elementary reaction cannot have fractional order.

STATEMENT-2: Stoichiometric coefficients in an elementary reaction can be fractional.

a) If both the statements are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1

b) If both the statements are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1

c) If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE

d) If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

A. If both the statements are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1

B. If both the statements are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1

C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE

D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

Answer: C

 [Watch Video Solution](#)

147. Each question contains STATEMENTS-1 (Assertion) and STATEMENT -2 (Reason).

Examine the statements carefully and mark the correct answer according to the instructions given below:

STATEMENT-1: Concentration of reactant in zero order reaction is constant.

STATEMENT-2: For zero order reaction $A \rightarrow B$, successive half life of reaction decrease with the progress of the reaction.

A. If both the statements are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1

- B. If both the statements are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1
- C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

Answer: D

 [Watch Video Solution](#)

148. Each question contains STATEMENTS-1 (Assertion) and STATEMENT -2 (Reason).

Examine the statements carefully and mark the correct answer according to the instructions given below:

STATEMENT-1: The order of reaction can have fractional value.

STATEMENT-2: For an elementary reaction, the partial orders are determined by the reaction stoichiometry

- A. If both the statements are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1
- B. If both the statement are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1
- C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

Answer: B



Watch Video Solution

149. Each question contains STATEMENTS-1 (Assertion) and STATEMENT -2 (Reason).

Examine the statements carefully and mark the correct answer according to the instructions given below:

STATEMENT-1: Catalyst may increase the rate constant to a large extent.

STATEMENT-2:By using suitable catalyst, we can significantly increase yield.

- a) If both the statements are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1
- b) If both the statement are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1
- c) If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- d) If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE
-
- A. If both the statements are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1
- B. If both the statement are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1
- C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

Answer: C



[Watch Video Solution](#)

150. Each question contains STATEMENTS-1 (Assertion) and STATEMENT -2 (Reason).

Examine the statements carefully and mark the correct answer according to the instructions given below:

STATEMENT-1: Product is formed only when the required orientation and energy conditions are met.

STATEMENT-2: All collisions between reactants yield the desired product

- a) If both the statements are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1
- b) If both the statements are TRUE and STATEMENT-2 is NOT the correct explanation of STATEMENT-1
- c) If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- d) If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

A. If both the statements are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1

B. If both the statement are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1

C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE

D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

Answer: C

 [Watch Video Solution](#)

151. Each question contains STATEMENTS-1 (Assertion) and STATEMENT -2 (Reason).

Examine the statements carefully and mark the correct answer according to the instructions given below:

STATEMENT-1: The plot of k versus $1/T$ is linear.

STATEMENT-2: $k = A \cdot e^{-E_a/RT}$

A. If both the statements are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1

B. If both the statement are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1

C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE

D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

Answer: D

 [Watch Video Solution](#)

152. Each question contains STATEMENTS-1 (Assertion) and STATEMENT -2 (Reason).

Examine the statements carefully and mark the correct answer according to the instructions given below:

STATEMENT-1: For exothermic reaction equilibrium constant decrease with increase in temperature.

STATEMETN-2: For exothermic reaction rate constant decrease with decrease in temperature.

A. If both the statements are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1

- B. If both the statements are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1
- C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

Answer: B

 [Watch Video Solution](#)

153. Each question contains STATEMENTS-1 (Assertion) and STATEMENT -2 (Reason).

Examine the statements carefully and mark the correct answer according to the instructions given below:

STATEMENT -1: If the activation energy of reaction is zero, temperature will have no effect on the rate constant.

STATEMENT-2: Lower the activation energy faster is the reaction.

- A. If both the statements are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1
- B. If both the statement are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1
- C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

Answer: B



[Watch Video Solution](#)

154. Each question contains STATEMENTS-1 (Assertion) and STATEMENT -2 (Reason).

Examine the statements carefully and mark the correct answer according to the instructions given below:

STATEMENT-1: Active complex is an intermediate product.

STATEMENT-2: Active complex is unstable with high vibrational energy.

- A. If both the statements are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1
- B. If both the statement are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1
- C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

Answer: D



[Watch Video Solution](#)

155. Each question contains STATEMENTS-1 (Assertion) and STATEMENT -2 (Reason).

Examine the statements carefully and mark the correct answer according to the instructions given below:

STATEMENT-1: The pre-exponential factor A has the same units for all reactions.

STATEMENT -2: $e^{-E_a/RT}$ has no unit.

- A. If both the statements are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1
- B. If both the statement are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1
- C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

Answer: D

 [Watch Video Solution](#)

156. Each question contains STATEMENTS-1 (Assertion) and STATEMENT -2 (Reason).

Examine the statements carefully and mark the correct answer according to the instructions given below:

STATEMENT -1: γ – rays have very high penetrating power.

STATEMENT-2: γ -rays are electromagnetic raidations of high energy.

- A. If both the statements are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1
- B. If both the statement are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1
- C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

Answer: A



[Watch Video Solution](#)

157. Each question contains STATEMENTS-1 (Assertion) and STATEMENT -2 (Reason).

Examine the statements carefully and mark the correct answer according to the instructions given below:

STATEMENT-1: Disintegration of ${}_{-1}^3\text{H}$ (tritium) is accompanied by β – emission.

STATEMENT-2: Tritium has high n/p ratio.

- A. If both the statements are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1
- B. If both the statement are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1
- C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

Answer: A



[Watch Video Solution](#)

158. Each question contains STATEMENTS-1 (Assertion) and STATEMENT -2 (Reason).

Examine the statements carefully and mark the correct answer according to the instructions given below:

STATEMENT-1: The life of radioactive object (organic origin) can found with the help of carbon dating.

STATEMENT-2: ${}_{6}^{14}\text{C}$ is a α and β -emitter.

- A. If both the statements are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1
- B. If both the statement are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1
- C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

Answer: C



[Watch Video Solution](#)

159. Each question contains STATEMENTS-1 (Assertion) and STATEMENT -2 (Reason).

Examine the statements carefully and mark the correct answer according to the instructions given below:

STATEMENT-1: Neutron are the best bombarding particles.

STATEMENT-2: Neutrons are neutral particles.

- A. If both the statements are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1
- B. If both the statement are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1
- C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

Answer: A



Watch Video Solution

160. Each question contains STATEMENTS-1 (Assertion) and STATEMENT -2 (Reason).

Examine the statements carefully and mark the correct answer according to the instructions given below:

STATEMENT-1: Nucleus does not contain free electrons, yet it emit beta-particles

STATEMENT-2: At high n/p ratio, one neutron supposed to give 1 proton and $1 e^{-}$ (β).

- A. If both the statements are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1
- B. If both the statement are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1
- C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

Answer: A



[Watch Video Solution](#)

161. Each question contains STATEMENTS-1 (Assertion) and STATEMENT -2 (Reason).

Examine the statements carefully and mark the correct answer according to the instructions given below:

STATEMENT-1: Rate of disintegration of thorium increases with the increase in moles of thorium.

STATEMENT-2: Rate of disintegration does not depend upon temperature, pressure

- A. If both the statements are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1
- B. If both the statement are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1
- C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

Answer: B



[Watch Video Solution](#)

162. The rate of decomposition of $NH_3(g)$ at 10 atm on platinum surface is zero order . What is rate of formation (in $M \min^{-1}$) of $H_2(g)$, if rate


constant of reaction $2NH_3(g) \rightarrow N_2(g) + 3H_2(g)$ is 2.0 M min^{-1} ?

 [Watch Video Solution](#)

163. In an elementary reaction $A(g) + 2B(g) \rightarrow C(g)$ the initial pressure of A and B are $P_A = 0.40 \text{ atm}$ and $P_B = 0.60 \text{ atm}$ respectively. After time T, if pressure of C is observed 0.1 atm, then find the value of $\frac{r_i(\text{ initial rate of reaction})}{r_t(\text{ rate of reaction after time t})}$.

 [Watch Video Solution](#)

164. Carbon monoxide reacts with O_2 to form CO_2 :



Informations about this reaction are given in the table below.

$[CO]$ mol/L	$[O_2]$ mol/L	Rate of reaction (mol/L.min)
0.02	0.02	4×10^{-5}
0.04	0.02	1.6×10^{-4}
0.02	0.04	8×10^{-5}

What is the value for the rate constant for the reaction in properly related unit?



Watch Video Solution

165. Half-life for the zero order reaction $A(g) \rightarrow B(g) + C(g)$ and half-life for the first order reaction $X(g) \rightarrow Y(g) + Z(g)$ are equal. If completion time for the zero order reaction is 13.86 min, then calculate the rate constant (in hr^{-1}) of the reaction $X(g) \rightarrow Y(g) + Z(g)$.



Watch Video Solution

166. For any acid catalysed reaction, $A \xrightarrow{H^+} B$

half-life period is independent of concentration of A at given pH. At definite concentration of A half-time is 10min at pH=2 and half-time is 100 min at pH=3. If the rate law expression of reaction is $r = k[A]^x [H^+]^y$ then calculate the value of (x+y).



Watch Video Solution

167. Iodine -131 is a radioactive isotope. If 1.0 mg of ^{131}I has an activity of 4.6×10^{12} Bq. What is the half-life of ^{131}I (in days)

 [Watch Video Solution](#)

168. The average life of a radioactive element is 7.2 min. Calculate the time interval (in min) between the stages of 33.33% and 66.66% decay

 [Watch Video Solution](#)

169. A, B and C are isodiaphers while C, D and E are isobars. Calculate the difference of protons between A and E. ${}_{82}^{206}\text{A} \rightarrow \text{B} \rightarrow \text{C} \rightarrow \text{D} \rightarrow \text{E}$


Given: Isodiaphers and isobars are formed in successive α and β – emission respectively.

 [Watch Video Solution](#)

170. ${}_{90}^{234}\text{Th}$ disintegrates to give ${}_{82}^{206}\text{Pb}$ as the final product. Total no. of α and β particle emitted out during this process are:

 [Watch Video Solution](#)

171. In the given radioactive disintegration series



Calculate difference between number of α and number of β particles emitted in this series.

 [Watch Video Solution](#)

Level 1 Q 3 To Q 32

1. The following data pertain to reaction between A and B

S.No	[A]	[B]	Rate
	molL^{-1}	molL^{-1}	$\text{molL}^{-1} \text{sec}^{-1}$
<i>I</i>	1×10^{-2}	2×10^{-2}	2×10^{-4}
<i>II</i>	2×10^{-2}	2×10^{-2}	4×10^{-4}
<i>III</i>	2×10^{-2}	4×10^{-2}	8×10^{-4}

Which of the following inference(s) can be drawn from the above data

- (a) Rate constant of the reaction 10^{-4}
- (b) Rate law of reaction is $k[A][B]$
- (c) Rate of reaction increases four times on doubling the concentration of both the reactant
- Select the correct answer codes:

a) a,b and C

b) a and b

c) b and c

d) c alone

A. a,b and C

B. a and b

C. b and c

D. c alone

Answer: C

 Watch Video Solution

Level 1 Q 33 To Q 62

1.

column I

P. Zero order reaction

Q. First order reaction

R. second order reactions

S. Pseudo unimolecular reaction

column II

1. $t_{1/2} \propto \frac{1}{[A]_0}$

2. $t_{100\%} = [A]_0 / k$

3. Involves at least two reactants

4. $[A] = [A]_0 e^{-kt}$

A.

<i>P</i>	<i>Q</i>	<i>R</i>	<i>S</i>
2	1	4	2

B.

<i>P</i>	<i>Q</i>	<i>R</i>	<i>S</i>
2	4	1	3

C.

<i>P</i>	<i>Q</i>	<i>R</i>	<i>S</i>
2	1	3	4


D.

<i>P</i>	<i>Q</i>	<i>R</i>	<i>S</i>
3	2	1	4

Answer: B

 Watch Video Solution

2. The decomposition of N_2O in carbon tetrachloride was followed by measuring the volume of O_2 gas evolved :



The maximum volume of O_2 gas obtained was 100 cm^3 . In 500 minutes, 90 cm of O_2 were evolved. The

first order rate constant (in min^{-1}) for the disappearance of N_2O is :

- a) $\frac{2.303}{500}$
- b) $\frac{2.303}{500} \log \frac{100}{90}$
- c) $\frac{2.303}{500} \log \frac{90}{100}$
- d) $\frac{100}{10 \times 500}$

- A. $\frac{2.303}{500}$
- B. $\frac{2.303}{500} \log \frac{100}{90}$
- C. $\frac{2.303}{500} \log \frac{90}{100}$
- D. $\frac{100}{10 \times 500}$

Answer: A



Watch Video Solution

Level 1 Q 63 To Q 92

1. A first order reaction is 50% completed in 20 minutes at 27°C and in 5 minutes at 47° . The energy of activation of the reaction is :

- a) 43.85 KJ/mol
- b) 55.14 KJ/mol
- c) 11.97 KJ/mol
- d) 6.65 KJ/mol

A. 43.85 KJ/mol

B. 55.14 KJ/mol

C. 11.97 KJ/mol

D. 6.65 KJ/mol

Answer: B



Watch Video Solution

2. A catalyst lowers the activation energy for a certain reaction from 83.314 to 75 KJ mol⁻¹ at 500 K . What will be the rate of reaction as compared to uncatalyst reaction ? Assume other things being equal

- a) Double
- b) 28 times
- c) 7.38 times
- d) 7.38×10^3 times

A. Double

B. 28 times

C. 7.38 times


D. 7.38×10^3 times

Answer: C



[Watch Video Solution](#)

1. In the radioactive decay



the sequence of the

radiation emitted is :

A. α, β, γ

B. γ, α, β

C. β, γ, α

D. β, α, γ

Answer: D



[Watch Video Solution](#)

2. A radioactive nuclide emits γ - rays due to the :

a) emission of an electron from its orbital

b) nuclear energy transition from a higher state to a lower state

c) presence of less neutrons than protons

d) presence of more neutrons than protons

- A. emission of an electron from its orbital
- B. nuclear energy transition from a higher state to a lower state
- C. presence of less neutrons than protons
- D. presence of more neutrons than protons

Answer: B

 [Watch Video Solution](#)

3. Consider the following decay ${}^A_Z X \rightarrow {}^A_{Z+1} Y + {}^0_{-1} e$, X is unstable because:

- A. its nucleus has excess energy
- B. $\frac{n}{p}$ ratio is high
- C. $\frac{n}{p}$ ratio is low
- D. none of these

Answer: B



Watch Video Solution

4. Consider the following decay ${}^A_Z X \rightarrow {}^A_{Z-1} Y + {}^0_{+1} e$, $(\beta^+)X$ is unstable because:

A. its nucleus has excess energy

B. $\frac{n}{p}$ ratio is high

C. $\frac{n}{p}$ ratio is low

D. none of these

Answer: C



Watch Video Solution

5. Which of the following processes causes the emission of X-ray?

a) α -emission

b) β -emission

c) β^+ (Positron) emission

d) electron capture

A. α -emission

B. β -emission

C. β^+ (Positron) emission

D. electron capture

Answer: D



Watch Video Solution

6. Which of the following processes result in an increase in the atomic number of a nuclide?

a) α -emission

b) electron capture

c) γ -emission

d) β^- emission

A. α -emission

B. electron capture

C. γ -emission

D. β -(Beta) emission

Answer: D



Watch Video Solution

7. Is produced when a positron and an electron collide.

a) X-ray

b) Neutron

c) γ -radiation

d) Neutrino

A. X-ray

B. Neutron

C. γ -radiation

D. Neutrino

Answer: C



Watch Video Solution

8. ${}_{67}^{165}\text{Ho}$ is stable isotope. ${}_{67}^{150}\text{Ho}$ is expected to distegrate by:

A. α -emission

B. β -emission

C. Positron emission

D. γ -emission

Answer: C



Watch Video Solution

9. ${}_1^1\text{H}$ is a stable isotope. ${}_1^3\text{H}$ is expected to disintegrated by :

- A. α -emission
- B. β -emission
- C. Positron emission
- D. proton emission

Answer: B



[Watch Video Solution](#)




10. Emission of β -particle is equivalent to:

- A. increase of one proton only
- B. decrease of one neutron only
- C. both (a) and (b)
- D. none of these

Answer: C

 [Watch Video Solution](#)

11. Pair of isobar is :



D. None of these

Answer: A

 [Watch Video Solution](#)

12. The 'group displacement law' was given by :

A. Bacquerel

B. Rutherford

C. Madam Curie

D. Soddy and Fajan

Answer: D

 [Watch Video Solution](#)

13. ${}^7_3\text{Li} + {}^1_1\text{P} \rightarrow X$, Identify X if reaction is (p, α) type.

A. ${}^8_4\text{Be}$

B. ${}^4_2\text{He}$


C. ${}^0_0\gamma$

D. none of these

Answer: B

 [Watch Video Solution](#)

14. Identify reaction type:



- a) (d,p)
- b) (p,p)
- c) (p,d)
- d) none of these

A. (d,p)

B. (p,p)

C. (p,d)

D. none of these



Answer: A






[Watch Video Solution](#)

15. ${}_{13}^{27}\text{Al} + {}_1^1\text{P} \rightarrow X + {}_0^0\gamma$, type artificial radioactive reaction.

a) ${}_{13}^{28}\text{Al}$



d) none of these






D. none of these


Answer: C



Watch Video Solution

16. What will be the product of reaction ${}_{101}^{255}\text{Md}(\alpha, 2n)$?









Answer: C



Watch Video Solution

17. Proton bombardment on Th^{230} followed by emission of two alpha particles will produce:











Answer: C



Watch Video Solution

18. ${}_{83}^{214}\text{Bi}$ decays to A by α -emission. A then decays to B by beta emission, which further decays to C by another beta emission. Element C decays to D by still another beta emission, and D decays by α -emission to form a stable isotope E. What is element E?



Answer: D



Watch Video Solution

19. The activity of a radioactive nuclide X^{100} is 6.023 curie at a certain time 't'. If its disintegration constant is $3.7 \times 10^4 s^{-1}$ the mass of X after t sec is :

a) $6.022 \times 10^6 \text{ g}$

b) 10^{-13} g

c) 10^{-15} g

d) 10^{-17} g

A. $6.022 \times 10^6 \text{ g}$

B. 10^{-13} g

C. 10^{-15} g

D. 10^{-17} g

Answer: C



Watch Video Solution

1. Activity of a radioactive substance is A_1 at time t_1 and A_2 at time t_2 ($t_2 > t_1$), then the ratio of $\frac{A_2}{A_1}$ is:

a) $e^{\lambda(t_2+t_1)}$

b) $e^{\lambda(t_1-t_2)}$

c) $e^{-\lambda(t_1+t_2)}$

d) $\frac{t_2}{t_1}$

A. $e^{\lambda(t_2+t_1)}$

B. $e^{\lambda(t_1-t_2)}$

C. $e^{-\lambda(t_1+t_2)}$

D. $\frac{t_2}{t_1}$

Answer: B



Watch Video Solution

2. The half-life of ${}_{6}^{14}\text{C}$ is 5730 year. What fraction of its original C^{14} would left after 22920 year of storage?

A. 0.5

B. 0.25

C. 0.125

D. 0.0625

Answer: D



[Watch Video Solution](#)

3. The amount of ${}_{6}^{14}\text{C}$ isotope in a piece of wood is found to one fourth (1/4) of that present in a fresh piece of wood. Calculate the age of the piece of wood ($t_{1/2}$ of ${}_{6}^{14}\text{C} = 5770$ years)

a) 7999 year

b) 11543 year

c) 16320 year

d) 23080 year

A. 7999 year

B. 11543 year

C. 16320 year

D. 23080 year

Answer: B



[Watch Video Solution](#)

4. A radioactive element undergoing decay is left 20% of its initial weight after certain period of time t . How many such periods should elapse from the start for the 50% of the element to be left over?

A. 3

B. 4

C. 5

D. none of these

Answer: D

 [Watch Video Solution](#)

5. In a sample of wood, the reading of a counter is 32 dpm and in a fresh sample of tree it is 122dpm . Due to error counter gives the reading 2 dpm in absence of ^{14}C . Half life of ^{14}C is 5770 years .

The approximate age (in years) of wood sample is :

A. 7997.2

B. 57570

C. 11540

D. 15140

Answer: C

 [Watch Video Solution](#)

6. A certain radioactive isotope ${}^A_Z X$ ($t_{1/2} = 100\text{days}$) decays to ${}^{A-8}_{Z-2} Y$.

If 1 mole of ${}^A_Z X$ is kept in sealed container, how much He gas will accumulate at STP in 200 days?

- A. 11.2 litres
- B. 33.6 litres
- C. 22.4 litres
- D. 44.8 litres

Answer: B



[Watch Video Solution](#)

7. Two radio isotopes A and B of atomic mass X and Y are mixed in equal amount by mass. After 20 days, their mass ratio is found to be 1:4.

Isotope A has a half-life of 1 day. The half-life of isotope B is :

- a) 1.11 days
- b) $0.11 \frac{X}{Y}$ day

c) 0.6237 day

d) 1.10 day

A. 1.11 days

B. $0.11 \frac{X}{Y}$ day

C. 0.6237 day

D. 1.10 day

Answer: D



Watch Video Solution

8. There are two radio nuclei A and B is a α -emitter and B is β -emitter. Their disintegration constant are in the ratio of 1:2 . What should be the number of atoms ratio of A and B at time $t=0$, so that initially probability of getting of α and β -particles are same.

A. 2: 1

B. 4: 1

C. 1:2


D. 1:4

Answer: A



Watch Video Solution

9. Ac^{227} has a half-life of 22 years . The decays follows two parallel paths



What are the decay constant (λ) for Th and Fr respectively ?

a) 0.03087,0.00063

b) 0.00063,0.03087

c) 0.02,0.98

d) None of these

A. 0.03087,0.00063

B. 0.00063,0.03087

C. 0.02,0.98

D. None of these

Answer: B

 [Watch Video Solution](#)

10. ${}_{84}^{218}\text{Po}$ ($t_{1/2} = 183$ sec) decay to ${}_{82}^{214}\text{Pb}$ ($t_{1/2} = 161$ sec) by α -emission, while ${}_{82}^{214}\text{Pb}$ is a β -emitter. In an experiment starting with 1 mole of pure ${}_{84}^{218}\text{Po}$, how many time would be required for the number of nuclei of ${}_{82}^{214}\text{Pb}$ to reach maximum ?

A. 147.5

B. 247.5

C. 182

D. 304


Answer: B



Watch Video Solution

Level 2

1. The forward rate constant for the elementary reversible gaseous reaction



What is the rate constant for the backward reaction at this temperature if 10^{-4} moles of CH_3 and 10 moles of C_2H_6 are present in a 10 litre vessel at equilibrium .

a) $1.57 \times 10^9 L \text{ mole}^{-1} s^{-1}$

b) $1.57 \times 10^{10} L \text{ mole}^{-1} s^{-1}$

c) $1.57 \times 10^{11} L \text{ mole}^{-1} s^{-1}$

d) $1.57 \times 10^7 L \text{ mole}^{-1} s^{-1}$

A. $1.57 \times 10^9 L \text{ mole}^{-1} s^{-1}$

B. $1.57 \times 10^{10} \text{ Lmole}^{-1} \text{ s}^{-1}$

C. $1.57 \times 10^{11} \text{ Lmole}^{-1} \text{ s}^{-1}$

D. $1.57 \times 10^7 \text{ Lmole}^{-1} \text{ s}^{-1}$

Answer: D

 [Watch Video Solution](#)

Level 2 Q 3 To Q 32

1. The reaction $A(g) \rightarrow B(g) + 2C(g)$ is a first order reaction with rate constant $2.772 \times 10^{-3} \text{ s}^{-1}$, Starting with 0.1 mole of A in 2 litre vessel, find the concentration of A after 250 sec when the reaction is allowed to take place at constant pressure and at 300 K?

a) $0.0125M$

b) $0.025M$

c) $0.05M$

d) none of these

A. $0.0125M$

B. $0.025M$

C. $0.05M$

D. none of these

Answer: A

 [Watch Video Solution](#)

2. $A(aq) \rightarrow B(aq) + C(aq)$ is first order reaction.

Time $t \quad \infty$

moles of reagent $n_1 \quad n_2$

Reaction progress is measured with the help of titration of reagent 'R'. If

all A, B and C react with reagent and have 'n' factors [n factor, eq. mass =

$\frac{\text{mol.mass}}{n}$ in the ratio of 1:2:3 with the reagent, the k in terms of

t, n_1 and n_2 is :

$$A. k = \frac{10}{t} \ln \left(\frac{n_2}{n_2 - n_1} \right)$$

$$B. k = \frac{1}{t} \ln. \left(\frac{2n_2}{n_2 - n_1} \right)$$

$$C. k = \frac{1}{t} \ln. \left(\frac{4n_2}{n_2 - n_1} \right)$$

$$D. k = \frac{1}{t} \ln. \left(\frac{4n_2}{5(n_2 - n_1)} \right)$$

Answer: D

 [Watch Video Solution](#)

3. The gaseous decomposition reaction, $A(g) \rightarrow 2B(g) + C(g)$ is observed to first order over the excess of liquid water at $25^\circ C$. It is found that after 10 minutes the total pressure of system is 188 torr and after very long time it is 388 torr. The rate constant of the reaction (in hr^{-1}) is : [Given : vapour pressure of H_2O at 25° is 28 torr ($\ln 2 = 0.7$, $\ln 3 = 1.1$, $\ln 10 = 2.3$)]
- a) 0.02
 - b) 1.2
 - c) 0.2
 - d) none of these

A. 0.02

B. 1.2

C. 0.2

D. none of these

Answer: B

 [Watch Video Solution](#)

4. The reaction, Sucrose $\xrightarrow{H^+}$ Glucose + Fructose, take place at certain temperature while the volume of solution is maintained at 1 litre. At time zero the initial rotation of the mixture is 34° . After 30 minutes the total rotation of solutions is 19° and after a very long time, the total rotation is $-11^\circ C$ Find the time when solution was optically inactive.

A. 135 min

B. 103.7 min



C. 38.7 min

D. 45 min

Answer: B

 [Watch Video Solution](#)

5. A compound A dissociates by two parallel first order paths at certain temperature



The reaction is started with 1 mole of pure 'A' litre closed container with initial pressure 2 atm. What is the pressure (in atm) developed in container after 50 minutes from start of experiment?

A. 1.25

B. 0.75

C. 1.50

D. 2.50

Answer: D



Watch Video Solution

6. The reaction $\text{cis-X} \xrightleftharpoons[k_b]{k_f} \text{trans-X}$ is first order in both directions. At 25°C , the equilibrium constant is 0.10 and the rate constant $k_f = 3 \times 10^{-4} \text{ s}^{-1}$. In an experiment starting with the pure cis-form, how long would it take for half of the equilibrium amount of the trans-isomer to be formed?

A. 150 sec

B. 200 sec

C. 240 sec


D. 210 sec

Answer: D



Watch Video Solution

7. Consider the reaction.



The rate constant for two parallel reactions were found to be $10^{-2} \text{ dm}^{-2} \text{ mol}^{-1} \text{ s}^{-1}$ and $4 \times 10^{-2} \text{ dm}^{-3} \text{ mol}^{-1} \text{ s}^{-1}$. If the corresponding energies of activation of the parallel reaction are 100 and 120 kJ/mol respectively, what is the net energy of activation (E_a) of A

- A. 100 kJ/mol
- B. 120 kJ/mol
- C. 116 kJ/mol
- D. 220 kJ/mol

Answer: C

 [Watch Video Solution](#)

8. A reaction takes place in various steps. The rate constant for first, second, third and fifth steps are k_1 , k_2 , k_3 and k_5 respectively. The overall rate constant is given by

$$k = \frac{k_2}{k_3} \left(\frac{k_1}{k_5} \right)^{1/2}$$

If activation energies are 40, 60, 50, and 10 kJ/mol respectively, the overall energy of activation (kJ/mol) is :

- A. 10
- B. 20
- C. 25
- D. none of these

Answer: C

 [Watch Video Solution](#)

9. For reaction $A \rightarrow B$, the rate constant $k_1 = A_1 e^{-E_{a1}/(RT)}$ and for the reaction $X \rightarrow Y$, the rate constant $k_2 = A_2 e^{-E_{a2}/(RT)}$. If $A_1 = 10^8$, $A_2 = 10^{10}$ and $E_{a1} = 600$ cal/mol, $E_{a2} = 1800$ cal/mol then the temperature at which $k_1 = k_2$ is (Given : $R=2$ cal/K-mol)

a) 1200 K

b) $1200 \times 4.606K$

c) $\frac{1200}{4.606}K$

d) $\frac{600}{4.606}K$

A. 1200 K

B. $1200 \times 4.606K$

C. $\frac{1200}{4.606}K$


D. $\frac{600}{4.606}K$

Answer: D



Watch Video Solution

10. For first order parallel reactions k_1 and k_2 are 4 and 2 min^{-1} respectively at 300 K. If the activation energies for the formation of B and C are respectively 30,000 and 38,314 joule/mol respectively, the temperature at which B and C will be obtained in equimolar ratio is :



A. 757.48 k

B. 378.74k

C. 600 k

D. none of these

Answer: B

11. In the series reaction

$A \xrightarrow{K_1} B \xrightarrow{K_2} C \xrightarrow{K_3} D$, if $K_1 > K_2 > K_3$ then the rate determining step of the reaction is

- a) $A \rightarrow B$
- b) $C \rightarrow D$
- c) $B \rightarrow C$
- d) Any step

A. $A \rightarrow B$

B. $C \rightarrow D$


C. $B \rightarrow C$

D. Any step



Answer: B



12. The mechanism of esterification in presence of acid catalyst (H_2SO_4)

is proposed as follows:



Which of the following potential energy V_s reaction co-ordinate diagram is consistent with given mechanism ?





Answer: A



Watch Video Solution

13. For the first order reaction $A \rightarrow B + C$, carried out at 27°C . If $3.8 \times 10^{-16} \%$ of the reactant molecules exists in the activated state, the E_a (activation energy) of the reaction is:

- a) 12 KJ/mol
- b) 831.4 KJ/mol
- c) 100 KJ/mol
- d) 88.57 KJ/mol

A. 12kJ/mol

B. 831.4kJ/mol

C. 100kJ/mol

D. 88.57kJ/mol

Answer: C



Watch Video Solution

14. Upon irradiating californium with neutrons, a scientist discovered a new nuclide having mass number of 250 and a half-life of 30 min. After 90 min. of irradiation, the observed radioactivity due to nuclide was 100 dis/min. How many atoms of the nuclide were prepared initially?

a) 2.4×10^4

b) 3.46×10^4

c) 1900

d) 800

A. 2.4×10^4

B. 3.46×10^4

C. 1900



D. 800

Answer: B



Watch Video Solution

15. The average (mean) life of a radio nuclide which decays by parallel path is



a) 52.63 sec

b) 500 sec

c) 50 sec

d) none of these

A. 52.63 sec

B. 500 sec

C. 50 sec

D. none of these

Answer: C



Watch Video Solution

16. The radioactive decay ${}_{83}^{211}\text{Bi} \rightarrow {}_{81}^{207}\text{Ti}$, takes place in 100L closed vessel at 27°C Starting with 2 mols of ${}_{83}^{211}\text{Bi}$ ($t_{1/2} = 130\text{sec}$), the pressure development in the vessel after 520 sec will be:

- a) 1.875 atm
- b) 0.2155atm
- c) 0.4618 atm
- d) 4.618 atm

A. 1.875 atm

B. 0.2155atm

C. 0.4618 atm

D. 4.618 atm

Answer: C

 [Watch Video Solution](#)

17. A fresh radioactive mixture contains short lived species A and B. Both emitting α -particles initially of 8000 α particles per minute. 20 minutes later, they emit at the rate of 3500 α -particles per minute. If the half-lives of the species A and B are 10 minutes and 500 hours respectively, then the ratio of activities of A: B the initial mixture was:

A. 4: 6

B. 6: 4

C. 3: 4

D. 3: 1

Answer: D

 [Watch Video Solution](#)

18. In order to determine the volume of blood in an animal, a 1.0mL sample of solution of 10^3 dpm of 3_1H is injected into the animal blood stream. After sufficient time for circulatory equilibrium to be established, 2mL of blood is found to have activity to 10 dpm. The volume of blood in animal is :

- a) 199 mL
- b) 198 mL
- c) 200 mL
- d) 20 mL

A. 199 mL

B. 198 mL

C. 200 mL

D. 20 mL

Answer: A



Watch Video Solution

Level 2 Q 33 To Q 35

1. In a sample of wood, the reading of a counter is 32 dpm and in a fresh sample of tree it is 122dpm . Due to error counter gives the reading 2 dpm in absence of ^{14}C . Half life of ^{14}C is 5770 years .

The approximate age (in years) of wood sample is :

A. 5.770×10^3

B. 16.87×10^3

C. 2488

D. none of these

Answer: A



[Watch Video Solution](#)

2. The isotopes ^{238}U and ^{235}U occur in nature in the mass ratio 140 : 1. It is assumed that initially they were found in equal mass. If half life ($t_{1/2}$) of $^{238}\text{U} = 4.5 \times 10^9$ and $t_{1/2}$ of $^{235}\text{U} = 7.13 \times 10^8$ years respectively then the age of earth is ($\log 7 = 0.846$, $\log 2 = 0.3$)


- A. 4.02×10^9 year
- B. 2.01×10^9 year
- C. 8.72×10^9 year
- D. none of these

Answer: A

 [Watch Video Solution](#)

Level 3 Passage


1. Two consecutive irreversible first order reactions can be represented by



The rate equation for A is readily interated to obtain

$$[A]_t = [A]_0 \cdot e^{-k_1 t}, \text{ and } [B]_t = \frac{k_1 [A]_0}{k_2 - k_1} \left[e^{-k_1(t)} - e^{-k_2(t)} \right]$$

When $k_1 = 1s^{-1}$ and $k_2 = 500s^{-1}$, select most appropriate graph



A. graph A

B. graph B

C. graph C

D. graph D

Answer: A



Watch Video Solution

1. Select the correct statement(s) :

a) When $T \rightarrow \infty$ or $E_a \rightarrow 0$ then $K=A$

b) A positive catalyst can change ΔH of the reaction

c) A mixture catalyst may be thermodynamically unstable by kinetically stable

d) A negative catalyst increases the activation energy of the reaction

A. When $T \rightarrow \infty$ or $E_a \rightarrow 0$ then $K=A$

B. A positive catalyst can change ΔH of the reaction

C. A mixture catalyst may be thermodynamically unstable by kinetically stable

D. A negative catalyst increases the activation energy of the reaction

Answer: A::C::D



Watch Video Solution

2. Consider a reaction $A + B \rightarrow C$, in which both reactants are in the same phase, may be

- a) unimolecular elementary reaction
- b) Exothermic
- c) Heterogeneous
- d) Photochemical

A. unimolecular elementary reaction

B. Exothermic

C. Heterogeneous


D. Photochemical

Answer: B::C::D



[Watch Video Solution](#)

3. In the following gaseous phase first order reaction



initial pressure was found to be 400 mm of Hg and it changed to 1000 mm of Hg after 20min. Then

a) half life for A is 10 min

b) rate constant is 0.0693 min^{-1}

c) partial pressure of C at 30 min is 350 mm of Hg

d) total pressure after 30 min is 1100 mm of Hg

A. half life for A is 10 min

B. rate constant is 0.0693 min^{-1}

C. partial pressure of C at 30 min is 350 mm of Hg

D. total pressure after 30 min is 1100 mm of Hg

Answer: A::B::C::D



[Watch Video Solution](#)

4. Identify the true statement(s)

a) A catalyst is chemically unchanged at the end of a reaction

b) A catalyst may appear in the kinetic rate equation of the reaction

c) A catalyst will not affect the composition of an equilibrium mixture
d) A catalyst cannot cause a non-spontaneous ($\Delta G > 0$) reaction to proceed

A. A catalyst is chemically unchanged at the end of a reaction

B. A catalyst may appear in the kinetic rate equation of the reaction

C. A catalyst will not affect the composition of an equilibrium mixture

D. A catalyst cannot cause a non-spontaneous ($\Delta G > 0$) reaction to proceed

Answer: A::B::C::D

 [Watch Video Solution](#)

5. Select the correct statement(s) :

A. α -particles are simply helium atoms

- B. γ -rays travel with higher speed as compared to α -particle and have higher ionization power as compare to β -particle
- C. Loss of β -particles results in the production of isobars
- D. β -particle are considered as the best bombarding particles

Answer: C



Watch Video Solution

6. Select the correct statement(s) :

- a) SI unit of radioactivity is becquerel (Bq)
- b) $1Ci = 3.7 \times 10^7$ Bq
- c) ${}^7_3Li + {}^1_1H \rightarrow {}^4_2He + (P, \alpha)$ type reaction
- d) The half -time of a particular radioactive isotope is a characteristics constant of that isotope

A. SI unit of radioactivity is becquerel (Bq)

B. $1Ci = 3.7 \times 10^7$ Bq

C. ${}^7_3\text{Li} + {}^1_1\text{H} \rightarrow {}^4_2\text{He} + \text{P}$ type reaction

D. The half -time of a particular radioactive isotope is a characteristics constant of that isotope

Answer: A::C::D



Watch Video Solution

7. Select the correct statement(s) :

- a) On bombarding ${}^{14}_7\text{N}$ nuclei with α -particle, the nuclei of the product formed after release of proton would be ${}^{17}_8\text{O}$
- b) Decay constant does not depend upon temperature
- c) Nuclide and its decay product after α -emission are called isodiaphers
- d) Half-life of radium is 1580 years . Its average life will be 1097.22 years

A. On bombarding ${}^{14}_7\text{N}$ nuclei with α -particle, the nuclide of the product formed after release of proton would be ${}^{17}_8\text{O}$

B. Decay constant does not depend upon temperature

C. Nuclide and its decay product after α -emission are called isodiaphers

D. Half-life of radium is 1580 years . Its average life will be 1097.22 years

Answer: A::B::C

 [Watch Video Solution](#)

8. In electron capture (radioactive process):

A. a neutron is formed

B. a proton is consumed

C. γ -ray emission take place

D. X-ray emission takes place

Answer: A::B::D

 [Watch Video Solution](#)

9. Select the correct statement(s) for positron emission by unstable nucleus.

- a) X-ray emission takes place
- b) a neutron is formed
- c) $\frac{n}{p}$ of daughter nucleus increases
- d) A neutron is consumed

A. X-ray emission takes place

B. a neutron is formed

C. $\frac{n}{p}$ of daughter nucleus increases

D. A neutron is consumed

Answer: B::C



Watch Video Solution

10. Select the correct statement(s):

- a) Mass number remains constant when positron emission takes place

- b) One neutron converts into proton in $\beta(^0_{-1}e)$ emission process
- c) Activity of a radioactive substance double when temp. increases from 300 K to 310 K
- d) Isodiaphers formed when one alpha particle emitted and isotopes formed when 2 beta particles emitted

A. Mass number remains constant when positron emission takes place

B. One neutron converts into proton in $\beta(^0_{-1}e)$ emission process

C. Activity of a radioactive substance double when temp. increases from 300 K to 310 K

D. Isodiaphers formed when one alpha particle emitted and isotopes formed when 2 beta particles emitted

Answer: A::B



Watch Video Solution

11. Match the following columns

Column-I

- (A) α -emission
- (B) β -emission
- (C) γ -emission
- (D) β^+ (Positron) emission

Column-II

- (P) Change in mass no.
- (Q) No change in atomic no. and mass no.
- (R) Atomic no. decreases
- (S) Atomic no. increases



[Watch Video Solution](#)

Level 3 Assertion Reason Type Questions

1. Each question contains STATEMENTS-1 (Assertion) and STATEMENT -2 (Reason).

Examine the statements carefully and mark the correct answer according to the instructions given below:

STATEMENTS-1: Acid catalysed hydrolysis of esters is pseudo first order reaction.

STATEMENT-2: Water is present in excess in given reaction.

a) If both the statements are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1

b) If both the statements are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1

c) If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE

d) If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

A. If both the statements are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1

B. If both the statements are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1

C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE

D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

Answer: A



[Watch Video Solution](#)

2. Each question contains STATEMENTS-1 (Assertion) and STATEMENT -2 (Reason).

Examine the statements carefully and mark the correct answer according to the instructions given below:

STATEMENT-1: For each 10°C rise of temperature the k is nearly double.

STATEMENT -2: Energy wise distribution of molecules in a gas sample is an exponential function of temperature so $e^{-E_a/RT}$ is doubled.

- A. If both the statements are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1
- B. If both the statement are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1
- C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

Answer: A



[Watch Video Solution](#)

3. Each question contains STATEMENTS-1 (Assertion) and STATEMENT -2 (Reason).

Examine the statements carefully and mark the correct answer according to the instructions given below:

STATEMENT-1: Nuclide $_{13}^{30}\text{Al}$ is less stable than $_{20}^{40}\text{Ca}$.

STATEMENT-2: Nuclide having odd number of protons and neutrons are generally unstable.

- A. If both the statements are TRUE and STATEMENT-2 is the correct explanation of STATEMENT-1
- B. If both the statement are TRUE but STATEMENT-2 is NOT the correct explanation of STATEMENT-1
- C. If STATEMENT-1 is TRUE and STATEMENT-2 is FALSE
- D. If STATEMENT-1 is FALSE and STATEMENT-2 is TRUE

Answer: B



Watch Video Solution

Level 3 Subjective Problems

1. $5A \rightarrow \text{Product}$

In above reaction, half-life period is directly proportional to initial concentration of reactant. The initial rate of reaction is $400 \text{ mol l}^{-1} \text{ min}^{-1}$

Then the order of reaction is



[Watch Video Solution](#)


2. For a reaction, $A \rightleftharpoons B$ equilibrium constant is 1.66 and

$$k_{\text{forward}} = 0.166 \text{ hr}^{-1}.$$

Calculate the time (in hour) when concentration of B is 80% of its equilibrium concentration. (Given : $\ln 25 = 3.20$)



[Watch Video Solution](#)



3. , At time $t=0$, initial mole of A is 1.

Overall half time of the reaction is 15 days. Then calculate the number of mole of C after 45 days if the ratio of $k_1 : k_2 : k_3$ is 4 : 2 : 1

[Watch Video Solution](#)