

MATHS

BOOKS - CENGAGE

BINOMIAL THEOREM

Examples

1. Prove that $\sum_{r=0}^n r^n C_r (-1)^r [i^r + i^{2r} + i^{3r} + i^{4r}] = 2^n + 2^{\frac{n}{2}+1} \cos(n\pi/4)$, where $i = \sqrt{-1}$.



Watch Video Solution

2. In the expansion of $\left(3\sqrt{\frac{a}{b}} + 3\sqrt{\frac{b}{a}}\right)^{21}$, the term containing same powers of a & b is



Watch Video Solution



3. Find a,b, and n in the expansion of $(a + b)^n$ if the first three terms of the expansion are 729, 7290 and 30375, respectively.



Watch Video Solution

4. If a and b are distinct integers, prove that a-b is a factor of $a^n - b^n$, whenever n is a positive integer.



Watch Video Solution

5. The number of terms in the expansion of $(a + b + c)^n$, where $n \in N$.



Watch Video Solution

6. Find the number of terms which are free from radical signs in the expansion of $(x^{1/5} + y^{1/10})^{55}$.



Watch Video Solution

7. Prove that $\sum_{k=1}^n \frac{1}{k(k+1)} = 1 - \frac{1}{n+1}$.



Watch Video Solution

8. Prove that $\sum_{r=0}^n nC_r \sin rx \cos(n-r)x = 2^{n-1} \sin(nx)$.



Watch Video Solution

9. Prove that $2 \leq \left(1 + \frac{1}{n}\right)^n < 3$ for all $n \in N$.



Watch Video Solution

10. The positive integer just greater than $(1 + .0001)^{10000}$ is



Watch Video Solution

11. Prove that $\sqrt{10} \left[(\sqrt{10} + 1)^{100} - (\sqrt{10} - 1)^{100} \right]$.



[View Text Solution](#)

12. If $9^7 - 7^9$ is divisible by 2^n , then find the greatest value of n , where $n \in N$.



[Watch Video Solution](#)

13. Find the degree of the polynomial

$$\frac{1}{\sqrt{4x+1}} \left\{ \left(\frac{1 + \sqrt{4x+1}}{2} \right)^7 - \left(\frac{1 - \sqrt{4x+1}}{2} \right)^7 \right\}$$



[Watch Video Solution](#)

14. If $T_0, T_1, T_2, \dots, T_n$ represent the terms in the expansion of $(x + a)^n$, then find the value of

$$(T_0 - T_2 + T_4 - \dots)^2 + (T_1 - T_3 + T_5 - \dots)^2 n \in N.$$



Watch Video Solution

15. If $n = 12m (m \in N)$, prove that

$$\hat{n}C_0 - \frac{\hat{n}C_2}{(2 + \sqrt{3})^2} + \frac{\hat{n}C_4}{(2 + \sqrt{3})^4} - \frac{\hat{n}C_6}{(2 + \sqrt{3})^6} + \dots = \left(\frac{2\sqrt{2}}{1 + \sqrt{3}} \right)^n.$$



Watch Video Solution

16. Prove that $\sum_{r=1}^k (-3)^{r-1} 3^n C_{2r-1} = 0$, where $k = 3n/2$ and n is an even integer.



View Text Solution

17. If the middle term in the expansion of $(x^2 + 1/x)^n$ is $924x^6$, then find the value of n .



Watch Video Solution

18. Show that the middle term in the expansion of $(1 + x)^{2n}$ is $\frac{1.3.5.....(2n - 1)}{n!} 2^n x^n$, where n is a positive integer.



Watch Video Solution

19. If the coefficient of the middle term in the expansion of $(1 + x)^{2n+2}$ is α and the coefficients of middle terms in the expansion of $(1 + x)^{2n+1}$ are β and γ then relate α , β and γ .



View Text Solution

20. Find the coefficient of a^3b^4c in the expansion of $(1 + a - b + c)^9$.



Watch Video Solution

21. Find the coefficient of $a^3b^4c^5$ in the expansion of $(bc + ca + ab)^6$.



Watch Video Solution

22. Find the coefficient of x^7 in the expansion of $(a + 3x - 2x^3)^{10}$.



Watch Video Solution

23. Find an approximation of $(0.99)^5$ using the first three terms of its expansion.



Watch Video Solution

24. Using the binomial theorem, evaluate $(102)^5$.



Watch Video Solution

25. Which is larger : $(99^{50} + 100^{50})$ or $(101)^{50}$.



View Text Solution

26. Find (i) the last digit, (ii) the last two digits, and (iii) the last three digits of 17^{256} .



Watch Video Solution

27. Using Binomial theorem, prove that $6^n - 5n$ always leaves remainder 1 when divided by 25 for all positive integer n .



Watch Video Solution

28. Using binomial theorem, prove that $2^{3n} - 7^n - 1$ is divisible by 49 , where $n \in N$.



Watch Video Solution

29. The remainder when 5^{99} is divided by 13 is



Watch Video Solution

30. Find the remainder when 27^{40} is divided by 12.



Watch Video Solution

31. Find the remainder when $1690^{2608} + 2608^{1690}$ is divided by 7.



Watch Video Solution

32. If $(2 + \sqrt{3})^n = I + f$, where I and n are positive integers and 0



Watch Video Solution

33. Statement 1: If p is a prime number ($p \neq 2$), then $[(2 + \sqrt{5})^p] - 2^{p+1}$ is always divisible by p (where $[.]$ denotes the greatest integer function). Statement 2: if n prime, then $\hat{n}C_1, {}^nC_2, {}^nC_2, {}^nC_{n-1}$ must be divisible by n .



Watch Video Solution

34. Find the coefficient of x^{20} in $\left(x^2 + 2 + \frac{1}{x^2}\right)^{-5} (1 + x^2)^{40}$.



Watch Video Solution

35. The coefficient of x^4 in the expansion of $\left(\frac{x}{2} - \frac{3}{x^2}\right)^{10}$ is



Watch Video Solution

36. Find the coefficient of x^{13} in the expansion of $(1 - x)^5 \times (1 + x + x^2 + x^3)^4$.



Watch Video Solution

37. Find the coefficient of x^{25} in expansion of expression

$$\sum_{r=0}^{50} \hat{(50)C_r(2x-3)^r(2-x)^{50-r}}.$$



Watch Video Solution

38. Find the coefficient of x^k in

$$1 + (1+x) + (1+x)^2 + \dots + (1+x)^n (0 \leq k \leq n).$$



Watch Video Solution

39. Find the coefficient of x^{50} in the expansion of

$$(1+x)^{101} \times (1-x+x^2)^{100}.$$



Watch Video Solution

40. Find the term independent of x in the expansion of

$$(2^x + 2^{-x} + \log_e e^{x+2}))^{30}.$$



Watch Video Solution

41. Find the coefficient of x^4 in the expansion of $(2 - x + 3x^2)^6$.



Watch Video Solution

42. Find the term independent of x in the expansion of $(1 + x + 2x^3)[(3x^2/2) - (1/3)]^9$



Watch Video Solution

43. Prove that in the expansion of $(1 + x)^n(1 + y)^n(1 + z)^n$, the sum of the coefficients of the terms of degree r is $s^{3n}C_r$.



Watch Video Solution

44. Find the coefficient of x^n in $\left(1 + \frac{x}{1!} + \frac{x^2}{2!} + \dots + \frac{x^n}{n!}\right)^2$.



Watch Video Solution

45. Find the sum $.^n C_1 + 2 \times .^n C_2 + 3 \times .^n C_3 + \dots + n \times .^n C_n$.



View Text Solution

46. Find the sum $\sum_{k=0}^{10} {}^{\wedge}(20)C_k$.



Watch Video Solution

47. Find the sum of the series ${}^{\wedge} 10C_0 + {}^{10} C_1 + {}^{10} C_2 + \dots + {}^{10} C_7$.



View Text Solution

48. If n is a positive integer, prove that

$$1 - 2n + \frac{2n(2n-1)}{2!} - \frac{2n(2n-1)(2n-2)}{3!} + + (-1)^{n-1} \frac{2n(2n-1)(2n-2)\dots(2n-n+1)}{(n-1)!}$$



Watch Video Solution

49. Prove that $\sum_{k=0}^n (-1)^k \cdot {}^{3n}C_k = (-1)^n \cdot {}^{3n-1}C_n$



Watch Video Solution

50. Find the sum of the coefficients of all the integral powers of x in the expansion of $(1 + 2\sqrt{x})^{40}$.



Watch Video Solution

51. Find the sum $\sum_{0 \leq i \leq n} \binom{n}{i} \cdot 2^i$



Watch Video Solution

52. Find the sum $\sum_{0 \leq i \leq n} \binom{n}{i} \cdot 2^i$



Watch Video Solution

53. Evaluate $\int(x^3 + 4x^2 + 5)dx$



Watch Video Solution

54. Find the following sums :

(i) $.^n C_0 - .^n C_2 + .^n C_4 - .^n C_6 + \dots$

(ii) $.^n C_1 - .^n C_3 + .^n C_5 - .^n C_7 + \dots$

(iii) $.^n C_0 + .^n C_4 + .^n C_8 + .^n C_{12} + \dots$

(iv) $.^n C_2 + .^n C_6 + .^n C_{10} + .^n C_{14} + \dots$

(v) $.^n C_1 + .^n C_5 + .^n C_9 + .^n C_{13} + \dots$

(vi) $.^n C_3 + .^n C_7 + .^n C_{11} + .^n C_{15} + \dots$



View Text Solution

55. Differentiate $y = \cos^3(2x)$.



Watch Video Solution

56. If the sum of the coefficient in the expansion of $(\alpha^2x^2 - 2\alpha x + 1)^{51}$ vanishes, then find the value of α

 Watch Video Solution

57. If $(1 + x - 2x^2)^{20} = a_0a_1x = a_2x^2 + a_3x^3 + \dots + a_{40}x^{40}$, then find the value of $a_1 + a_3 + a_5 + \dots + a_{39}$.

 Watch Video Solution

58. If $(1 + x + x^2)^n = a_0 + a_1x + a_2x^2 + \dots + a_{2n}x^{2n}$, find the value of $a_0 + a_6 + \dots, n \in N$.

 Watch Video Solution

59. Prove that $\frac{\sum_{\alpha+\beta+\gamma=10}^{10!}}{\alpha!\beta!\gamma!} = 3^{10}$.

 Watch Video Solution

60. If the binomial coefficients of three consecutive terms in the expansion of $(a + x)^n$ are in the ratio 1:7 :42 , then find n .



Watch Video Solution

61. In the coefficients of rth, $(r + 1)th$, and $(r + 2)th$ terms in the binomial expansion of $(1 + y)^m$ are in A.P., then prove that $m^2 - m(4r + 1) + 4r^2 - 2 = 0$.



Watch Video Solution

62. If $C_r = {}^n C_r$ then prove that $(C_0 + C_1)(C_1 + C_2) \dots \dots (C_{n-1} + C_n) = (C_1 C_2 \dots C_{n-1} C_n)(n + 1)^n / n!$



Watch Video Solution

63. If a_1, a_2, a_3, a_4 be the coefficient of four consecutive terms in the expansion of $(1 + x)^n$, then prove that:

$$\frac{a_1}{a_1 + a_2} + \frac{a_3}{a_3 + a_4} = \frac{2a_2}{a_2 + a_3}.$$



[Watch Video Solution](#)

64. Find the sum of $\sum_{r=1}^n \frac{r^n C_r}{n C_{r-1}}$.



[Watch Video Solution](#)

65. Find the numerically Greatest Term In the expansion of $(3 - 5x)^{15}$ when $x=1/5$



[Watch Video Solution](#)

66. Find the value of the second largest term in the expansion of $(4 + 5x)^{20}$ when $x = 1/3$



[View Text Solution](#)

67. Find the greatest coefficient in the expansion of $(1 + 2x/3)^{15}$.



[Watch Video Solution](#)

68. Given that the 4th term in the expansion of $[2 + (3/8x)]^{10}$ has the maximum numerical value. Then find the range of value of x .



[Watch Video Solution](#)

69. Prove that $.^n C_1 + 2.^n C_2 + 3.^n C_3 + \dots + n.^n C_n = n2^{n-1}$.



[Watch Video Solution](#)

70. Find the sum $1.^n C_0 + 3.^n C_1 + 5.^n C_2 + \dots + (2n+1).^n C_n$.



[Watch Video Solution](#)

71. If $(1 + x + x^2 + \dots + x^p)^n = a_0 + a_1x + a_2x^2 + \dots + a_{np}x^{np}$, then find the value of $a_1 + 2a_2 + 3a_3 + \dots + npa_{np}$.



Watch Video Solution

72. Evaluate $\int (1 + x)^2 dx$



Watch Video Solution

73. If $n > 2$, then prove that $C_1(a - 1) - C_2 \times (a - 2) + \dots + (-1)^{n-1}C_n(a - n) = a$, where $C_r = {}^n C_r$



Watch Video Solution

74. Find the sum $3^n C_0 - 8^n C_1 + 13^n C_2 - {}^n C_3 + \dots$



Watch Video Solution

75. If $x + y = 1$, prove that $\sum_{r=0}^n r \cdot {}^n C_r x^r y^{n-r} = nx$.



[Watch Video Solution](#)

76. Prove that $\frac{C_1}{2} + \frac{C_3}{3} + \frac{C_5}{6} + \dots = \frac{2^{n+1} - 1}{n + 1}$.



[Watch Video Solution](#)

77. Prove that $\frac{C_1}{2} + \frac{C_3}{3} + \frac{C_5}{6} + \dots = \frac{2^{n+1} - 1}{n + 1}$.



[Watch Video Solution](#)

78. Find the sum

$$2 \cdot {}^{10} C_0 + \frac{2^2}{2} \cdot {}^{10} C_1 + \frac{2^3}{3} \cdot {}^{10} C_2 + \frac{2^4}{4} \cdot {}^{10} C_3 + \dots + \frac{2^{11}}{11} \cdot {}^{10} C_{10}.$$



[View Text Solution](#)

79.

Prove

that

$$\frac{C_1}{1} - \frac{C_2}{2} + \frac{C_3}{3} - \frac{C_4}{4} + \dots + \frac{(-1)^{n-1}}{n} C_n = 1 + \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{n}.$$



Watch Video Solution

80. Prove that $\sum_{r=1}^n (-1)^{r-1} \left(1 + \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{r} \right)^n C_r = \frac{1}{n}.$



Watch Video Solution

81. Prove that $\frac{3!}{2(n+3)} = \sum_{r=0}^n (-1)^r \left(\frac{\hat{n}C_r}{\hat{(r+3)}C_r} \right)$



Watch Video Solution

82. There are two bags can each of which contains n balls. A man has to select an equal number of balls from both the bags. Prove that the number of ways in which a man can choose at least one ball from each bag is $\hat{2n}C_n - 1$.



Watch Video Solution

83. Find the sum $\sum_{i=0}^r {}^n C_r {}^{n_2}_{r-i} C_i$.



Watch Video Solution

84. Prove that $\sum_{r=0}^{2n} r \left({}^{2n} C_r \right)^2 = n^{4n} C_{2n}$.



Watch Video Solution

85. Using binomial theorem (without using the formula for ${}^n C_r$), prove that

$${}^n C_4 + {}^m C_2 - {}^m C_1 {}^n C_2 = {}^m C_4 - {}^{m+n} C_1 {}^m C_3 + {}^{m+n} C_2 {}^m C_2 - {}^{m+n} C_3 {}^m C_1 + {}^{m+n} C_0$$



Watch Video Solution

86. Evaluate $\int (x^2 + 8x + 3) dx$



Watch Video Solution

87.

Prove

that

$${}^{\wedge} 100 C_2^{100} C_2 + {}^{100} C_2^{100} C_4 + {}^{100} C_4^{100} C_6 + \dots + {}^{100} C_{98}^{100} C_{100} = \frac{1}{2} [{}^{\wedge} (200) C_{98}^{100}]$$



View Text Solution

88. Let $m, n \in \mathbb{N}$ and $C_r = {}^n C_r$, for $0 \leq r \leq n$

Statement-1:

$$\begin{aligned} & \frac{1}{m!} C_0 + \frac{n}{(m+1)!} C_1 + \frac{n(n-1)}{(m+2)!} C_2 + \dots + \frac{n(n-1)(n-2)\dots 2.1}{(m+n)!} C_n \\ &= \frac{(m+n+1)(m+n+2)\dots(m+2n)}{(m+n)!} \end{aligned}$$

Statement-2: For $r \leq 0$

$${}^m C_r {}^n C_0 + {}^m C_{r-1} {}^n C_1 + {}^m C_{r-2} {}^n C_2 + \dots + {}^m C_0 {}^n C_r = {}^{m+n} C_r$$



Watch Video Solution

89. Statement1: if $n \in N$ and n is not a multiple of 3 and

$$(1 + x + x^2) = \sum_{r=0}^{2n} a_r x^r, \text{ then the value of } \sum_{r=0}^n (-1)^r a_r n C_r \text{ is zero}$$

Statement 2: The coefficient of x^n in the expansion of $(1 - x^3)$ is zero, if

$n = 3k + 1$ or $n = 3k + 2$.



[Watch Video Solution](#)

90. Find the sum $\sum_{r=0}^n (n+r) C_r$.



[Watch Video Solution](#)

91.

Prove

that

$${}^n C_0^{2n} C_n - {}^n C_1^{2n-1} C_n + {}^n C_2 \times {}^{2n-2} C_n + \dots + (-1)^n {}^n C_n^n C_n = 1.$$



[Watch Video Solution](#)

92. Prove that ${}^m C_1^n C_m - {}^m C_2^{2n} C_m + {}^m C_3^{3n} C_m \equiv (-1)^{m-1} n^m$.



Watch Video Solution

93. If $(18x^2 + 12x + 4)^n = a_0 + a_{1x} + a_2x^2 + \dots + a_{2n}x^{2n}$, prove that $a_r = 2^n 3^r \left({}^n C_r + {}^n C_1^{2n-2} C_r + {}^n C_2^{2n-4} C_r + \dots \right)$.



Watch Video Solution

94. Find the value of x , for which $1/(\sqrt{5+4x})$ can be expanded as infinite series.



Watch Video Solution

95. Prove that $\lim_{x \rightarrow 0} \frac{(1+x)^n - 1}{x} = n$.



Watch Video Solution

96. Find the fourth term in the expansion of $(1 - 2x)^{3/2}$.



Watch Video Solution

97. Prove that the coefficient of x^r in the expansion of $(1 - 2x)^{1/2}$ is $(2r)! / [2^r(r!)^2]$.



View Text Solution

98. Find the sum $1 - \frac{1}{8} + \frac{1}{8} \times \frac{3}{16} - \frac{1 \times 3 \times 5}{8 \times 16 \times 24} + \dots$



Watch Video Solution

99. Find the coefficient of x^n in the expansion of $(1 - 9x + 20x^2)^{-1}$.



View Text Solution

100. Assuming x to be so small that x^2 and higher power of x can be neglected, prove that



Watch Video Solution

101. If x is very large as compare to y , then prove that

$$\sqrt{\frac{x}{x+y}} \sqrt{\frac{x}{x-y}} = 1 + \frac{y^2}{2x^2}.$$



Watch Video Solution

102. Prove that the coefficient of x^n in the expansion of

$$\frac{1}{(1-x)(1-2x)(1-3x)} \text{ is } \frac{1}{2}(3^{n+2} - 2^{n+3} + 1).$$



Watch Video Solution

103. Prove that

$${}^n C_0 - {}^n C_1 + {}^n C_2 - {}^n C_3 + \dots + (-1)^r + {}^n C_r + \dots = (-1)^r \times {}^{n-1}$$

.



Watch Video Solution

104. Find the value of

$$\cdot^{20} C_0 \times \cdot^{13} C_{10} - \cdot^{20} C_1 \times \cdot^{12} C_9 + \cdot^{20} C_2 \times \cdot^{11} C_8 - \dots + \cdot^{20} C_{10}.$$



Watch Video Solution

105. If $U_n = (\sqrt{3} + 1)^{2n} + (\sqrt{3} - 1)^{2n}$, then prove that $U_{n+1} = 8U_n - 4U_{n-1}$.



Watch Video Solution

106. Evaluate $\int(x^{10} - 6x^5 + 2x^3)dx$



Watch Video Solution

107. Find the coefficients of x^{50} in the expression $(1+x)^{1000} + 2x(1+x)^{999} + 3x^2(1+x)^{998} + \dots + 1001x^{1000}$.



Watch Video Solution

108.

Given,

$$s_n = 1 + q + q^2 + \dots + q^n, S_n = 1 + \frac{q+1}{2} + \left(\frac{q+1}{2}\right)^2 + \dots + \left(\frac{q+1}{2}\right)^n,$$

prove that $\hat{n+1}C_1 +^{n+1}C_2 s_1 +^{n+1}C_3 s_2 + \dots +^{n+1}C_{n+1} s_n 2^n S_n$.



Watch Video Solution

109. Find the derivative of $y = 3 \sin^3(2x^4 + 1)$.



Watch Video Solution

110. Evaluate $\int \frac{x+1}{x^2+2x+1} dx$



Watch Video Solution

111. Evaluate $\int \frac{x^4+x-4}{x^2+2} dx$



Watch Video Solution

112. Solve the differential equation $\frac{dx}{dy} = xy + 3x + y + 3$



[Watch Video Solution](#)

113. Prove that $\frac{.^n C_1 \sin 2x + .^n C_2 \sin 4x + .^n C_3 \sin 6x + \dots}{1 + .^n C_1 \cos 2x + .^n C_2 \cos 4x + .^n C_3 \cos 6x + \dots} = \frac{(a^x + b^x)^2}{a^x b^x}$



[Watch Video Solution](#)

114. Evaluate $\int \frac{(a^x + b^x)^2}{a^x b^x} dx$



[Watch Video Solution](#)

Exercise 8.1

1. The first three terms in the expansion of $(1 + ax)^n (n \neq 0)$ are 1, $6x$ and $16x^2$. Then find the value of a and n .



[Watch Video Solution](#)

2. about to only mathematics

 Watch Video Solution

3. The coefficients of three consecutive terms in the expansion of $(1 + a)^n$ are in the ratio 1: 7: 42 Find n.

 Watch Video Solution

4. If the number of terms in the expansion of $(x + y + z)^n$ are 36, then find the value of n.

 Watch Video Solution

5. Find the value of

$$\frac{1}{81^n} - \frac{10}{(81^n)^{2n}} C_1 + \frac{10^2}{(81^n)^{2n}} C_2 - \frac{10^3}{(81^n)^{2n}} C_3 + \dots + \frac{10^{2n}}{81^n}$$



Watch Video Solution

6.

Prove

that

$$\sum_{r=0}^n (-1)^r \cdot nC_r \left[\frac{1}{2^r} + \frac{3}{2^{2r}} + \frac{7}{2^{3r}} + \frac{15}{2^{4r}} + \text{up to } m \text{ terms} \right] = \frac{2^{mn} - 1}{2^{mn}(2^n - 1)}$$



Watch Video Solution

7. In $\left(33 + \frac{1}{33}\right)^n$ if the ratio of 7th term from the beginning to the 7th term from the end is $1/6$, then find the value of n .



Watch Video Solution

8. If the coefficients of $(r-5)^t h$ and $(2r-1)^t h$ terms in the expansion of $(1+x)^{34}$ are equal, find r .



Watch Video Solution

9. Find the number of irrational terms in the expansion of

$$\left(5^{1/6} + 2^{1/8}\right)^{100}.$$



Watch Video Solution

10. Represent $\cos 6\theta$ in terms of $\cos \theta$.



Watch Video Solution

11. Find the number of nonzero terms in the expansion of

$$(1 + 3\sqrt{2}x)^9 + (1 - 3\sqrt{2}x)^9.$$



Watch Video Solution

12. Find the value of $(\sqrt{2} + 1)^6 + (\sqrt{2} - 1)^6$.



Watch Video Solution

13.

If

$$\frac{1}{\sqrt{4x+1}} \left\{ \left(\frac{1 + \sqrt{4x+1}}{2} \right)^n - \left(\frac{1 - \sqrt{4x+1}}{2} \right)^n \right\} = a_0 + a_1 x$$

then find the possible value of n .



Watch Video Solution

14. Let $R = (5\sqrt{5} + 11)^{2n+1}$ and $f = R - [R]$ where $[]$ denotes the greatest integer function, prove that $Rf = 4^{2n+1}$



Watch Video Solution

15. If the middle term in the binomial expansion of $\left(\frac{1}{x} + x \sin x\right)^{10}$ is equal to $\frac{63}{8}$, find the value of x .



Watch Video Solution

16. Find the middle term in the expansion of $\left(x^2 + \frac{1}{x^2} + 2\right)^n$.



Watch Video Solution

17. If the number of terms in the expansion $(1 + 2x - 3y + 4z)^n$ is 286, then find the coefficient of term containing xyz .



Watch Video Solution

Exercise 8.2

1. Let n be an odd natural number greater than 1. Then , find the number of zeros at the end of the sum $99^n + 1$.



Watch Video Solution

2. By mathematical induction prove that $2^{3n}-1$ is divisible by 7.



Watch Video Solution

3. Find the last three digits of the number 27^{27} .



Watch Video Solution

4. If 10^m divides the number $101^{100} - 1$ then, find the greatest value of m .



Watch Video Solution

5. Show that $9^n + 1 - 8n - 9$ is divisible by 64, whenever n is a positive integer.



Watch Video Solution

6. Show that $2^{4n+4} - 15n - 16$, where $n \in N$ is divisible by 225.



Watch Video Solution

7. Find the remainder when 27^{40} is divided by 12.



Watch Video Solution

8. Find the value of $\{3^{2003} / 28\}$, where $\{\cdot\}$ denotes the fractional part.



Watch Video Solution

9. Statement 1: Remainder when 3456^{2222} is divided by 7 is 4. Statement 2: Remainder when 5^{2222} is divided by 7 is 4
option1: BOTH the statement are TRUE and STATEMENT 2 is the correct explanation
option2: BOTH the statement are TRUE and STATEMENT 2 is NOT the correct explanation
option3: STATEMENT 1 is TRUE and STATEMENT 2 is FALSE
option4: STATEMENT 1 is FALSE and STATEMENT 2 is TRUE



Watch Video Solution

10. Show that the integer next above $(\sqrt{3} + 1)^{2m}$ contains 2^{m+1} as a factor.



Watch Video Solution

Exercise 8.3

1. If x^4 occurs in the rth term in the expansion of $\left(x^4 + \frac{1}{x^3}\right)^{15}$, then find the value of r .



Watch Video Solution

2. If x^p occurs in the expansion of $(x^2 + 1/x)^{2n}$, prove that its coefficient is $\frac{(2n)!}{\left[\frac{1}{3}(4n - p)\right]! \left[\frac{1}{3}(2n + p)\right]!}$.



Watch Video Solution

3. Find the coefficient of t^8 in the expansion of $(1 + 2t^2 - t^3)^9$.



Watch Video Solution

4. If $|x| < 1$, then find the coefficient of x^n in the expansion of $(1 + 2x + 3x^2 + 4x^3 + \dots)^{1/2}$.



Watch Video Solution

5. The coefficient of the term independent of x in the expansion of

$$\left(\frac{x+1}{x^{2/3} - x^{1/3} + 1} - \frac{x-1}{x - x^{1/2}} \right)^{10}$$



Watch Video Solution

6. In the expansion of $(1 + 3x + 2x^2)^6$, the coefficient of x^{11} is a. 144 b. 288 c. 216 d. 576



Watch Video Solution

7. Find the coefficient of x^{12} in expansion of $(1 - x^2 + x^4)^3(1 - x)^7$.



Watch Video Solution

Exercise 8.4

1. In the expansion of $(1 + x)^{50}$ the sum of the coefficients of the odd power of x is



Watch Video Solution

2. Find the following sum: $\frac{1}{n!} + \frac{1}{2!(n-2)!} + \frac{1}{4!(n-4)!} + \dots$



Watch Video Solution

3. Find the sum of the coefficients in the expansion of $(1 + 2x + 3x^2 + nx^n)^2$.



Watch Video Solution

4. Find the sum $\sum_{j=0}^n \left({}^{4n+1}C_j + {}^{4n+1}C_{2n-j} \right)$.



Watch Video Solution

5. Find the remainder when $\left(\sum_{r=1}^5 {}^{20}C_{2r-1} \right)^6$ is divided by 11.



Watch Video Solution

6. Prove that

$$.{}^n C_0 + 5 \times .{}^n C_1 + 9 \times .{}^n C_2 + \dots + (4n+1) \times .{}^n C_n = (2n+1)2^n.$$



Watch Video Solution

7. Prove that $nC_0 + {}^n C_3 + {}^n C_6 + \dots = \frac{1}{3} \left(2^n + 2 \frac{\cos(n\pi)}{3} \right)$.



Watch Video Solution



Watch Video Solution

8. Find the value of ${}^{\wedge} 4nC_0 + {}^{4n} C_4 + {}^{4n} C_8 + \dots + {}^{4n} C_{4n}$.



Watch Video Solution

9. Prove that $\sum_{r=0}^s \sum_{\substack{s=1 \\ r \leq s}}^n {}^n C_s {}^s C_r = 3^n - 1$.



Watch Video Solution

10. Find the sum of coefficients in $(1 + x - 3x^2)^{4163}$.



Watch Video Solution

11. Find the sum of coefficients in the expansion of $(x - 2y + 3z)^n$ is 128, then find the greatest coefficients in the expansion of $(1 + x)^n$.



Watch Video Solution

12. Find the sum of the coefficients in the expansion of $(1 + 2x + 3x^2 + nx^n)^2$.



Watch Video Solution

13. If $(1 + x - 2x^2)^6 = 1 + a_1x + a_2x^2 + a_3x^3 + \dots + a_{12}x^{12}$, then the value of $a_2 + a_4 + a_6 + \dots + a_{12}$ will be



Watch Video Solution

Exercise 8.5

1. In the expansion of $(1 + x)^n$, 7th and 8th terms are equal. Find the value of $(7/x + 6)^2$.



Watch Video Solution

2. Find the sum $\sum_{r=1}^n r^2 \frac{\hat{n}C_r}{\hat{n}C_{r-1}}$.



[Watch Video Solution](#)

3. Statement 1: Three consecutive binomial coefficients are always in A.P.

Statement 2: Three consecutive binomial coefficients are not in H.P. or G.P.



[Watch Video Solution](#)

4. If the 3rd, 4th, 5th and 6th term in the expansion of $(x + \alpha)^n$ be,

respectively, a, b, c and d , prove that $\frac{b^2 - ac}{c^2 - bd} = \frac{5a}{3c}$.



[Watch Video Solution](#)

5. Find the largest term in the expansion of $(3 + 2x)^{50}$, where $x = 1/5$.



[Watch Video Solution](#)

6. If $x = 1/3$, find the greatest term in the expansion of $(1 + 4x)^8$.



Watch Video Solution

7. If n is an even positive integer, then find the value of x if the greatest term in the expansion of $1 + x^n$ may have the greatest coefficient also.



Watch Video Solution

8. If in the expansion of $2x + 5^{10}$, the numerically greatest term is equal to the middle term, then find the values of x .



Watch Video Solution

Exercise 8.6

1. Find the value of
 $(.^{10} C_{10}) + (.^{10} C_0 + .^{10} C_1) + (.^{10} C_0 + .^{10} C_1 + .^{10} C_2) + \dots + (.^{10} C_0 + .^{10} C_1 + .^{10} C_2 + \dots + .^{10} C_{10})$



Watch Video Solution

2. Prove that

$$\frac{1^2}{3} \cdot {}^n C_1 + \frac{1^2 + 2^2}{7} \cdot {}^n C_2 + \frac{1^2 + 2^2 + 3^2}{7} \cdot {}^n C_3 + \dots + \frac{1^2 + 2^3 + \dots + n^2}{2n+1} \cdot {}^n C_n = \frac{n(n+3)}{6} 2^{n-2}.$$



Watch Video Solution

3. If $p + q = 1$, then show that $\sum_{r=0}^n r^2 \wedge {}^n C_r p^r q^{n-r} = npq + n^2 p^2$.



Watch Video Solution

4. Prove that $1 - {}^n C_1 \frac{1+x}{1+nx} + {}^n C_2 \frac{1+2x}{(1+nx)^2} - {}^n C_3 \frac{1+3x}{(1+nx)^3} + \dots \dots (n+1) terms = 0$.



Watch Video Solution

5. Prove that

$$\frac{\hat{n}C_0}{1} + \frac{\hat{n}C_2}{3} + \frac{\hat{n}C_4}{5} + \frac{\hat{n}C_6}{7} + \dots + = \frac{2^n}{n+1}.$$



Watch Video Solution

6. If $(1+x)^{15} = C_0 + C_1x + C_2x^2 + \dots + C_{15}x^{15}$, then find the sum of $C_1 + 2C_3 + 3C_4 + \dots + 14C_{15}$.



Watch Video Solution

7. Find the coefficient of x^n in the polynomial $(x + {}^n C_0)(x + 3^n C_1) \times (x + 5^n C_2) [x + (2n+1)^n C_n]$.



Watch Video Solution

8. Find the value of $\hat{^20}C_0 - \frac{\hat{^20}C_1}{2} + \frac{\hat{^20}C_2}{3} - \frac{\hat{^20}C_3}{4} + \dots$.



Watch Video Solution

9. Prove that ${}^{10}C_2 + 2 \times {}^{10}C_3 + {}^{10}C_4 = {}^{12}C_4$



Watch Video Solution

10. If $\frac{dy}{dx} = \frac{1}{x}$ then $y = \dots$



Watch Video Solution

Exercise 8.7

1. Prove that $\sum_{r=0}^{2n} r \left({}^{2n}C_r \right)^2 = n^{4n} C_{2n}$.



Watch Video Solution

2.

Prove

that

$$(2nC_0)^2 + (2nC_1)^2 + (2nC_2)^2 - \dots + (2nC_{2n})^2 = (-1)^n 2nC_n.$$



Watch Video Solution

3. Find the sum of the series

$$\cdot^{84} C_4 + 6 \times \cdot^{84} C_5 + 15 \times \cdot^{84} C_6 + 20 \times \cdot^{84} C_7 + 15 \times \cdot^{84} C_8 + 6 \times \cdot^{84} C_9 + \dots$$



Watch Video Solution

4. Find an anti derivative (or integral) of the given function by the method of inspection. $(ax + b)$.



Watch Video Solution

5.

Prove

that

$$C_0 - 2^2 C_1 + 3^2 C_2 - 4^2 C_3 + \dots + (-1)^n (n+1)^2 \times C_n = 0 \text{ where } C_r = {}^n C_r$$



Watch Video Solution

6. Find the value of $\sum_{1 \leq i}^n$



Watch Video Solution

7. Prove that $n C_0^n C_0 - {}^{n+1} C_1^n C_1 + {}^{n+2} C_2^n C_2 \equiv (-1)^n$.



Watch Video Solution

8. Prove that $n C_0^{2n} C_n - {}^n C_1^{2n-2} C_n + {}^n C_2^{2n-4} C_n \equiv 2^n$.



Watch Video Solution

9. Find the value of $\sum_{p=1}^n \left(\sum_{m=p}^n {}^n C_m {}^m C_p \right)$. And hence, find the value of $(\lim)_{n \rightarrow \infty} \frac{1}{3^n} \sum_{p=1}^n \left(\sum_{m=p}^n {}^n C_m {}^m C_p \right)$.



Watch Video Solution

Exercise 8.8

1. If the third term in the expansion of $(1 + x)^m$ is $-\frac{1}{8}x^2$, then find the value of m .



Watch Video Solution

2. Find the cube root of 217, correct to two decimal places.



Watch Video Solution

3. Show that $\sqrt{3} = 1 + (1/3) + (1/3)(3/6) + (1/3)((3/6)(5/9)) + \dots$



Watch Video Solution

4. Find the coefficient of x^2 in $\left(\frac{a}{a+x}\right)^{1/2} + \left(\frac{a}{a-x}\right)^{1/2}$



Watch Video Solution

5. If $|x| < 1$, then find the coefficient of x^n in the expansion of $(1 + x + x^2 + x^3 + \dots)^2$.



Watch Video Solution

6. If $|x| > 1$, then expand $(1 + x)^{-2}$.



Watch Video Solution

7. If $|x| < 1$, then find the coefficient of x^n in the expansion of $(1 + 2x + 3x^2 + 4x^3 + \dots)^{1/2}$.



Watch Video Solution

8. If $(r + 1)th$ term is the first negative term in the expansion of $(1 + x)^{7/2}$, then find the value of r .



Watch Video Solution

9. Prove that the coefficient of x^n in the expansion of $\frac{1}{(1 - x)(1 - 2x)(1 - 3x)}$ is $\frac{1}{2}(3^{n+2} - 2^{n+3} + 1)$.



Watch Video Solution

10. Prove that

$$(r + 1) \times {}^n C_0 - r \times {}^n C_1 + (r - 1) \times {}^n C_2 - (r - 2) \cdot {}^n C_3 + \dots + (-1)^r \cdot {}^n C_r = 0$$

.



View Text Solution

Exercise (Single)

1. If the coefficients of 5th, 6th , and 7th terms in the expansion of $(1 + x)^n$ are in A.P., then $n =$ a. 7 only b. 14 only c. 7 or 14 d. none of these

A. 7 only

B. 14 only

C. 7 or 14

D. none of these

Answer: C



Watch Video Solution

2. The coefficient of the middle term in the binomial expansion in powers of x of $(1 + \alpha x)^4$ and of $(1 - \alpha x)^6$ is the same if α equals

A. $-\frac{5}{3}$

B. $\frac{10}{3}$

C. $-\frac{3}{10}$

D. $\frac{3}{5}$

Answer: C



Watch Video Solution

3. If $(1 + x)^5 = a_0 + a_1x + a_2x^2 + a_3x^3 + a_4x^4 + a_5x^5$, then the value of $(a_0 - a_2 + a_4)^2 + (a_1 - a_3 + a_5)^2$ is equal to
a. 243 b. 32 c. 1 d. 2^{10}

A. 243

B. 32

C. 1

D. 2^{10}

Answer: B



Watch Video Solution

4.

The

expression

$$\left(\sqrt{2x^2 + 1} + \sqrt{2x^2 - 1} \right)^6 \left(\frac{2}{\left(\sqrt{2x^2 + 1} + \sqrt{2x^2 - 1} \right)^\square} \right)^6 \quad \text{is}$$

polynomial of degree 6 b. 8 c. 10 d. 12

A. 6

B. 8

C. 10

D. 12

Answer: A



Watch Video Solution

5. If the 6th term in the expansion of $\left(\frac{1}{x^{\frac{8}{3}}} + x^2 (\log)_{10} x \right)^8$ is 5600, then x equals

A. 1

B. $\log_e 10$

C. 10

D. x does not exist

Answer: C



Watch Video Solution

6. If in the expansion of $(a - 2b)^n$, the sum of 5^{th} and 6^{th} terms is 0, then
the values of $a/b =$

A. $\frac{n - 4}{5}$

B. $\frac{2(n - 4)}{5}$

C. $\frac{5}{n - 4}$

D. $\frac{5}{2(n - 4)}$

Answer: B



Watch Video Solution

7. The number of real negative terms in the binomial expansion of $(1 + ix)^{4n-2}$, $n \in N$, $x > 0$ is

A. n

B. n+1

C. n-1

D. 2n

Answer: A



Watch Video Solution

8. The sum of rational term in $(\sqrt{2} + \sqrt[3]{3} + \sqrt[6]{5})^{10}$ is equal to 12632 b.

1260 c. 126 d. none of these

A. 12632

B. 1260

C. 126

D. 11792

Answer: D



Watch Video Solution

9. The value of x for which the sixth term in the expansion of

$$\left[2^{\log} - 2^{\sqrt{9^{(x-1)} + 7}} + \frac{1}{2^{\frac{1}{5}} (\log)_2 (3^{(x-1) + 1})} \right]^7 \text{ is } 84 \text{ is}$$

A. 4

B. 1 or 2

C. 0 or 1

D. 3

Answer: B



Watch Video Solution

10. The number of distinct terms in the expansion of $\left(x + \frac{1}{x} + \frac{1}{x^2}\right)^{15}$ is/are (with respect to different power of x)
a. 255 b. 61 c. 127 d. none of these

A. 255

B. 61

C. 127

D. none of these

Answer: B



Watch Video Solution

11. The value of $2 \times {}^n C_1 + 2^3 \times {}^n C_3 + 2^5 + \dots$ is

A. $\frac{3^n + (-1)^n}{2}$

B. $\frac{3^n - (-1)^n}{2}$

C. $\frac{3^n + 1}{2}$

D. $\frac{3^n - 1}{2}$

Answer: B



Watch Video Solution

12. If $(4x^2 + 1)^n = \sum_{r=0}^n a_r (1 + x^2)^{n-r} x^{2r}$, then the value of

A. 3^n

B. 4^n

C. 5^n

D. 6^n

Answer: B



Watch Video Solution

13. The fractional part of $2^{4n}/15$ is ($n \in N$)
a. $\frac{1}{15}$ b. $\frac{2}{15}$ c. $\frac{4}{15}$ d. none of these

A. $\frac{1}{15}$

B. $\frac{2}{15}$

C. $\frac{4}{15}$

D. none of these

Answer: A



Watch Video Solution

14. If $p = (8 + 3\sqrt{7})^n$ and $f = p - [p]$, where $[.]$ denotes the greatest integer function, then the value of $p(1 - f)$ is equal to
a. 1 b. 2 c. 2^n d. 2^{2n}

A. 1

B. 2

C. 2^n

D. 2^{2n}

Answer: A



Watch Video Solution

15. The remainder when the number $3^{256} - 3^{12}$ is divided by 8 is

A. 0

B. 3

C. 4

D. 7

Answer: A



Watch Video Solution

16. The smallest integer larger than $(\sqrt{3} + \sqrt{2})^6$ is

A. 969

B. 970

C. 971

D. 972

Answer: B



Watch Video Solution

17. Find the coefficient of x^5 in the expansion of $(1 + x^2)^5(1 + x)^4$

A. 12

B. 5

C. 4

D. 56

Answer: C



Watch Video Solution

18. Coefficient of x^2 in the expansion of $(x^3 + 2x^2 + x + 4)^{15}$ is

- A. Prime
- B. Composite
- C. 0
- D. Perfect square

Answer: D



Watch Video Solution

19. If the coefficients of r th and $(r + 1)$ th terms in the expansion of $(3 + 7x)^{29}$ are equal, then r equals a. 15 b. 21 c. 14 d. none of these

- A. 15
- B. 21
- C. 14

D. none of these

Answer: B



Watch Video Solution

20. In the expansion of $\left(x^3 - \frac{1}{x^2}\right)^n$, $n \in N$, if the sum of the coefficients of x^5 and x^{10} , then n is a. 25 b. 20 c. 15 d. none of these

A. 25

B. 20

C. 15

D. None of these

Answer: C



Watch Video Solution

21. If $(1 + 2x + x^2)^n = \sum_{r=0}^{2n} a_r x^r$, then $a =$ $({}^n C_2)^2$ b.

${}^n C_r$ c. ${}^n C_{r+1}$ d. ${}^n C_r$

A. $({}^n C_r)^2$

B. ${}^n C_r \cdot {}^n C_{r+1}$

C. ${}^{2n} C_r$

D. ${}^{2n} C_{r+1}$

Answer: C



Watch Video Solution

22. If the term independent of x in the $\left(\sqrt{x} - \frac{k}{x^2}\right)^{10}$ is 405, then k equals 2, -2 b. 3, -3 c. 4, -4 d. 1, -1

A. 2, -2

B. 3, -3

C. 4, - 4

D. 1, - 1

Answer: B



Watch Video Solution

23. The coefficient of x^{53} in the expansion

$\sum_{m=0}^{100} {}^{\wedge}(100)C_m(x-3)^{100-m}2^m$ is ^ 100C₄₇ b. ^ 100C₅₃ c. -¹⁰⁰C₅₃ d.

none of these

A. .¹⁰⁰C₄₇

B. .¹⁰⁰C₅₃

C. -.¹⁰⁰C₅₃

D. -.⁵³C₁₀₀

Answer: C



Watch Video Solution

24. If the coefficient of x^7 in $\left[ax^2 + \left(\frac{1}{bx}\right)\right]^{11}$ equals the coefficient of x^{-7} in $\left[ax - \left(\frac{1}{bx^2}\right)\right]^{11}$, then a and b satisfy the relation

A. $a + b = 1$

B. $a - b = 1$

C. $ab = 1$

D. $\frac{a}{b} = 1$

Answer: C



Watch Video Solution

25. Find a positive value of m for which the coefficient of x^2 in the expansion $(1 + x)^m$ is 6.

A. -83

B. 0

C. $.^{30} C_{10}$

D. none of these

Answer: D



Watch Video Solution

26. The term independent of a in the expansion of

$$\left(1 + \sqrt{a} + \frac{1}{\sqrt{a} - 1}\right)^{-30}$$

is

a. $.^{30} C_{20}$ b. 0 c. $.^{\wedge} 30 C_{10}$ d. none of these

A. $.^{30} C_{20}$

B. 0

C. $.^{30} C_{10}$

D. none of these

Answer: B



Watch Video Solution

27. The coefficient of x^{10} in the expansion of $(1 + x^2 - x^3)^8$ is

a. 476 b. 496

c. 506 d. 528

A. 476

B. 496

C. 506

D. 528

Answer: A



Watch Video Solution

28. The coefficient of x^n in $(1 + x)^{101}(1 - x + x^2)^{100}$ is nonzero, then n

cannot be of the form 3r + 1 b. 3r c. 3r + 2 d. none of these

A. 3r + 1

B. 3r

C. 3r + 2

D. none of these

Answer: C



Watch Video Solution

29. The coefficient of x^{28} in the expansion of $(1 + x^3 - x^6)^{30}$ is 1 b. 0 c.

30C₆ d. ^ 30C₃

A. 1

B. 0

C. .³⁰ C₆

D. .³⁰ C₃

Answer: B



Watch Video Solution

30. The coefficient of $a^8b^4c^9d^9$ in $(abc + abd + acdd + bcd)^{10}$ is 10! b.

$$\frac{10!}{8!4!9!9!} \text{ c. } 2520 \text{ d. none of these}$$

A. 10!

B. $\frac{10!}{8!4!9!9!}$

C. 2520

D. none of these

Answer: C



Watch Video Solution

31. In the expansion of $\left(1 + x + \frac{7}{x}\right)^{11}$ find the term not containing x.

A. $\sum_{r=0}^5 .^{11}C_r .^{11-r} C_{2r} 7^r$

B. $\sum_{r=0}^5 .^{11}C_r .^{11-r} C_{11-2r} 7^r$

C. $\sum_{r=0}^5 .^{11}C_r .^{11-r} C_{11-2r} 7^r$

D. none of these

Answer: C



Watch Video Solution

32. The coefficient of x^7 in the expansion of $(1 - x - x^3 + x^4)^8$ is equal to

A. - 648

B. 792

C. - 792

D. 648

Answer: C



Watch Video Solution

33. Sum of the coefficients of terms of degree 13 in the expansion of

$$(1+x)^{11}(1+y^2-z)^{10} \text{ is}$$

A. $.^{10} C_3$

B. $.^{10} C_4$

C. $.^{11} C_3$

D. $.^{11} C_4$

Answer: B



Watch Video Solution

34. The coefficient of x^2y^3 in the expansion of $(1-x+y)^{20}$ is $\frac{20!}{213!}$ b.

- $\frac{20!}{213!}$ c. $\frac{20!}{15!2!3!}$ d. none of these

A. $\frac{20!}{2!3!}$

B. - $\frac{20!}{2!3!}$

C. $\frac{20!}{5!2!3!}$

$$D. \frac{20!}{15!2!3!}$$

Answer: D



Watch Video Solution

35. If coefficient of $a^2b^3c^4 \in (a + b + c)^m$ (where $m \in N$) is L ($L \neq 0$) , then in same expansion coefficient of $a^4b^4c^1$ will be L b. $\frac{L}{3}$ c. $\frac{mL}{4}$ d. $\frac{L}{2}$

A. L

B. $\frac{L}{3}$

C. $\frac{mL}{4}$

D. $\frac{L}{2}$

Answer: D



Watch Video Solution

36. The coefficient of x^r [$0 \leq r \leq (n - 1)$] in the expansion of $(x + 3)^{n-1} + (x + 3)^{n-2}(x + 2) + (x + 3)^{n-3}(x + 2)^2 + \dots + (x + 2)^{n-1}$ is

a. ${}^n C_r (3^r - 2^n)$ b. ${}^n C_r (3^{n-r} - 2^{n-r})$ c. ${}^n C_r (3^r + 2^{n-r})$ d. none of these

A. ${}^n C_r (3^r - 2^n)$

B. ${}^n C_r (3^{n-r} - 2^{n-r})$

C. ${}^n C_r (3^r + 2^{n-r})$

D. none of these

Answer: B



Watch Video Solution

37. If $(1 + 2x + 3x^2)^{10} = a_0 + a_1x + a_2x^2 + \dots + a_{20}x^{20}$, then a_1 equals

10 b. 20 c. 210 d. none of these

A. 10

B. 20

C. 210

D. none of these

Answer: B



Watch Video Solution

38. If $f(x) = 1 - x + x^2 - x^3 + \dots + x^{15} + x^{16} - x^{17}$, then the coefficient of $x^2 \in f(x - 1)$ is 826 b. 816 c. 822 d. none of these

A. 826

B. 816

C. 822

D. none of these

Answer: B



Watch Video Solution

39. Let $f(x) = a_0 + a_1x + a_2x^2 + \dots + a_nx^n$ and

$$\frac{f(x)}{1-x} = b_0 + b_1x + b_2x^2 + \dots + b_nx^n, \text{ then } b_n + b_{n-1} = a_n \quad \text{b.}$$

$b_n - b_{n-1} = a_n$ c. $b_n / b_{n-1} = a_n$ d. none of these

A. $b_n + b_{n-1} = a_n$

B. $b_n - b_{n-1} = a_n$

C. $b_n / b_{n-1} = a_n$

D. none of these

Answer: B



Watch Video Solution

40. Statement 1: The coefficient of x^n is $\left(1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots + \frac{x^n}{n!}\right)^3$

is $\frac{3^n}{n!}$ Statement 2: The coefficient of $x^n \in e^{3x}$ is $\frac{3^n}{n!}$

A. $\frac{3^n}{n!}$

B. $\frac{3^{n-1}}{n!}$

C. $\frac{3^n}{(n-1)!}$

D. $\frac{1}{3^n \cdot n!}$

Answer: A



Watch Video Solution

41. In the expansion of $\left(3^{-x/4} + 3^{5x/4}\right)^n$ the sum of binomial coefficient is 64 and term with the greatest binomial coefficient exceeds the third by $(n-1)$, the value of x must be

A. 0

B. 1

C. 2

D. 3

Answer: A



42. The sum of the coefficients of even power of x in the expansion of $(1 + x + x^2 + x^3)^5$ is a. 256 b. 128 c. 512 d. 64

A. 256

B. 128

C. 512

D. 64

Answer: C



43. Maximum sum of coefficient in the expansion of $(1 - x \sin \theta + x^2)^n$ is 1 b. 2^n c. 3^n d. 0

A. 1

B. 2^n

C. 3^n

D. 0

Answer: C



Watch Video Solution

44. If the sum of the coefficients in the expansion of $(a + b)^n$ is 4096, then the greatest coefficient in the expansion is

A. 924

B. 792

C. 1594

D. none of these

Answer: A



Watch Video Solution

45. The value of $\hat{^20}C_0 + ^{20}C_1 + ^{20}C_2 + ^{20}C_3 + ^{20}C_4 + ^{20}C_{12} + ^{20}C_{13} + ^{20}C_{14} + ^{20}C_{15}$ is $2^{19} - \frac{(\hat{^20}C_{10} + ^{20}C_9)}{2}$ b. $2^{19} - \frac{(\hat{^20}C_{10} + 2 \times ^{20}C_9)}{2}$ c. $2^{19} - \frac{\hat{^20}C_{10}}{2}$ d. none of these
- A. $2^{19} - \frac{(.^{20}C_{10} + .^{20}C_9)}{2}$
- B. $2^{19} - \frac{(.^{20}C_{10} + 2 \times .^{20}C_9)}{2}$
- C. $2^{19} - \frac{.^{20}C_{10}}{2}$
- D. none of these

Answer: B



Watch Video Solution

46. The sum of series $\hat{^20}C_0 - ^{20}C_1 + ^{20}C_2 - ^{20}C_3 + \dots + ^{20}C_{10}$ is

A. $\frac{1}{2} \cdot ^{20}C_{10}$

B. 0

C. $.^{20} C_{10}$

D. $- .^{20} C_{10}$

Answer: A



Watch Video Solution

47. If $(3 + x^{2008} + x^{2009})^{2010} = a_0 + a_1x + a_2x^2 + \dots + a_nx^n$, then the value of $a_0 - \frac{1}{2}a_1 - \frac{1}{2}a_2 + a_3 - \frac{1}{2}a_4 - \frac{1}{2}a_5 + a_6 - \dots$ is 3^{2010} b. 1 c. 2^{2010} d. none of these

A. 3^{2010}

B. 1

C. 2^{2010}

D. none of these

Answer: C



48. Value of $\sum_{k=1}^{\infty} \sum_{r=0}^k \frac{1}{3^k} (^k C_r)$ is **a. $\frac{2}{3}$** b. $\frac{4}{3}$ c. 2 d. 1

A. $\frac{2}{3}$

B. $\frac{4}{3}$

C. 2

D. 1

Answer: C



Watch Video Solution

49. The value of $\sum_{r=0}^{10} (r)^{20} C_r$ is equal to $20(2^{18} + {}^{19} C_{10})$ b. $10(2^{18} + {}^{19} C_{10})$ c. $20(2^{18} + {}^{19} C_{11})$ d. $10(2^{18} + {}^{19} C_{11})$

A. $20(2^{18} + {}^{19} C_{10})$

B. $10(2^{18} + {}^{18} C_{10})$

C. $20(2^{16} + .^{19} C_{11})$

D. $10(2^{18} + .^{19} C_{11})$

Answer: A



Watch Video Solution

50.

$$\left[({}^n nC_0 + {}^n C_3 +)1/2({}^n nC_1 + {}^n C_2 + {}^n C_4 + {}^n C_5)^2 + 3/4({}^n nC_1 - {}^n C_2 \right]$$

3 b. 4 c. 2 d. 1

A. 3

B. 4

C. 2

D. 1

Answer: D



Watch Video Solution

51. The value of $\sum_{r=1}^{n+1} \left(\sum_{k=1}^n {}^k C_{r-1} \right)$ (wherer, $k, n \in N$) is equal to

a. $2^{n+1} - 2$ b. $2^{n+1} - 1$ c. 2^{n+1} d. none of these

A. $2^{n+1} - 2$

B. $2^{n+1} - 1$

C. 2^{n+1}

D. none of these

Answer: A



Watch Video Solution

52. The sum $\sum_{0 \leq i \leq j \leq 10} \left({}^{10} C_j \right) \left({}^j C_{r-1} \right)$ is equal to

A. $2^{10} - 1$

B. 2^{10}

C. $3^{10} - 1$

D. 3^{10}

Answer: C



Watch Video Solution

53. The value of ${}^{1000}C_{50} + {}^{999}C_{49} + {}^{998}C_{48} + \dots + {}^{950}C_0$ is

A. ${}^{1001}C_{50}$

B. ${}^{1002}C_{951}$

C. ${}^{1001}C_{950}$

D. ${}^{1002}C_{50}$

Answer: A



Watch Video Solution

54. If $\sum_{r=0}^n \{a_r(x - \alpha + 2)^r - b_r(\alpha - x - 1)^r\} = 0$, then prove that $b_n - (-1)^n a_n = 0$.

A. $b_n = 1 + a_n$

B. $b_n = (-1)^n \times a_n$

C. $b_n = (-1)^{n-1} \times a_n$

D. $b_n + 1 = a_n$

Answer: B



Watch Video Solution

55. If $\sum_{r=0}^{2n} a_r(x - 2)^r = \sum_{r=0}^{2n} b_r(x - 3)^r$ and $a_k = 1$ for all $k \geq n$, then

show that $b_n = {}^{2n+1} C_{n+1}$.

A. ${}^{2n+1} C_{n-1}$

B. ${}^{2n} C_{n+1}$

C. $\dots^{2n} C_n$

D. $\dots^{2n+1} C_{n+1}$

Answer: D



Watch Video Solution

56. The value of $\sum_{r=0}^{40} r \cdot {}^{40} C_r \cdot {}^{30} C_r$ is

A. 10460

B. 11240

C. 11520

D. 12640

Answer: C



Watch Video Solution

57. If ${}^n C_{r+1} : {}^n C_r : {}^{n-1} C_{r-1} = 11 : 6 : 3$, then $nr =$

A. 20

B. 30

C. 40

D. 50

Answer: D



Watch Video Solution

58. If the three consecutive terms in the expansion of $(1 + x)^n$ are 28, 56, and 70, then the value of n is.

A. $\frac{ac + ab + bc}{b^2 + ac}$

B. $\frac{2ac + ab + bc}{b^2 - ac}$

C. $\frac{ab + ac}{b^2 - ac}$

D. none of these

Answer: B



Watch Video Solution

59. Which term in the expansion of $(2 - 3x)^{19}$ has algebraically the last coefficients ?

A. 10^{th}

B. 11^{th}

C. 12^{th}

D. 13^{th}

Answer: C



Watch Video Solution

60. The value of $\frac{{}^nC_0}{n} + \frac{{}^nC_1}{n+1} + \frac{{}^nC_2}{n+2} + \dots + \frac{{}^nC_n}{2n}$ is equal to

- A. $\int_0^1 x^{n-1} (1-x)^n dx$
- B. $\int_1^2 x^n (x-1)^{n-1} dx$
- C. $\int_1^2 (1+x)^n dx$
- D. $\int_0^1 (1-x)^n x^{n-1} dx$

Answer: B



Watch Video Solution

61. The value of $\sum_{r=1}^n (-1)^{r+1} \frac{\wedge nCr}{r+1}$ is equal to - $\frac{1}{n+1}$ b. $\frac{1}{n}$ c. $\frac{1}{n+1}$ d. $\frac{n}{n+1}$

A. $-\frac{1}{n+1}$

B. $-\frac{1}{n}$

C. $\frac{1}{n+1}$

D. $\frac{n}{n+1}$

Answer: D



Watch Video Solution

62. If $\sum_{r=0}^{10} \left(\frac{r+2}{r+1} \right) \wedge nC_r = \frac{2^8 - 1}{6}$, then n is 8 b. 4 c. 6 d. 5

A. 8

B. 4

C. 6

D. 5

Answer: D



Watch Video Solution

63. The value of $\sum_{r=0}^{10} (-1)^r \cdot 4^{10-r} 30C_r {}^{30-r} C_{10-r}$ is equal to

A. ${}^{30} C_{10} \times 2^{10}$

B. $.^{30}C_9 \times 4^{10}$

C. $.^{30}C_{10} \times 3^{10}$

D. $.^{30}C_9 \times 4^{10}$

Answer: C



Watch Video Solution

64. The value of $\sum_{r=0}^{10} r^{10} C_r, 3^r \cdot (-2)^{10-r}$ is -

A. 20

B. 10

C. 300

D. 30

Answer: D



Watch Video Solution

65. ${}^{15}C_0 =$

A. 15

B. -15

C. 0

D. 51

Answer: C



Watch Video Solution

66.

If

$(1 + x)^n = C_0 + C_1x + C_2x^2 + \dots + C_nx^n$, then $C_0C_2 + C_1C_3 + C_2C_4 + \dots +$

a. $\frac{(2n)!}{(n!)^2}$ b. $\frac{(2n)!}{(n-1)!(n+1)!}$ c. $\frac{(2n)!}{(n-2)!(n+2)!}$ d. none of these

A. $\frac{(2n)!}{(n!)^2}$

B. $\frac{(2n)!}{(n-1)!(n+1)!}$

$$\text{C. } \frac{(2n)!}{(n-2)!(n+2)!}$$

D. none of these

Answer: C



Watch Video Solution

$$67. {}^{\wedge} 404C_4 - {}^{303}C_4. {}^4C_1 + {}^{202}C_4. {}^4C_2 - {}^{101}C_4. {}^4C_3 =$$

A. $(401)^4$

B. $(101)^4$

C. 0

D. $(201)^4$

Answer: B



Watch Video Solution

68. The value of $\sum_{r=0}^{40} r^{40} C_r {}^{30}C_r$ is

A. ${}^{40}C_{29}$

B. ${}^{40}C_{30}$

C. ${}^{60}C_{29}$

D. ${}^{70}C_{30}$

Answer: A



Watch Video Solution

69. The value of $\sum_{r=0}^{50} (-1)^r \frac{\hat{C}_r}{r+2}$ is equal to $\frac{1}{50 \times 51}$ b. $\frac{1}{52 \times 50}$ c. $\frac{1}{52 \times 51}$ d. none of these

A. $\frac{(17)! - 2^{16}}{(17)!}$

B. $\frac{(18)! 2^{17}}{(18)!}$

C. $\frac{(16)! - 2^{15}}{(16)!}$

$$D. \frac{(15)! - 2^{14}}{(15)!}$$

Answer: A



Watch Video Solution

70. $(n + 2)nC_0(2^{n+1}) - (n + 1)nC_1(2^n) + (n)nC_2(2^{n-1}) - \dots$ is equal to

A. 4

B. $4n$

C. $4(n+1)$

D. $2(n+2)$

Answer: C



Watch Video Solution

71. The value of $\sum_{r=0}^{50} (-1)^r \frac{\hat{(50)}C_r}{r+2}$ is equal to $\frac{1}{50 \times 51}$ b. $\frac{1}{52 \times 50}$ c. $\frac{1}{52 \times 51}$ d. none of these

A. $\frac{1}{50 \times 51}$

B. $\frac{1}{52 \times 50}$

C. $\frac{1}{52 \times 51}$

D. none of these

Answer: C



Watch Video Solution

72. $\sum_{r=0}^{300} a_r x^r = (1 + x + x^2 + x^3)^{100}$. If $a = \sum_{r=0}^{300} a_r$, then $\sum_{r=0}^{300} r a_r$ is equal to 300a b. 100a c. 150a d. 75a

A. ${}^n C_r$

B. ${}^n C_r 3^r$

C. $.^2n C_r$

D. $.^n C_r 2^r$

Answer: D



Watch Video Solution

73. $\sum_{r=0}^{300} a_r x^r = (1 + x + x^2 + x^3)^{100}$. If $a = \sum_{r=0}^{300} a_r$, then $\sum_{r=0}^{300} r a_r$ is

equal to 300a b. 100a c. 150a d. 75a

A. 300a

B. 100a

C. 150a

D. 75a\

Answer: C



Watch Video Solution

74. The value of $\sum_{r=0}^{20} r(20-r)(\hat{(20)}C_r)^2$ is equal to $400^{39}C_{20}$ b.
400⁴⁰C₁₉ c. 400³⁹C₁₉ d. 400³⁸C₂₀

A. 400.³⁹ C₂₀

B. 400.⁴⁰ C₁₉

C. 400.³⁹ C₁₉

D. 400.³⁸ C₂₀

Answer: D



Watch Video Solution

75.

If $f(x) = .^{40}C_1 \cdot x(1-x)^{39} + 2 \cdot .^{40}C_2 x^2(1-x)^{38} + 3 \cdot .^{40}C_3 x^3(1-x)^{37} + \dots$, then the value of $f(3)$ is

A. 120

B. 150

C. 200

D. 240

Answer: A



Watch Video Solution

76. If $a_n = \sum_{r=0}^n \frac{1}{{}^n C_r}$, then the value of $\left(\sum \sum \right)_{0 \leq i < j \leq n} \left(\frac{i}{{}^n C_i} + \frac{j}{{}^n C_j} \right)$

A. an^2

B. $\frac{a^2 n}{2}$

C. $a^2 n$

D. $\frac{n^2 a}{2}$

Answer: D



Watch Video Solution

77. In the expansion of $[(1+x)/(1-x)]^2$, the coefficient of x^n will be

a. $4n$ b. $4n - 3$ c. $4n + 1$ d. none of these

A. $4n$

B. $4n - 3$

C. $4n + 1$

D. none of these

Answer: A



Watch Video Solution

78. Show that $x^n = 1 + n\left(1 - \frac{1}{x}\right) + \frac{n(n+1)}{1.2}\left(1 - \frac{1}{x}\right)^2 + \dots$

A. x^n

B. x^{-n}

C. $\left(1 - \frac{1}{x}\right)^n$

D. none of these

Answer: A



Watch Video Solution

$$79. \sum_{k=1}^{\infty} k \left(1 - \frac{1}{n}\right)^{k-1} = n(n-1) \text{ b. } n(n+1) \text{ c. } n^2 \text{ d. } (n+1)^2$$

A. $n(n-1)$

B. $n(n+1)$

C. n^2

D. $(n+1)^2$

Answer: C



Watch Video Solution

80. The coefficient of x^4 in the expansion of $\left\{ \sqrt{1+x^2} - x \right\}^{-1}$ in ascending powers of x , when $|x| < 1$, is
a. 0 b. $\frac{1}{2}$ c. $-\frac{1}{2}$ d. $-\frac{1}{8}$

A. 0

B. $\frac{1}{2}$

C. $-\frac{1}{2}$

D. $-\frac{1}{8}$

Answer: D



Watch Video Solution

81. $1 + \frac{1}{3}x + \frac{1 \times 4}{3 \times 6}x^2 + \frac{1 \times 4 \times 7}{3 \times 6 \times 9}x^3 + \dots$ is equal to

A. x

B. $(1 + x)^{1/3}$

C. $(1 - x)^{1/3}$

D. $(1 - x)^{-1/3}$

Answer: D



Watch Video Solution

82. $1 + \frac{1}{4} + \frac{1 \times 3}{4 \times 8} + \frac{1 \times 3 \times 5}{4 \times 8 \times 12} + \dots \dots$ is equal to

A. $\sqrt{2}$

B. $\frac{1}{\sqrt{2}}$

C. $\sqrt{3}$

D. $\frac{1}{\sqrt{3}}$

Answer: A



Watch Video Solution

83. If $|x| < 1$, then $1 + n\left(\frac{2x}{1+x}\right) + \frac{n(n+1)}{2!}\left(\frac{2x}{1+x}\right)^2 + \dots \dots$ is equal to

- A. $\left(\frac{2x}{1+x}\right)^n$
- B. $\left(\frac{1+x}{2x}\right)^n$
- C. $\left(\frac{1-x}{1+x}\right)^n$
- D. $\left(\frac{1+x}{1-x}\right)^n$

Answer: D



Watch Video Solution

84. Find the coefficient of x^5 in $(1 + 2x + 3x^2 \dots \dots \dots)^{-\frac{3}{2}}$

- A. 21
- B. 25
- C. 26
- D. none of these

Answer: D



Watch Video Solution

85. If $|x| < 1$, then find the coefficient of x^n in the expansion of $(1 + x + x^2 + x^3 + \dots)^2$.

- A. n
- B. n-1
- C. n+2
- D. n+1

Answer: D



Watch Video Solution

86. If x is positive, the first negative term in the expansion of $(1 + x)^{\frac{27}{5}}$ is

- A. 5th term
- B. 8th term
- C. 6th term

D. 7th term

Answer: B



Watch Video Solution

87. If x is so small that x^3 and higher power of x may neglected, then

$$\frac{(1+x)^{\frac{3}{2}} - \left(1 + \frac{1}{2}x\right)^3}{(1-x)^{\frac{1}{2}}} \text{ may be approximated as}$$

A. $3x + \frac{3}{x^2}$

B. $1 - \frac{3}{8}x^2$

C. $\frac{x}{2} - \frac{3}{x}x^2$

D. $-\frac{3}{8}x^2$

Answer: D



Watch Video Solution

88. If the expansion in power of x of the function

$$\frac{1}{(1-ax)(1-bx)} \text{ is } a_0 + a_1x + a_2x^2 + a_3x^3 + \dots, \text{ then } a_n \text{ is}$$

A. $\frac{b^n - a^n}{b - a}$

B. $\frac{a^n - b^n}{b - a}$

C. $\frac{a^{n+1} - b^{n+1}}{b - a}$

D. $\frac{b^{n+1} - a^{n+1}}{b - a}$

Answer: D



Watch Video Solution

89. If $\frac{x^2 + x + 1}{1 - x} = a_0 + a_1x + a_2x^2 + \dots$, the $b \sum_{r=1}^{50} a_r$ equal to

A. 148

B. 146

C. 149

D. none of these

Answer: C



Watch Video Solution

90.

If

$(1 - x)^{-n} = a_0 + a_1x + a_2x^2 + \dots + a_rx^r + , \text{ then } a_0 + a_1 + a_2 + \dots + a_r$

is equal to (a) $\frac{n(n+1)(n+2)(n+r)}{r!}$ (b) $\frac{(n+1)(n+2)(n+r)}{r!}$ (c)

$\frac{n(n+1)(n+2)(n+r-1)}{r!}$ (d) none of these

A. $\frac{n(n+1)(n+2)\dots(n+r)}{r!}$

B. $\frac{(n+1)(n+2)\dots(n+r)}{r!}$

C. $\frac{n(n+1)(n+2)\dots(n+r-1)}{r!}$

D. none of these

Answer: B



Watch Video Solution

Exercise (Multiple)

1. The value of x in the expression $\left(x + x^{(\log)_{10}}\right)^5$ if third term in the expansion is 10,00,000 is/are a. 10 b. 100 c. $10^{-5/2}$ d. $10^{-3/2}$

A. 10

B. 100

C. $10^{-5/2}$

D. $10^{-3/2}$

Answer: A::C



Watch Video Solution

2. The value of x for which the sixth term in the expansion of

$$\left[2^{\log} - 2^{\sqrt{9^{(x-1)} + 7}} + \frac{1}{2^{\frac{1}{5}} (\log)_2 (3^{(x-1)+1})} \right]^7 \text{ is } 84 \text{ is}$$

A. 4

B. 3

C. 2

D. 1

Answer: C::D



Watch Video Solution

3. If the 4th term in the expansion of $(ax + 1/x)^n$ is $5/2$, then (a) $a = \frac{1}{2}$
b. $n = 8$ c. $a = \frac{2}{3}$ d. $n = 6$

A. $a = \frac{1}{2}$

B. $n = 8$

C. $a = \frac{2}{3}$

D. $n = 6$

Answer: A::D



Watch Video Solution

4. In the expansion of $(x + a)^n$ if the sum of odd terms is P and the sum of even terms is Q , then (a) $P^2 - Q^2 = (x^2 - a^2)^n$ (b)

$$4PQ = (x + a)^{2n} - (x - a)^{2n} \quad (c)$$

$$2(P^2 + Q^2) = (x + a)^{2n} + (x - a)^{2n} \quad (d)\text{all of these}$$

A. $P^2 - Q^2 = (x^2 - a^2)^n$

B. $4PQ = (x + a)^{2n} - (x - a)^{2n}$

C. $2(P^2 + Q^2) = (x + a)^{2n} + (x - a)^{2n}$

D. none of these

Answer: A::B::C



Watch Video Solution

5. If $(4 + \sqrt{15})^n = I + f$, where n is an odd natural number, I is an integer and '0

A. I is an odd integer

B. I is an even nteger

C. $(I + f)(I - f) = 1$

D. $I - f = (4 - \sqrt{15})^n$

Answer: A::C::D



Watch Video Solution

6. If the coefficients of x^{39} and x^{40} are equal in the expansion of $(p + qx)^{49}$. then the possible values of p and q are

A. 4,16

B. 1,4

C. 5,20

D. 3,9

Answer: A::B::C



Watch Video Solution

7. The sum of coefficient in the expansion of $(1 + ad - 2x^2)^n$ is positive, when $a < 1$ and $n = 2k, k \in N$ negative, when $a < 1$ and $n = 2k + 1, k \in N$ positive, when $a < 1$ and $n \in N$ zero, when $a = 1$

A. positive, when $a < 1$ and $n = 2k, k \in N$

B. negative, when $a < 1$ and $n = 2k + 1, k \in N$

C. positive, when $a > 1$ and $n \in N$

D. zero, when $a = 1$

Answer: A::B::C::D



Watch Video Solution

8. Let $(1 + x^2)^2(1 + x)^n = \sum_{k=0}^{n+4} a_k x^k$. If a_1, a_2 and a_3 are in arithmetic progression, then the possible value/values of n is/are a. 5 b. 4 c. 3 d. 2

A. 5

B. 4

C. 3

D. 2

Answer: B::C::D



Watch Video Solution

9.

For natural numbers

$m, n,$ if $(1 - y)^m(1 + y)^n = 1 + a_1y + a_2y^2 + , and a_1 = a_2 = 10, then$

' $m \neq n. m+n=80$ and $m-n=20$ '

A. $m < n$

B. $m > n$

C. $m + n = 80$

D. $m - n = 20$

Answer: A::C



Watch Video Solution

10. The middle term in the expansion of $(x/2 + 2)^8$ is 1120, then $x \in R$ is equal to a. -2 b. 3 c. -3 d. 2

A. -2

B. 3

C. -3

D. 2

Answer: A::D



Watch Video Solution

11. In the expansion of $\left(3 - \sqrt{\frac{17}{4} + 3\sqrt{2}}\right)^{15}$ the 11th term is a

A. an irrational number

B. a rational number

C. a positive integer

D. a negative integer

Answer: A



Watch Video Solution

12. For the expansion $(x \sin p + x^{-1} \cos p)^{10}$, ($p \in R$),

A. the greatest value of the term independent of x is $10! / 2^5 (5!)^2$

B. the least value of sum of coefficient is zero

C. the greatest value of sum of coefficient is 12

D. the last value of the term independent of x occurs when

$$p = (2n + 1)' \frac{\pi}{4}, n \in Z$$

Answer: A::B::C



Watch Video Solution

13. For which of the following values of x , 5th term is the numerically greatest term in the expansion of $(1 + x/3)^{10}$, a. -2 b. 1.8 c. 2 d. -1. 9

A. -2

B. 1.8

C. 2

D. -19

Answer: A::B::C::D



Watch Video Solution

14. Which of the following is/are true ?

A. $1000^{1000} > 1002^{999}$

B. $1000^{1000} < 1002^{999}$

C. $1000^{1002} < 1002^{1000}$

D. $1000^{1002} > 1002^{1000}$

Answer: A::D



Watch Video Solution

15. If $\sum_{r=0}^n \frac{r}{nC_r} = \sum_{r=0}^n \frac{n^2 - 3n + 3}{2 \cdot {}^n C_r}$, then a. $n = 1$ b. $n = 2$ c. $n = 3$

d. none of these

A. $n = 1$

B. $n = 2$

C. $n = 3$

D. none of these

Answer: A::C



Watch Video Solution

16. The value of $\hat{n}C_1 + \hat{n+1}C_2 + \hat{n+2}C_3 + \dots + \hat{n+m-1}C_m$ is equal to

$$\hat{m} + nC_{n-1} + \hat{m+n-1}$$

$$\hat{m+n-1}C_{m-1}$$

A. $\hat{m+n}C_n - 1$

B. $\hat{m+n}C_{n-1}$

C. $\hat{m}C_1 + \hat{m+1}C_2 + \hat{m+2}C_3 + \dots + \hat{m+n-1}C_n$

D. $\hat{m+n}C_m - 1$

Answer: A::C::D



Watch Video Solution

17. If the number of terms in the expansion of $\left(1 - \frac{2}{x} + \frac{4}{x^2}\right)x \neq 0$, is

28, then the sum of coefficient of all the terms in this expansion, is

A. number of terms is $2n + 1$

B. constant term is 2^{n-1}

C. coefficient of x^{2n-2} is n

D. coefficient of x^2 in n

Answer: A::C



Watch Video Solution

18. In the expansion of $(7^{1/3} + 11^{1/9})^{6561}$,

A. there are exactly 730 rational terms

B. there are exactly 5832 irrational terms

C. the term which involves greatest binomial coefficients is irrational

D. the term which involves greatest binomial coefficients is rational

Answer: A::B::C



Watch Video Solution

19. If $(1 - x)^n = C_0 + C_1x + C_2x^2 + \dots + C_nx^n$, then the value of

1. $C_1 + 2. C_3 + 3. C_5 + \dots + n. C_n =$

A. a positive value

B. a negative value

C. divisible by 2^{n-1}

D. divisible by 2^n

Answer: B::C



Watch Video Solution

20. In the expansion of $(a + b)^n$, if two consecutive terms are equal, then

which of the following is/are always integer ?

A. $\frac{(n + 1)b}{a + b}$

B. $\frac{(n + 1)a}{a + b}$

C. $\frac{na}{a - b}$

$$D. \frac{na}{a+b}$$

Answer: A::B



Watch Video Solution

21. If for z as real or complex,

$$(1 + z^2 + z^4)^8 = C_0 + C_1 z^2 + C_2 z^4 + \dots + C_{16} z^{32} \text{ then}$$

$$C_0 - C_1 + C_2 - C_3 + \dots + C_{16} = 1$$

$$C_0 + C_3 + C_6 + C_9 + C_{12} + C_{15} = 3^7$$

$$C_2 + C_5 + C_8 + C_{11} + C_{14} = 3^6$$

$$C_1 + C_4 + C_7 + C_{10} + C_{13} + C_{16} = 3^7$$

A. $C_0 - C_1 + C_2 - C_3 + \dots + C_{16} = 1$

B. $C_0 + C_3 + C_6 + C_{12} + C_{15} = 3^7$

C. $C_2 + C_5 + C_8 + C_{11} + C_{14} = 3^6$

D. $C_1 + C_4 + C_7 + C_{10} + C_{13} + C_{16} = 3^7$

Answer: A::B::D



Watch Video Solution

22. If $f(m) = \sum_{i=0}^{30} - (m) \binom{30}{30-i} \binom{20}{m-i}$ where $\binom{p}{q} = {}^p C_q$, then

- A. maximum value of $f(m)$ is $.{}^{50} C_{25}$
- B. $f(0) + f(1) + \dots + f(50) = 2^{50}$
- C. $f(m)$ is always divisible by 50 ($1 \leq m \leq 49$)
- D. The value of $\sum_{m=0}^{50} (f(m))^2 = .{}^{100} C_{50}$

Answer: A::B::D



Watch Video Solution

23.

$$(1+x)^n = C_0 + C_1x + C_2x^2 + \dots + C_nx^n, n \in N, \text{ then } C_0 - C_1 + C_2 - \dots$$

is equal to '(m

A. $\frac{(n-1)(n-2)\dots(n-m+1)}{(m-1)!} (-1)^{m-1}$

B. $.^{n-1}C_{m-1}(-1)^{m-1}$

C. $\frac{(n-1)(n-2)\dots(n-m)}{(m-1)!}(-1)^{m-1}$

D. $.^{n-1}C_{n-m}(-1)^{m-1}$

Answer: A::B::D



Watch Video Solution

24. If $\sum_{r=0}^n (pr + 2) \cdot {}^nC_r = (25)(64)$ where $n, p \in N$, then

A. $p = 3$

B. $p = 4$

C. $n = 7$

D. $n = 6$

Answer: A::C



Watch Video Solution

25.

If

$$\left(x + \frac{1}{x} + 1\right)^6 = a_0 + \left(a_1x + \frac{b_i}{x}\right) + \left(a_2x^2 + \frac{b_2}{x^2}\right) + \dots + \left(a_6x^6 + \frac{b_6}{x^6}\right)$$

, then

A. $a = 141$

B. $a_5 = 6$

C. $\sum_{i=1}^6 a_i + b_i = 588$

D. $\sum_{i=1}^6 a_i + b_i = 3^6$

Answer: A::B::C



Watch Video Solution

26.

The value of

$${}^n C_0 \times {}^{2n} C_r - {}^n C_1 \times {}^{2n-2} C_r + {}^n C_2 \times {}^{2n-4} C_r + \dots \text{ is equal to}$$

A. ${}^n C_{r-n} \times 2^{2n-r}$ if $r \geq n$

B. 0, if $r < n$

C. $.^n C_{r-n} \times 2^{n-r}$ if $r \geq n$

D. $.^{-n} C_{r-n} \times 2^{2n-r}$ if $r < n$

Answer: A::B



Watch Video Solution

27.

The

sum

$$2 \times .^{40} C_2 + 6 \times .^{40} C_3 + 12 \times .^{40} C_4 + 20 \times .^{40} C_5 + \dots + 1560 \times .^{40} C_{40}$$

is divisible by

A. 3

B. 5

C. 13

D. 2^{41}

Answer: A::B::C::D



Watch Video Solution

Exercise (Comprehension)

1. The sixth term in the expansion of $\left(\sqrt{2^{\log(10 - 3^x)}} + \left(2^{(x-2)\log 3} \right)^{\frac{1}{5}} \right)^m$

is equal to 21, if it is known that the binomial coefficient of the 2nd 3rd and 4th terms in the expansion represent, respectively, the first, third and fifth terms of an A.P. (the symbol log stands for logarithm to the base 10)

The value of m is

A. 6

B. 7

C. 8

D. 9

Answer: B



Watch Video Solution

2. The sixth term in the expansion of $\left(\sqrt{2^{\log(10 - 3^x)}} + \left(2^{(x-2)\log 3} \right)^{\frac{1}{5}} \right)^m$ is equal to 21, if it is known that the binomial coefficient of the 2nd 3rd and 4th terms in the expansion represent, respectively, the first, third and fifth terms of an A.P. (the symbol log stands for logarithm to the base 10) The sum of possible value of x is

A. 1

B. 3

C. 4

D. 2

Answer: D



Watch Video Solution

3. The sixth term in the expansion of $\left(\sqrt{2^{\log(10 - 3^x)}} + \left(2^{(x-2)\log 3} \right)^{\frac{1}{5}} \right)^m$ is equal to 21, if it is known that the

binomial coefficient of the 2nd 3rd and 4th terms in the expansion represent, respectively, the first, third and fifth terms of an A.P. (the symbol log stands for logarithm to the base 10) The value of m is

A. 6

B. 7

C. 8

D. 9

Answer: C



Watch Video Solution

4. If the 2nd, 3rd and 4th terms in the binomial expansion of $(x + a)^n$ are 240, 720 and 1080 for a suitable values of x, a and n.



Watch Video Solution

5. If the 2nd, 3rd and 4th terms in the binomial expansion of $(x + a)^n$ are 240, 720 and 1080 for a suitable values of x, find x, a and n.

A. 16

B. 160

C. 32

D. 81

Answer: C



Watch Video Solution

6. If the 2nd, 3rd and 4th terms in the binomial expansion of $(x + a)^n$ are 240, 720 and 1080 for a suitable values of x, find x, a and n.

A. 1664

B. 2376

C. 1562

D. 1486

Answer: C



Watch Video Solution

7. An equation $a_0 + a_1x + a_2x^2 + \dots + a_{99}x^{99} + x^{100} = 0$ has roots $99C_0, 99C_1, 99C_2, \dots, 99C_{99}$. Find the value of a_{99}



Watch Video Solution

8. An equation $a_0 + a_1x + a_2x^2 + \dots + a_{99}x^{99} + x^{100} = 0$ has roots $.^{99}C_0, .^{99}C_1, C_{99}C_2, \dots, .^{99}C_{99}$

The value of a_{99} is equal to



Watch Video Solution

9. An equation $a_0 + a_2x^2 + \dots + a_{99}x^{99} + x^{100} = 0$ has roots $.^{99}C_0, .^{99}C_1, C_{99}C_2, \dots, .^{99}C_{99}$

The value of $(.^{99}C_0)^2 + (.^{99}C_1)^2 + \dots + (.^{99}C_{99})^2$ is equal to

A. $2a_{98} - a_{99}^2$

B. $a_{99}^2 - a_{98}$

C. $a_{99}^2 - 2a_{98}$

D. none of these

Answer: C



Watch Video Solution

10.

If

$$a = .^{20}C_0 + .^{20}C_3 + .^{20}C_6 + .^{20}C_9 + \dots, b = .^{20}C_1 + .^{20}C_4 + .^{20}C_7 + \dots$$

$$\text{and } c = .^{20}C_2 + .^{20}C_5 + .^{20}C_8 + \dots, \text{ then}$$

Value of $a^3 + b^3 + c^3 - 3abc$ is



Watch Video Solution

11.

If

$$a = .^{20} C_0 + .^{20} C_3 + .^{20} C_6 + .^{20} C_9 + \dots, b = .^{20} C_1 + .^{20} C_4 + .^{20} C_7 + \dots$$

$$\text{and } c = .^{20} C_2 + .^{20} C_5 + .^{20} C_8 + \dots, \text{ then}$$

Value of $(a - b)^2 + (b - c)^2 + (c - a)^2$ is

A. 1

B. 2

C. 2^{20}

D. 2^{40}

Answer: B



Watch Video Solution

- 12.** The sum of all the coefficients of the terms in the expansion of $(x + y + z + w)^6$ which contain x but not y , is

A. 4096

B. 1560

C. 3367

D. 670

Answer: B



Watch Video Solution

13. Consider the expansion of $(a + b + c + d)^6$. Then the sum of all the coefficients of the term

Which contains all of $a, b, c,$ and d is

A. 729

B. 3367

C. 665

D. 1024

Answer: C



Watch Video Solution

14. Consider the expansion of $(a + b + c + d)^6$. Then the sum of all the coefficients of the term

Which contains all of $a, b, c,$ and d is

A. 2884

B. 4032

C. 1974

D. 2702

Answer: D



Watch Video Solution

15.

Let

$$P = \sum_{r=1}^{50} \frac{^{50+r}C_r(2r-1)}{^{50}C_r(50+r)}, Q = \sum_{r=0}^{50} (r+1)^{50} ({}^{50}C_r)^2, R = \sum_{r=0}^{100} (-1)^r ({}^{100}C_r)^2$$

The value of $P - Q$ is equal to



Watch Video Solution

16.

Let

$$P = \sum_{r=1}^{50} \frac{^{50+r}C_r(2r-1)}{^{50}C_r(50+r)}, Q = \sum_{r=0}^{50} ({}^{50}C_r)^2, R = \sum_{r=0}^{100} (-1)^r ({}^{100}C_r)^2$$

The value of $P - R$ is equal to

A. 1

B. -1

C. 2^{50}

D. 2^{100}

Answer: B



Watch Video Solution

17.

Let

$$P = \sum_{r=1}^{50} \frac{^{50+r}C_r(2r-1)}{^{50}C_r(50+r)}, Q = \sum_{r=0}^{50} (^{50}C_r)^2, R = \sum_{r=0}^{100} (-1)^r (^{100}C_r)^2$$

The value of $Q + R$ is equal to

A. $2P + 1$

B. $2P - 1$

C. $2P + 2$

D. $2P - 2$

Answer: C



Watch Video Solution

18. If $(1 + x + x^2)^{20} = a_0 + a_1x^2 + \dots + a_{40}x^{40}$, then following questions.

The value of $a_0 + a_1 + a_2 + \dots + a_{19}$ is



Watch Video Solution

19. If $(1 + x + x^2)^{20} = a_0 + a_1x^2 + \dots + a_{40}x^{40}$, then following questions.

The value of $a_0^2 - a_1^2 + a_2^2 - \dots - a_{19}^2$ is



Watch Video Solution

20. If $(1 + x + x^2)^{20} = a_0 + a_1x^2 + \dots + a_{40}x^{40}$, then following questions.

The value of $a_0 + 3a_1 + 5a_2 + \dots + 81a_{40}$ is

A. 161×3^{20}

B. 41×3^{40}

C. 41×3^{20}

D. none of these

Answer: C



Watch Video Solution

Matrix

1. Evaluate $\int_1^3 (2x - 3)dx$

 Watch Video Solution

2. Evaluate $\int \frac{x - 1}{x^2 - 2x + 1} dx$

 Watch Video Solution

3. Evaluate $\int_0^3 (x^2 - 6x + 5)dx$

 Watch Video Solution

4. If the second term of the expansion $\left[a^{\frac{1}{13}} + \frac{a}{\sqrt{a^{-1}}} \right]^n$ is $14a^{5/2}$, then the value of $\frac{\hat{n}C_3}{\hat{n}C_2}$ is.



Watch Video Solution

Exercise (Numerical)

1. Evaluate $\int_0^3 (x^2 - 6x + 5) dx$



Watch Video Solution

2. If the three consecutive in the expansion of $(1 + x)^n$ are 28, 56, and 70, then the value of n is.



Watch Video Solution

3. Least positive integer just greater than $(1 + 0.00002)^{50000}$ is.



Watch Video Solution

4. If the second term of the expansion $\left[a^{\frac{1}{13}} + \frac{a}{\sqrt{a^{-1}}} \right]^n$ is $14a^{5/2}$, then the value of $\frac{\hat{n}C_3}{\hat{n}C_2}$ is.



Watch Video Solution

5. If the constant term in the binomial expansion of $\left(x^2 - \frac{1}{x} \right)^n$, $n \in N$ is 15, then the value of n is equal to.



Watch Video Solution

6. The largest value of x for which the fourth tem in the expansion $\left(\frac{5^2}{3} (\log)_5 \sqrt{4^{x+44}} + \frac{1}{5^{\log} - 52^{(x-1)+73}} \right)$ is 336 is.



Watch Video Solution

7. Let a and b be the coefficients of x^3 in $(1 + x + 2x^2 + 3x^3)^4$ and $(1 + x + 2x^2 + 3x^3 + 4x^4)^4$, then respectively. Then the value of $4a/b$ is.

 Watch Video Solution

8. If R is remainder when $6^{83} + 8^{83}$ is divided by 49, then the value of $R/5$ is.

 Watch Video Solution

9. The remainder, if $1 + 2 + 2^2 + \dots + 2^{1999}$ is divided by 5 is.

 Watch Video Solution

10. Given $(1 - 2x + 5x^2 - 10x^3)(1 + x)^n = 1 + a_1x + a_2x^2 + \dots$ and that $a_1 = 2a_2$ then the value of n is.

 Watch Video Solution

11. The largest real value of x such that $\sum_{k=0}^4 \left(\frac{3^{4-k}}{(4-k)!} \right) \left(\frac{x^k}{k!} \right) = \frac{32}{3}$ is.



Watch Video Solution

12. Coefficient of x^{2009} in $(1 + x + x^2 + x^3 + x^4)^{1001}(1 - x)^{1002}$ is



Watch Video Solution

13. The total number of different terms in the product $(.^{101} C_0 - .^{101} C_1 x + .^{101} C_2 x^2 - \dots - .^{101} C_{101} x^{101}) (1 + x + x^2 + \dots +$ is ____.



Watch Video Solution

14. The constant term in the expansion of $\left(x - \frac{2}{x^2} \right)^9$ is



Watch Video Solution

15. The value of $\sum_{r=0}^3 {}^8C_r ({}^5C_{r+1} - {}^4C_r)$ is ____.



Watch Video Solution

16. The value of $(.^{21}C_1 - .^{10}C_1) + (.^{21}C_2 - .^{10}C_2) + (.^{21}C_3 - .^{10}C_3) + (.^{21}C_4 - .^{10}C_4)$ is



Watch Video Solution

17. Let $a = 3^{\frac{1}{223}} + 1$ and for all $t \geq 3, t \leq n$

$$f(n) = {}^nC_0 a^{n-1} - {}^nC_1 a^{n-2} + {}^nC_2 a^{n-3} - \dots + (-1)^{n-1} \dot{n}C_{n-1} a^0.$$

If the value of $f(2007) + f(2008) = 3^k$ where $k \in N$, then the value of k is.



Watch Video Solution

18. Let $1 + \sum_{r=1}^{10} (3^r 10C_r + r 10C_r) = 2^{10}(\alpha \cdot 4^5 + \beta)$ where $\alpha, \beta \in N$ and $f(x) = x^2 - 2x - k^2 + 1$. If α, β lies between the roots of $f(x) = 0$, then find the smallest positive integral value of k .



Watch Video Solution

19. The value of $(\lim)_{n \rightarrow \infty} \sum_{r=1}^{r-1} \left(\sum_{t=0}^{r-1} \frac{1}{5^n} \dot{+} nC_r \dot{+} rC_t \cdot 3^t \right)$ is equal to



Watch Video Solution

20. If $\sum_{r=0}^{10} \left(\frac{r+2}{r+1} \right) \wedge nC_r = \frac{2^8 - 1}{6}$, then n is 8 b. 4 c. 6 d. 5



Watch Video Solution

21. If $S_n = (.^n C_0)^2 + (.^n C_1)^2 + (.^n C_n)^n$, then maximum value of $\left[\frac{S_{n+1}}{S_n} \right]$ is ____.

(where $[\cdot]$ denotes the greatest integer function)



Watch Video Solution

22. The value of $.^{40} C_0 \times .^{100} C_{40} - .^{40} C_1 \times .^{99} C_{40} + .^{40} C_2 \times .^{98} C_{40} - \dots + .^{40} C_{40} \times .^{60}$ is equal to ____.



Watch Video Solution

23. The value of $\sum_{0 \leq i < j \leq 5} \sum ({}^5 C_j) ({}^j C_i)$ is equal to ____



Watch Video Solution

24. If $(1 - x - x^2)^{20} = \sum_{r=0}^{40} a_r \cdot x^r$, then value of $a_1 + 3a_3 + 5a_5 + \dots + 39a_{39}$ is



[Watch Video Solution](#)

25. The value of $\sum_{r=1}^{11} \tan^2\left(\frac{r\pi}{24}\right)$ is



[Watch Video Solution](#)

JEE Previous Year

1. The remainder left out when $8^{2n}(62)^{2n+1}$ is divided by 9 is (1) 0 (2) 2 (3) 7 (4) 8

A. 0

B. 2

C. 7

Answer: B**Watch Video Solution**

2. Let $S_1 = \sum_{j=1}^{10} (j-1)^{10} j(j-1)^{10} C_j$, $S_2 = \sum_{j=1}^{10} {}^{10}C_j$, and $S_3 = \sum_{j=1}^{10} j^2 \cdot {}^{10}C_j$.

Statement 1: $S_3 \times 2^9$.

Statement 2: $S_1 = 90 \times 2^8$ and $S_2 = 10 \times 2^8$.

- A. Statement 1 is false, statement 2 is true.
- B. Statement 1 is true, statement 2 is true, statement 2 is a correct explanation for statement 1.
- C. Statement 1 is true, statement 2 is true, statement 2 is not a correct explanation for statement 2.
- D. statement 1 is true, statement 2 is false.

Answer: B



Watch Video Solution

3. The coefficient of x^7 in the expansion of $(1 - x - x^2 + x^3)^6$ is :

A. 132

B. 144

C. -132

D. -144

Answer: D



Watch Video Solution

4. If n is a positive integer, then $(\sqrt{3} + 1)^{2n} - (\sqrt{3} - 1)^{2n}$ is (1) an irrational number (2) an odd positive integer (3) an even positive integer (4) a rational number other than positive integers

A. an irrational number

B. an odd positive integer

C. an even positive integer

D. a rational number other than positive integers

Answer: A



Watch Video Solution

5. The coefficient of the term independent of x in the expansion of

$$\left(\frac{x+1}{x^{2/3} - x^{1/3} + 1} - \frac{x-1}{x - x^{1/2}} \right)^{10}$$
 is 210 b. 105 c. 70 d. 112



Watch Video Solution

6. If the coefficients of x^3 and x^4 in the expansion of

$(1+ax+bx^2)(1-2x)^{18}$ in powers of x are both zero, then (a, b) is

equal to (1) $\left(16, \frac{251}{3}\right)$ (3) $\left(14, \frac{251}{3}\right)$ (2) $\left(14, \frac{272}{3}\right)$ (4) $\left(16, \frac{272}{3}\right)$

- A. $\left(16, \frac{251}{3}\right)$
- B. $\left(14, \frac{251}{3}\right)$
- C. $\left(14, \frac{272}{3}\right)$
- D. $\left(16, \frac{272}{3}\right)$

Answer: D



Watch Video Solution

7. The sum of coefficients of integral powers of x in the binomial expansion of $(1 - 2\sqrt{x})^{50}$ is: (1) $\frac{1}{2}(3^{50} + 1)$ (2) $\frac{1}{2}(3^{50})$ (3) $\frac{1}{2}(3^{50} - 1)$ (4) $\frac{1}{2}(2^{50} + 1)$

A. $\frac{1}{2}(3^{50} + 1)$

B. $\frac{1}{2}(3^{50})$

C. $\frac{1}{2}(3^{50} - 1)$

D. $\frac{1}{2}(2^{10} + 1)$

Answer: A



Watch Video Solution

8. If the number of terms in the expansion of $\left(1 - \frac{2}{x} + \frac{4}{x^2}\right)^n$, $x \neq 0$, is 28, then the sum of the coefficients of all the terms in this expansion, is : (1) 64 (2) 2187 (3) 243 (4) 729

A. 2187

B. 243

C. 729

D. 64

Answer: C



Watch Video Solution

9. The value of

$$(.^{21} C_1 - .^{10} C_1) + (.^{21} C_2 - .^{10} C_2) + (.^{21} C_3 - .^{10} C_3) + (.^{21} C_4 - .^{10} C_4)$$

is

A. $2^{20} - 2^{10}$

B. $2^{21} - 2^{11}$

C. $2^{21} - 2^{10}$

D. $2^{20} - 2^9$

Answer: A



Watch Video Solution

10. The sum of the co-efficients of all odd degree terms in the expansion

$$\text{of } (x+\sqrt{x^3-1})^5 + (x-\sqrt{x^3-1})^5, (x > 1)$$

A. 2

B. -1

C. 0

D. 1

Answer: A



Watch Video Solution

11. For $r = 0, 1, \dots, 10$, let A_r , B_r , and C_r denote, respectively, the coefficient of x^r in the expansion of $(1 + x)^{10}$, $(1 + x)^{20}$ and $(1 + x)^{30}$.

Then $\sum_{r=1}^{10} A_r(B_{10}B_r - C_{10}A_r)$ is equal to

A. $B_{10} - C_{10}$

B. $A_1(B_{10}^2 - C_{10}A_{10})$

C. 0

D. $C_{10} - B_{10}$

Answer: D



Watch Video Solution

12. Coefficient of x^{11} in the expansion of $(1 + x^2)(1 + x^3)^7(1 + x^4)^{12}$ is
1051 b. 1106 c. 1113 d. 1120

A. 1051

B. 1106

C. 1113

D. 1120

Answer: C



Watch Video Solution

13. The coefficients of three consecutive terms of $(1 + x)^{n+5}$ are in the ratio 5:10:14. Then $n = \underline{\hspace{2cm}}$.



Watch Video Solution

14. The coefficient of x^9 in the expansion of $(1 + x)(16x^2)(1 + x^3)(1 + x^{100})$ is

 Watch Video Solution

15. Let m be the smallest positive integer such that the coefficient of x^2 in the expansion of $(1 + x)^2 + (1 + x)^3 + (1 + x)^4 + \dots + (1 + x)^{49} + (1 + mx)^{50}$ is $(3n + 1)^{51} C_3$ for some positive integer n . Then the value of n is

 Watch Video Solution

16. Let $X = ({}^{10}C_1)^2 + 2({}^{10}C_2)^2 + 3({}^{10}C_3)^2 + \dots + 10({}^{10}C_{10})^2$, where ${}^{10}C_r$, $r \in \{1, 2, \dots, 10\}$ denote binomial coefficients. Then, the value of $\frac{1}{1430} X$ is _____.

 Watch Video Solution

Single correct Answer

1. Coefficient of x^6 in $(1 + x)(1 + x^2)^2(1 + x^3)^3 \dots (1 + x^n)^n$ is

A. 26

B. 28

C. 30

D. 35

Answer: B



Watch Video Solution

2. The term independent of x in the product $(4 + x + 7x^2)\left(x - \frac{3}{x}\right)^{11}$

is

A. $7 \cdot {}^{11}C_6$

B. $36 \cdot {}^{11}C_6$

C. $3^5 \cdot {}^{11}C_5$

D. $-12 \cdot 2^{11}$

Answer: B



Watch Video Solution

3. The 13^{th} term in the expansion of $(x^2 + 2/x)^n$ is independent of x

then the sum of the divisors of n is

A. 36

B. 37

C. 39

D. 51

Answer: D



Watch Video Solution

4. Coefficient of x^{2009} in $(1 + x + x^2 + x^3 + x^4)^{1001}(1 - x)^{1002}$ is

- A. 0
- B. $4 \cdot {}^{1001}C_{501}$
- C. -2009
- D. none of these

Answer: A



Watch Video Solution

5. If the constant term in the binomial expansion of $\left(x^2 - \frac{1}{x}\right)^n$, $n \in N$

is 15, then the value of n is equal to.

- A. 6
- B. 9
- C. 12
- D. 15

Answer: A



Watch Video Solution

6. If $p^4 + q^3 = 2(p > 0, q > 0)$, then the maximum value of term independent of x in the expansion of $\left(px^{\frac{1}{12}} + qx^{-\frac{1}{9}}\right)^{14}$ is

A. ${}^{14}C_4$

B. ${}^{14}C_6$

C. ${}^{14}C_7$

D. ${}^{14}C_{12}$

Answer: B



Watch Video Solution

7. In the expansion of $\left(x^3 - \frac{1}{x^2}\right)^n$, $n \in N$, if the sum of the coefficients of x^5 and x^{10} , then n is a. 25 b. 20 c. 15 d. none of these

A. 25

B. 20

C. 15

D. None of these

Answer: C



Watch Video Solution

8. Find the coefficient of t^8 in the expansion of $(1 + 2t^2 - t^3)^9$.

A. 1680

B. 2140

C. 2520

D. 2730

Answer: C



Watch Video Solution

9. Find the 13^{th} term in the expansion of $\left(9x - \frac{1}{3\sqrt{x}}\right)^{18}, x \neq 0$.

A. 3

B. $\frac{1}{3}$

C. $-\frac{1}{3}$

D. 1

Answer: D



Watch Video Solution

10. In the expansion of $\left(\frac{x}{\cos \theta} + \frac{1}{x \sin \theta}\right)^{16}$, if l_1 is the least value of the term independent of x when $\frac{\pi}{8} \leq \theta \leq \frac{\pi}{4}$ and l_2 is the least value of the term independent of x when $\frac{\pi}{16} \leq \theta \leq \frac{\pi}{8}$, then the value of $\frac{l_2}{l_1}$ is

A. 8

B. 32

C. 16

D. 64

Answer: C



Watch Video Solution

11. If $A_{i,j}$ be the coefficient of $a^i b^j c^{2010-i-j}$ in the expansion of $(a + b + c)^{2010}$, then

A. $A_{i,i}$ is defined for $i \geq 1010$

B. $A_{i,j} = A_{j,i}$

C. $A_{2i,3i}$ is defined for $i \geq 405$

D. $A_{0,1} = 2000$

Answer: B



Watch Video Solution

12. The coefficient of x^{301} in the expansion of $(1+x)^{500} + x(1+x)^{499} + x^2(1+x)^{498} + \dots + x^{500}$ is
- A. ${}^{501}C_{301}$
- B. ${}^{500}C_{301}$
- C. ${}^{501}C_{300}$
- D. none of these

Answer: A



Watch Video Solution

13. The coefficient of x^{70} in the product $(x-1)(x^2-2)(x^3-3)\dots(x^{12}-12)$ is
- A. 4
- B. 6

C. 8

D. 12

Answer: A



Watch Video Solution

14. Given $(1 - x^3)^n = \sum_{k=0}^n a_k x^k (1 - x)^{3n-2k}$ then the value of $3 \cdot a_{k-1} + a_k$ is

A. ${}^n C_k \cdot 3^k$

B. ${}^{n+1} C_k \cdot 3^k$

C. ${}^{n+1} C_k \cdot 3^{k-1}$

D. ${}^n C_{k-1} \cdot 3^k$

Answer: B



Watch Video Solution

15. Find the sum of the roots (real or complex) of the equation

$$x^{2001} + \left(\frac{1}{2} - x\right)^{2001} = 0.$$

A. 2000

B. 2001

C. 1000

D. 500

Answer: D



Watch Video Solution

16. If the 4^{th} term of $\left\{ \sqrt[x^{\frac{1}{1+\log_{10}x}}]{ } + \sqrt[12]{x} \right\}^6$ is equal to 200, $x > 1$ and the

logarithm is common logarithm, then x is not divisible by

A. 2

B. 5

C. 10

D. 4

Answer: D



Watch Video Solution

17. The number of distinct terms in the expansion of $(x + y^2)^{13} + (x^2 + y)^{14}$ is

A. 27

B. 29

C. 28

D. 25

Answer: C



Watch Video Solution

18. The value of $\sum_{r=1}^n \left(\sum_{p=0}^{r-1} nC_r^r C_p 2^p \right)$ is equal to

A. $4^n - 3^n + 1$

B. $4^n - 3^n - 1$

C. $4^n - 3^n + 2$

D. $4^n - 3^n$

Answer: D



Watch Video Solution

19. If in the expansion of $\left(x^3 - \frac{2}{\sqrt{x}} \right)^n$ a term like x^2 exists and 'n' is a double digit number, then least value of 'n' is

A. 10

B. 11

C. 12

D. 13

Answer: A



Watch Video Solution

20. In $\left(33 + \frac{1}{33}\right)^n$ if the ratio of 7th term from the beginning to the 7th term from the end is $1/6$, then find the value of n .

A. 6

B. 9

C. 12

D. 15

Answer: B



Watch Video Solution

21. The number of distinct terms in the expansion of is

$$\left(x^3 + \frac{1}{x^3} + 1\right)^{200} \text{ is}$$

A. 201

B. 400

C. 401

D. 500

Answer: C



Watch Video Solution

22. If r^{th} and $(r + 1)^{th}$ term in the expansion of $(p + q)^n$ are equal, then

$$\frac{(n + 1)q}{r(p + q)} \text{ is}$$

A. $\frac{1}{2}$

B. $\frac{1}{4}$

C. 1

D. 0

Answer: C



Watch Video Solution

23. If $(3 + a\sqrt{2})^{100} + (3 + b\sqrt{2})^{100} = 7 + 5\sqrt{2}$ number of pairs (a, b) for which the equation is true is, (a, b are rational numbers)

A. 1

B. 6

C. 0

D. infinite

Answer: C



Watch Video Solution

24. The coefficient of the middle term in the expansion of $(x + 2y)^6$ is

A. $\frac{(6n)!x^n}{(3n)!(3n)!}$

B. $\frac{(6n)!x^{3n}}{(3n)!}$

C. $\frac{(6n)}{(3n)!(3n)!}(-x)^{3n}$

D. None of these

Answer: C



Watch Video Solution

25. The algebraically second largest term in the expansion of $(3 - 2x)^{15}$

at $x = \frac{4}{3}$.

A. 5

B. 7

C. 9

D. 11

Answer: B



Watch Video Solution

26. If 6^{th} term in the expansion of $\left(\frac{3}{2} + \frac{x}{3}\right)^n$ is numerically greatest, when $x = 3$, then the sum of possible integral values of ' n ' is

A. 23

B. 24

C. 25

D. 26

Answer: C



Watch Video Solution

27. Let $(5 + 2\sqrt{6})^n = p + f$, where ' n ' in N and ' p ' in N and ' 0

- A. a natural number
- B. a negative integer
- C. a prime number
- D. are irrational number

Answer: B



Watch Video Solution

28. The sum of last 3 digits of 3^{100} is

- A. 1
- B. 2
- C. 3
- D. 4

Answer: A



Watch Video Solution

29. The remainder when $27^{10} + 7^{51}$ is divided by 10

A. 4

B. 6

C. 9

D. 2

Answer: D



Watch Video Solution

30. Consider the sequence $({}^nC_0)/(1.2.3), ({}^nC_1)/(2.3.4), ({}^nC_2)/(3.4.5), \dots$, if $n=50$ then greatest term is

A. 30^{th}

B. 24^{th}

C. 26^{th}

D. 27^{th}

Answer: B



Watch Video Solution

31. If P_n denotes the product of all the coefficients of $(1 + x)^n$ and $9!P_{n+1} = 10^9 P_n$ then n is equal to

A. 10

B. 9

C. 19

D. none of these

Answer: B



Watch Video Solution

32. If N is a prime number which divides

$S = {}^{39}P_{19} + {}^{38}P_{19} + {}^{37}P_{19} + \dots + {}^{20}P_{19}$, then the largest possible value of N among following is

A. 41

B. 31

C. 37

D. 19

Answer: A



Watch Video Solution

33. If $\sum_{r=0}^n \left\{ \frac{{}^nC_{r-1}}{{}^nC_r + {}^nC_{r-1}} \right\}^3 = \frac{25}{24}$, then n is equal to

A. 3

B. 4

C. 5

D. 6

Answer: C



Watch Video Solution

34. If a, b, c, d be four consecutive coefficients in the binomial expansion of $(1 + x)^n$, then value of the expression $\left(\left(\frac{b}{b+c} \right)^2 - \frac{ac}{(a+b)(c+d)} \right)$ (where $x > 0$ and $n \in N$) is
- A. positive
 - B. negative
 - C. zero
 - D. depends on n

Answer: A



Watch Video Solution

35.

$$({}^m C_0 + {}^m C_1 - {}^m C_2 - {}^m C_3) + ({}^m C_4 + {}^m C_5 - {}^m C_6 - {}^m C_7) + \dots = 0$$

if and only if for some positive integer k , $m =$

A. $4k$

B. $4k + 1$

C. $4k - 1$

D. $4k + 2$

Answer: C



Watch Video Solution

36. The value of $\sum_{r=0}^{3n-1} (-1)^r 6nC_{2r+1} 3^r$ is

A. 2^{3n}

B. 2^{2n-1}

C. 2^{6n-1}

D. 0

Answer: D



Watch Video Solution

37. The coefficient of x^{50} in $(x + {}^{101}C_0)(x + {}^{101}C_1) \dots (x + {}^{101}C_{50})$ is

A. 4^{50}

B. 2^{50}

C. $2^{101} - 1$

D. 2^{101}

Answer: A



Watch Video Solution

38. In the expansion of $(1 + x)^{70}$, the sum of coefficients of odd powers of x is

A. 0

B. 2^{69}

C. 2^{70}

D. 2^{71}

Answer: B



Watch Video Solution

39. The sum of all the coefficients of the terms in the expansion of $(x + y + z + w)^6$ which contain x but not y , is

A. 3^6

B. 2^6

C. $3^6 - 2^6$

D. none of these

Answer: C



Watch Video Solution

40. The value of ${}^{12}C_2 + {}^{13}C_3 + {}^{14}C_4 + \dots + {}^{999}C_{989}$ is

A. ${}^{1000}C_{11} - 12$

B. ${}^{1000}C_{11} + 12$

C. ${}^{900}C_{11} - 12$

D. ${}^{1000}C_{989}$

Answer: A



Watch Video Solution

41. If $(1 + x + x^2)^{25} = a_0 + a_1x + a_2x^2 + \dots + a_{50}x^{50}$ then

$a_0 + a_2 + a_4 + \dots + a_{50}$ is :

- A. even
- B. odd and of the form $3n$
- C. odd and of the form $(3n - 1)$
- D. odd and of the form $(3n + 1)$

Answer: A



Watch Video Solution

42. If the sum of the coefficients in the expansion of $(q+r)^{20}(1+(p-2)x)^{20}$ is equal to square of the sum of the coefficients in the expansion of $[2rqx - (r+q) \cdot y]^{10}$, where p, r, q are positive constants, then

- A. $\leq P$

B. $\frac{r+q}{2} \geq p$

C. r, p and q are in $G.P.$

D. $1/r, 1/p$ and $1/q$ are in $H.P.$

Answer: B



Watch Video Solution

43. The sum $S_n = \sum_{k=0}^n (-1)^k \cdot {}^{3n}C_k$, where $n = 1, 2, \dots$ is

A. $(-1)^n \cdot {}^{3n-1}C_{n-1}$

B. $(-1)^n \cdot {}^{3n-1}C_n$

C. $(-1)n \cdot {}^{3n-1}C_{n+1}$

D. None of these

Answer: B



Watch Video Solution

44.

If

for

$$n \in I, n > 10; 1 + (1+x) + (1+x)^2 + \dots + (1+x)^n = \sum_{k=0}^n a_k \cdot x^k, x \neq 0$$

then

A. $a_{n-2} = \frac{n(n+1)}{2}$

B. $a_9^2 - a_8^2 = {}^{n+2}C_{10}({}^{n+1}C_{10} - {}^{n+1}C_9)$

C. $a_p > a_{p-1}$ for $p < \frac{n}{2}$

D. $\sum_{k=0}^n a_k = 2^{n+1}$

Answer: D



Watch Video Solution

45.

Given

$${}^8C_1 x(1-x)^7 + 2 \cdot {}^8C_2 x^2(1-x)^6 + 3 \cdot {}^8C_3 x^3(1-x)^5 + \dots + 8 \cdot x^8 = ax$$

, then $a + b$ is

A. 4

B. 6

C. 8

D. 10

Answer: C



[View Text Solution](#)

46. The value of

$$99^{50} - \frac{99}{1} \cdot 98^{50} + \frac{99.98}{1.2} 97^{50} - \dots - \frac{99.98}{1.2} \cdot 2^{50} + \frac{99}{1} \cdot 1^{50}$$
 is

A. 0

B. -1

C. -2

D. -3

Answer: A



[Watch Video Solution](#)

47. Let $f(n) = \sum_{k=1}^n k^2(nC_k)^2$ then the value of $f(5)$ equals

A. 1000

B. 1250

C. 1750

D. 2500

Answer: C



Watch Video Solution

48. The value of $\sum_{r=1}^n (-1)^{r-1} \left(\frac{r}{r+1} \right) \cdot {}^n C_r$ is

A. $\frac{1}{n+1}$

B. $\frac{1}{n}$

C. $\frac{1}{n-1}$

D. 0

Answer: A



Watch Video Solution

49. The value of

$$\binom{100}{0} \binom{200}{150} + \binom{100}{1} \binom{200}{151} + \dots + \binom{100}{50} \binom{200}{200} \quad \text{equals}$$

(where $\binom{n}{r} = {}^n C_r$)

A. $\binom{300}{50}$

B. $\binom{100}{50} \binom{200}{150}$

C. $\left(\binom{100}{50}\right)^2$

D. $\left(\binom{300}{50}\right)^2$

Answer: A



Watch Video Solution

50. Let $t_{100} = \sum_{r=0}^{100} \frac{1}{\binom{100}{r}^5}$ and $S_{100} = \sum_{r=0}^{100} \frac{r}{\binom{100}{r}^5}$, then the value of $\frac{100t_{100}}{S_{100}}$ is

A. 1

B. 2

C. 3

D. 4

Answer: B



[View Text Solution](#)

51. Let $S_1 = \sum_{0 \leq i < j \leq 100} C_i C_j$, $S_2 = \sum_{0 \leq j < i \leq 100} C_i C_j$ and $S_3 = \sum_{0 \leq i=j \leq 100} C_i C_j$ where C_r represents coefficient of x^r in the binomial expansion of $(1+x)^{100}$

If $S_1 + S_2 + S_3 = a^b$ where $a, b \in N$, then the least value of $(a+b)$ is

A. 66

B. 72

C. 46

D. 52

Answer: A



Watch Video Solution

52. ${}^{74}C_{37} - 2$ is divisible by

A. 37^2

B. 38

C. 36

D. none of these

Answer: A



Watch Video Solution

53. If ${}^nC_0 - {}^nC_1 + {}^nC_2 - {}^nC_3 + \dots + (-1)^r \cdot {}^nC_r = 28$, then n is equal to

A. 7

B. 8

C. 9

D. 11

Answer: C



Watch Video Solution

54. If the value of ${}^nC_0 + 2 \cdot {}^nC_1 + 3 \cdot {}^nC_2 + \dots + (n+1) \cdot {}^nC_n = 576$, then n is

A. 7

B. 5

C. 6

D. 9

Answer: A



Watch Video Solution

55. The value of

$$\binom{50}{6} - \binom{5}{1}\binom{40}{6} + \binom{5}{2}\binom{30}{6} - \binom{5}{3}\binom{20}{6} + \binom{5}{4}\binom{10}{6}$$

where $\binom{n}{r}$ denotes nC_r , is

A. 15625

B. 0

C. 1000000

D. 2250000

Answer: D



Watch Video Solution

56. The value of the expansion

$$\left(\sum \sum \right)_{0 \leq i < j \leq n} (-1)^{i+j-1} {}^n C_i \cdot {}^n C_j =$$

A. ${}^{2n-1} C_n$

B. ${}^{2n} C_n$

C. ${}^{2n+1} C_n$

D. None of these

Answer: A



Watch Video Solution

57. $\sum_{m=1}^n \left(\sum_{k=1}^m \left(\sum_{p=1}^m {}^n C_m \cdot {}^m C_p \cdot {}^p C_k \right) \right) =$

A. $3^n - 2^n$

B. $4^n - 3^n$

C. $3^n + 2^n$

D. $4^n - 1$

Answer: B



Watch Video Solution

58. If $\frac{(1 - 3x)^{1/2} + (1 - x)^{5/3}}{\sqrt{4 - x}}$ is approximately equal to $a + bx$ for small values of x , then $(a, b) =$

- A. $\left(1, \frac{35}{24}\right)$
- B. $\left(1, -\frac{35}{24}\right)$
- C. $\left(2, \frac{35}{12}\right)$
- D. $\left(2, -\frac{35}{12}\right)$

Answer: B



Watch Video Solution

59. The sum of the series

$$1 + \frac{1}{3^2} + \frac{1 \cdot 4}{1 \cdot 2} \frac{1}{3^4} + \frac{1 \cdot 4 \cdot 7}{1 \cdot 2 \cdot 3} \frac{1}{3^6} + \dots, \text{ is}$$

A. $\left(\frac{3}{2}\right)^{\frac{1}{3}}$

B. $\left(\frac{5}{4}\right)^{\frac{1}{3}}$

C. $\left(\frac{3}{2}\right)^{\frac{1}{6}}$

D. None of these

Answer: A



Watch Video Solution

60. Coefficient of $x^{2^{m+1}}$ in the expansion of

$$\frac{1}{(1+x)(1+x^2)(1+x^4)(1+x^8)\dots\cdot(1+x^{2^m})} \quad (|x| < 1)$$

A. 0

B. 1

C. 2^m

D. none of these

Answer: B



Watch Video Solution

61. Let $(2x^2 + 3x + 4)^{10} = \sum_{r=0}^{20} a_r x^r$, then the value of $\frac{a_7}{a_{13}}$ is

A. 6

B. 8

C. 12

D. 16

Answer: B



Watch Video Solution

62. ${}^{30}C_0 \cdot {}^{20}C_{10} + {}^{31}C_1 \cdot {}^{19}C_{10} + {}^{32}C_2 \cdot 18C_{10} + \dots {}^{40}C_{10} \cdot {}^{10}C_{10}$ is equal to

A. ${}^{51}C_{41}$

B. ${}^{50}C_{40}$

C. ${}^{51}C_{21}$

D. ${}^{50}C_{40}$

Answer: A



Watch Video Solution

Multiple Correct Answer

1. Prove that ${}^nC_0^{2n}C_n - {}^nC_1^{2n-2}C_n + {}^nC_2^{2n-4}C_n \equiv 2^n$.

A. $\binom{n}{m-n}2^{2n-m}$ if $m \geq n$

B. 0 if $m < n$

C. $\binom{n}{m-n}2^{2n+m}$ if $m \geq n$

D. 1 if $m < n$


Answer: A::B



Watch Video Solution

2. Which of the following is/are correct ?

- (I) Density of all the nuclei is same.
- (II) Heavy elements exhibit radioactivity.
- (III) Amongst alpha, beta and gamma rays, α -particle has minimum penetrating power.
- (IV) Isobars are the atoms of same elements having the different mass number.



Answer: A:D



Watch Video Solution

3. The value of $\sum_{k=0}^7 \left[\frac{\binom{7}{k}}{\binom{14}{k}} \sum_{r=k}^{14} \binom{r}{k} \binom{14}{r} \right]$, where $\binom{n}{r}$ denotes

${}^n C_r$ is

A. 6^7

B. greater than 7^6

C. 8^7

D. greater than 7^8

Answer: A::B



Watch Video Solution

4. The value of ${}^{1000}C_{50} + {}^{999}C_{49} + {}^{998}C_{48} + \dots + {}^{950}C_0$ is

A. ${}^{1001}C_{50}$

B. ${}^{1002}C_{951} - {}^{1001}C_{51}$

C. ${}^{1001}C_{951}$

D. ${}^{1002}C_{51} - {}^{1001}C_{95}$

Answer: A::B::C::D



Watch Video Solution

Comprehension

1. Consider a G. P. with first term $(1 + x)^n$, $|x| < 1$, common ratio $\frac{1+x}{2}$ and number of terms $(n + 1)$. Let ' S ' be sum of all the terms of the G. P. , then

The coefficient of x^n is ' S ' is

A. 2^n

B. 2^{n+1}

C. 2^{2n}

D. 2^{2n+1}

Answer: A



Watch Video Solution

2. Consider a G. P. with first term $(1 + x)^n$, $|x| < 1$, common ratio $\frac{1+x}{2}$ and number of terms $(n + 1)$. Let ' S ' be sum of all the terms of the G. P., then

$$\sum_{r=0}^n {}^{n+r}C_r \left(\frac{1}{2}\right)^r \text{ equals}$$

A. $(3/4)$

B. 1

C. 2^n

D. 3^n

Answer: C



Watch Video Solution

3. A path of length n is a sequence of points $(x_1, y_1), (x_2, y_2), \dots, (x_n, y_n)$ with integer coordinates such that for all i between 1 and $n - 1$ both inclusive,

either $x_{i+1} = x_i + 1$ and $y_{i+1} = y_i$ (in which case we say the i^{th} step is rightward)

or $x_{i+1} = x_i$ and $y_{i+1} = y_i + 1$ (in which case we say that the i^{th} step is upward).

This path is said to start at (x_1, y_1) and end at (x_n, y_n) . Let $P(a, b)$, for a and b non-negative integers, denotes the number of paths that start at $(0, 0)$ and end at (a, b) .

The value of $\sum_{i=0}^{10} P(i, 10 - i)$ is

A. 1024

B. 512

C. 256

D. 128

Answer: A



Watch Video Solution

4. A path of length n is a sequence of points $(x_1, y_1), (x_2, y_2), \dots, (x_n, y_n)$ with integer coordinates such that for all i between 1 and $n - 1$ both inclusive,

either $x_{i+1} = x_i + 1$ and $y_{i+1} = y_i$ (in which case we say the i^{th} step is rightward)

or $x_{i+1} = x_i$ and $y_{i+1} = y_i + 1$ (in which case we say that the i^{th} step is upward).

This path is said to start at (x_1, y_1) and end at (x_n, y_n) . Let $P(a, b)$, for a and b non-negative integers, denotes the number of paths that start at $(0, 0)$ and end at (a, b) .

Number of ordered pairs (i, j) where $i \neq j$ for which $P(i, 100 - i) = P(i, 100 - j)$ is

A. 50

B. 99

C. 100

Answer: C**Watch Video Solution**

5. A path of length n is a sequence of points $(x_1, y_1), (x_2, y_2), \dots, (x_n, y_n)$ with integer coordinates such that for all i between 1 and $n - 1$ both inclusive,

either $x_{i+1} = x_i + 1$ and $y_{i+1} = y_i$ (in which case we say the i^{th} step is rightward)

or $x_{i+1} = x_i$ and $y_{i+1} = y_i + 1$ (in which case we say that the i^{th} step is upward).

This path is said to start at (x_1, y_1) and end at (x_n, y_n) . Let $P(a, b)$, for a and b non-negative integers, denotes the number of paths that start at $(0, 0)$ and end at (a, b) .

The sum $P(43, 4) + \sum_{j=1}^5 P(49 - j, 3)$ is equal to

- A. $P(4, 48)$

B. $P(3, 49)$

C. $P(4, 47)$

D. $P(5, 47)$

Answer: A



Watch Video Solution

6.

The

expansion

$1 + x, 1 + x + x^2, 1 + x + x^2 + x^3, \dots, 1 + x + x^2 + \dots + x^{20}$ are multiplied together and the terms of the product thus obtained are arranged in increasing powers of x in the form of $a_0 + a_1x + a_2x^2 + \dots$, then,

Number of terms in the product

A. 200

B. 211

C. 231

D. none of these

Answer: B



Watch Video Solution

7.

The

expressions

$1 + x, 1 + x + x^2, 1 + x + x^2 + x^3, \dots, 1 + x + x^2 + \dots, + x^c$

are multiplied together and the terms of the product thus obtained are

arranged in increasing powers of x in the from of

$a_0 + a_1x + a_2x^2 + \dots, \text{ then sum of even coefficients?}$

A. $20!$

B. $21!$

C. $\frac{21!}{2}$

D. $19!$

Answer: C



Watch Video Solution

8.

The

expansion

$1 + x, 1 + x + x^2, 1 + x + x^2 + x^3, \dots, 1 + x + x^2 + \dots + x^{20}$ are multiplied together and the terms of the product thus obtained are arranged in increasing powers of x in the form of $a_0 + a_1x + a_2x^2 + \dots$, then,

The value of $\frac{a_r}{a_{n-r}}$, where n is the degree of the product.

A. (a)2

B. (b)1

C. (c)1/2

D. (d)depends on r

Answer: B



Watch Video Solution

9. If $(1 + px + x^2)^n = 1 + a_1x + a_2x^2 + \dots + a_{2n}x^{2n}$.

Which of the following is true for $1 < r < 2n$

- A. $(np + pr)a_r = (r + 1)a_{r+1} + (r - 1)a_{r-1}$
- B. $(np - pr)a_r = (r + 1)a_{r+1} + (r - 1 - 2n)a_{r-1}$
- C. $(np - pr)a_r = (r + 1)a_{r+1} + (r - 1 - n)a_{r-1}$
- D. $(2np + pr)a_r = (r + 1 + n)a_{r+1} + (r + 1 - n)a_{r-1}$

Answer: B



Watch Video Solution

10. If $(1 + px + x^2)^n = 1 + a_1x + a_2x^2 + \dots + a_{2n}x^{2n}$.

The remainder obtained when $a_1 + 5a_2 + 9a_3 + 13a_4 + \dots + (8n - 3)a_{2n}$ is divided by $(p + 2)$ is

A. 1

B. 2

C. 3

D. 0

Answer: C



Watch Video Solution

11. If $(1 + px + x^2)^n = 1 + a_1x + a_2x^2 + \dots + a_{2n}x^{2n}$.

The value of $a_1 + 3a_2 + 5a_3 + 7a_4 + \dots + (4n - 1)a_{2n}$ when $p = -3$ and $n \in \text{even}$ is

A. n

B. $2n - 1$

C. $2n - 2$

D. $2n$

Answer: D



Watch Video Solution

