



## MATHS

### BOOKS - CENGAGE

### Differential Equations

#### Solved Examples And Exercises

1. If  $y = \frac{1}{\sqrt{a^2 - b^2}} \cos^{-1} \left( \frac{a \cos x + b}{a + b \cos x} \right)$ , then  $\frac{d^2 y}{dx^2} =$  (i)

$$\frac{b \sin x}{(a + b \cos x)^2} \quad \text{(ii)} \quad - \frac{b \sin x}{(a + b \cos x)^2} \quad \text{(iii)} \quad \frac{b \cos x}{(a + b \cos x)^2} \quad \text{(iii)}$$
$$- \frac{b \cos x}{(a + b \cos x)^2}$$



Watch Video Solution

2. Find the differential equation of all non-vertical lines in a plane.



Watch Video Solution

3. Form the differential equation of all circles touching the x-axis at the origin and center on y-axis.

 [Watch Video Solution](#)

4. Form the differential equation of the family of parabolas with focus at the origin and the axis of symmetry along the axis.

 [Watch Video Solution](#)

5. Find the differential equation of all parabolas whose axis are parallel to the x-axis.

 [Watch Video Solution](#)

6. A body at a temperature of  $50^{\circ}F$  is placed outdoors where the temperature is  $100^{\circ}F$ . If the rate of change of the temperature of a body is proportional to the temperature difference between the body and its surrounding medium. If after 5 min the temperature of the body is  $60^{\circ}F$ , find (a) how long it will take the body to reach a temperature of  $75^{\circ}F$  and (b) the temperature of the body after 20 min.

 [Watch Video Solution](#)

7. Solve  $\frac{dy}{dx} = \frac{2x - y + 1}{x + 2y - 3}$

 [Watch Video Solution](#)

8. Find the degree of the differential equation satisfying the relation

$$\sqrt{1+x^2} + \sqrt{1+y^2} = \lambda \left( x\sqrt{1+y^2} - y\sqrt{1+x^2} \right)$$

 [Watch Video Solution](#)

9. Find the particular solution of the differential equation

$$(1 + e^{2x})dy + (1 + y^2)e^x dx = 0. \text{ Given that } y = 1 \text{ when } x = 0.$$

 [Watch Video Solution](#)

10. Find the time required for a cylindrical tank of radius 2.5 m and height 3 m to empty through a round hole of 2.5 cm with a velocity  $2.5\sqrt{h}$  m/s,  $h$  being the depth of the water in the tank.

 [Watch Video Solution](#)

11. Solve:  $(dy/dx) + (y/x) = y^3$

 [Watch Video Solution](#)

12. Solve the differential equation  $xy \frac{dy}{dx} = \frac{1 + y^2}{1 + x^2} (1 + x + x^2)$

 [Watch Video Solution](#)

13. Solve the equation  $\frac{dy}{dx} + x \sin 2y = x^3 \cos^2 y$

 [Watch Video Solution](#)

14. Solve the equation  $\frac{dy}{dx} + \frac{xy}{(1-x^2)} = x\sqrt{y}$

 [Watch Video Solution](#)

15. Solve the equation  $\frac{dy}{dx} + (2x \tan^{-1} y - x^3)(1 + y^2) = 0$

 [Watch Video Solution](#)

16. Find the orthogonal trajectory of  $y^2 = 4ax$  (a being the parameter).

 [Watch Video Solution](#)

17.  $\left[ x \sin^2 \left( \frac{y}{x} \right) - y \right] dx + x dy = 0, y = \frac{\pi}{4}$  when  $x = 1$



Watch Video Solution

18. Find the orthogonal trajectories of family of curves  $x^2 + y^2 = cx$



Watch Video Solution

19. Show that the differential equation  $y^3 dy + (x + y^2) dx = 0$  can be reduced to a homogeneous equation.



Watch Video Solution

20. Solve  $\left[2\sqrt{xy} - x\right] dy + y dx = 0$



Watch Video Solution

21. Find the equation of the curve which is such that the area of the rectangle constructed on the abscissa of any point and the intercept of

the tangent at this point on the y-axis is equal to 4.

 [Watch Video Solution](#)

22. Solve  $x \left( \frac{dy}{dx} \right) = y(\log y - \log x + 1)$

 [Watch Video Solution](#)

23. If the population of country double in 50 years, in how many years will it triple under the assumption that the rate of increase is proportional to the number of inhabitants.

 [Watch Video Solution](#)

24. Find the equation of the curve passing through the origin if the middle point of the segment of its normal from any point of the curve to the x-axis lies on the parabola  $2y^2 = x$

 [Watch Video Solution](#)

25. Find the real value of  $m$  for which the substitution  $y = u^m$  will transform the differential equation  $2x^4y \frac{dy}{dx} + y^4 = 4x^6$  into a homogenous equation.

 [Watch Video Solution](#)

26. A curve  $y = f(x)$  passes through the origin. Through any point  $(x, y)$  on the curve, lines are drawn parallel to the co-ordinate axes. If the curve divides the area formed by these lines and co-ordinates axes in the ratio  $m:n$ , find the curve.

 [Watch Video Solution](#)

27. Find the equation of a curve passing through  $(0, 1)$  and having gradient  $\frac{-(y + y^3)}{1 + x + xy}$  at  $(x, y)$

 [Watch Video Solution](#)



28. Solve:  $\frac{dy}{dx} = \frac{x + 2y + 5}{2x + y - 1}$

 [Watch Video Solution](#)

29. Solve  $xdy = \left( y + x \frac{f\left(\frac{y}{x}\right)}{f'\left(\frac{y}{x}\right)} \right) dx$

 [Watch Video Solution](#)

30. Solve the equation  $ydx + (x - y^2)dy = 0$

 [Watch Video Solution](#)

31. Find the orthogonal trajectory of  $y^2 = 4ax$  (a being the parameter).

 [Watch Video Solution](#)

32. Solve  $e^{\frac{dy}{dx}} = x + 1$ , given that when  $x = 0, y = 3$ .

 [Watch Video Solution](#)

33. Solve  $\log\left(\frac{dy}{dx}\right) = 4x - 2y - 2$ , given that  $y = 1$  when  $x = 1$ .

 [Watch Video Solution](#)

34. Form the differential equation of all concentric circles at the origin.

 [Watch Video Solution](#)

35. Solve  $\frac{d^2y}{dx^2} = \left(\frac{dy}{dx}\right)^2$

 [Watch Video Solution](#)

36. Solve  $\frac{dy}{dx} \sqrt{1 + x + y} = x + y - 1$

 [Watch Video Solution](#)

37. The solution of  $\sec^2 x \tan y dx + \sec^2 y \tan x dy = 0$  is

 [Watch Video Solution](#)

38. Solve the differential equation  $ye^{\frac{x}{y}} dx = \left(xe^{\frac{x}{y}} + y^2\right) dy (y \neq 0)$ .

 [Watch Video Solution](#)

39. Find the order and degree (if defined) of the equation:

$$\frac{d^3 y}{dx^3} = x \ln \left( \frac{dy}{dx} \right)$$

 [Watch Video Solution](#)

40. Find the order and degree (if defined) of the equation:

$$\frac{d^2 y}{dx^2} = \left\{ 1 + \left( \frac{dy}{dx} \right)^4 \right\}^{\frac{5}{3}}$$



Watch Video Solution

41. What constant interest rate is required if an initial deposit placed into an account accrues interest compounded continuously is to double its value in six years? ( $\ln|x| = 0.6930$ )



Watch Video Solution

42. For each of the differential equations given below, indicate its order and degree ( if defined ).

$$\frac{d^4y}{dx^4} - \sin\left(\frac{d^3y}{dx^3}\right) = 0$$



Watch Video Solution

43. Find the order and degree (if defined) of the equation:

$$a = \frac{1 \left[ 1 + \left( \frac{dy}{dx} \right)^2 \right]^{\frac{3}{2}}}{\frac{d^2y}{dx^2}}, \text{ where } a \text{ is constant}$$



Watch Video Solution

44. Determine its order, degree (if exists)

$$\left(\frac{d^3y}{dx^3}\right)^{\frac{2}{3}} - 3\frac{d^2y}{dx^2} + 5\frac{dy}{dx} + 4 = 0$$



Watch Video Solution

45. Find the order and degree (if defined) of the equation:

$$\frac{d^4y}{dx^4} + 3\left(\frac{d^2y}{dx^2}\right)^6 + \sin x = 2 \cos x$$



Watch Video Solution

46. Solve  $(1 - x^2) \frac{dy}{dx} + 2xy = x \sqrt{1 - x^2}$



Watch Video Solution

47. Solve the equation  $(x + y + 1) \left(\frac{dy}{dx}\right) = 1$



Watch Video Solution

48. Solve the equation  $\frac{dy}{dx} + y \cot x = \sin x$



Watch Video Solution

49. Find the  $\sum_{0 \leq i < j \leq n} \sum 1$ .



Watch Video Solution

50. Solve the differential equation:  $(1 + y^2) + (x - e^{\tan^{-1} y}) \frac{dy}{dx} = 0$



Watch Video Solution

51. Solve:

$$ydx - xdy + \log x dx = 0$$



Watch Video Solution

52. Solve the differential equation  $\left[ \frac{e^{-(2\sqrt{x})}}{\sqrt{x}} - \frac{y}{\sqrt{x}} \right] \frac{dx}{dy} = 1 (x \neq 0)$ .

 [Watch Video Solution](#)

53. Find the order and degree of the following differential equation:

$$\sin^{-1} \left( \frac{dy}{dx} \right) = x + y$$

 [Watch Video Solution](#)

54. Find the order and degree of the following differential equation:

$$e^{\frac{d^3y}{dx^3}} - x \frac{d^2y}{dx^2} + y = 0$$

 [Watch Video Solution](#)

55. Find the order and degree of the following differential equation:

$$\frac{dy}{dx} + y = \frac{1}{\frac{dy}{dx}}$$

 [Watch Video Solution](#)

56. Find the curve for which the perpendicular from the foot of the ordinate to the tangent is of constant length.

 [Watch Video Solution](#)

57. Find the order and degree of the following differential equation:

$$\frac{d^2y}{dx^2} = \left[ y + \left( \frac{dy}{dx} \right)^6 \right]^{\frac{1}{4}}$$

 [Watch Video Solution](#)



58. Find the order and degree of the following differential equation:

$$\ln\left(\frac{dy}{dx}\right) = ax + by$$

 [Watch Video Solution](#)

59. Solve  $(xy^2 - e^{\frac{1}{x^3}}) dx - x^2ydy = 0$

 [Watch Video Solution](#)

60. The rate at which a substance cools in moving air is proportional to the difference between the temperatures of the substance and that of the air. If the temperature of the air is 290 K and the substance cools from 370 K to 330 K in 10 min, when will the temperature be 295 K?

 [Watch Video Solution](#)

61. Find the equation of the curve in which the subnormal varies as the square of the ordinate.

 [Watch Video Solution](#)

62. Solve  $\frac{dy}{dx} = \frac{\sin y + x}{\sin 2y - x \cos y}$

 [Watch Video Solution](#)

63. Solve:  $(x - 1)dy + ydx = x(x - 1)y^{1/3}dx$ .

 [Watch Video Solution](#)

64. A normal is drawn at a point  $P(x, y)$  of a curve. It meets the x-axis and the y-axis in point  $A$  AND  $B$ , respectively, such that  $\frac{1}{OA} + \frac{1}{OB} = 1$ , where  $O$  is the origin. Find the equation of such a curve passing through  $(5, 4)$



[Watch Video Solution](#)

65. Find the curve for which the length of normal is equal to the radius vector.



[Watch Video Solution](#)

66. The integrating factor of the differential equation

$$(1 - y^2) \frac{dy}{dx} + yx = ay \quad (-1 < y' < 1) \text{ is}$$



[Watch Video Solution](#)

67. The solution of the differential equation

$$\left\{1 + x\sqrt{(x^2 + y^2)}\right\}dx + \left\{\sqrt{(x^2 + y^2)} - 1\right\}ydy = 0 \text{ is equal to}$$



[Watch Video Solution](#)

68. The solution of the differential equation  $\frac{dy}{dx} = \frac{3x^2y^4 + 2xy}{x^2 - 2x^3y^3}$  is

 [Watch Video Solution](#)

69. Solution of the differential equation

$$\left\{ \frac{1}{x} - \frac{y^2}{(x-y)^2} \right\} dx + \left\{ \frac{x^2}{(x-y)^2} - \frac{1}{y} \right\} dy = 0 \text{ is}$$

 [Watch Video Solution](#)

70. The solution of the differential equation

$$2x^2y \frac{dy}{dx} = \tan(x^2y^2) - 2xy^2, \text{ given } x = 1, y = \frac{\pi}{2}, \text{ is}$$

 [Watch Video Solution](#)

71. The solution of the differential equation

$$x^2 \frac{dy}{dx} \cos\left(\frac{1}{x}\right) - y \sin\left(\frac{1}{x}\right) = -1, \text{ where } y \rightarrow -1 \text{ as } x \rightarrow \infty \text{ is}$$

 [Watch Video Solution](#)

Watch Video Solution

72. The solution of the differential equation  $(x + 2y^3) \frac{dy}{dx} = y$  is

 Watch Video Solution

73. Find the curve satisfying the equation  $\frac{dy}{dx} = \frac{y(x + y^3)}{x(y^3 - x)}$  and passing through the point  $(4, -2)$  is

 Watch Video Solution

74. Find the solution of  $ye^{-\frac{x}{y}} dx - \left( xe^{-\frac{x}{y}} + y^3 \right) dy = 0$  is

 Watch Video Solution

75. The solution of differential equation

$(2y + xy^3) dx + (x + x^2y^2) dy = 0$  is

 Watch Video Solution

76. if  $y + x \frac{dy}{dx} = x \frac{\phi(xy)}{\phi'(xy)}$  then  $\phi(xy)$  is

 Watch Video Solution

77. Let  $y'(x) + y(x)g'(x) = g(x)g'(x)$ ,  $y(0)$ ,  $x \in R$ , where  $f'(x)$  denotes  $\frac{dy(x)}{dx}$ , and  $g(x)$  is a given non-constant differentiable function on  $R$  with  $g(0) = g(2) = 0$ . Then the value of  $y(2)$  is \_\_\_\_\_

 Watch Video Solution

78. Let  $f: [1, \infty]$  be a differentiable function such that  $f(1) = 2$ . If  $6 \int_1^x f(t) dt = 3x f(x) - x^3$  for all  $x \geq 1$ , then the value of  $f(2)$  is

 Watch Video Solution

79. If  $y(x)$  satisfies the differential equation  $y' - y \tan x = 2x \sec x$  and  $y(0) = 0$ , then

 [Watch Video Solution](#)

80. Consider the family of all circles whose centers lie on the straight line  $y = x$ . If this family of circles is represented by the differential equation

$P y'' + Q y' + 1 = 0$ , where  $P, Q$  are functions of  $x, y$  and  $y'$  (here  $y' = \frac{dy}{dx}, y'' = \frac{d^2y}{dx^2}$ ), then which of the following statements is

(are) true? (a)  $P = y + x$  (b)  $P = y - x$  (c)

$P + Q = 1 - x + y + y' + (y')^2$  (d)  $P - Q = x + y - y' - (y')^2$

 [Watch Video Solution](#)

81. Let  $y(x)$  be a solution of the differential equation  $(1 + e^x)y' + ye^x = 1$ . If  $y(0) = 2$ , then which of the following statements is (are) true? (a)  $y(-4) = 0$  (b)  $y(-2) = 0$  (c)  $y(x)$  has a

critical point in the interval  $(-1, 0)$  (d)  $y(x)$  has no critical point in the interval  $(-1, 0)$



[Watch Video Solution](#)

**82.** Let  $f$  be a real-valued differentiable function on  $R$  (the set of all real numbers) such that  $f(1) = 1$ . If the  $y$ -intercept of the tangent at any point  $P(x, y)$  on the curve  $y = f(x)$  is equal to the cube of the abscissa of  $P$ , then the value of  $f(-3)$  is equal to \_\_\_\_\_



[Watch Video Solution](#)

**83.** The function  $y = f(x)$  is the solution of the differential equation

$$\frac{dy}{dx} + \frac{xy}{x^2 - 1} = \frac{x^4 + 2x}{\sqrt{1 - x^2}} \quad \text{in } (-1, 1) \quad \text{satisfying } f(0) = 0. \quad \text{Then}$$

$$\int_{\frac{\sqrt{3}}{2}}^{\frac{\sqrt{3}}{2}} f(x) dx \text{ is}$$



[Watch Video Solution](#)



84. The order of the differential equation whose general solution is given by  $y = (C_1 + C_2)\cos(x + C_3) - C_4e^{x+C_5}$ , where  $C_1, C_2, C_3, C_4, C_5$ , are arbitrary constants, is (a) 5 (b) 4 (c) 3 (d) 2



Watch Video Solution

85. The differential equation representing the family of curves  $y^2 = 2c(x + \sqrt{c})$ , where  $c$  is a positive parameter, is of (A) order 1 (B) order 2 (C) degree 3 (D) degree 4



Watch Video Solution

86. Find the equation of a curve passing through  $\left(2, \frac{7}{2}\right)$  and having gradient  $1 - \frac{1}{x^2}$  at  $(x, y)$  is



Watch Video Solution

87. The equation of the curve which is such that the portion of the axis of  $x$  cut off between the origin and tangent at any point is proportional to the ordinate of that point is

 [Watch Video Solution](#)

88. A normal at  $P(x, y)$  on a curve meets the  $x$ -axis at  $Q$  and  $N$  is the foot of the ordinate at  $P$ . If  $NQ = \frac{x(1 + y^2)}{1 + x^2}$ , then the equation of curve given that it passes through the point  $(3, 1)$  is

 [Watch Video Solution](#)

89. The differential equation of the family of all non-vertical lines in a plane is

 [Watch Video Solution](#)

90. The curve in the first quadrant for which the normal at any point  $(x, y)$  and the line joining the origin to that point form an isosceles triangle with the x-axis as base is (a) an ellipse (b) a rectangular hyperbola (c) a circle (d) None of these

 [Watch Video Solution](#)

91. The differential equation of the curve for which the initial ordinate of any tangent is equal to the corresponding subnormal (a) is linear (b) is homogeneous of second degree (c) has separable variables (d) is of second order

 [Watch Video Solution](#)

92. Orthogonal trajectories of family of the curve  $x^{\frac{2}{3}} + y^{\frac{2}{3}} = a \left(\frac{2}{3}\right)$ , where  $a$  is any arbitrary constant, is

 [Watch Video Solution](#)

93. Which of the following is not the differential equation of family of curves whose tangent form an angle of  $\frac{\pi}{4}$  with the hyperbola  $xy = c^2$ ?

(a)  $\frac{dy}{dx} = \frac{x - y}{x + y}$  (b)  $\frac{dy}{dx} = \frac{x}{x - y}$  (c)  $\frac{dy}{dx} = \frac{x + y}{x - y}$  (d) N.O.T.



Watch Video Solution

94. Tangent to a curve intercepts the y-axis at a point  $P$ . A line perpendicular to this tangent through  $P$  passes through another point  $(1, 0)$ . The differential equation of the curve is



Watch Video Solution

95. The solution of the differential equation  $(x \cot y + \log \cos x)dy + (\log \sin y - y \tan x)dx = 0$  is



Watch Video Solution

96. The curve with the property that the projection of the ordinate on the normal is constant and has a length equal to  $a$  is

 [Watch Video Solution](#)

97. The differential equation of all parabolas each of which has a latus rectum  $4a$  and whose axes are parallel to the  $x$ -axis is (a) of order 1 and degree 2 (b) of order 2 and degree 3 (c) of order 2 and degree 1 (d) of order 2 and degree 2

 [Watch Video Solution](#)

98. The solution of differential equation  $\frac{y(2x^4 + y)dy}{dx} = (1 - 4xy^2)x^2$  is given by

 [Watch Video Solution](#)

99. The normal to a curve at  $P(x, y)$  meet the x-axis at  $G$ . If the distance of  $G$  from the origin is twice the abscissa of  $P$ , then the curve is a (a) parabola (b) circle (c) hyperbola (d) ellipse



Watch Video Solution

100. A normal at any point  $(x, y)$  to the curve  $y = f(x)$  cuts a triangle of unit area with the axis, the differential equation of the curve is



Watch Video Solution

101. If  $a, b$  are two fixed positive integers such that  $f(a + x) = b + \left[ b^3 + 1 - 3b^2 f(x) + 3b\{f(x)\}^2 - \{f(x)\}^3 \right]^{\frac{1}{3}}$  for all real  $x$ , then prove that  $f(x)$  is periodic and find its period.



Watch Video Solution

**102.** The differential equation whose general solution is given by  $y = c_1 \cos(x + c_2) - c_3 e^{(-x + c_4)} + (c_5 \sin x)$ , where  $c_1, c_2, c_3, c_4, c_5$  are arbitrary constants, is

 [Watch Video Solution](#)

**103.** The solution to the differential equation  $y \log y + xy' = 0$ , where  $y(1) = e$ , is

 [Watch Video Solution](#)

**104.** The form of the differential equation of the central conics  $ax^2 + by^2 = 1$  is

 [Watch Video Solution](#)

**105.** The differential equation for the family of curve  $x^2 + y^2 - 2ay = 0$ , where  $a$  is an arbitrary constant, is

 [Watch Video Solution](#)

**106.** If  $\frac{dy}{dx} = (e^y - x)^{-1}$ , where  $y(0) = 0$ , then  $y$  is expressed explicitly as

 [Watch Video Solution](#)

**107.** If  $y = \frac{x}{\log|cx|}$  (where  $c$  is an arbitrary constant) is the general solution of the differential equation  $\frac{dy}{dx} = \frac{y}{x} + \varphi\left(\frac{x}{y}\right)$ , then the function  $\varphi\left(\frac{x}{y}\right)$  is

 [Watch Video Solution](#)



**108.** Differential equation of the family of circles touching the line  $y = 2$  at  $(0, 2)$  is

 [Watch Video Solution](#)

**109.** The differential equation of all parabolas whose axis are parallel to the  $y$ -axis is

 [Watch Video Solution](#)

**110.** A differential equation associated to the primitive  $y = a + be^{5x} + ce^{-7x}$  is (where  $y_n$  is  $n$ th derivative w.r.t.  $x$ )

(a)  $y_3 + 2y_2 - y_1 = 0$  (b)  $4y_3 + 5y_2 - 20y_1 = 0$  (c)  $y_3 + 2y_2 - 35y_1 = 0$

(d) none of these

 [Watch Video Solution](#)

111. The order and degree of the differential equation of all tangent lines to the parabola  $y = x^2$  is (a) 1,2 (b) 2,3 (c) 2,1 (d) 1,1

 [Watch Video Solution](#)

112. A solution of the differential equation,  $\left(\frac{dy}{dx}\right)^2 - x\frac{dy}{dx} + y = 0$

 [Watch Video Solution](#)

113. A curve  $y = f(x)$  passes through point  $P(1, 1)$ . The normal to the curve at  $P$  is a  $(y - 1) + (x - 1) = 0$ . If the slope of the tangent at any point on the curve is proportional to the ordinate of the point, then the equation of the curve is

 [Watch Video Solution](#)

114. A spherical rain drop evaporates at a rate proportional to its surface area at any instant  $t$ . The differential equation giving the rate of change of the radius of the rain drop is \_\_\_\_\_



[Watch Video Solution](#)

115. If length of tangent at any point on the curve  $y = f(x)$ . Intercepted between the point and the x-axis is of length 1. Find the equation of the curve.



[Watch Video Solution](#)

116. For the primitive integral equation  $ydx + y^2dy = xdy; x \in R, y > 0, y(1) = 1$ , then  $y(-3)$  is (a) 3 (b) 2 (c) 1 (d) 5



[Watch Video Solution](#)

117. The solution of the primitive integral equation  $(x^2 + y^2)dy = xydx$  is  $y = y(x)$ . If  $y(1) = 1$  and  $y(x_0) = e$ , then  $x_0$  is

 [Watch Video Solution](#)

118. if  $y = y(x)$  and  $\frac{2 + \sin x}{y + 1} \left( \frac{dy}{dx} \right) = -\cos x$ ,  $y(0) = 1$ , then  $y\left(\frac{\pi}{2}\right) =$

 [Watch Video Solution](#)

119. If  $y(t)$  is a solution of  $(1 + t) \frac{dy}{dt} - ty = 1$  and  $y(0) = -1$  then  $y(1)$  is

 [Watch Video Solution](#)

120. A curve passes through the point  $\left(1, \frac{\pi}{6}\right)$ . Let the slope of the curve at each point  $(x, y)$  be  $\frac{y}{x} + \sec\left(\frac{y}{x}\right)$ ,  $x > 0$ . Then the equation of the curve is



[Watch Video Solution](#)

121. The differential equation  $\frac{dy}{dx} = \frac{\sqrt{1-y^2}}{y}$  determines a family of circle with (a) variable radii and a fixed centre at  $(0, 1)$  (b) variable radii and a fixed centre at  $(0, -1)$  (c) Fixed radius 1 and variable centres along the x-axis. (d) Fixed radius 1 and variable centres along the y-axis.



[Watch Video Solution](#)

122. A spherical rain drop evaporates at a rate proportional to its surface area at any instant  $t$ . The differential equation giving the rate of change of the radius of the rain drop is \_\_\_\_\_



[Watch Video Solution](#)

123. Water is drained from a vertical cylindrical tank by opening a valve at the base of the tank. It is known that the rate at which the water level

drops is proportional to the square root of water depth  $y$ , where the constant of proportionality  $k > 0$  depends on the acceleration due to gravity and the geometry of the hole. If  $t$  is measured in minutes and  $k = \frac{1}{15}$ , then the time to drain the tank if the water is 4 m deep to start with is (a) 30 min (b) 45 min (c) 60 min (d) 80 min

 [Watch Video Solution](#)

**124.** The population of a country increases at a rate proportional to the number of inhabitants.  $f$  is the population which doubles in 30 years, then the population will triple in approximately. (a) 30 years (b) 45 years (c) 48 years (d) 54 years

 [Watch Video Solution](#)

**125.** An object falling from rest in air is subject not only to the gravitational force but also to air resistance. Assume that the air resistance is proportional to the velocity with constant of proportionality

as  $k > 0$ , and acts in a direction opposite to motion  $\left(g = 9.8 \frac{m}{s^2}\right)$ .

Then velocity cannot exceed.

 [Watch Video Solution](#)

**126.** Find the solution of differential equation

$$x^2 = 1 + \left(\frac{x}{y}\right)^{-1} \frac{dy}{dx} + \frac{\left(\frac{x}{y}\right)^{-2} \left(\frac{dy}{dx}\right)^2}{2!} + \dots$$

 [Watch Video Solution](#)

**127.** The solution of the differential equation  $y' y'''' = 3(y'')^2$  is

 [Watch Video Solution](#)

**128.** Number of values of  $m \in N$  for which  $y = e^{mx}$  is a solution of the

differential equation  $\frac{d^3y}{dx^3} - 3\frac{d^2y}{dx^2} - 4\frac{dy}{dx} + 12y = 0$  (a) 0 (b) 1 (c) 2 (d)

More than 2



 [Watch Video Solution](#)

129. A curve passing through  $(2, 3)$  and satisfying the differential equation  $\int_0^x ty(t)dt = x^2y(x)$ ,  $(x > 0)$  is

 [Watch Video Solution](#)

130. The solution of the differential equation  $\frac{d^2y}{dx^2} = \sin 3x + e^x + x^2$  when  $y_1(0) = 1$  and  $y(0) = 0$  is

 [Watch Video Solution](#)

131. The solution of the differential equation

$$\frac{x + \frac{x^3}{3!} + \frac{x^5}{5!} + \dots}{1 + \frac{x^2}{2!} + \frac{x^4}{4!} + \dots} = \frac{dx - dy}{dx + dy} \text{ is}$$

 [Watch Video Solution](#)



132.  $A$  and  $B$  are two separate reservoirs of water. Capacity of reservoir  $A$  is double the capacity of reservoir  $B$ . Both the reservoirs are filled completely with water, their inlets are closed and then the water is released simultaneously from both the reservoirs. The rate of flow of water out of each reservoir at any instant of time is proportional to the quantity of water in the reservoir at the time. One hour after the water is released, the quantity of water in reservoir  $A$  is  $1\frac{1}{2}$  times the quantity of water in reservoir  $B$ . After how many hours do both the reservoirs have the same quantity of water?

 [Watch Video Solution](#)

133. If  $\int_a^x ty(t) dt = x^2 + y(x)$ , then find  $y(x)$

 [Watch Video Solution](#)

134. Given a function '  $g$  ' which has a derivative  $g'(x)$  for every real  $x$  and satisfies  $g'(0) = 2$  and  $g(x + y) = e^y g(x) + e^x g(y)$  for all  $x$  and  $y$  then:



Watch Video Solution

135. Solve  $y \left( \frac{dy}{dx} \right)^2 + 2x \frac{dy}{dx} - y = 0$  given that  $y(0) = \sqrt{5}$



Watch Video Solution

136. If  $y + \frac{d}{dx}(xy) = x(\sin x + \log x)$ ,  $f \in dy(x)$ .



Watch Video Solution

137. Show that the given differential equation is homogeneous and solve each of them.

$$\left\{ x \cos\left(\frac{y}{x}\right) + y \sin\left(\frac{y}{x}\right) \right\} y dx = \left\{ y \sin\left(\frac{y}{x}\right) - x \cos\left(\frac{y}{x}\right) \right\} x dy$$



Watch Video Solution

138. Solve  $\frac{dy}{dx} = \frac{(x+y)^2}{(x+2)(y-2)}$



Watch Video Solution

139. Solve  $\frac{x + y \frac{dy}{dx}}{y - x \frac{dy}{dx}} = x^2 + 2y^2 + \frac{y^4}{x^2}$



Watch Video Solution

140. For the differential equation whose solution is  $(x - h)^2 + (y - k)^2 = a^2$  ( $a$  is a constant), its (a) order is 2 (b) order is 3 (c) degree is 2 (d) degree is 3



Watch Video Solution

141. Which one of the following function(s) is/are homogeneous?

(a)  $f(x, y) = \frac{x - y}{x^2 + y^2}$  (b)  $f(x, y) = x^{\frac{1}{3}} y^{-\frac{2}{3}} \tan^{-1} \left( \frac{x}{y} \right)$  (c)  $f(x, y) = x \left( \ln \frac{x}{y} \right)$

(d) none of these



Watch Video Solution

**142.** The solution of the differential equation

$$(e^{x^2} + e^{y^2})y \frac{dy}{dx} + e^{x^2}(xy^2 - x) = 0 \text{ is}$$

 [Watch Video Solution](#)

**143.** The solution of the differential equation  $y'''' - 8y'' = 0$ , where

$$y(0) = \frac{1}{8}, y'(0) = 0, y''(0) = 1, \text{ is}$$

 [Watch Video Solution](#)

**144.** Differential equation of the family of curves  $v = \frac{A}{r} + B$ , where  $A$  and  $B$  are arbitrary constants, is

 [Watch Video Solution](#)

**145.** The function  $f(\theta) = \frac{d}{d(\theta)} \int_0^\theta \frac{dx}{1 - \cos \theta \cos x}$  satisfies the differential equation



Watch Video Solution

146. The differential equation of the curve  $\frac{x}{c-1} + \frac{y}{c+1} = 1$  is



Watch Video Solution

147. The solution of the differential equation

$$x = 1 + xy \frac{dy}{dx} + \frac{x^2 y^2}{2!} \left( \frac{dy}{dx} \right)^2 + \frac{x^3 y^3}{3!} \left( \frac{dy}{dx} \right)^3 + \dots \text{is}$$



Watch Video Solution

148. Which of the following equation(s) is/are linear?

(a)  $\frac{dy}{dx} + \frac{y}{x} = \ln x$  (b)  $y \left( \frac{dy}{dx} \right) + 4x = 0$  (c)  $(2x + y^3) \left( \frac{dy}{dx} \right) = 3y$  (d)  $N. O$



Watch Video Solution

**149.** The equation of the curve satisfying the differential equation

$$y\left(\frac{dy}{dx}\right)^2 + (x - y)\frac{dy}{dx} - x = 0 \text{ can be a (a) circle (b) Straight line (c)}$$

Parabola (d) Ellipse



**Watch Video Solution**

**150.** Find a pair of curves such that (a) the tangents drawn at points with equal abscissas intersect on the y-axis. (b) the normal drawn at points with equal abscissas intersect on the x-axis. (c) one curve passes through (1,1) and other passes through (2, 3).



**Watch Video Solution**

**151.** If  $y_1$  and  $y_2$  are two solutions to the differential equation

$$\frac{dy}{dx} + P(x)y = Q(x) . \text{ Then prove that } y = y_1 + c(y_1 - y_2) \text{ is the}$$

general solution to the equation where  $c$  is any constant.



**Watch Video Solution**

152. If  $y_1$  and  $y_2$  are the solution of the differential equation  $\frac{dy}{dx} + Py = Q$ , where  $P$  and  $Q$  are functions of  $x$  alone and  $y_2 = y_1 z$ , then prove that  $z = 1 + c \cdot e^{-\int \frac{Q}{y_1} dx}$ , where  $c$  is an arbitrary constant.

 [Watch Video Solution](#)

153. Let the function  $\ln f(x)$  is defined where  $f(x)$  exists for  $x \geq 2$  and  $k$  is fixed positive real numbers prove that if  $\frac{d}{dx}(x \cdot f(x)) \geq -kf(x)$  then  $f(x) \geq Ax^{-1-k}$  where  $A$  is independent of  $x$ .

 [Watch Video Solution](#)

154. The degree of the differential equation satisfying  $\sqrt{1-x^2} + \sqrt{1-y^2} = a(x-y)$  is (a) 1 (b) 2 (c) 3 (d) none of these

 [Watch Video Solution](#)

**155.** The force of resistance encountered by water on a motor boat of mass  $m$  going in still water with velocity  $v$  is proportional to the velocity  $v$ . At  $t = 0$  when its velocity is  $v_0$ , then engine shuts off. Find an expression for the position of motor boat at time  $t$  and also the distance travelled by the boat before it comes to rest. Take the proportionality constant as  $k > 0$ .



[Watch Video Solution](#)

**156.** A cyclist moving on a level road at 4 m/s stops pedalling and lets the wheels come to rest. The retardation of the cycle has two components: a constant  $0.08 \text{ m/s}^2$  due to friction in the working parts and a resistance of  $0.02v^2/m$ , where  $v$  is speed in meters per second. What distance is traversed by the cycle before it comes to rest? (consider  $\ln 5=1.61$ ).



[Watch Video Solution](#)



157. Given two curves:  $y = f(x)$  passing through the point  $(0, 1)$  and  $g(x) = \int_{-\infty}^x f(t)dt$  passing through the point  $\left(0, \frac{1}{n}\right)$ . The tangents drawn to both the curves at the points with equal abscissas intersect on the x-axis. Find the curve  $y = f(x)$ .

 [Watch Video Solution](#)

158. The differential equation of the family of curves  $y = e^x(A \cos x + B \sin x)$ , where  $A$  and  $B$  are arbitrary constants is

 [Watch Video Solution](#)

159. The differential equation whose solution is  $Ax^2 + By^2 = 1$ , where  $A$  and  $B$  are arbitrary constants, is of (a) second order and second degree (b) first order and second degree (c) first order and first degree (d) second order and first degree

 [Watch Video Solution](#)

160. Solve:  $(x-4)(x-7)(x-2)(x+1)=16$

 [Watch Video Solution](#)

161. The solution of the equation  $\frac{dy}{dx} = \frac{x(2 \log x + 1)}{\sin y + y \cos y}$  is

 [Watch Video Solution](#)

162. The solution of  $\log \left( \frac{dy}{dx} \right) = ax + by$  is \_\_\_\_\_.

 [Watch Video Solution](#)

163. if  $y = y(x)$  and  $\frac{2 + \sin x}{y + 1} \left( \frac{dy}{dx} \right) = -\cos x$ ,  $y(0) = 1$ , then  $y\left(\frac{\pi}{2}\right) =$  (a)  $\frac{1}{e}$  (b)  $\frac{2}{m}$  (c)  $\frac{1}{u}$  (d) 1

 [Watch Video Solution](#)

164. The equation of the curves through the point (1, 0) and whose slope is  $\frac{y - 1}{x^2 + x}$  is

 [Watch Video Solution](#)

165. The solution of  $\frac{dv}{dt} + \frac{k}{m}v = -g$  is

 [Watch Video Solution](#)

166. The solution of the equation  $\frac{dy}{dx} = \cos(x - y)$  is

 [Watch Video Solution](#)

167. The solution of the equation  $(x^2y + x^2)dx + y^2(x - 1)dy = 0$  is given by

 [Watch Video Solution](#)

168. Solution of differential equation  $dy - \sin x \sin y dx = 0$  is

 [Watch Video Solution](#)

169. Solution of  $\frac{dy}{dx} + 2xy = y$  is

 [Watch Video Solution](#)

170. The general solution of the differential equation

$\frac{dy}{dx} + \sin\left(\frac{x+y}{2}\right) = \sin\left(\frac{x-y}{2}\right)$  is

 [Watch Video Solution](#)

171. The solution of  $\frac{xdx + ydy}{xdy - ydx} = \sqrt{\frac{1 - x^2 - y^2}{x^2 + y^2}}$  is

 [Watch Video Solution](#)

172. The curve for which the length of the normal is equal to the length of the radius vector is/are (a) circles (b) rectangular hyperbola (c) ellipses (d) straight lines

 [Watch Video Solution](#)

173.  $f(x, y) = \sin^{-1}(xy) + \tan^{-1}(yx)$  is homogeneous function of degree:

 [Watch Video Solution](#)

174. The graph of the function  $y = f(x)$  passing through the point (0,1) and satisfying the differential equation  $\frac{dy}{dx} + y \cos x = \cos x$  is such that (a) it is a differential function for all  $x \in \mathbb{R}$ . (b) it is continuous for all  $x \in \mathbb{R}$ . (c) it is periodic. (d) it is passing through  $(\pi, 1)$

 [Watch Video Solution](#)

175. The solution of  $\frac{dy}{dx} = \frac{ax + h}{by + k}$  represent a parabola when (a)  $a = 0, b \neq 0$  (b)  $a \neq 0, b \neq 0$  (c)  $b = 0, a \neq 0$  (d)  $a = 0, b \in R$

 [Watch Video Solution](#)

176. The equation of the curve satisfying the differential equation  $y_2(x^2 + 1) = 2xy_1$  passing through the point (0,1) and having slope of tangent at  $x = 0$  as 3 (where  $y_2$  and  $y_1$  represent 2nd and 1st order derivative), then (a)  $y = f(x)$  is a strictly increasing function (b)  $y = f(x)$  is a non-monotonic function (c)  $y = f(x)$  has a three distinct real roots (d)  $y = f(x)$  has only one negative root.

 [Watch Video Solution](#)

177.  $y = ae^{-\frac{1}{x}} + b$  is a solution of  $\frac{dy}{dx} = \frac{y}{x^2}$ , then (a)  $a \in R$  (b)  $b = 0$  (c)  $b = 1$  (d)  $a$  takes finite number of values

 [Watch Video Solution](#)

**178.** For equation of the curve whose subnormal is constant, then (a) its eccentricity is 1 (b) its eccentricity is  $\sqrt{2}$  (c) its axis is the x-axis (d) its axis is the y-axis.



[Watch Video Solution](#)

**179.** If  $f(x)$ ,  $g(x)$  be twice differentiable functions on  $[0,2]$  satisfying  $f''(x) = g''(x)$ ,  $f'(1) = 2g'(1) = 4$  and  $f(2) = 3g(2) = 9$ , then  $f(x) - g(x)$  at  $x = 4$  equals (A) 0 (B) 10 (C) 8 (D) 2



[Watch Video Solution](#)

**180.** The solution of the differential equation  $(x^2y^2 - 1)dy + 2xy^3dx = 0$  is



[Watch Video Solution](#)

181. The solution of  $\frac{x^2 dy}{dx} - xy = 1 + \cos\left(\frac{y}{x}\right)$  is

 [Watch Video Solution](#)

182. Solve the equation  $(x + y + 1)\left(\frac{dy}{dx}\right) = 1$

 [Watch Video Solution](#)

183. Solve  $x\left(\frac{dy}{dx}\right) = y(\log y - \log x + 1)$

 [Watch Video Solution](#)

184. The slope of the tangent at  $(x, y)$  to a curve passing through  $\left(1, \frac{\pi}{4}\right)$  is given by  $\frac{y}{x} - \cos^2\left(\frac{y}{x}\right)$ , then the equation of the curve is

 [Watch Video Solution](#)



185. The solution of  $(x^2 + xy)dy = (x^2 + y^2)dx$  is

 [Watch Video Solution](#)

186. The solution of differential equation  $yy' = x \left( \frac{y^2}{x^2} + \frac{f\left(\frac{y^2}{x^2}\right)}{f'\left(\frac{y^2}{x^2}\right)} \right)$  is

 [Watch Video Solution](#)

187. The solution of  $(x^2 + xy)dy = (x^2 + y^2)dx$  is (a)

(b)  $\log x = \log((d)(e)x - y(f)) + (g)\frac{y}{h}x(i)(j) + c(k)$  (l) (m)

(n)  $\log x = 2\log((p)(q)x - y(r)) + (s)\frac{y}{t}x(u)(v) + c(w)$  (x) (y)

*[Math Processing Error]* (jj) (kk) None of these

 [Watch Video Solution](#)

188. The solution of the differential equation

$$\left\{1 + x\sqrt{(x^2 + y^2)}\right\}dx + \left\{\sqrt{(x^2 + y^2)} - 1\right\}ydy = 0$$
 is equal to

 [Watch Video Solution](#)

189. The slope of the tangent at  $(x, y)$  to a curve passing through a point

$(2, 1)$  is  $\frac{x^2 + y^2}{2xy}$ , then the equation of the curve is

 [Watch Video Solution](#)

190. If  $x \frac{dy}{dx} = x^2 + y - 2$ ,  $y(1) = 1$ , then  $y(2)$  equals .....

 [Watch Video Solution](#)

191. If  $y = y(x)$  and it follows the relation  $4xe^{xy} = y + 5\sin^2 x$ , then

$y'(0)$  is equal to \_\_\_\_\_

 [Watch Video Solution](#)

**192.** A right circular cone with radius  $R$  and height  $H$  contains a liquid which evaporates at a rate proportional to its surface area in contact with air (proportionality constant  $k$  is positive). Suppose that  $r(t)$  is the radius of the liquid cone at time  $t$ . The time after which the cone is empty is

 [Watch Video Solution](#)

**193.** A curve  $C$  passes through  $(2,0)$  and the slope at  $(x, y)$  as  $\frac{(x+1)^2 + (y-3)}{x+1}$ . Find the equation of the curve.

 [Watch Video Solution](#)

**194.** The equation of motion is given by  $s(t) = 2t^3 - 6t^2 + 6$ . At what time, the velocity and accelerations are zero?

 [Watch Video Solution](#)

**195.** Statement 1 : The differential equation of all circles in a plane must be of order 3. Statement 2 : There is only one circle passing through three non-collinear points.



**Watch Video Solution**

**196.** In which of the following differential equation degree is not defined?

(a)  $\frac{d^2y}{dx^2} + 3\left(\frac{dy}{dx}\right)^2 = x \log\left(\frac{d^2y}{dx^2}\right)$  (b)

(c)  $x = \sin\left(\left(\frac{dy}{dx}\right) - 2y\right), |x| < 1$

(d)  $x - 2y = \log\left(\frac{dy}{dx}\right)$



**Watch Video Solution**

**197.** Statement 1 : Degree of the differential equation  $2x - 3y + 2 = \log\left(\frac{dy}{dx}\right)$  is not defined. Statement 2 : In the given differential equation, the power of highest order derivative when expressed as the polynomials of derivatives is called degree.



Watch Video Solution

**198.** Statement 1 : The differential equation of the family of curves represented by  $y = Ae^x$  is given by  $\frac{dy}{dx} = y$  Statement 2 :  $\frac{dy}{dx} = y$  is valid for every member of the given family.



Watch Video Solution

**199.** What is the order of the differential equation whose general solution is  $y = c_1 \cos 2x + c_2 \sin^2 x + c_3 \cos^2 x + c_4 e^{2x} + c_5 e^{2x+c_6}$ ?



Watch Video Solution

**200.** Statement 1 : Order of a differential equation represents the number of arbitrary constants in the general solution. Statement 2 : Degree of a differential equation represents the number of family of curves.



Watch Video Solution

**201.** Solve the differential equation  $x \frac{dy}{dx} = x^2 + y$



[Watch Video Solution](#)

**202.** Let  $f(x), x \geq 0$ , be a non-negative continuous function, and let

$$F(x) = \int_0^x f(t)dt, x \geq 0, \text{ if for some } c > 0, f(x) \leq cF(x) \text{ for all}$$

$x \geq 0$ , then show that  $f(x) = 0$  for all  $x \geq 0$ .



[Watch Video Solution](#)

**203.** A particle moves so that the distance moved is according to the law

$$s(t) = \frac{t^3}{3} - t^2 + 3. \text{ At what time the velocity and acceleration are zero}$$

respectively?



[Watch Video Solution](#)

**204.** A curve  $C$  has the property that if the tangent drawn at any point  $P$  on  $C$  meets the co-ordinate axis at  $A$  and  $B$ , then  $P$  is the mid-point of  $AB$ . The curve passes through the point  $(1,1)$ . Determine the equation of the curve.



[Watch Video Solution](#)

**205.** A country has a food deficit of 10%. Its population ear. Its annual food production every year is 4% more than that of the last year. Assuming that the average food requirement per person remains constant, prove that the country will become self-sufficient in food after  $n$  years, where  $n$  is the smallest integer bigger than or equal to

$$\frac{\log_e 10 - \log_e 9}{(\log_e 1.04) - 0.03}$$



[Watch Video Solution](#)

**206.** Integrating factor of the differential equation

$$\cos x \frac{dy}{dx} + y \sin x = 1$$
 is



Watch Video Solution

207. Solution of the equation  $\cos^2 x \left( \frac{dy}{dx} \right) - (\tan 2x)y = \cos^4 x$ , when  $y(\pi/6) = (3\sqrt{3})/8$  is



Watch Video Solution

208. If integrating factor of  $x(1 - x^2)dy + (2x^2y - y - ax^3)dx = 0$  is  $e^{\int p dx}$ , then  $P$  is equal to



Watch Video Solution

209. A function  $y = f(x)$  satisfies  $(x + 1)f'(x) - 2(x^2 + x)f(x) = \frac{e^x - 2}{(x + 1)}$ ,  $\forall x > -1$ . If  $f(0) = 5$ , then  $f(x)$  is



Watch Video Solution



210. The general solution of the differential equation,  $y' + y\varphi'(x) - \varphi'(x)\varphi(x) = 0$ , where  $\varphi(x)$  is a known function, is

 [Watch Video Solution](#)

211. The solution of  $\frac{dy}{dx} = \frac{x^2 + y^2 + 1}{2xy}$  satisfying  $y(1) = 1$  is

 [Watch Video Solution](#)

212. The integrating factor of the differential equation  $\frac{dy}{dx}(x(\log)x) + y = 2(\log)x$  is given by

 [Watch Video Solution](#)

213. The solution of the differential equation  $x(x^2 + 1)\left(\frac{dy}{dx}\right) = y(1 - x^2) + x^3 \log x$  is

 [Watch Video Solution](#)

**214.** The solution of the differential equation  $\frac{dy}{dx} = \frac{1}{xy[x^2\sin y^2 + 1]}$  is

 [Watch Video Solution](#)

**215.** The general solution of the equation  $\frac{dy}{dx} = 1 + xy$  is

 [Watch Video Solution](#)

**216.** A normal is drawn at a point  $P(x, y)$  of a curve. It meets the x-axis at  $Q$ . If  $PQ$  has constant length  $k$ , then show that the differential equation describing such curves is  $y\frac{dy}{dx} = \pm\sqrt{k^2 - y^2}$ . Find the equation of such a curve passing through  $(0, k)$ .

 [Watch Video Solution](#)

217. If the solution of the differential equation  $\frac{dy}{dx} - y = 1 - e^{-x}$  and  $y(0) = y_0$  has a finite value, when  $x \rightarrow \infty$ , then the value of  $\left| \frac{2}{y_0} \right|$  is \_

 [Watch Video Solution](#)

218. Let  $y = y(t)$  be a solution to the differential equation  $y' + 2ty = t^2$ , then  $16(\lim)_{t \rightarrow \infty} \frac{y}{t}$  is \_\_\_\_\_

 [Watch Video Solution](#)

219. If the dependent variable  $y$  is changed to  $z$  by the substitution  $y = \tan z$  and the differential equation  $\frac{d^2y}{dx^2} = 1 + \frac{2(1+y)}{1+y^2} \left( \frac{dy}{dx} \right)^2$  is changed to  $\frac{d^2z}{dx^2} = \cos^2 z + k \left( \frac{dz}{dx} \right)^2$ , then the value of  $k$  equal to \_\_\_\_\_

 [Watch Video Solution](#)

**220.** If the independent variable  $x$  is changed to  $y$ , then the differential

equation  $x \frac{d^2y}{dx^2} + \left(\frac{dy}{dx}\right)^3 - \frac{dy}{dx} = 0$  is changed to  $x \frac{d^2x}{dy^2} + \left(\frac{dx}{dy}\right)^2 = k$  where  $k$  equals \_\_\_\_



[Watch Video Solution](#)

**221.** If the solution of the differential equation  $\frac{dy}{dx} = \frac{1}{x \cos y + \sin 2y}$  is

$x = ce^{\sin y} - k(1 + \sin y)$ , then the value of  $k$  is \_\_\_\_\_



[Watch Video Solution](#)

**222.** Tangent is drawn at the point  $(x_i, y_i)$  on the curve  $y = f(x)$ , which intersects the x-axis at  $(x_{i+1}, 0)$ . Now, again a tangent is drawn at

$(x_{i+1}, y_{i+1})$  on the curve which intersect the x-axis at  $(x_{i+2}, 0)$  and the

process is repeated  $n$  times, i.e.  $i = 1, 2, 3, \dots, n$ . If  $x_1, x_2, x_3, \dots, x_n$  form an arithmetic progression with common difference equal to  $(\log)_2 e$  and

curve passes through  $(0, 2)$ . Now if curve passes through the point  $(-2, k)$ , then the value of  $k$  is \_\_\_\_

 [Watch Video Solution](#)

**223.** The curve passing through the point  $(1, 1)$  satisfies the differential equation  $\frac{dy}{dx} + \frac{\sqrt{(x^2 - 1)(y^2 - 1)}}{xy} = 0$ . If the curve passes through the point  $(\sqrt{2}, k)$ , then the value of  $[k]$  is (where  $[.]$  represents greatest integer function) \_\_\_\_

 [Watch Video Solution](#)

**224.** If the eccentricity of the curve for which tangent at point  $P$  intersects the  $y$ -axis at  $M$  such that the point of tangency is equidistant from  $M$  and the origin is  $e$ , then the value of  $5e^2$  is \_\_\_\_

 [Watch Video Solution](#)

225. The perpendicular from the origin to the tangent at any point on a curve is equal to the abscissa of the point of contact. Also curve passes through the point (1,1). Then the length of intercept of the curve on the x-axis is \_\_\_\_\_



Watch Video Solution

## Examples

1. Find the order and degree of the following differential equations. i)

$$\frac{dy}{dx} + y = \frac{1}{\frac{dy}{dx}}, \text{ ii) } e^{\frac{d^3y}{dx^3}} - x \frac{d^2y}{dx^2} + y = 0, \text{ iii) } \sin^{-1}\left(\frac{dy}{dx}\right) = x + y, \text{ iv)}$$

$$\log_e\left(\frac{dy}{dx}\right) = ax + by$$

$$\text{v) } y \frac{d^2y}{dx^2} + x \left(\frac{dy}{dx}\right)^2 - 4y \frac{dy}{dx} = 0$$



Watch Video Solution

2. Form the differential equation of family of lines concurrent at the origin.

 [Watch Video Solution](#)

3. Form the differential equation of all circle touching the x-axis at the origin and centre on the y-axis.

 [Watch Video Solution](#)

4. Form the differential equation of family of lines situated at a constant distance  $p$  from the origin.

 [Watch Video Solution](#)

5. From the differential equation of the family of parabolas with focus at the origin and axis of symmetry along the x-axis. Find the order and

degree of the differential equation.



[Watch Video Solution](#)

6. The differential equation of all parabolas whose axis are parallel to the y-axis is (a)

(b) (c) (d)  $\frac{(e)(f)d^{(g)3(h)}(i)y}{j} \left( (k)d(l)x^{(m)3(n)}(o) \right) (p)(q) = 0(r) (s) (b)$

(t)(u)(v)  $\frac{(w)(x)d^{(y)2(z)}(aa)x}{bb} \left( (cc)d(dd)y^{(ee)2(ff)}(gg) \right) (hh)(ii) = C(jj)$

(kk) (c) *[Math Processing Error]* (ii) (d) *[Math Processing Error]* (ggg)



[View Text Solution](#)

7. From the differential equation of the family curves having equation

$$y = (\sin^{-1} x)^2 + A \cos^{-1} x + B.$$



[Watch Video Solution](#)



8. What is the order of the differential equation whose general solution is

$$y = c_1 \cos 2x + c_2 \sin^2 x + c_3 \cos^2 x + c_4 e^{2x} + c_5 e^{2x+c_6}$$

 [Watch Video Solution](#)

9. Find the particular solution of the differential equation

$$(1 + e^{2x})dy + (1 + y^2)e^x dx = 0. \text{ Given that } y = 1 \text{ when } x = 0.$$

 [Watch Video Solution](#)

10. Solve  $\log\left(\frac{dy}{dx}\right) = 4x - 2y - 2$ , given that  $y = 1$  when  $x = 1$ .

 [Watch Video Solution](#)

11. Solve the differential equation  $xy \frac{dx}{ydx} = \frac{1 + y^2}{1 + x^2} (1 + x + x^2)$

 [Watch Video Solution](#)

12. Solve the following differential equation:

$$\sqrt{1 + x^2 + y^2 + x^2 y^2} + xy \frac{dy}{dx} = 0$$

 [Watch Video Solution](#)

13. Solve  $\frac{d^2y}{dx^2} = \left(\frac{dy}{dx}\right)^2$

 [Watch Video Solution](#)

14. Solve  $\frac{dy}{dx} = (x + y)^2$

 [Watch Video Solution](#)

15. Solve  $\frac{dy}{dx} \sqrt{1 + x + y} = x + y - 1$

 [Watch Video Solution](#)

16. Show that the given differential equation is homogeneous and solve each of them.

$$(x^2 + xy)dy = (x^2 + y^2)dx$$

 [Watch Video Solution](#)

17.  $xdy - ydx = \sqrt{x^2 + y^2}dx$

 [Watch Video Solution](#)

18. Solve  $x \sin\left(\frac{y}{x}\right)dy = \left(y \sin\left(\frac{y}{x}\right) - x\right)dx$ .

 [Watch Video Solution](#)

19.  $\left(1 + \frac{e^x}{y}\right)dx + \frac{e^x}{y}\left(1 - \frac{x}{y}\right)dy = 0$

 [Watch Video Solution](#)

20. Solve  $x dy = \left( y + x \frac{f\left(\frac{y}{x}\right)}{f'\left(\frac{y}{x}\right)} \right) dx$

 [Watch Video Solution](#)

21. Find the real value of  $m$  for which the substitution  $y = u^m$  will transform the differential equation  $2x^4y \frac{dy}{dx} + y^4 = 4x^6$  in to a homogeneous equation.

 [Watch Video Solution](#)

22. Solve  $\frac{dy}{dx} = \frac{x + 2y + 3}{2x + 3y + 4}$

 [Watch Video Solution](#)

23. Solve:

$$x dy + y dx = \frac{x dy - y dx}{x^2 + y^2}$$



 [Watch Video Solution](#)

24. Solve:  $\left[ (x + 1) \frac{y}{x} + \sin y \right] dx + [x + \log_e x + x \cos y] dy = 0$

 [Watch Video Solution](#)

25. Solve:  $(x \cos x - \sin x) dx = \frac{x}{y} \sin x dy$

 [Watch Video Solution](#)

26. Solve:  $y^4 dx + 2xy^3 dy = \frac{y dx - x dy}{x^3 y^3}$

 [Watch Video Solution](#)

27. Solve:

$$\frac{dy}{dx} = \frac{yf'(x) - y^2}{f(x)}$$

 [Watch Video Solution](#)

28. Solve the differential equation  $ye^{\frac{x}{y}} dx = \left(xe^{\frac{x}{y}} + y^2\right) dy (y \neq 0)$ .

 [Watch Video Solution](#)

29. Solve  $\frac{dy}{dx} = \frac{x - 2y + 5}{2x + y - 1}$

 [Watch Video Solution](#)

30. Solve  $\frac{x + y \frac{dy}{dx}}{y - x \frac{dy}{dx}} = x^2 + 2y^2 + \frac{y^4}{x^2}$

 [Watch Video Solution](#)

31. Solve:

$$1 \left( \frac{1}{y} \frac{\sin x}{y} - \frac{y}{x^2} \frac{\cos y}{x} + 1 \right) dx + \left( \frac{1}{x} \frac{\cos y}{x} - \frac{x}{y^2} \frac{\sin y}{x} + \frac{1}{y^2} \right) dy = 0$$

 [View Text Solution](#)

32. Solve  $x^2 \left( \frac{dy}{dx} \right) + y = 1$

 [Watch Video Solution](#)

33. Solve:

$$ydx - xdy + \log x dx = 0$$

 [Watch Video Solution](#)

34. Solve  $(1 + y + x^2y)dx + (x + x^3)dy = 0$

 [Watch Video Solution](#)

35. Solve :  $(x + 2y^3) \frac{dy}{dx} = y$

 [Watch Video Solution](#)

36. Solve the differential equation:  $(1 + y^2) + (x - e^{\tan^{-1}y}) \frac{dy}{dx} = 0$



Watch Video Solution

37. Let  $u(x)$  and  $v(x)$  satisfy the differential equation  $\frac{du}{dx} + p(x)u = f(x)$  and  $\frac{dv}{dx} + p(x)v = g(x)$  are continuous functions. If  $u(x_1) > v(x_1)$  for some  $x_1$  and  $f(x) > g(x)$  for all  $x > x_1$ , prove that any point  $(x, y)$ , where  $x > x_1$ , does not satisfy the equations  $y = u(x)$  and  $y = v(x)$ .



View Text Solution

38. Solve  $\left(\frac{dy}{dx}\right) + \left(\frac{y}{x}\right) = y^3$



Watch Video Solution

39. Solve  $\left(\frac{dy}{dx}\right) = e^{x-y}(e^x - e^y)$ .





Watch Video Solution

40. Solve  $(x - 1)dy + ydx = x(x - 1)y^{\frac{1}{3}}dx$ .



Watch Video Solution

41. The solution of the differential equation,  $e^x(x + 1)dx + (ye^y - xe^x)dy = 0$  with initial condition  $f(0) = 0$ , is



Watch Video Solution

42. The slope of a curve, passing through (3,4) at any point is the reciprocal of twice the ordinate of that point. Show that it is parabola.



Watch Video Solution

**43.** Find the equation of a curve passing through the point  $(0, 2)$ , given that the sum of the coordinates of any point on the curve exceeds the magnitude of the slope of the tangent to the curve at that point by 5.

 [Watch Video Solution](#)

**44.** Find the equation of the curve passing through  $(2,1)$  which has constant sub-tangent.

 [Watch Video Solution](#)

**45.** If length of tangent at any point on the curve  $y = f(x)$  intercepted between the point and the x-axis is of length 1. Find the equation of the curve.

 [Watch Video Solution](#)

**46.** Find the equation of a curve passing through the point (1,1) if the perpendicular distance of the origin from the normal at any point  $P(x, y)$  of the curve is equal to the distance of  $P$  from the  $x$ -axis.

 [View Text Solution](#)

**47.** Find the equation of the curve such that the portion of the  $x$ -axis cut off between the origin and the tangent at a point is twice the abscissa and which passes through the point (1,2).

 [View Text Solution](#)

**48.** Find the equation of the curve passing through the point, (5,4) if the sum of reciprocal of the intercepts of the normal drawn at any point  $P(x,y)$  on it is 1.

 [View Text Solution](#)

49. Find the equation of the curve passing through the origin if the middle point of the segment of its normal from any point of the curve to the x-axis lies on the parabola  $2y^2 = x$

 [Watch Video Solution](#)

50. Find the equation of family of curves which intersect the family of curves  $xy=c$  at an angle  $45^\circ$ .

 [View Text Solution](#)

51. Find the orthogonal trajectory of  $y^2 = 4ax$  (a being the parameter).

 [Watch Video Solution](#)

52. The population of a certain is known to increase at a rate proportional to the number of people presently living in the country. If after two years

the population has doubled, and after three years the population is 20,000 estimates the number of people initially living in the country.

 [View Text Solution](#)

**53.** What constant interest rate is required if an initial deposit placed into an account accrues interest compounded continuously is to double its value in six years? ( $\ln|x| = 0.6930$ )

 [Watch Video Solution](#)

**54.** Suppose that a mothball loses volume by evaporation at a rate proportional to its instantaneous area. If the diameter of the ball decreases from 2cm to 1cm in 3 months, how long will it take until the ball has practically gone?

 [View Text Solution](#)

55. A body at a temperature of  $50^{\circ}F$  is placed outdoors where the temperature is  $100^{\circ}F$ . If the rate of change of the temperature of a body is proportional to the temperature difference between the body and its surrounding medium. If after 5 min the temperature of the body is  $60^{\circ}F$ , find (a) how long it will take the body to reach a temperature of  $75^{\circ}F$  and (b) the temperature of the body after 20 min.

 [Watch Video Solution](#)

56. Find the time required for a cylindrical tank of radius  $r$  and height  $H$  to empty through a round hole of area  $a$  at the bottom. The flow through the hole is according to the law  $v(t) = k\sqrt{2gh(t)}$ , where  $v(t)$  and  $h(t)$ , are respectively, the velocity of flow through the hole and the height of the water level above the hole at time  $t$ , and  $g$  is the acceleration due to gravity.

 [View Text Solution](#)

57. A country has a food deficit of 10%. Its population grows continuously at a rate of 3% per year. Its annual food production every year is 4% more than that of the last year. Assuming that the average food requirement per person remains constant, prove that the country will become self-sufficient in food after  $n$  years, where  $n$  is the smallest integer bigger than or equal to  $\frac{\ln 10 - \ln 9}{\ln(1.04) - 0.03}$



[View Text Solution](#)

58. A hemi-spherical tank of radius 2 m is initially full of water and has an outlet of  $12\text{cm}^2$  cross-sectional area at the bottom. The outlet is opened at some instant. The flow through the outlet is according to the law  $v(t) = \sqrt{0.62gh(t)}$ , where  $v(t)$  and  $h(t)$  are, respectively, the velocity of the flow through the outlet and the height of water level above the outlet and the height of water level above the outlet at time  $t$ , and  $g$  is the acceleration due to gravity. Find the time it takes to empty the tank.



[View Text Solution](#)

59. Solve  $\frac{x + y \frac{dy}{dx}}{y - x \frac{dy}{dx}} = x^2 + 2y^2 + \frac{y^4}{x^2}$

 [Watch Video Solution](#)

60. Show that the given differential equation is homogeneous and solve each of them.

$$\left\{ x \cos\left(\frac{y}{x}\right) + y \sin\left(\frac{y}{x}\right) \right\} y dx = \left\{ y \sin\left(\frac{y}{x}\right) - x \cos\left(\frac{y}{x}\right) \right\} x dy$$

 [Watch Video Solution](#)

61. Solve  $\frac{dy}{dx} = \frac{(x + y)^2}{(x + 2)(y - 2)}$

 [Watch Video Solution](#)

62. Solve  $y \left( \frac{dy}{dx} \right)^2$

 [View Text Solution](#)



63. If  $y + \frac{d}{dx}(xy) = x(\sin x + \log x)$ ,  $f \in dy(x)$ .

 [Watch Video Solution](#)

64. If  $\int_a^x ty(t)dt = x^2 + y(x)$ , then find  $y(x)$

 [Watch Video Solution](#)

65. about to only mathematics

 [Watch Video Solution](#)

66. Solve  $\frac{dy}{dx} = \frac{s \in}{\sin 2y - x \cos y}$

 [Watch Video Solution](#)

67. Solve:

$$\frac{dy}{dx} + \frac{3y}{x} = g(x), \text{ where } g(x) = \begin{cases} 1 & \text{if } 0 \leq x \leq 1 \\ \frac{1}{x} & \text{if } x > 1 \end{cases}$$

$$y\left(\frac{1}{2}\right) = \frac{1}{8} \text{ and } y(x) \text{ is continuous on } [0, \infty].$$



[View Text Solution](#)

68. Solve:  $(x \cos y - y \sin y)dy + (x \sin y + y \cos y)dx = 0$



[View Text Solution](#)

69. If  $y_1$  and  $y_2$  are the solution of the differential equation

$$\frac{dy}{dx} + Py = Q, \text{ where } P \text{ and } Q \text{ are functions of } x \text{ alone and } y_2 = y_1 z,$$

then prove that  $z = 1 + c \cdot e^{-\int \frac{Q}{y_1} dx}$ , where  $c$  is an arbitrary constant.



[Watch Video Solution](#)

70. If  $y_1$  and  $y_2$  are two solutions to the differential equation  $\frac{dy}{dx} + P(x)y = Q(x)$ . Then prove that  $y = y_1 + c(y_1 - y_2)$  is the general solution to the equation where  $c$  is any constant.



[Watch Video Solution](#)

71. Let  $f(x), x \geq 0$ , be a non-negative continuous function, and let  $f(x) = \int_0^x f(t)dt, x \geq 0$ , if for some  $c > 0, f(x) \leq cF(x)$  for all  $x \geq 0$ , then show that  $f(x) = 0$  for all  $x \geq 0$ .



[Watch Video Solution](#)

72. Find a pair of curves such that (a) the tangents drawn at points with equal abscissas intersect on the y-axis. (b) the normal drawn at points with equal abscissas intersect on the x-axis. (c) one curve passes through (1,1) and other passes through (2, 3).



[Watch Video Solution](#)

**73.** Given two curves:  $y = f(x)$  passing through the point  $(0, 1)$  and  $g(x) = \int_{-\infty}^x f(t)dt$  passing through the point  $\left(0, \frac{1}{n}\right)$ . The tangents drawn to both the curves at the points with equal abscissas intersect on the x-axis. Find the curve  $y = f(x)$ .

 [Watch Video Solution](#)

**74.** A cyclist moving on a level road at 4 m/s stops pedalling and lets the wheels come to rest. The retardation of the cycle has two components: a constant  $0.08 \text{ m/s}^2$  due to friction in the working parts and a resistance of  $0.02v^2/m$ , where  $v$  is speed in meters per second. What distance is traversed by the cycle before it comes to rest? (consider  $\ln 5=1.61$ ).

 [Watch Video Solution](#)

**75.** The force of resistance encountered by water on a motor boat of mass  $m$  going in still water with velocity  $v$  is proportional to the velocity  $v$ . At  $t = 0$  when its velocity is  $v_0$ , then engine shuts off. Find an expression

for the position of motor boat at time  $t$  and also the distance travelled by the boat before it comes to rest. Take the proportionality constant as  $k > 0$ .

 [Watch Video Solution](#)

**76.**  $A$  and  $B$  are two separate reservoirs of water. Capacity of reservoir  $A$  is double the capacity of reservoir  $B$ . Both the reservoirs are filled completely with water, their inlets are closed and then the water is released simultaneously from both the reservoirs. The rate of flow of water out of each reservoir at any instant of time is proportional to the quantity of water in the reservoir at the time. One hour after the water is released, the quantity of water in reservoir  $A$  is  $1\frac{1}{2}$  times the quantity of water in reservoir  $B$ . After how many hours do both the reservoirs have the same quantity of water?

 [Watch Video Solution](#)

1. Find the order and degree (if defined) of the following differential equations:

$$\frac{d^2y}{dx^2} = \left\{ 1 + \left( \frac{dy}{dx} \right)^4 \right\}^{5/3}$$



[View Text Solution](#)

2. Find the order and degree (if defined) of the equation:

$$\frac{d^3y}{dx^3} = x \ln \left( \frac{dy}{dx} \right)$$



[Watch Video Solution](#)

$$3. \left( \frac{d^4y}{dx^4} \right)^3 + 3 \left( \frac{d^2y}{dx^2} \right)^6 + \sin x = 2 \cos x$$



[View Text Solution](#)

4. Determine its order, degree (if exists)

$$\left(\frac{d^3y}{dx^3}\right)^{\frac{2}{3}} - 3\frac{d^2y}{dx^2} + 5\frac{dy}{dx} + 4 = 0$$



[Watch Video Solution](#)

5. Find the order and degree (if defined) of the equation:

$$a = \frac{1\left[1 + \left(\frac{dy}{dx}\right)^2\right]^{\frac{3}{2}}}{\frac{d^2y}{dx^2}}, \text{ where } a \text{ is constant}$$



[Watch Video Solution](#)

6. For each of the differential equations given below, indicate its order and degree (if defined).

$$\frac{d^4y}{dx^4} - \sin\left(\frac{d^3y}{dx^3}\right) = 0$$



[Watch Video Solution](#)

## Exercise 10.2

1. The differential equation of the family of all non-horizontal lines in a plane is

 [Watch Video Solution](#)

2. Find the differential equation of the family of circles with their centre at the origin.

 [Watch Video Solution](#)

3. Find the differential equation of all parabolas whose axes are parallel to the x-axis and having latus rectum  $a$ .

 [Watch Video Solution](#)



4. Find the differential equation of the family of curves  $y = Ae^{2x} + Be^{-2x}$ , where A and B are arbitrary constants.

 [Watch Video Solution](#)

5. Find the degree of the differential equation satisfying the relation

$$\sqrt{1+x^2} + \sqrt{1+y^2} = \lambda \left( x\sqrt{1+y^2} - y\sqrt{1+x^2} \right)$$

 [Watch Video Solution](#)

6. Find the differential equation whose general solution is given by

$y = (c_1 + c_2)\cos(x + c_3) - c_4e^{x+c}$ , where  $c_1, c_2, c_3, c_4, c_5$  are arbitrary constants.

 [Watch Video Solution](#)

1. Solve  $e^{\frac{dy}{dx}} = x + 1$ , given that when  $x = 0, y = 3$ .

 [Watch Video Solution](#)

2. Solve  $(x - y^2x)dx = (y - x^2y)dy$ .

 [Watch Video Solution](#)

3. The solution of  $\sec^2 x \tan y dx + \sec^2 y \tan x dy = 0$  is

 [Watch Video Solution](#)

4.  $e^x \tan y dx + (1 - e^x) \sec^2 y dy = 0$

 [Watch Video Solution](#)

5. Solve the following differential equations:  $\frac{dy}{dx} = 1 + x + y + xy$  (ii)

$$y - x \frac{dy}{dx} = a \left( y^2 + \frac{dy}{dx} \right)$$

 [Watch Video Solution](#)

6. Find a particular solution of the differential equation

$(x - y)(dx + dy) = dx - dy$ . Given that  $y = -1$ , when  $x = 0$ .

 [Watch Video Solution](#)

7. Solve  $\frac{dy}{dx} + yf'(x) = f(x)f'(x)$ , where  $f(x)$  is a given integrable function of  $x$ .

 [Watch Video Solution](#)

8. Solve  $\frac{dy}{dx} = \cos(x + y) - \sin(x + y)$ .

 [Watch Video Solution](#)

9. If a function 'f' satisfies the relation

$$f(x)f''(x) - f(x)f'(x) - f'(x)^2 = 0 \forall x \in R \text{ and } f(0) = 1 = f'(0).$$

Then find  $f(x)$ .

 [Watch Video Solution](#)

## Exercise 10.4

1. Solve the following differential equation:  $x \frac{dy}{dx} - y = 2 \sqrt{y^2 - x^2}$

 [Watch Video Solution](#)

2. Solve  $[2\sqrt{xy} - x]dy + ydx = 0$

 [Watch Video Solution](#)

3. Solve  $x \left( \frac{dy}{dx} \right) = y(\log y - \log x + 1)$

 [Watch Video Solution](#)

4.  $\left[ x \sin^2 \left( \frac{y}{x} \right) - y \right] dx + x dy = 0, y = \frac{\pi}{4}$  when  $x = 1$

 [Watch Video Solution](#)

5. Show that the differential equation  $y^3 dy - (x + y^2) dx = 0$  can be reduced to a homogenous equation.

 [Watch Video Solution](#)

6. Solve  $\frac{dy}{dx} = \frac{2x - y + 1}{x + 2y - 3}$

 [Watch Video Solution](#)

## Exercise 10.5

1. The solution of  $ydx - xdy + 3x^2y^2e^{x^3} dx = 0$  is

 [Watch Video Solution](#)

2. 
$$\frac{dy}{dx} = \frac{2xy}{x^2 - 1 - 2y}$$

 [Watch Video Solution](#)

3. The solution of the differential equation  $ydx + (x + x^2y)dy = 0$  is

 [Watch Video Solution](#)

4. The solution of the differential equation  $(xy^4 + y)dx - xdy = 0$ , is

 [Watch Video Solution](#)

$$5. y(x^2y + e^x)dx - e^x dy = 0$$

 [Watch Video Solution](#)

$$6. \frac{dy}{dx} = -\frac{y + \sin x}{x} \text{ satisfying condition } y(0) = 1$$

 [Watch Video Solution](#)

$$7. y(xy + 1)dx + x(1 + xy + x^2y^2)dy = 0$$

 [Watch Video Solution](#)

## Exercise 10.6

1. The integrating factor of the differential equation  $(1 - y^2) \frac{dy}{dx} + yx = ay$  ( $-1 < y' < 1$ ) is

 [Watch Video Solution](#)

2. Solve  $\frac{dy}{dx} + y \cot x = \sin x$

 [Watch Video Solution](#)

3. Solve the equation  $(x + y + 1) \left( \frac{dy}{dx} \right) = 1$

 [Watch Video Solution](#)

4. Solve  $(1 - x^2) \frac{dy}{dx} + 2xy = x \sqrt{1 - x^2}$

 [Watch Video Solution](#)

5. Solve the equation  $\frac{dy}{dx} = \frac{y}{2y \ln y + y - x}$

 [Watch Video Solution](#)



6. Solve the equation  $ydx + (x - y^2)dy = 0$



Watch Video Solution

7. Find the equation of a curve passing through  $(0, 1)$  and having gradient  $\frac{-(y + y^3)}{1 + x + xy}$  at  $(x, y)$



Watch Video Solution

## Exercise 10.7

1. Solve the equation  $\frac{dy}{dx} + \frac{1}{x} = \frac{e^y}{x^2}$



Watch Video Solution

2. Solve the equation  $\frac{dy}{dx} = +x \sin 2y = x^3 \cos^2 y$



Watch Video Solution

3. Solve the equation  $\frac{dy}{dx} + \frac{xy}{(1-x^2)}x\sqrt{y}$

 [Watch Video Solution](#)

4. Solve the equation  $\frac{dy}{dx} + (2x \tan^{-1} y - x^3)(1 + y^2) = 0$

 [Watch Video Solution](#)

5.  $\frac{dy}{dx} - \frac{\tan x}{1+x} = (1+x)e^x \sec y$

 [Watch Video Solution](#)

## Exercise 10.8

1. Find the equation of the curve in which the subnormal varies as the square of the ordinate.



[Watch Video Solution](#)

2. The curve for which the length of the normal is equal to the length of the radius vector is/are (a) circles (b) rectangular hyperbola (c) ellipses (d) straight lines



[Watch Video Solution](#)

3. Find the curve for which the perpendicular from the foot of the ordinate to the tangent is of constant length.



[Watch Video Solution](#)

4. A curve  $y = f(x)$  passes through the origin. Through any point  $(x, y)$  on the curve, lines are drawn parallel to the co-ordinate axes. If the curve divides the area formed by these lines and co-ordinates axes in the ratio  $m:n$ , find the curve.



[Watch Video Solution](#)

 Watch Video Solution

5. A normal is drawn at a point  $P(x, y)$  of a curve. It meets the x-axis at  $Q$ . If  $PQ$  has constant length  $k$ , then show that the differential equation describing such curves is  $y \frac{dy}{dx} = \pm \sqrt{k^2 - y^2}$ . Find the equation of such a curve passing through  $(0, k)$ .

 Watch Video Solution

6. Find the orthogonal trajectories of family of curves  $x^2 + y^2 = cx$

 Watch Video Solution

7. A curve  $C$  has the property that if the tangent drawn at any point  $P$  on  $C$  meets the co-ordinate axis at  $A$  and  $B$ , then  $P$  is the mid-point of  $AB$ . The curve passes through the point  $(1,1)$ . Determine the equation of the curve.

 Watch Video Solution

## Exercise 10.9

1. A person places Rs 500 in an account that interest compounded continuously. Assuming no additional deposits or withdrawals, how much will be in the account after seven years if the interest rate is a constant 8.5 percent for the first four years and a constant 9.25 percent for the last three years

 [View Text Solution](#)

2. Find the time required for a cylindrical tank of radius 2.5 m and height 3 m to empty through a round hole of 2.5 cm with a velocity  $2.5\sqrt{h}$  m/s,  $h$  being the depth of the water in the tank.

 [Watch Video Solution](#)

3. If the population of country double in 50 years, in how many years will it triple under the assumption that the rate of increase is proportional to the number of inhabitants.



[Watch Video Solution](#)

4. The rate at which a substance cools in moving air is proportional to the difference between the temperatures of the substance and that of the air. If the temperature of the air is 290 K and the substance cools from 370 K to 330 K in 10 min, when will the temperature be 295 K?



[Watch Video Solution](#)

5. about to only mathematics



[Watch Video Solution](#)

1. The degree of the differential equation satisfying

$$\sqrt{1-x^2} + \sqrt{1-y^2} = a(x-y), \text{ is}$$

- A. 1
- B. 2
- C. 3
- D. None of these

**Answer: A**



[View Text Solution](#)

2. Number of values of  $m \in \mathbb{N}$  for which  $y = e^{mx}$  is a solution of the

differential equation  $\frac{d^3y}{dx^3} - 3\frac{d^2y}{dx^2} - 4\frac{dy}{dx} + 12y = 0$  (a) 0 (b) 1 (c) 2 (d)

More than 2

- A. 0

B. 1

C. 2

D. More than 2

**Answer: C**



**Watch Video Solution**

3. The order of differential equation of family of circles passing through intersection of  $3x + 4y - 7 = 0$  and  $S = -x^2 + y^2 - 2x + 1 = 0$  is

A. 1

B. 2

C. 3

D. 4

**Answer: A**



**View Text Solution**



4. The differential equation of the family of all non-horizontal lines in a plane is

A.  $\frac{d^2y}{dx^2}$

B.  $\frac{d^2x}{dy^2} = 0$

C.  $\frac{dy}{dx} = 0$

D.  $\frac{dx}{dy} = 0$

Answer: B



Watch Video Solution

5. The differential equation of the family of curves

$y = e^x(A \cos x + B \sin x)$ , where  $A$  and  $B$  are arbitrary constants is (a)

(b)(c)(d)  $\frac{(e)(f)d^{(g)2(h)}(i)y}{j} \left( (k)d(l)x^{(m)2(n)}(o) \right) (p)(q) - 2(r) \frac{(s)dy}{t} ((u,$

(y) (z) [Math Processing Error] (xx) (yy) [Math Processing Error] (eeee) (ffff)

[Math Processing Error] (ddddd)

A.  $\frac{d^2y}{dx^2} - 2\frac{dy}{dx} + 2y = 0$

B.  $\frac{d^2y}{dx^2} + 2\frac{dy}{dx} - 2y = 0$

C.  $\frac{d^2y}{dx^2} + \left(\frac{dy}{dx}\right)^2 + y = 0$

D.  $\frac{d^2y}{dx^2} - 7\frac{dy}{dx} + 2y = 0$

Answer: A



Watch Video Solution

6. Differential equation of the family of circles touching the line  $y = 2$  at

$(0, 2)$  is (a)

(b)  $x^2 + (y - 2)^2 = r^2$  (c)  $x^2 + (y - 2)^2 = r^2$  (d)  $x^2 + (y - 2)^2 = r^2$  (e)  $x^2 + (y - 2)^2 = r^2$  (f)  $x^2 + (y - 2)^2 = r^2$  (g)  $x^2 + (y - 2)^2 = r^2$  (h)  $x^2 + (y - 2)^2 = r^2$  (i)  $x^2 + (y - 2)^2 = r^2$  (j)  $x^2 + (y - 2)^2 = r^2$  (k)  $x^2 + (y - 2)^2 = r^2$  (l)  $x^2 + (y - 2)^2 = r^2$  (m)  $x^2 + (y - 2)^2 = r^2$  (n)  $x^2 + (y - 2)^2 = r^2$  (o)  $x^2 + (y - 2)^2 = r^2$  (p)  $x^2 + (y - 2)^2 = r^2$  (q)  $x^2 + (y - 2)^2 = r^2$  (r)  $x^2 + (y - 2)^2 = r^2$  (s)  $x^2 + (y - 2)^2 = r^2$  (t)  $x^2 + (y - 2)^2 = r^2$  (u)  $x^2 + (y - 2)^2 = r^2$  (v)  $x^2 + (y - 2)^2 = r^2$  (w)  $x^2 + (y - 2)^2 = r^2$  (x)  $x^2 + (y - 2)^2 = r^2$  (y)  $x^2 + (y - 2)^2 = r^2$  (z)  $x^2 + (y - 2)^2 = r^2$

(aa) **[Math Processing Error]** (uu) (vv)

(ww)  $x^2 + (y - 2)^2 = r^2$  (xx)  $x^2 + (y - 2)^2 = r^2$  (yy)  $x^2 + (y - 2)^2 = r^2$  (zz)  $x^2 + (y - 2)^2 = r^2$  (aaa)  $x^2 + (y - 2)^2 = r^2$  (bbb)  $x^2 + (y - 2)^2 = r^2$  (ccc)  $x^2 + (y - 2)^2 = r^2$  (ddd)  $x^2 + (y - 2)^2 = r^2$  (eee)  $x^2 + (y - 2)^2 = r^2$  (fff)  $x^2 + (y - 2)^2 = r^2$  (ggg)  $x^2 + (y - 2)^2 = r^2$  (hhh)  $x^2 + (y - 2)^2 = r^2$  (iii)  $x^2 + (y - 2)^2 = r^2$  (jjj)  $x^2 + (y - 2)^2 = r^2$  (kkk)  $x^2 + (y - 2)^2 = r^2$  (lll)  $x^2 + (y - 2)^2 = r^2$  (mmm)  $x^2 + (y - 2)^2 = r^2$  (nnn)  $x^2 + (y - 2)^2 = r^2$  (ooo)  $x^2 + (y - 2)^2 = r^2$  (ppp)  $x^2 + (y - 2)^2 = r^2$  (qqq)  $x^2 + (y - 2)^2 = r^2$  (rrr)  $x^2 + (y - 2)^2 = r^2$  (sss)  $x^2 + (y - 2)^2 = r^2$  (ttt)  $x^2 + (y - 2)^2 = r^2$  (uuu)  $x^2 + (y - 2)^2 = r^2$  (vvv)  $x^2 + (y - 2)^2 = r^2$  (www)  $x^2 + (y - 2)^2 = r^2$  (xxx)  $x^2 + (y - 2)^2 = r^2$  (yyy)  $x^2 + (y - 2)^2 = r^2$  (zzz)  $x^2 + (y - 2)^2 = r^2$

$x^2 + (y - 2)^2 = r^2$

(xxx) (yyy) None of these

A.  $x^2 + (y - 2)^2 + \frac{dy}{dx}(y - 2) = 0$

B.  $x^2 + (y - 2) \left( 2 - 2x \frac{dx}{dy} - y \right) = 0$

C.  $x^2 + (y - 2)^2 + \left( \frac{dx}{dy} + y - 2 \right) (y - 2) = 0$

D. None of these

**Answer: B**



[Watch Video Solution](#)

7. The order and degree of the differential equation of all tangent lines to the parabola  $y = x^2$  is (a) 1,2 (b) 2,3 (c) 2,1 (d) 1,1

A. 1,2

B. 2,3

C. 2,1

D. 1,1

**Answer: A**



[Watch Video Solution](#)

8. The differential equation for the family of curve  $x^2 + y^2 - 2ay = 0$ ,

where  $a$  is an arbitrary constant, is (a)

(b)  $2\left(\frac{d}{dx}(x^2 + y^2 - 2ay) - (y^2)' - (2a)'\right) = xy$

(t) (u)

(v)  $2\left(\frac{d}{dx}(x^2 + y^2 - 2ay) + (y^2)'\right) = xy$

(nn) (oo) **[Math Processing Error]** (hhh) (iii)

(jjj)  $\left(\frac{d}{dx}(x^2 + y^2 - 2ay) + (y^2)'\right) = xy$

(bbbb)

A.  $2(x^2 - y^2)y' = xy$

B.  $2(x^2 + y^2)y' = xy$

C.  $(x^2 - y^2)y' = 2xy$

D.  $(x^2 + y^2)y' = 2xy$

Answer: C

 Watch Video Solution

9. The differential equation whose general solution is given by

$$y = c_1 \cos(x + c_2) - c_3 e^{(-x + c_4)} + (c_5 \sin x), \quad \text{where } c_1, c_2, c_3, c_4, c_5$$

are arbitrary constants, is



Watch Video Solution

10. If  $y = \frac{x}{\ln|cx|}$  (where  $c$  is an arbitrary constant) is the general solution

of the differential equation  $\frac{dy}{dx} = \frac{y}{x} + \phi\left(\frac{x}{y}\right)$  then function  $\phi\left(\frac{x}{y}\right)$  is:

A.  $x^2 / y^2$

B.  $-x^2 / y^2$

C.  $y^2 / x^2$

D.  $-y^2 / x^2$

Answer: D



Watch Video Solution

11. The differential equation of the curve  $\frac{x}{c-1} + \frac{y}{c+1} = 1$  is (a) [Math

Processing

Error]

(cc)

(dd)

(ee)(ff)  $\left( (gg)(hh)(ii) \frac{(jj)dy}{kk} ((ll)dx)(mm)(n) + 1(oo) \right) \left( (pp)(qq)y - x \right)$

(fff) (ggg)

(hhh)(iii)  $\left( (jjj)(kkk)(lll) \frac{(mmm)dy}{nnn} ((ooo)dx)(ppp)(qqq) + 1(rrr) \right) \left( (sss) \right)$

$$= 2(bbbb) \frac{(cccc)dy}{dddd} ((eeee)dx)(ffff)(gggg)(hhhh)$$

(iiii)

A.  $\frac{dy}{dx} - 1 \left( y + x \frac{dy}{dx} \right) = 2 \frac{dy}{dx}$

B.  $\left( \frac{dy}{dx} + 1 \right) \left( y - x \frac{dy}{dx} \right) = \frac{dy}{dx}$

C.  $\left( \frac{dy}{dx} + 1 \right) \left( y - x \frac{dy}{dx} \right) = 2 \frac{dy}{dx}$

D.  $y^2 / x^2$

Answer: C



Watch Video Solution

12. if  $y = y(x)$  and  $\frac{2 + \sin x}{y + 1} \left( \frac{dy}{dx} \right) = -\cos x$ ,  $y(0) = 1$ , then  $y\left(\frac{\pi}{2}\right) =$  (a)  $\frac{1}{e}$  (b)  $\frac{2}{m}$  (c)  $\frac{1}{3}$  (d)  $\frac{2}{3}$  (e)  $\frac{1}{3}$  (f)  $\frac{2}{3}$  (g)  $\frac{1}{3}$  (h)  $\frac{2}{3}$  (i)  $\frac{1}{3}$  (j)  $\frac{2}{3}$  (k)  $\frac{1}{3}$  (l)  $\frac{2}{3}$  (m)  $\frac{1}{3}$  (n)  $\frac{2}{3}$  (o)  $\frac{1}{3}$  (p)  $\frac{2}{3}$  (q)  $\frac{1}{3}$  (r)  $\frac{2}{3}$  (s)  $\frac{1}{3}$  (t)  $\frac{2}{3}$  (u)  $\frac{1}{3}$  (v)  $\frac{2}{3}$  (w)  $\frac{1}{3}$  (x)  $\frac{2}{3}$  (y)  $\frac{1}{3}$  (z)  $\frac{2}{3}$

A.  $1/3$

B.  $2/3$

C.  $-1/3$

D.  $1$

Answer: A



Watch Video Solution

13. Find the equation of the curve whose slope is  $\frac{y - 1}{x^2 + x}$  and which passes through the point (1,0).

A.  $(y - 1)(x + 1) + 2x = 0$

B.  $2x(y - 1) + x + 1 = 0$

C.  $x(y - 1)(x + 1) + 2 = 0$

D. None of these

**Answer: A**



**Watch Video Solution**

14. The solution of the equation  $\frac{dy}{dx} = \left( x \frac{2 \log x + 1}{\sin y + y \cos y} \right)$  is (a)
- (b)  $y \sin y = (d)x^{(e)2(f)}(g) \log x + (h) \frac{(i)(j)x^{(k)2(l)}(m)}{n} 2(o)(p) + c(q)$
- (r) (s) (t)  $y \cos y = (v)x^{(w)2(x)}(y)(\log x + 1) + c(z)$  (aa) (bb)
- (cc)  $y \cos y = (ee)x^{(ff)2(gg)}(hh) \log x + (ii) \frac{(jj)(kk)x^{(ll)2(mm)}(nn)}{oo} 2$
- (ss) (tt)  $(uu)(\vee) y \sin y = (ww)x^{(xx)2(yy)}(zz) \log x + c(aaa) (bbb)$

A.  $y \sin y = x^2 \log x + \frac{x^2}{2} + c$

B.  $y \cos y = x^2(\log x + 1) + c$

C.  $y \cos y = x^2 \log x + \frac{x^2}{2} + c$

D.  $y \sin y = x^2 \log x + c$



Answer: D



Watch Video Solution

15. The solution of the equation  $(x^2y + x^2)dx + y^2(x - 1)dy = 0$  is given by (a)

(b)  $x^{(e)2(f)} + (h)y^{(i)2(j)} + 2((l)(m)x - y(n)) + 21n(o) \frac{(p)}{c}$

(aa) (bb)

(cc)  $x^{(ff)2(gg)} + (ii)y^{(jj)2(kk)} + 2((mm)(\cap)x - y(\infty))$

(bbb) (ccc)

(ddd)  $(eee)(fff)x^{(ggg)2(hhh)} + (jjj)y^{(kkk)2(lll)} + 2((nnn)(c) = 0$  (bbbb)

(cccc) (dddd) None of these

A.  $x^2 + y^2 + 2(x - y) + 2\ln \frac{(x - 1)(y + 1)}{c} = 0$

B.  $x^2 + y^2 + 2(x - y) + \ln \frac{(x - 1)(y + 1)}{c} = 0$

C.  $x^2 + y^2 + 2(x - y) - 2\ln \frac{(x - 1)(y + 1)}{c} = 0$

D. None of these

Answer: A



Watch Video Solution

16. Solve the following differential equations

$$\frac{dy}{dx} = \sin x \cdot \sin y$$



Watch Video Solution

17. The solution of  $\frac{dv}{dt} + \frac{k}{m}v = -g$  is (a)

(b)  $v = c + \frac{mg}{k} e^{-\frac{k}{m}t}$  (s)

(b) [Math Processing Error] (kk) (c)

(d)  $v = c - \frac{mg}{k} e^{-\frac{k}{m}t}$  (u)

(d) [Math Processing Error] (mm)

A.  $v = ce^{-\frac{k}{m}t} - \frac{mg}{k}$

B.  $v = c - \frac{mg}{k} e^{-\frac{k}{m}t}$

C.  $ve^{-\frac{k}{m}t} = c - \frac{mg}{k}$

$$D. ve^{\frac{k}{m}t} = c - \frac{mg}{k}$$

Answer: A



Watch Video Solution

18. The general solution of the differential equation

$$\frac{dy}{dx} + \frac{\sin(x+y)}{2} = \frac{\sin(x-y)}{2} \quad \text{is} \quad (a)$$

$$(b)(c) \log \tan \left( (d)(e)(f) \frac{y}{g} 2(h)(i)(j) \right) = c - 2 \sin x (k) \quad (l) \quad (m) \quad \text{[Math$$

*Processing Error]* (ee) (ff) *[Math Processing Error]* (uu) (vv)

$$(ww)(\times) \log \tan \left( (yy)(zz)(aaa) \frac{y}{bbb} 4(ccc)(ddd) + (eee) \frac{\pi}{fff} 4(ggg)(hhh) \right)$$

(rrr)

A.  $\log \tan \left( \frac{y}{2} \right) = c - 2 \sin x$

B.  $\log \tan \left( \frac{y}{2} \right) = c - 2 \sin \left( \frac{x}{2} \right)$

C.  $\log \tan \left( \frac{y}{2} + \frac{\pi}{4} \right) = c - 2 \sin x$

D.  $\log \tan \left( \frac{y}{4} + \frac{\pi}{4} \right) = c - 2 \sin \left( \frac{x}{2} \right)$

**Answer: B**

 **Watch Video Solution**

19. if  $y + x \frac{dy}{dx} = x \frac{\phi(xy)}{\phi'(xy)}$  then  $\phi(xy)$  is

 **Watch Video Solution**

20. Find the solution of differential equation

$$x^2 = 1 + \left(\frac{x}{y}\right)^{-1} \frac{dy}{dx} + \frac{\left(\frac{x}{y}\right)^{-2} \left(\frac{dy}{dx}\right)^2}{2!} + \dots$$

 **Watch Video Solution**

21. The solution of the differential equation

$$\frac{x + \frac{x^3}{3!} + \frac{x^5}{5!} + \dots}{1 + \frac{x^2}{2!} + \frac{x^4}{4!} + \dots} = \frac{dx - dy}{dx + dy} \quad (a)$$

$$(b)(c)2y(d)e^{(e)(f)2x(g)}(h) = C(i)e^{(j)(k)2x(l)}(m) + 1(n) \quad (o) \quad (p)$$

$$(q)(r)2y(s)e^{(t)(u)2x(v)}(w) = C(x)e^{(y)(z)2x(aa)}(bb) - 1(cc) \quad (dd) \quad (ee)$$

$$(ff)(gg)y(hh)e^{(ii)(jj)2x(kk)}(ll) = C(mm)e^{(nn)(\infty)2x(pp)}(qq) + 2(rr)$$

(ss) (d) None of these

A.  $2ye^{2x} = Ce^{2x} + 1$

B.  $2ye^{2x} = Ce^{2x} - 1$

C.  $ye^{2x} = Ce^{2x} + 2$

D. None of these

**Answer: B**



**Watch Video Solution**

22. The solution of the differential equation

$$x = 1 + xy \frac{dy}{dx} + \frac{x^2 y^2}{2!} \left( \frac{dy}{dx} \right)^2 + \frac{x^3 y^3}{3!} \left( \frac{dy}{dx} \right)^3 + \dots \quad (a)$$

$$(b)(c)y = \ln((d)x(e)) + c(f) \quad (g) \quad (b)$$

$$(h)(i)(j)y^{(k)2(l)}(m) = (n)(o)((p)(q)\ln x(r))^{(s)2(t)}(u) + c(v) \quad (w) \quad (c)$$

$$(d)(e)y = \log x + xy(f) \quad (g) \quad (d) \quad (h)(i)xy = (j)x^{(k)y(l)}(m) + c(n) \quad (o)$$

A.  $y = \ln(x) + C$

B.  $y^2 = (\ln x)^2 + c$

C.  $y = \log x + xy$

D.  $xy = x^y + c$

**Answer: B**



**Watch Video Solution**

**23.** A curve passing through  $(2, 3)$  and satisfying the differential equation

$\int_0^x ty(t)dt = x^2y(x), (x > 0)$  is (a)

(b)  $(c)(d)x^{(e)2(f)}(g) + (h)y^{(i)2(j)}(k) = 13(l)$  (m) (b)

(n)(o)(p)y^{(q)2(r)}(s) = (t)\frac{9}{u}2(v)(w)x(x) (y) (c)

(d)(e)(f)\frac{(g)(h)x^{(i)2(j)}(k)}{l}8(m)(n) + (o)\frac{(p)(q)y^{(r)2(s)}(t)}{u}((v)18)(w)(x)

(z) (d) *[Math Processing Error]* (dd)

A.  $x^2 + y^2 = 13$

B.  $y^2 = \frac{9}{2}x$

C.  $\frac{x^2}{8} + \frac{y^2}{18} = 1$

D.  $xy = 6$

Answer: D

 [Watch Video Solution](#)

24. The solution of the differential equation  $\frac{d^2y}{dx^2} = \sin 3x + e^x + x^2$  when  $y_1(0) = 1$  and  $y(0) = 0$  is

A.  $y = \frac{-\sin 3x}{9} + e^x + \frac{x^4}{12} + \frac{1}{3}x - 1$

B.  $y = \frac{-\sin 3x}{9} + e^x + \frac{x^4}{12} + \frac{1}{3}x$

C.  $y = \frac{-\cos 3x}{3} + e^x + \frac{x^4}{12} + \frac{1}{3}x + 1$

D. None of these

Answer: A

 [Watch Video Solution](#)

25. Find the solution of the differential equation  $y' y'''' = 3(y'')^2$  is

 [Watch Video Solution](#)

26. The solution of the differential equation  $y'''' - 8y'' = 0$ , where  $y(0) = \frac{1}{8}$ ,  $y'(0) = 0$ ,  $y''(0) = 1$ , is

 [Watch Video Solution](#)

27. The slope of the tangent at  $(x, y)$  to a curve passing through  $(1, \frac{\pi}{4})$  is given by  $\frac{y}{x} - \cos^2\left(\frac{y}{x}\right)$ , then the equation of the curve is (a) (b) (c)  $y = (d)(e)\tan^{(f)(g)-1(h)}(i)\left((j)(k)\log\left((l)(m)(n)\frac{e}{o}x(p)(q)(r)\right)\right)(s)$  (u) (v) *[Math Processing Error]* (pp) (qq) *[Math Processing Error]* (kkk) (d) none of these

A.  $y = \tan^{-1} \log\left(\frac{e}{x}\right)$

B.  $y = x \tan^{-1} \log\left(\frac{x}{e}\right)$

C.  $y = x \tan^{-1} \log\left(\frac{e}{x}\right)$



D. None of these

**Answer: C**

 [Watch Video Solution](#)

28. Solve  $x \left( \frac{dy}{dx} \right) = y(\log y - \log x + 1)$

A.  $\frac{\log x}{y} = cy$

B.  $\frac{\log y}{x} = cy$

C.  $\frac{\log y}{x} = cx$

D. None of these

**Answer: C**

 [Watch Video Solution](#)

29. The solution of differential equation  $xy' = x \left( \frac{y^2}{x^2} + \frac{f\left(\frac{y^2}{x^2}\right)}{f'\left(\frac{y^2}{x^2}\right)} \right)$  is

(a)

$$(b)(c) f \left( (d)(e)(f) \frac{(g)(h)y^{(i)2(j)}(k)}{l} \left( (m)(n)x^{(o)2(p)}(q) \right) (r)(s)(t) \right) = c$$

(z) (b) **[Math Processing Error]** (ggg) (c)

$$(d)(e)(f)x^{(g)2(h)}(i) f \left( (j)(k)(l) \frac{(m)(n)y^{(o)2(p)}(q)}{r} \left( (s)(t)x^{(u)2(v)}(w) \right) \right)$$

(bb) (d) **[Math Processing Error]** (bbb)

A.  $f(y^2/x^2) = cx^2$

B.  $x^2 f(y^2/x^2) = c^2 y^2$

C.  $x^2 f(y^2/x^2) = c$

D.  $f(y^2/x^2) = cy/x$

**Answer: A**



**Watch Video Solution**

30. The solution of  $(x^2 + xy)dy = (x^2 + y^2)dx$  is (a)

(b)  $\log x = \log((d)(e)x - y(f)) + (g)\frac{y}{h}x(i)(j) + c(k)$  (l) (m)

(n)  $\log x = 2\log((p)(q)x - y(r)) + (s)\frac{y}{t}x(u)(v) + c(w)$  (x) (y)

*[Math Processing Error]* (jj) (kk) None of these

A.  $\log x = \log(x - y) + \frac{y}{x} + c$

B.  $\log x = 2\log(x - y) + \frac{y}{x} + c$

C.  $\log x = \log(x - y) + \frac{x}{y} + c$

D. None of these

**Answer: B**



**Watch Video Solution**

31. The solution of  $(y + x + 5)dy = (y - x + 1)dx$  is

A.  $\log\left((y + 3)^2 + (x + 2)^2\right) + \frac{\tan^{-1}(y + 3)}{y + 2} + C$

B.  $\log\left((y + 3)^2 + (x - 2)^2\right) + \frac{\tan^{-1}(y - 3)}{x - 2} = C$

$$C. \log\left((y+3)^2 + (x+2)^2\right) + 2\frac{\tan^{-1}(y+3)}{x+2} = C$$

$$D. \log\left((y+3)^2 + (x+2)^2\right) - 2\frac{\tan^{-1}(y+3)}{x+2} = C$$

**Answer: A**



**View Text Solution**

**32.** The slope of the tangent at  $(x, y)$  to a curve passing through a point

$(2, 1)$  is  $\frac{x^2 + y^2}{2xy}$ , then the equation of the curve is (a)

(b)(c)  $2\left((d)(e)(f)x^{(g)2(h)}(i) - (j)y^{(k)2(l)}(m)(n)\right) = 3x(o)$  (p) (b)

*[Math Processing Error]* (ee) (c)

(d)(e)  $x\left((f)(g)(h)x^{(i)2(j)}(k) - (l)y^{(m)2(n)}(o)(p)\right) = 6(q)$  (r) (d)

(s)(t)  $x\left((u)(v)(w)x^{(x)2(y)}(z) + (aa)y^{(bb)2(cc)}(dd)(ee)\right) = 10(ff)$

(gg)

A.  $2(x^2 - y^2) = 3x$

B.  $2(x^2 - y^2) = 6y$

C.  $x(x^2 - y^2) = 6$

D.  $x(x^2 + y^2) = 10$

Answer: A



Watch Video Solution

33. Solution of the differential equation  $ydx - xdy + x\sqrt{xy}dy = 0$  is



Watch Video Solution

34. The solution of  $\frac{x^2 dy}{dx} - xy = 1 + \frac{\cos y}{x}$  is (a)

(b)  $\tan\left(\frac{y}{2x}\right) = c - \frac{1}{m} \left(2x^{(p)2(q)}\right)$

(v) (w) **[Math Processing Error]** (ii) (jj)

(kk)  $\cos\left(\frac{y}{x}\right) = 1 + \frac{c}{uu} x^{(v)}(ww)$

(yy) (d) **[Math Processing Error]** (rrr)

A.  $\tan\left(\frac{y}{2x}\right) = c - \frac{1}{2x^2}$

B.  $\frac{\tan y}{x} = c + \frac{1}{x}$

$$C. \cos\left(\frac{y}{x}\right) = 1 + \frac{c}{x}$$

$$D. x^2 = (c + x^2) \frac{\tan y}{x}$$

**Answer: A**



**Watch Video Solution**

**35.** The solution of the differential equation

$$2x^2y \frac{dy}{dx} = \tan(x^2y^2) - 2xy^2, \text{ given } x = 1, y = \frac{\pi}{2}, \text{ is}$$

A.  $\sin^2 y^2 = e^{x+1}$

B.  $\sin(x^2y^2) = x$

C.  $\cos x^2y^2 + x = 0$

D.  $\sin(x^2y^2) = e^{x-1}$

**Answer: D**



**Watch Video Solution**

36. Solution of the differential equation

$$\left\{ \frac{1}{x} - \frac{y^2}{(x-y)^2} \right\} dx + \left\{ \frac{x^2}{(x-y)^2} - \frac{1}{y} \right\} dy = 0 \quad \text{is} \quad \text{(a)}$$

$$(b)(c) \ln \left| \frac{x}{y} \right| + (d)(e) \frac{x}{f} y (g)(h) + (i) \frac{(j)dy}{k} ((l)x - y)(m)(n) = c(o) \quad \text{(p)} \quad \text{(b)}$$

[Math Processing Error] (ii) (c)

$$(d)(e) \ln |(f)xy| = c + (g) \frac{(h)xy}{i} ((j)x - y)(k)(l)(m) \quad \text{(n)} \quad \text{(d) None of}$$

these

A.  $\ln \left| \frac{x}{y} \right| + \frac{xy}{x-y} = c$

B.  $\frac{xy}{x-y} = ce^{x/y}$

C.  $\ln |xy| + \frac{x^4 y^4}{4} = C$

D. None of these

Answer: A



Watch Video Solution

37. The solution of differential equation

$$(2y + xy^3) dx + (x + x^2 y^2) dy = 0 \quad \text{is} \quad \text{(a)}$$

$$(b)(c)(d)x^{(e)2(f)}(g)y + (h)\frac{(i)(j)x^{(k)3(l)}(m)(n)y^{(o)3(p)}(q)}{r}3(s)(t) = c$$

(v) (b)

$$(w)(x)x(y)y^{(z)2(aa)}(bb) + (cc)\frac{(dd)(ee)x^{(ff)3(gg)}(hh)(ii)y^{(jj)3(kk)}(ll)}{mm}3$$

(qq) (c)

$$(d)(e)(f)x^{(g)2(h)}(i)y + (j)\frac{(k)(l)x^{(m)4(n)}(o)(p)y^{(q)4(r)}(s)}{t}4(u)(v) = c$$

(x) (d) None of these

A.  $x^2y + \frac{x^3y^3}{3} = c$

B.  $xy^2 + \frac{x^3y^3}{3} = c$

C.  $x^2y + \frac{x^4y^4}{4} = c$

D. None of these

**Answer: A**

 [Watch Video Solution](#)

38. Find the solution of  $ye^{-\frac{x}{y}}dx - \left(xe^{-\frac{x}{y}} + y^3\right)dy = 0$  is

 [Watch Video Solution](#)



39. Find the curve satisfying the equation  $\frac{dy}{dx} = \frac{y(x + y^3)}{x(y^3 - x)}$  and passing through the point  $(4, -2)$  is

 [Watch Video Solution](#)

40. Find the solution of differential equation

$$\frac{x + y \frac{dy}{dx}}{y - x \frac{dy}{dx}} = \left( \frac{x}{y^3} \right) (\cos^2(x^2 + y^2))$$

 [Watch Video Solution](#)

41. The solution of the differential equation  $\frac{dy}{dx} = \frac{3x^2y^4 + 2xy}{x^2 - 2x^3y^3}$  is (a)

(b)(c)(d)  $\frac{(e)(f)y^{(g)2(h)}(i)}{j}x(k)(l) - (m)x^{(n)3(o)}(p)(q)y^{(r)2(s)}(t) = c(u)$

(v) (w) **[Math Processing Error]** (ww) (xx)

(yy)(zz)(aaa)  $\frac{(bbb)(c)x^{(ddd)2(eee)}(fff)}{ggg}y(hhh)(iii) + (jjj)x^{(kkk)3(lll)}(m)$

(sss) (d)

(ttt)(U)(V)  $\frac{(www)(\times x)x^{(yyy)^2(zzz)}(aaaa)}{bbbb}$  ((ccc)3y)(dddd)(eeee) -

(oooo)

A.  $\frac{y^2}{x} - x^3y^2 = c$

B.  $\frac{x^2}{y^2} + x^3y^3 = c$

C.  $\frac{x^2}{y} + x^3y^2 = c$

D.  $\frac{x^2}{3y} - 2x^3y^2 = c$

**Answer: C**



**Watch Video Solution**

**42.** The solution of the differential equation

$\left\{1 + x\sqrt{(x^2 + y^2)}\right\}dx + \left\{\sqrt{(x^2 + y^2)} - 1\right\}ydy = 0$  is equal to

A.  $x^2 + \frac{y^2}{2} + \frac{1}{3}(x^2 + y^2)^{3/2} = c$

B.  $x - \frac{y^3}{3} + \frac{1}{2}(x^2 + y^2)^{1/2} = c$

C.  $x - \frac{y^2}{2} + \frac{1}{3}(x^2 + y^2)^{3/2} = c$

D. None of these

Answer: C



Watch Video Solution

43. The solution of differential equation  $\frac{y(2x^4 + y)dy}{dx} = (1 - 4xy^2)x^2$  is given by (a)

(b)  $3(d)(e) \left( (f)(g)(h)x^{(i)2(j)}(k)y(l) \right)^{(m)2(n)}(o) + (p)y^{(q)3(r)}(s) - (t)$

(y) (z) [Math Processing Error] (zz) (aaa) [Math Processing Error] (nnnn)

(d) None of these

A.  $3(x^2y^2)^2 + y^3 - x^3 = c$

B.  $xy^2 + \frac{y^3}{3} - \frac{x^3}{3} + c = 0$

C.  $\frac{2}{5}yx^5 + \frac{y^3}{3} = \frac{x^3}{3} - \frac{4xy^3}{3} + c$

D. None of these

Answer: A



Watch Video Solution

44. The solution of the differential equation

$$(x \cot y + \log \cos x)dy + (\log \sin y - y \tan x)dx = 0 \quad \text{is} \quad (a)$$

$$(b)(c)(d)(e)((f)(g)\sin x(h))^{(i)y(j)}(k)(l)(m)((n)(o)\cos y(p))^{(q)x(r)}(s) =$$

(u) (v) *[Math Processing Error]* (pp) (qq) *[Math Processing Error]* (kkk) (d)

None of these

A.  $(\sin x)^y(\cos y)^x = c$

B.  $(\sin y)^x(\cos x)^y = c$

C.  $(\sin x)^x(\cos y)^y = c$

D. None of these

Answer: B



Watch Video Solution

45. If  $y = (e^y - x)^{-1}$ , where  $y(0) = 0$ , then  $y$  is expressed explicitly as

(a)  $\frac{1}{e} \ln \left( (h)(i)1 + (j)x^{(k)2(l)}(m)(n) \right) (o) (p) (b)$

(q)(r)  $\ln \left( (s)(t)1 + (u)x^{(v)2(w)}(x)(y) \right) (z) (aa) (c)$

(d)(e)  $\ln \left( (f)(g)x + \sqrt{(h)(i)1 + (j)x^{(k)2(l)}(m)(n)(o)(p)} \right) (q) (r) (d)$

*[Math Processing Error]* (gg)

A.  $\frac{1}{2} \log_e (1 + x^2)$

B.  $\log_e (1 + x^2)$

C.  $\log_e \left( x + \sqrt{1 + x^2} \right) = c$

D. None of these

**Answer: C**

 **Watch Video Solution**

46. The general solution of the differential equation,

$y' + y\varphi'(x) - \varphi'(x)\varphi(x) = 0$ , where  $\varphi(x)$  is a known function, is

A.  $y = ce^{-\phi(x)} + \phi(x) - 1$

B.  $y = ce^{+\phi(x)} + \phi(x) - 1$

C.  $y = ce^{-\phi(x)} - \phi(x) + 1$

D.  $y = ce^{-\phi(x)} + \phi(x) + 1$

**Answer: A**



**Watch Video Solution**

**47.** The integrating factor of the differential equation

$$\frac{dy}{dx}(x(\log)x) + y = 2(\log)x \text{ is given by}$$

A.  $x$

B.  $e^x$

C.  $\log(x)$

D.  $\log(\log(x))$

**Answer: C**



Watch Video Solution

48. The solution of the differential equation

$$x(x^2 + 1) \left( \frac{dy}{dx} \right) = y(1 - x^2) + x^3 \log x$$
 is

A.  $y(x^2 + 1) / x = \frac{1}{4}x^2 \log x + \frac{1}{2}x^2 + c$

B.  $y^2(x^2 - 1) / x = \frac{1}{2}x^2 \log x - \frac{1}{4}x^2 + c$

C.  $y(x^2 + 1)x = \frac{1}{2}x^2 \log x - \frac{1}{4}x^2 + c$

D. None of these

Answer: C



Watch Video Solution

49. Integrating factor of the differential equation  $\cos x \frac{dy}{dx} + y \sin x = 1$

is

A.  $\cos x$

B.  $\tan x$

C.  $\sec x$

D.  $\sin x$

**Answer: C**



**Watch Video Solution**

50. Solution of the equation  $\cos^2 x \left( \frac{dy}{dx} \right) - (\tan 2x)y = \cos^4 x$ , when  $y(\pi/6) = (3\sqrt{3})/8$  is

A.  $y = \tan 2x \cos^2 x$

B.  $y = \cot 2x \cos^2 x$

C.  $y = \frac{1}{2} \tan 2x \cos^2 x$

D.  $y = \frac{1}{2} \cot 2x \cos^2 x$

**Answer: C**



**Watch Video Solution**



51. If integrating factor of  $x(1 - x^2)dy + (2x^2y - y - ax^3)dx = 0$  is  $e^{\int p dx}$ , then  $P$  is equal to

A.  $\frac{2x^2 - ax^2}{x(1 - x^2)}$

B.  $2x^3 - 1$

C.  $\frac{2x^2 - a}{ax^3}$

D.  $\frac{2x^2 - 1}{x(1 - x^2)}$

**Answer: D**



**Watch Video Solution**

52. A function  $y = f(x)$  satisfies

$$(x + 1)f'(x) - 2(x^2 + x)f(x) = \frac{e^{x^2} - 2}{(x + 1)}, \forall x > -1. \text{ If } f(0) = 5,$$

then  $f(x)$  is

A.  $\left(\frac{3x + 5}{x + 1}\right)e^{x^2}$

B.  $\left(\frac{6x + 5}{x + 1}\right)e^{x^2}$

C.  $\left(\frac{6x + 5}{(x + 1)^2}\right)e^{x^2}$

D.  $\left(\frac{5 - 6x}{x + 1}\right)e^{x^2}$

**Answer: B**



**Watch Video Solution**

53. The general solution of the equation  $\frac{dy}{dx} = 1 + xy$  is (a)

(b)(c)  $y = c(d)e^{e(f)} - (g) \left( (h)(i) \frac{x^{((j)2(k)) (l)}}{m} - 2(n)(o)(p) \right) (q)(r)$  (s)

(b)

(t)(u)  $y = c(v)e^{w(x)}(y) \left( (z)(aa) \frac{x^{((bb)2(cc)) (dd)}}{ee} - 2(ff)(gg)(hh) \right) (ii)(jj)$

(kk)

(c)

(d)(e)  $y = ((f)(g)x + c(h)), (i)e^j(k) - (l) \left( (m)(n) \frac{x^{((o)2(p)) (q)}}{r} - 2(s)(t) \right)$

(x) (d) None of these

A.  $y = ce^{-x^2/2}$

B.  $y = ce^{x^2/2}$

C.  $y = (x + c), e^{-x^2/2}$

D. None of these

**Answer: D**

 [Watch Video Solution](#)

54. The solution of the differential equation  $(x + 2y^3) \frac{dy}{dx} = y$  is

A.  $\frac{x}{y^2} = y + c$

B.  $\frac{x}{y} = y^2 + c$

C.  $\frac{x^2}{y} = y^2 + c$

D.  $\frac{y}{x} = x^2 + c$

**Answer: B**

 [Watch Video Solution](#)

55. The solution of the differential equation

$$x^2 \frac{dy}{dx} \cos\left(\frac{1}{x}\right) - y \sin\left(\frac{1}{x}\right) = -1, \text{ where } y \rightarrow -1 \text{ as } x \rightarrow \infty \text{ is}$$

A.  $y = \sin\left(\frac{1}{x}\right) - \cos\left(\frac{1}{x}\right)$

B.  $y = \frac{x+1}{x \sin\left(\frac{1}{x}\right)}$

C.  $y = \cos\left(\frac{1}{x}\right) + \sin\left(\frac{1}{x}\right)$

D.  $y = \frac{x+1}{x \cos(1/x)}$

**Answer: A**



**Watch Video Solution**

56. The solution of  $\frac{dy}{dx} = \frac{x^2 + y^2 + 1}{2xy}$  satisfying  $y(1) = 1$  is

A. a system of parabolas

B. a system of circles

C.  $y^2 = x(1+x) - 1$

D.  $(x - 2)^2 + (y - 3)^2 = 5$

Answer: C



Watch Video Solution

57. The solution of the differential equation  $\frac{dy}{dx} = \frac{1}{xy[x^2 \sin y^2 + 1]}$  is

(a)

(b)  $\cos y - \sin y - 2C e^{-y^2}$

(c)  $\cos y^2 - \sin y^2 - 2C e^{-y^2}$  (aa) *[Math Processing Error]* (zz) (aaa)

(d)  $\cos x^2 - \sin x^2 - 2C e^{-x^2}$  (bbb)  $\cos(hhh)y^{(iii)^2} - \sin(lll)y^{(mmm)}$

(eee) (zzz) (aaaa) None of these

A.  $x^2 (\cos y^2 - \sin y^2 - 2C e^{-y^2}) = 2$

B.  $y^2 (\cos x^2 - \sin y^2 - 2C e^{-y^2}) = 4C$

C. None of these

D. a system of circles

Answer: A



Watch Video Solution

58. Find the equation of a curve passing through  $\left(2, \frac{7}{2}\right)$  and having gradient  $1 - \frac{1}{x^2}$  at  $(x, y)$  is

A.  $y = x^2 + x + 1$

B.  $xy = x^2 + x + 1$

C.  $xy = x + 1$

D. None of these

Answer: B



Watch Video Solution

59. Which of the following is not the differential equation of family of curves whose tangent form an angle of  $\frac{\pi}{4}$  with (a)

(b) (c) (d)  $\frac{(e)dy}{f} = (g)dx$  (h) (i)  $= (j) \frac{(k)x - y}{l} = ((m)x + y)(n)$  (o) (p) (q)

$$(b) \frac{(u)dy}{v}((w)dx)(x)(y) = (z) \frac{x}{aa}((bb)x - y)(cc)(dd)(ee) (ff)$$

$$(c) \frac{(g)dy}{h}((i)dx)(j)(k) = (l) \frac{(m)x + y}{n}((o)y - x)(p)(q)(r)$$

(s) (d) None of these

A.  $\frac{dy}{dx} = \frac{x - y}{x + y}$

B.  $\frac{dy}{dx} = \frac{x}{x - y}$

C.  $\frac{dy}{dx} = \frac{x + y}{y - x}$

D. None of these

**Answer: B**



**Watch Video Solution**

**60.** Tangent to a curve intercepts the y-axis at a point  $P$ . A line perpendicular to this tangent through  $P$  passes through another point  $(1,0)$ . The differential equation of the curve is (a)

$$(b)(c)y(d) \frac{(e)dy}{f}((g)dx)(h)(i) - x(j)(k) \left( (l)(m)(n) \frac{(o)dy}{p}((q)dx)(r)(s)(t) \right)$$

(d)(e)y(f)  $\frac{(g)dx}{h}$  ((i)dy)(j)(k) + x = 1(l) (m) (d) None of these

A.  $y \frac{dy}{dx} - x \left( \frac{dy}{dx} \right)^2 = 1$

B.  $\frac{xd^2y}{dx^2} + \left( \frac{dy}{dx} \right)^2 = 0$

C.  $y \frac{dy}{dx} + x = 1$

D. None of these

**Answer: A**



**Watch Video Solution**

61. Orthogonal trajectories of family of the curve  $x^{\frac{2}{3}} + y^{\frac{2}{3}} = a \left( \frac{2}{3} \right)$ ,

where  $a$  is any arbitrary constant, is

A.  $x^{2/3} - y^{2/3} = c$

B.  $x^{4/3} - y^{4/3} = c$

C.  $x^{4/3} + y^{4/3} = c$



D.  $x^{\frac{1}{3}} - y^{1/3} = c$

**Answer: B**



[Watch Video Solution](#)

62. The curve in the first quadrant for which the normal at any point  $(x, y)$  and the line joining the origin to that point form an isosceles triangle with the x-axis as base is (a) an ellipse (b) a rectangular hyperbola (c) a circle (d) None of these

A. an ellipse

B. a rectangular hyperbola

C. a circle

D. None of these

**Answer: B**



[Watch Video Solution](#)

63. The equation of the curve which is such that the portion of the axis of  $x$  cut off between the origin and tangent at any point is proportional to the ordinate of that point is (a)  $(b)(c)x = y(a - b \log x)(d)$  (e) (f)  $(g)(h)\log x = b(i)y^{(j)2(k)}(l) + a(m)$  (n) (o)  $(p)(q)(r)x^{(s)2(t)}(u) = y(a - b \log y)(v)$  (w) (d) None of these

A.  $x = y(a - b \log y)$

B.  $\log_x = by^2 + a$

C.  $x^2 = y(a - b \log y)$

D. None of these

**Answer: A**



**Watch Video Solution**

64. The family of curves represented by  $\frac{dy}{dx} = \frac{x^2 + x + 1}{y^2 + y + 1}$  and the family represented by  $\frac{dy}{dx} + \frac{y^2 + y + 1}{x^2 + x + 1} = 0$

- A. Touch each other
- B. Are orthogonal
- C. Are one and the same
- D. None of these

**Answer: B**

 [Watch Video Solution](#)

65. A normal at  $P(x, y)$  on a curve meets the x-axis at  $Q$  and  $N$  is the foot of the ordinate at  $P$ . If  $NQ = \frac{x(1 + y^2)}{1 + x^2}$ , then the equation of curve given that it passes through the point  $(3, 1)$  is (a)

(b)  $(c)(d)x^{(e)2(f)}(g) - (h)y^{(i)2(j)}(k) = 8(l)$  (m) (b)

(n)(o)(p)x<sup>(q)2(r)</sup>(s) + 2(t)y<sup>(u)2(v)</sup>(w) = 11(x) (y) (c)

(d)(e)(f)x<sup>(g)2(h)</sup>(i) - 5(j)y<sup>(k)2(l)</sup>(m) = 4(n) (o) (d) None of these

A.  $x^2 - y^2 = 8$

B.  $x^2 + 2y^2 = 11$

C.  $x^2 - 5y^2 = 4$

D. None of these

**Answer: C**



**Watch Video Solution**

**66.** A curve is such that the mid-point of the portion of the tangent intercepted between the point where the tangent is drawn and the point where the tangent meets the y-axis lies on the line  $y = x$ . If the curve passes through  $(1, 0)$ , then the curve is (a)

(b)  $2y = (d)x^{(e)2(f)}(g) - x(h)$  (i) (b)

(j)  $(k)y = (l)x^{(m)2(n)}(o) - x(p)$  (q) (c)

(d)  $(e)y = x - (f)x^{(g)2(h)}(i)(j)$  (k) (d)

(l)  $(m)y = 2\left((n)(o)x - (p)x^{(q)2(r)}(s)(t)\right)(u)(v)$

A.  $2y = x^2 - x$

B.  $y = x^2 - x$

C.  $y = x - x^2$

D.  $y = 2(x - x^2)$

**Answer: C**



**Watch Video Solution**

67. The normal to a curve at  $P(x, y)$  meet the x-axis at  $G$ . If the distance of  $G$  from the origin is twice the abscissa of  $P$ , then the curve is a (a) parabola (b) circle (c) hyperbola (d) ellipse

A. parabola

B. circle

C. hyperbola

D. ellipse

**Answer: C**



**Watch Video Solution**

68. The x-intercept of the tangent to a curve is equal to the ordinate of the point of contact. The equation of the curve through the point (1,1) is

A.  $ye^{x/y} = e$

B.  $xe^{x/y} = e$

C.  $xe^{y/x} = e$

D.  $ye^{y/x} = e$

Answer: A



Watch Video Solution

69. The equation of a curve passing through (1,0) for which the product of the abscissa of a point  $P$  and the intercept made by a normal at  $P$  on the x-axis equal twice the square of the radius vector of the point  $P$  is (a)

(b)(c)(d)  $x^{(e)2(f)}(g) + (h)y^{(i)2(j)}(k) = (l)x^{(m)4(n)}(o)(p)$  (q) (b)

(r)(s)(t)  $x^{(u)2(v)}(w) + (x)y^{(y)2(z)}(aa) = 2(bb)x^{(cc)4(dd)}(ee)(ff)$

(gg) (c) (d)(e)(f)  $x^{(g)2(h)}(i) + (j)y^{(k)2(l)}(m) = 4(n)x^{(o)4(p)}(q)(r)$

(s) (d) None of these

A.  $x^2 + y^2 = x^4$

B.  $x^2 + y^2 = 2x^4$

C.  $x^2 + y^2 = 4x^4$

D. None of these

Answer: A



Watch Video Solution

70. The curve with the property that the projection of the ordinate on the normal is constant and has a length equal to  $a$  is (a)

(b)  $a \ln \left( \sqrt{(f)(g)(h)y^{(i)2(j)}(k) - (l)a^{(m)2(n)}(o)(p)(q) + y(r)} \right)$

(t) (u)

(v)  $(w)x + \sqrt{(x)(y)(z)a^{(aa)2(bb)}(cc) - (dd)y^{(ee)2(ff)}(gg)(hh)(ii)} = c(j)$

(kk) (ll)  $(mm)(\cap)(\infty)(pp)((qq)(rr)y - a(ss))^{(tt)2(uu)}(vv) = cx(ww)$

(xx) (yy)

(zz)  $(aaa)ay = (bbb)(c)\tan^{(ddd)(eee)-1}(fff)(ggg)((hhh)(iii)x + c(jjj))(l)$

(III)

A.  $a \ln \left( \sqrt{y^2 - a^2} \right) = x + c$

B.  $x + \sqrt{a^2 - y^2} = c$

C.  $(y - a)^2 = cx$

D.  $ay = \tan^{-1}(x + c)$

**Answer: A**

 **Watch Video Solution**

71. Spherical rain drop evaporates at a rate proportional to its surface area. The differential equation corresponding to the rate of change of the radius of the rain drop if the constant of proportionality is  $K > 0$  is (a)

(b)  $(c) \frac{(e)dy}{f} ((g)dt)(h)(i) + K = 0(j)$  (k) (b)

(l)(m)(n)  $\frac{(o)dr}{p} ((q)dt)(r)(s) - K = 0(t)$  (u) (c)

(d)(e)(f)  $\frac{(g)dr}{h} ((i)dt)(j)(k) = Kr(l)$  (m) (d) None of these

A.  $\frac{dr}{dt} + K = 0$

B.  $\frac{dr}{dt} - K = 0$



C.  $\frac{dr}{dt} = Kr$

D. None of these

**Answer: A**



**Watch Video Solution**

72. Water is drained from a vertical cylindrical tank by opening a valve at the base of the tank. It is known that the rate at which the water level drops is proportional to the square root of water depth  $y$ , where the constant of proportionality  $k > 0$  depends on the acceleration due to gravity and the geometry of the hole. If  $t$  is measured in minutes and  $k = \frac{1}{15}$ , then the time to drain the tank if the water is 4 m deep to start with is (a) 30 min (b) 45 min (c) 60 min (d) 80 min

A. 30 min

B. 45 min

C. 60 min

D. 80 min

**Answer: C**



[Watch Video Solution](#)

**73.** The population of a country increases at a rate proportional to the number of inhabitants.  $f$  is the population which doubles in 30 years, then the population will triple in approximately. (a) 30 years (b) 45 years (c) 48 years (d) 54 years

A. 30 years

B. 45 years

C. 48 years

D. 54 years

**Answer: C**



[Watch Video Solution](#)

74. An object falling from rest in air is subject not only to the gravitational force but also to air resistance. Assume that the air resistance is proportional to the velocity with constant of proportionality as  $k > 0$ , and acts in a direction opposite to motion  $\left(g = 9.8 \frac{m}{s^2}\right)$ . Then velocity cannot exceed. (c) (d)(e)  $9.8/km/s$ (f) (g) (b) (h)(i)  $98/km/s$ (j) (k) (c) (d)(e)(f)  $\frac{k}{g}$ ((h)  $9.8$ )(i)(j)  $m/s$ (k) (l) (d) None of these

A.  $9.8/km/s$

B.  $98/k m/s$

C.  $\frac{k}{9.8}m/x$

D. None of these

**Answer: A**



**Watch Video Solution**

**Exercise (Multiple)**

1. Which one of the following function(s) is/are homogeneous?

A.  $f(x, y) = \frac{x - y}{x^2 + y^2}$

B.  $f(x, y) = x^{\frac{1}{3}}y^{-\frac{2}{3}}\frac{\tan^{-1} x}{y}$

C.  $f(x, y) = x \left( \ln \sqrt{x^2 + y^2} \right) - \ln y + ye^{x/y}$

D.  $f(x, y) = x \left[ \ln (2x^2 + y^2)x - \ln (x + y) \right] + y^2 \frac{\tan(x + 2y)}{3x - y}$

Answer: A::B::C



Watch Video Solution

2. For the differential equation whose solution is  $(x - h)^2 + (y - k)^2 = a^2$  ( $a$  is a constant), its (a) order is 2 (b) order is 3 (c) degree is 2 (d) degree is 3

A. order is 2

B. order is 3

C. degree is 2

D. degree is 3

**Answer: A::C**



[Watch Video Solution](#)

3. The equation of the curve satisfying the differential equation

$$y \left( \frac{dy}{dx} \right)^2 + (x - y) \frac{dy}{dx} - x = 0 \text{ can be a (a) circle (b) Straight line (c)}$$

Parabola (d) Ellipse

A. Circle

B. Straight line

C. Parabola

D. Ellipse

**Answer: A::B**



[Watch Video Solution](#)

4. Which of the following equation(s) is/are linear?

A.  $\frac{dy}{dx} + \frac{y}{x} = \log x$

B.  $y\frac{dy}{dx} + 4x = 0$

C.  $(2x + y^3)\frac{dy}{dx} = 3y$

D. None of these

Answer: A:C



Watch Video Solution

5. The solution of  $\frac{dy}{dx} = \frac{ax + h}{by + k}$  represents a parabola when

A.  $a = 0, b \neq 0$

B.  $a \neq 0, b \neq 0$

C.  $b = 0, a \neq 0$

D.  $a = 0, b \in R$

Answer: A::C



Watch Video Solution

6. The equation of the curve satisfying the differential equation  $y_2(x^2 + 1) = 2xy_1$  passing through the point (0,1) and having slope of tangent at  $x = 0$  as 3 (where  $y_2$  and  $y_1$  represent 2nd and 1st order derivative), then (a) (b)(c) $y = f((d)x(e))(f)$  (g) is a strictly increasing function (h) (i)(j) $y = f((k)x(l))(m)$  (n) is a non-monotonic function (o) (p)(q) $y = f((r)x(s))(t)$  (u) has a three distinct real roots (v) (w)(x) $y = f((y)x(z))(aa)$  (bb) has only one negative root.

A.  $y = f(x)$  is a strictly increasing function

B.  $y = f(x)$  is a non-monoatomic function

C.  $y = f(x)$  has three distinct real root

D.  $y = f(x)$  has only one negative root

Answer: A::D



7. Identify the statement(s) which is/are true.

A.  $f(x, y) = e^{y/x} = e^{\frac{y}{x}} + \frac{\tan y}{x}$  is a homogeneous of degree zero.

B.  $x \ln \frac{y}{x} dx + \frac{y^2 \sin^{-1} y}{x} dx = 0$  is a homogenous differential equation.

C.  $f(x, y) = x^2 + \sin x \cos y$  is a non-homogenous

D.  $(x^2 + y^2) dx - (xy^2 - y^3) dy = 0$  is a homogenous differential equationn.

Answer: A::B::C



8. The graph of the function  $y = f(x)$  passing through the point  $(0, 1)$  and satisfying the differential equation  $\frac{dy}{dx} + y \cos x = \cos x$  is such that



- A. It is constant function
- B. It is periodic
- C. it is neither an even nor an odd function.
- D. it is continuous and differentiable for all x

**Answer: A::B::D**

 [Watch Video Solution](#)

9. If  $f(x)$ ,  $g(x)$  be twice differentiable functions on  $[0,2]$  satisfying  $f''(x) = g''(x)$ ,  $f'(1) = 2g'(1) = 4$  and  $f(2) = 3g(2) = 9$ , then  $f(x) - g(x)$  at  $x = 4$  equals (A) 0 (B) 10 (C) 8 (D) 2

- A.  $f(4) - g(4) = 10$
- B.  $|f(x) - g(x)| < 2 \Rightarrow -2 < x < 0$
- C.  $f(2) = g(2) \Rightarrow x = -1$
- D.  $f(x) - g(x) = 2x$  has real root

Answer: A::B::C



Watch Video Solution

10. The solution of the differential equation  $(x^2y^2 - 1)dy + 2xy^3dx = 0$

is (a) (b)(c)  $1 + (d)x^{(e)2(f)}(g)(h)y^{(i)2(j)}(k) = cx(l)$  (m) (b)

(n)(o)  $1 + (p)x^{(q)2(r)}(s)(t)y^{(u)2(v)}(w) = cy(x)$  (y) (c) (d)(e)  $y = 0(f)$

(g) (d) (h)(i)  $y = - (j)\frac{1}{k} \left( (l)(m)x^{(n)2(o)}(p) \right) (q)(r)(s) (t)$

A.  $1 + x^2y^2 = cx$

B.  $1 + x^2y^2 = cy$

C.  $y = 0$

D.  $y = -\frac{1}{x^2}$

Answer: B



Watch Video Solution

11.  $y = ae^{-\frac{1}{x}} + b$  is a solution of  $\frac{dy}{dx} = \frac{y}{x^2}$ , then (a)  $a \in \mathbb{R}$  (b)  $a = 0$  (c)  $b = 1$  (d)  $b = 0$  (e)  $a$  takes finite number of values (f)  $b = 1$  (g)  $b = 0$  (h)  $a$  takes finite number of values (i)  $a \in \mathbb{R}$  (j)  $a = 0$

A.  $x \in \mathbb{R} - \{0\}$

B.  $b = 0$

C.  $b = 1$

D.  $a$  takes finite number of values

**Answer: A:B**



**Watch Video Solution**

12. For equation of the curve whose subnormal is constant, then (a) its eccentricity is 1 (b) its eccentricity is  $\sqrt{2}$  (c) its axis is the x-axis (d) its axis is the y-axis (e) its eccentricity is  $\sqrt{2}$  (f) its axis is the x-axis (g) its axis is the y-axis (h) its eccentricity is 1 (i) its axis is the x-axis (j) its axis is the y-axis (k) its axis is the x-axis (l) its axis is the y-axis

A. its eccentricity is 1

B. its eccentricity is  $\sqrt{2}$

C. its axis is the x-axis

D. its axis is the y-axis

**Answer: B**

 [Watch Video Solution](#)

13. The solution of  $\frac{xdx + ydy}{xdy - ydx} = \sqrt{\frac{1 - x^2 - y^2}{x^2 + y^2}}$  is

A.  $\sqrt{x^2 + y^2} = \sin\{(\tan^{-1} y/x) + C\}$

B.  $\sqrt{x^2 + y^2} = \cos\{(\tan^{-1} y/x) + C\}$

C.  $\sqrt{x^2 + y^2} = (\tan(\sin^{-1} y/x) + C)$

D.  $y = x \tan\left(c + \sin^{-1} \sqrt{x^2 + y}\right)$

**Answer: A:D**

 [Watch Video Solution](#)

14. The curve for which the length of the normal is equal to the length of the radius vector is/are (a) circles (b) rectangular hyperbola (c) ellipses (d) straight lines

A. circles

B. rectangular hyperbola

C. ellipses

D. straight lines

**Answer: A::B**



**Watch Video Solution**

15. In which of the following differential equation degree is not defined?

A.  $\frac{d^2y}{dx^2} + 3\left(\frac{dy}{dx}\right)^2 = x \log\left(\frac{d^2y}{dx^2}\right)$

B.  $\frac{d^2y}{dx^2} + \left(\frac{dy}{dx}\right)^2 = x \sin\left(\frac{d^2y}{dx^2}\right)$

C.  $x = \sin\left(\frac{dy}{dx} - 2y\right), |x| < 1$

$$D. x - 2y = \log\left(\frac{dy}{dx}\right)$$

**Answer: A&B**



**Watch Video Solution**

16. If  $y = f(x)$  is the solution of equation  $yx + dy = -e^x y^2 dy$ ,  $f(0)=1$  and area bounded by the curve  $y = f(x)$ ,  $y = e^x$  and  $x=1$  is A, then

A. curve  $y=f(x)$  is passing through  $(-2, e)$ .

B. Curve  $y = f(x)$  is passing through  $(1, 1/e)$

C. curve  $y = f(x)$  is passing through  $(1, 1/3)$

$$D. A = e + \frac{2}{\sqrt{e}} - 3$$

**Answer: A::D**



**Watch Video Solution**

17. A particle falls in a medium whose resistance is proportional to the square of the velocity of the particles. If the differential equation of the free fall is  $\frac{dv}{dt} = g - kv^2$  ( $k$  is constant) then

A.  $v = 2\sqrt{\frac{g}{k}} \frac{e^{2t\sqrt{g/k}} + 1}{e^{2t\sqrt{g/k}} - 1}$

B.  $v = \sqrt{\frac{g}{k}} \frac{e^{2t\sqrt{gk}} - 1}{e^{2t\sqrt{gk}} + 1}$

C.  $v \rightarrow 0$  as  $t \rightarrow \infty$

D.  $v \rightarrow \sqrt{\frac{g}{k}}$  as  $t \rightarrow \infty$

Answer: B::D



Watch Video Solution

### Exercise (Comprehension)

1. For certain curves  $y = f(x)$  satisfying  $\frac{d^2y}{dx^2} = 6x - 4$ ,  $f(x)$  has local minimum value 5 when  $x=1$ . find Number of critical point for  $y=f(x)$  for  $x \in [0,2]$  (a) 0 (b)1. c).2 d) 3

A. 0

B. 1

C. 2

D. 3

**Answer: C**



**Watch Video Solution**

2. For certain curves  $y = f(x)$  satisfying  $\frac{d^2y}{dx^2} = 6x - 4$ ,  $f(x)$  has local minimum value 5 when  $x=1$ . 9. Number of critical point for  $y=f(x)$  for  $x \in [0,2]$  (a) 0 (b)1. c).2 d) 3 10. Global minimum value  $y = f(x)$  for  $x \in [0,2]$  is (a)5 (b)7 (c)8 d) 9 11 Global maximum value of  $y = f(x)$  for  $x \in [0,2]$  is (a) 5 (b) 7 (c) 8 (d) 9

A. 5

B. 7

C. 8



D. 9

**Answer: A**



[Watch Video Solution](#)

3. For certain curve  $y = f(x)$  satisfying  $\frac{d^2y}{dx^2} = 6x - 4$ ,  $f(x)$  has local minimum value 5 when  $x = 1$

Global minimum value of  $y = f(x)$  for  $x \in [0, 2]$  is

A. 5

B. 7

C. 8

D. 9

**Answer: B**



[Watch Video Solution](#)

4. The differential equation  $y = px + f(p)$ , .....(i) where  $p = \frac{dy}{dx}$ , is known as Clairout's equation. To solve equation i) differentiate it with respect to  $x$ , which gives either  $\frac{dp}{dx} = 0 \Rightarrow p = c$ .....(ii) or  $x + f'(p) = 0$ .....(iii) Which of the following is true about solutions of differential equation  $y = xy' + \sqrt{1 + y'^2}$ ?

- A. the general solution of equation is family of parabolas
- B. the general solution of equation is family of circles
- C. the singular solution of equation is circle
- D. the singular solution of equation is ellipse

**Answer: C**

 [Watch Video Solution](#)

5. The differential equation  $y = px + f(p)$ , .....(i) where  $p = \frac{dy}{dx}$ , is known as Clairout's equation. To solve equation i) differentiate it with respect to  $x$ , which gives either

$$\frac{dp}{dx} = 0 \Rightarrow p = c \dots \dots \dots (ii)$$

$$\text{or } x + f^i(p) = 0 \dots \dots \dots (iii)$$

The number of solution of the equation  $f(x) = -1$  and the singular

solution of the equation  $y = x \frac{dy}{dx} + \left( \frac{dy}{dx} \right)^2$  is

A. 1

B. 2

C. 4

D. 0

**Answer: B**



[Watch Video Solution](#)

6. The differential equation  $y = px + f(p)$ , .....(i)

where  $p = \frac{dy}{dx}$ , is known as Clairout's equation. To solve equation i)

differentiate it with respect to x, which gives either

$$\frac{dp}{dx} = 0 \Rightarrow p = c \dots \dots \dots (ii)$$

or  $x + f^i(p) = 0$ .....(iii)

The singular solution of the differential equation  $y = mx + m - m^3$ ,

where  $m = \frac{dy}{dx}$ , passes through the point.

A. (0,0)

B. (0,1)

C. (1,0)

D. (-1,0)

**Answer: D**



[Watch Video Solution](#)

7. Let  $f(x)$  be a non-positive continuous function and

$F(x) = \int_0^x f(t)dt \forall x \geq 0$  and  $f(x) \geq cF(x)$  where  $c < 0$  and let

$g: [0, \infty) \rightarrow R$  be a function such that  $\frac{dg(x)}{dx} < g(x) \forall x > 0$  and

$g(0) = 0$

The total number of root(s) of the equation  $f(x) = g(x)$  is/ are

A.  $\infty$

B. 1

C. 2

D. 0

**Answer: B**



**Watch Video Solution**

8. Let  $f(x)$  be a non-positive continuous function and  $F(x) = \int_0^x f(t)dt \forall x \geq 0$  and  $f(x) \geq cF(x)$  where  $c < 0$  and let  $g: [0, \infty) \rightarrow R$  be a function such that  $\frac{dg(x)}{dx} < g(x) \forall x > 0$  and  $g(0) = 0$

The number of solution(s) of the equation

$$|x^2 + x - 6| = f(x) + g(x) \text{ is/are}$$

A. 2

B. 1

C. 0

D. 3

**Answer: C**



**Watch Video Solution**

9. Let  $f(x)$  be a non-positive continuous function and  $F(x) = \int_0^x f(t)dt \forall x \geq 0$  and  $f(x) \geq cF(x)$  where  $c < 0$  and let  $g: [0, \infty) \rightarrow R$  be a function such that  $\frac{dg(x)}{dx} < g(x) \forall x > 0$  and  $g(0) = 0$

The solution set of inequation  $g(x)(\cos^{-1} x - \sin^{-1} x) \leq 0$

A.  $\left[ -1, \frac{1}{\sqrt{2}} \right]$

B.  $\left[ \frac{1}{\sqrt{2}}, 1 \right]$

C.  $\left[ 0, \frac{1}{\sqrt{2}} \right]$

D.  $\left[ 0, \frac{1}{\sqrt{2}} \right]$

**Answer: A**



**Watch Video Solution**

10. A curve 'C' with negative slope through the point(0,1) lies in the I Quadrant. The tangent at any point 'P' on it meets the x-axis at 'Q'. Such that  $PQ = 1$ . Then

The curve in parametric form is

A.  $x = \cos \theta + \log_e \tan(\theta/2), y = \sin \theta$

B.  $x = -\cos \theta + \log_e \tan(\theta/2), y = \sin \theta$

C.  $x = -\cos \theta - \log_e \tan \theta/2, y = \sin \theta$

D. None of these

**Answer: C**



**Watch Video Solution**

11. A curve 'C' with negative slope through the point(0,1) lies in the I Quadrant. The tangent at any point 'P' on it meets the x-axis at 'Q'. Such that  $PQ = 1$ . Then

The area bounded by 'C' and the co-ordinate axes is

A. 1

B.  $\log_e 2$

C.  $\pi / 4$

D.  $\pi / 2$

**Answer: C**



[Watch Video Solution](#)

12. A curve 'C' with negative slope through the point(0,1) lies in the I Quadrant. The tangent at any point 'P' on it meets the x-axis at 'Q'. Such that  $PQ = 1$ . Then

The orthogonal trajectories of 'C' are



A. Circles

B. Parabolas

C. Ellipses

D. Hyperbolas

**Answer: A**



**Watch Video Solution**

13. Let  $y = f(x)$  satisfies the equation

$$f(x) = (e^{-x} + e^x) \cos x - 2x - \int_0^x (x-t)f'(t)dt$$
  $y$  satisfies the

differential equation

A.  $\frac{dy}{dx} + y = e^x(\cos x - \sin x) - e^{-x}(\cos x + \sin x)$

B.  $\frac{dy}{dx} - y = e^x(\cos x - \sin x) + e^{-x}(\cos x + \sin x)$

C.  $\frac{dy}{dx} + y = e^x(\cos x + \sin x) - e^{-x}(\cos x - \sin x)$

D.  $\frac{dy}{dx} - y = e^x(\cos x - \sin x) + e^{-x}(\cos x - \sin x)$

**Answer: A**



**Watch Video Solution**

**14.** Let  $y = f(x)$  satisfies the equation

$$f(x) = (e^{-x} + e^x)\cos x - 2x + \int_0^x (x - t)f'(t)dt$$

The value of  $f(0) + f'(0)$  equal

A.  $-1$

B.  $0$

C.  $1$

D.  $1$

**Answer: B**



**Watch Video Solution**

15. Let  $y = f(x)$  satisfies the equation

$$f(x) = (e^{-x} + e^x) \cos x - 2x + \int_0^x (x - t) f'(t) dt$$

$y$  satisfies the differential equation

A.  $e^{-x}(\cos x - \sin x) + \frac{e^x}{5}(3 \cos x + \sin x) + \frac{2}{5}e^{-x}$

B.  $e^{-x}(\cos x + \sin x) + \frac{e^x}{5}(3 \cos x - \sin x) - \frac{2}{5}e^{-x}$

C.  $e^{-x}(\cos x - \sin x) + \frac{e^x}{5}(3 \cos x - \sin x) + \frac{2}{5}e^{-x}$

D.  $e^{-x}(\cos x + \sin x) + \frac{e^x}{5}(3 \cos x - \sin x) - \frac{2}{5}e^{-x}$

**Answer: C**



**Watch Video Solution**

16. A certain radioactive material is known to decay at a rate proportional to the amount present. If initially there is 50 kg of the material present and after two hours it is observed that the material has lost 10% of its original mass, find (i) an expression for the mass of the material remaining at any time  $t$ , (ii) the mass of the material after four hours and

(iii) the time at which the material has decayed to one half of its initial mass.

A.  $N = 50e^{(1/2)(\ln 9)t}$

B.  $50e^{(1/4)(\ln 9)t}$

C.  $N = 50e^{-(\ln 0.9)t}$

D. None of these

**Answer: A**



**Watch Video Solution**

**17.** A certain radioactive material is known to decay at a rate proportional to the amount present. Initially there is 50 kg of the material present and after 2h it is observed that the material has lost 10 percent of its original mass. Based on these data answer the following questions.

The time at which the material has decayed to one half of its initial mass is

A.  $50^{-0.5 \ln 9}$

B.  $50e^{-2 \ln 9}$

C.  $50e^{-2 \ln 0.9}$

D. None of these

**Answer: C**

 [Watch Video Solution](#)

**18.** A certain radioactive material is known to decay at a rate proportional to the amount present. If initially there is 50 kg of the material present and after two hours it is observed that the material has lost 10% of its original mass, find (i) an expression for the mass of the material remaining at any time  $t$ , (ii) the mass of the material after four hours and (iii) the time at which the material has decayed to one half of its initial mass.

A.  $\frac{\ln 0.25}{\ln 0.9} h$

B.  $\frac{\ln 0.5}{\ln 0.81}h$

C.  $\frac{\ln 0.25}{\ln 0.81}h$

D. None of these

**Answer: C**



**Watch Video Solution**

**19.** Consider a tank which initially holds  $V_0$  liter of brine that contains a lb of salt. Another brine solution, containing  $b$  lb of salt per liter is poured into the tank at the rate of  $eL/\text{min}$ . The problem is to find the amount of salt in the tank at any time  $t$ .

Let  $Q$  denote the amount of salt in the tank at any time. The time rate of change of  $Q$ ,  $\frac{dQ}{dt}$ , equals the rate at which salt enters the tank at the rate at which salt leaves the tank. Salt enters the tank at the rate of  $be$  lb/min.

To determine the rate at which salt leaves the tank, we first calculate the volume of brine in the tank at any time  $t$ , which is the initial volume  $V_0$  plus the volume of brine added and minus the volume of brine removed.

Thus, the volume of brine at any time is

$$V_0 + et - ft$$

The concentration of salt in the tank at any time is  $Q/(V_0 + et - ft)$ ,

from which it follows that salt leaves the tank at the rate of

$f\left(\frac{Q}{V_0 + et - ft}\right)$  lb/min. Thus,

$$\frac{dQ}{dt} = be - f\left(\frac{Q}{V_0 + et - ft}\right)Q = be$$

A tank initially holds 100 L of a brine solution containing 20 lb of salt. At

$t=0$ , fresh water is poured into the tank at the rate of 5 L/min, while the

well-stirred mixture leaves the tank at the same rate. Then the amount of

salt in the tank after 20 min.

A.  $20/e$

B.  $10/e$

C.  $40/e^2$

D.  $5/e$  L

**Answer: A**



**Watch Video Solution**

20. A 50 L tank initially contains 10 L of fresh water, At  $t=0$ , a brine solution containing 1 lb of salt per gallon is poured into the tank at the rate of 4 L/min, while the well-stirred mixture leaves the tank at the rate of 2 L/min. Then the amount of time required for overflow to occur in

- A. 30 min
- B. 20 min
- C. 10 min
- D. 40 min

**Answer: B**



[Watch Video Solution](#)

21. A 50 L tank initially contains 10 L of fresh water, At  $t=0$ , a brine solution containing 1 lb of salt per gallon is poured into the tank at the rate of 4 L/min, while the well-stirred mixture leaves the tank at the rate of 2 L/min. Then the amount of time required for overflow to occur in



A. 20 lb

B. 50 lb

C. 30 lb

D. None of these

**Answer: D**



[Watch Video Solution](#)

## Matrix Match Type

1. A bag contains 15 balls, some are red and others are white. If a ball is drawn at random from the bag, the probability that it is red is  $\frac{1}{5}$ . Find the number of white balls in the bag.



[Watch Video Solution](#)

2. Find the derivative of  $y = \log 3x$



Watch Video Solution

3. Find the derivative of  $y = \ln x^3$



Watch Video Solution

## Exercise (Numerical)

1. If  $y = y(x)$  and it follows the relation  $4xe^{xy} = y + 5\sin^2 x$ , then  $y'(0)$  is equal to \_\_\_\_\_



Watch Video Solution

2. If  $x \frac{dy}{dx} = x^2 + y - 2$ ,  $y(1) = 1$ , then  $y(2)$  equal \_\_\_\_\_



Watch Video Solution

3. If the dependent variable  $y$  is changed to  $z$  by the substitution  $y = \tan z$  and the differential equation  $\frac{d^2y}{dx^2} = 1 + \frac{2(1+y)}{1+y^2} \left(\frac{dy}{dx}\right)^2$  is changed to  $\frac{d^2z}{dx^2} = \cos^2 z + k \left(\frac{dz}{dx}\right)^2$ , then the value of  $k$  equal to \_\_\_\_\_

 [Watch Video Solution](#)

4. Let  $y=y(t)$  be a solution to the differential equation  $y' + 2ty = t^2$ , then  $\lim_{t \rightarrow \infty} \frac{y}{t}$  is.....

 [Watch Video Solution](#)

5. If the solution of the differential equation  $\frac{dy}{dx} = \frac{1}{x \cos y + \sin 2y}$  is  $x = ce^{\sin y} - k(1 + \sin y)$ , then the value of  $k$  is \_\_\_\_\_

 [Watch Video Solution](#)

6. If the independent variable  $x$  is changed to  $y$ , then the differential

equation  $x \frac{d^2y}{dx^2} + \left(\frac{dy}{dx}\right)^3 - \frac{dy}{dx} = 0$  is changed to  $x \frac{d^2x}{dy^2} + \left(\frac{dx}{dy}\right)^2 = k$  where  $k$  equals \_\_\_\_



[Watch Video Solution](#)

7. Let  $y_1$  and  $y_2$  be two different solutions of the equation

$\frac{dy}{dx} + P(x).y = Q(x)$ . Then  $\alpha y_1 + \beta y_2$  will be solution of the given equation if  $\alpha + \beta = \dots\dots\dots$



[Watch Video Solution](#)

8. Tangent is drawn at the point  $(x_i, y_i)$  on the curve  $y = f(x)$ , which intersects the x-axis at  $(x_{i+1}, 0)$ . Now, again a tangent is drawn at  $(x_{i+1}, y_{i+1})$  on the curve which intersect the x-axis at  $(x_{i+2}, 0)$  and the process is repeated  $n$  times, i.e.  $i = 1, 2, 3, \dots, n$ . If  $x_1, x_2, x_3, \dots, x_n$  form an arithmetic progression with common difference equal to  $(\log)_2 e$  and

curve passes through  $(0, 2)$ . Now if curve passes through the point  $(-2, k)$ , then the value of  $k$  is \_\_\_\_

 [Watch Video Solution](#)

9. The perpendicular from the origin to the tangent at any point on a curve is equal to the abscissa of the point of contact. Also curve passes through the point  $(1,1)$ . Then the length of intercept of the curve on the x-axis is \_\_\_\_\_

 [Watch Video Solution](#)

10. If the eccentricity of the curve for which tangent at point  $P$  intersects the y-axis at  $M$  such that the point of tangency is equidistant from  $M$  and the origin is  $e$ , then the value of  $5e^2$  is \_\_\_\_

 [Watch Video Solution](#)

11. If the solution of the differential equation  $\frac{dy}{dx} - y = 1 - e^{-x}$  and  $y(0) = y_0$  has a finite value, when  $x \rightarrow \infty$ , then the value of  $\left| \frac{2}{y_0} \right|$  is \_\_

 [Watch Video Solution](#)

12. Let  $f$  be a function defined on the interval  $[0, 2\pi]$  such that  $\int_0^x (f'(t) - \sin 2t) dt = \int_x^0 f(t) \tan t dt$  and  $f(0) = 1$ . Then the maximum value of  $f(x)$  is.....

 [Watch Video Solution](#)

13. Let  $y(x)$  be a function satisfying  $\frac{d^2y}{dx^2} - \frac{dy}{dx} + e^{2x} = 0$ ,  $y(0) = \alpha$  and  $y'(0) = 1$ . If maximum value of  $y(x)$  is  $y(\alpha)$ , then integral part of  $2\alpha$  is.....

 [Watch Video Solution](#)

14. if the differential equation of a curve, passing through  $(0, -\frac{\pi}{4})$  and  $(t, 0)$  is  $\cos y \left( \frac{dy}{dx} + e^{-x} \right) + \sin y \left( e^{-x} - \frac{dy}{dx} \right) = e^{e^{-x}}$  then find the value of  $t$ .  $e^{e^{-1}}$

 [Watch Video Solution](#)

15. Let  $f$  be a continuous function satisfying the equation  $\int_0^x f(t)dt + \int_0^x tf(x-t)dt = e^{-x} - 1$ , then find the value of  $e^9 f(9)$  is equal to.....

 [Watch Video Solution](#)

## JEE Main Previous Year

1. The differential equation which represents the family of curves  $y = C_1 e^{C_2 x}$ , where  $C_1$  and  $C_2$  are arbitrary constants, is

A.  $y' = y^2$

B.  $y'' = y'y$

C.  $yy'' = y'$

D.  $yy'' = (y')^2$

**Answer: D**

 [Watch Video Solution](#)

**2. Solution of the differential equation**

$$\cos x dy = y(\sin x - y) dx, 0 < x < \frac{\pi}{2} \text{ is}$$

A.  $\tan x = (\sec x + c)y$

B.  $\sec x = (\tan x + c)y$

C.  $y \sec x = \tan x + c$

D.  $y \tan x = \sec x + c$

**Answer: B**

 [Watch Video Solution](#)



3. Let  $I$  be the purchase value of an equipment and  $V(t)$  be the value after it has been used for  $t$  years. The value  $V(t)$  depreciates at a rate given by differential equation  $\left( dV \frac{t}{dt} = -k(T-t) \right)$ , where  $k > 0$  is a constant and  $T$  is the total life in years of the equipment. Then the scrap value  $V(T)$  of the equipment is : (1)  $T^2 - \frac{1}{k}$  (2)  $I - \frac{kT^2}{2}$  (3)  $I - \frac{k(T-t)^2}{2}$  (4)  $e^{-kT}$

A.  $e^{-kT}$

B.  $T^2 - \frac{I}{k}$

C.  $I - \frac{kT^2}{2}$

D.  $I - \frac{k(T-t)^2}{2}$

**Answer: C**



**Watch Video Solution**

4. If  $\frac{dy}{dx} = y + 3$  and  $y(0) = 2$ , then  $y(\ln 2)$  is equal to

A.  $-2$

B.  $7$

C.  $5$

D.  $13$

**Answer: B**



**Watch Video Solution**

5. A spherical balloon is filled with  $4500\pi$  cubic meters of helium gas. If a leak in the balloon causes the gas to escape at the rate of  $72\pi$  cubic meters per minute, then the rate (in meters per minute) at which the radius of the balloon decreases 49 minutes after the leakage began is (1)

$\frac{9}{7}$  (2)  $\frac{7}{9}$  (3)  $\frac{2}{9}$  (4)  $\frac{9}{2}$

A.  $6/7$

B.  $4/9$

C.  $2/9$

D. None of these

**Answer: C**



[Watch Video Solution](#)

6. The population  $p(t)$  at time  $t$  of a certain mouse species satisfies the differential equation  $\left( dp \frac{t}{dt} = 0.5p(t) - 450 \right)$  If  $p(0) = 850$ , then the time at which the population becomes zero is (1)  $2 \ln 18$  (2)  $\ln 9$  (3)  $\frac{1}{2} \ln 18$  (4)  $\ln 18$

A.  $2 \ln 18$

B.  $\ln 9$

C.  $\frac{1}{2} \ln 18$

D.  $\ln 18$

**Answer: A**



[Watch Video Solution](#)

7. At present, a firm is manufacturing 2000 items. It is estimated that the rate of change of production  $P$  w.r.t. additional number of workers  $x$  is given by  $\frac{dP}{dx} = 100 - 12\sqrt{x}$ . If the firm employs 25 more workers, then the new level of production of items is (1) 3000 (2) 3500 (3) 4500 (4) 2500

A. 2500

B. 3000

C. 3500

D. 4500

**Answer: C**



[Watch Video Solution](#)

8. Let the population of rabbits surviving at a time  $t$  be governed by the differential equation  $\left( dp \frac{t}{dt} = \frac{1}{2}p(t) - 200 \right)$ . If  $p(0) = 100$ , then  $p(t)$  equals (1)  $400 - 300e^{t/2}$  (2)  $300 - 200e^{-t/2}$  (3)  $600 - 500e^{t/2}$  (4)  $400 - 300e^{-t/2}$

A.  $40 - 300e^{t/2}$

B.  $200 - 200e^{-t/2}$

C.  $600 - 500e^{t/2}$

D.  $400 - 300e^{-t/2}$

**Answer: A**



**Watch Video Solution**

9. Let  $y(x)$  be the solution of the differential equation

$(x \log x) \frac{dy}{dx} + y = 2x \log x, (x \geq 1)$ , Then  $y(e)$  is equal to

A.  $e$

B.  $0$

C.  $2$

D.  $2e$

**Answer: C**



Watch Video Solution

10. If a curve  $y = f(x)$  passes through the point  $(1, -1)$  and satisfies the differential equation  $y(1 + xy)dx = xdy$ , then  $f\left(-\frac{1}{2}\right)$  is equal to: (1)  $-\frac{2}{5}$  (2)  $-\frac{4}{5}$  (3)  $\frac{2}{5}$  (4)  $\frac{4}{5}$

A.  $-\frac{4}{5}$

B.  $\frac{2}{5}$

C.  $\frac{4}{5}$

D.  $-\frac{2}{5}$

Answer: C



Watch Video Solution

11. If  $(2 + \sin x) \frac{dy}{dx} + (y + 1)\cos x = 0$  and  $y(0) = 1$ , then  $y\left(\frac{\pi}{2}\right)$  is equal to

A.  $\frac{4}{3}$

B.  $\frac{1}{3}$

C.  $-\frac{3}{8}$

D.  $-\frac{1}{3}$

**Answer: B**



**Watch Video Solution**

**12.** Let  $y = g(x)$  be the solution of the differential equation

$\sin x \left( \frac{dy}{dx} \right) + y \cos x = 4x$ , If  $y\left(\frac{\pi}{2}\right) = 0$ , then  $y(\pi/6)$  is equal to

A.  $-\frac{4}{9}\pi^2$

B.  $\frac{4}{9\sqrt{3}}\pi^2$

C.  $-\frac{8}{9\sqrt{3}}\pi^2$

D.  $-\frac{8}{9}\pi^2$

**Answer: D**

[Watch Video Solution](#)

## JEE Advanced Previous Year

1. A curve passes through the point  $\left(1, \frac{\pi}{6}\right)$ . Let the slope of the curve at each point  $(x, y)$  be  $\frac{y}{x} + \sec\left(\frac{y}{x}\right)$ ,  $x > 0$ . Then the equation of the curve is

A.  $\sin\left(\frac{y}{x}\right) = \log x + \frac{1}{2}$

B.  $\operatorname{cosec}\left(\frac{y}{x}\right) = \log x + 2$

C.  $\frac{\sec(2y)}{x} = \log x + 2$

D.  $\frac{\cos(2y)}{x} = \log x + \frac{1}{2}$

**Answer: A**

[Watch Video Solution](#)



2. The function  $y = f(x)$  is the solution of the differential equation

$$\frac{dy}{dx} + \frac{xy}{x^2 - 1} = \frac{x^4 + 2x}{\sqrt{1 - x^2}} \text{ in } (-1, 1) \text{ satisfying } f(0) = 0. \text{ Then}$$

$$\int_{\frac{\sqrt{3}}{2}}^{\frac{\sqrt{3}}{2}} f(x) dx \text{ is (a) (b)(c)(d) } \frac{\pi}{e} 3(f)(g) - (h) \frac{(i)\sqrt{(j)3(k)(l)}}{m} 2(n)(o)(p)$$

$$(q) (b) (r)(s)(t) \frac{\pi}{u} 3(v)(w) - (x) \frac{(y)\sqrt{(z)3(aa)(bb)}}{cc} 4(dd)(ee)(ff) (gg)$$

$$(c) (d)(e)(f) \frac{\pi}{g} 6(h)(i) - (j) \frac{(k)\sqrt{(l)3(m)(n)}}{o} 4(p)(q)(r) \quad (s) \quad (d)$$

$$(t)(u)(v) \frac{\pi}{w} 6(x)(y) - (z) \frac{(aa)\sqrt{(bb)3(cc)(dd)}}{ee} 2(ff)(gg)(hh) \text{ (ii)}$$

A.  $\frac{\pi}{3} - \frac{\sqrt{3}}{2}$

B.  $\frac{\pi}{3} - \frac{\sqrt{3}}{4}$

C.  $\frac{\pi}{6} - \frac{\sqrt{3}}{4}$

D.  $\frac{\pi}{6} - \frac{\sqrt{3}}{2}$

Answer: B



Watch Video Solution

3. If  $y = y(x)$  satisfies the differential equation  $8\sqrt{x}(\sqrt{9 + \sqrt{x}}) dy = (\sqrt{4 + \sqrt{9 + \sqrt{x}}})^{-1} dx, x > 0$  and  $y(0) = \sqrt{7}$ ,

then  $y(256) = 16$  (b) 80 (c) 3 (d) 9

A. 3

B. 9

C. 16

D. 80

**Answer: A**



**Watch Video Solution**

4. If  $y(x)$  satisfies the differential equation  $y' - y \tan x = 2x \sec x$  and  $y(0) = 0$ , then

A.  $y\left(\frac{\pi}{4}\right) = \frac{\pi^2}{8\sqrt{2}}$

B.  $y'\left(\frac{\pi}{4}\right) = \frac{\pi^2}{18}$

$$C. y\left(\frac{\pi}{3}\right) = \frac{\pi^2}{9}$$

$$D. y'\left(\frac{\pi}{3}\right) = \frac{4\pi}{3} + \frac{2\pi^2}{3\sqrt{3}}$$

**Answer: A::D**



**Watch Video Solution**

5. Consider the family of all circles whose centers lie on the straight line  $y = x$ . If this family of circles is represented by the differential equation

$P y' + Q y' + 1 = 0$ , where  $P, Q$  are functions of  $x, y$  and  $y'$  (here  $y' = \frac{dy}{dx}$ ,  $y'' = \frac{d^2y}{dx^2}$ ), then which of the following statements is

(are) true? (a) (b) (c)  $P = y + x$  (d) (e) (b) (f) (g)  $P = y - x$  (h) (i) (c)

(d) (e)  $P + Q = 1 - x + y + y + (f) (g) \left( (h) (i) y^{(j)'} (k) (l) (m) \right)^{(n)2(o)} (p)$

(r) (s)

(t) (u)  $P - Q = x + y - y - (v) (w) \left( (x) (y) y^{(z)'} (aa) (bb) (cc) \right)^{(dd)2(ee)} (f)$

(hh)

A.  $P = y + x$

B.  $P = y - x$

$$C. P + Q = 1 - x + y + y' + (y')^2$$

$$D. P - Q = x + y - y' - (y')^2$$

Answer: B::C



Watch Video Solution

6. Let  $y(x)$  be a solution of the differential equation  $(1 + e^x)y' + ye^x = 1$ . If  $y(0) = 2$ , then which of the following statements is (are) true? (a)  $y(-4) = 0$  (b)  $y(-2) = 0$  (c)  $y(x)$  has a critical point in the interval  $(-1, 0)$  (d)  $y(x)$  has no critical point in the interval  $(-1, 0)$

A.  $y(-4) = 0$

B.  $y(-2) = 0$

C.  $y(x)$  has a critical point in the interval  $(-1, 0)$

D.  $y(x)$  has no critical point in the interval  $(-1, 0)$

**Answer: A:C**



**Watch Video Solution**

7. A solution curve of the differential equation  $(x^2 + xy + 4x + 2y + 4) \left( \frac{dy}{dx} \right) - y^2 = 0$  passes through the point  $(1, 3)$ . Then the solution curve is

A. intersects  $y = x + 2$  exactly at one point

B. intersects  $y = x + 2$  exactly at two points

C. intersects  $y = (x + 2)^2$

D. does NOT intersect  $y = (x + 3)^2$

**Answer: A:D**



**View Text Solution**

8. Let  $f: R \rightarrow R$  and  $g: R \rightarrow R$  be two non-constant differentiable functions. If

$$f'(x) = \left( e^{(f(x) - g(x))} \right) g'(x) \text{ for all } x \in R \text{ and } f(1) = g(2) = 1,$$

then which of the following statement(s) is (are) TRUE?

A.  $f(2) < 1 - \log_e 2$

B.  $f(2) > 1 - \log_e(2)$

C.  $g(1) < 1 - \log_e 2$

D.  $g(1) < 1 - \log_e 2$

**Answer: B::C**



[Watch Video Solution](#)

9. Let  $f$  be a real-valued differentiable function on  $R$  (the set of all real numbers) such that  $f(1) = 1$ . If the  $y$ -intercept of the tangent at any point  $P(x, y)$  on the curve  $y = f(x)$  is equal to the cube of the abscissa of  $P$ , then the value of  $f(-3)$  is equal to \_\_\_\_\_



[Watch Video Solution](#)

10. Let  $y'(x) + y(x)g'(x) = g(x)g'(x), y(0), x \in R,$  where  $f'(x)$  denotes  $\frac{dy(x)}{dx}$ , and  $g(x)$  is a given non-constant differentiable function on  $R$  with  $g(0) = g(2) = 0$ . Then the value of  $y(2)$  is\_\_\_\_\_

 Watch Video Solution

11. Let  $f: [1, \infty]$  be a differentiable function such that  $f(1) = 2$ . If  $\int_1^x f(t)dt = 3xf(x) - x^3$  for all  $x \geq 1$ , then the value of  $f(2)$  is

 Watch Video Solution

12. Let  $f: R \rightarrow R$  be a differentiable function with  $f(0) = 0$ . If  $y = f(x)$  satisfies the differential equation

$\frac{dy}{dx} = (2 + 5y)(5y - 2)$ , then the value of  $\lim_{x \rightarrow \infty} f(x)$  is.....

 View Text Solution

1. The order of the differential equation whose general solution is

$$y = c_1 \cos 2x + c_2 \cos^2 x + c_3 \sin^2 x + c_4 \text{ is}$$

A. 2

B. 4

C. 3

D. None of these

**Answer: A**



[View Text Solution](#)

2. Order of the differential equation whose general solution is

$$y = \frac{ax}{bx + c}, \text{ where } a, b, c \text{ are arbitrary constants is}$$

A. 1



B. 2

C. 3

D. None of these

**Answer: B**



[View Text Solution](#)

3. If  $p$  and  $q$  are order and degree of differential equation

$$y^2 \left( \frac{d^2 y}{dx^2} \right)^2 + 3x \left( \frac{dy}{dx} \right)^{\frac{1}{3}} + x^2 y^2 = \sin x, \text{ then}$$

A.  $p > q$

B.  $\frac{p}{2} = \frac{q}{2}$

C.  $p = q$

D.  $p < q$

**Answer: D**



[Watch Video Solution](#)

4. If  $m$  and  $n$  are the order and degree of the differential equation

$$(y'')^5 + 4 \cdot \frac{(y'')^3}{y'''} + y''' = \sin x, \text{ then}$$

A.  $m = 3, n = 5$

B.  $m = 3, n = 1$

C.  $m = 3, n = 3$

D.  $m = 3, n = 2$

**Answer: D**



[Watch Video Solution](#)

5. If the differential equation  $\frac{dx}{3y + f} + \frac{dy}{px + g} = 0$  represents a family of circle, then  $p =$

A.  $g$

B.  $f$

C. 4

D. 3

**Answer: D**



**Watch Video Solution**

6. The general solution of  $\frac{dy}{dx} = 1 - x^2 - y^2 + x^2y^2$  is

A.  $2 \sin^{-1} y = x\sqrt{1 - y^2} + c$

B.  $\sin^{-1} y = \frac{1}{2} \sin^{-1} x + c$

C.  $\cos^{-1} y = x \cos^{-1} x + c$

D.  $\frac{1}{2} \log\left(\frac{1 + y}{1 - y}\right) = x - \frac{x^3}{3} + c$

**Answer: D**



**Watch Video Solution**

7. The solution of the differential equation  $e^{-x}(y + 1)dy + (\cos^2 x - \sin 2x)y(dx) = 0$  subjected to the condition that  $y = 1$  when  $x = 0$  is

A.  $(y + 1) + e^x \cos^2 x = 2$

B.  $y + \log y = e^x \cos^2 x$

C.  $\log(y + 1) + e^x \cos^2 x = 1$

D.  $y + \log y + e^x \cos^2 x = 2$

**Answer: D**



**Watch Video Solution**

8. The solution of the differential equation

$$(x^2 - yx^2) \frac{dy}{dx} + y^2 + xy^2 = 0 \text{ is}$$

A.  $\log\left(\frac{x}{y}\right) = \frac{1}{x} + \frac{1}{y} + c$

B.  $\log\left(\frac{y}{x}\right) = \frac{1}{x} + \frac{1}{y} + c$

$$C. \log(xy) = \frac{1}{x} + \frac{1}{y} + c$$

$$D. \log(xy) + \frac{1}{x} + \frac{1}{y} = c$$

**Answer: A**



**Watch Video Solution**

9. The family of curves passing through  $(0, 0)$  and satisfying the differential equation  $\frac{y_2}{y_1} = 1$  (where,  $y_n = \frac{d^n y}{dx^n}$ ) is (A)  $y = k$  (B)  $y = kx$  (C)  $y = k(e^x + 1)$  (D)  $y = k(e^x - 1)$

A.  $y = k$

B.  $y = kx$

C.  $y = k(e^x + 1)$

D.  $y = k(e^x - 1)$

**Answer: D**



**View Text Solution**

10. The solution of the differential equation

$$y^2 dx + (x^2 - xy + y^2) dy = 0 \text{ is}$$

A.  $\tan^{-1}\left(\frac{x}{y}\right) + \ln y + C = 0$

B.  $2 \tan^{-1}\left(\frac{x}{y}\right) + \ln x + C = 0$

C.  $\ln\left(y + \sqrt{x^2 + y^2}\right) + \ln y + C = 0$

D.  $\ln\left(x + \sqrt{x^2 + y^2}\right) + C = 0$

**Answer: A**



**Watch Video Solution**

11. The solution of differential equation  $(1 - xy + x^2y^2) dx = x^2 dy$  is

A.  $\tan xy = \log |cx|$

B.  $\tan (y/x) = \tan \log |cx|$

C.  $xy = \tan \log |cx|$

D. None of these

**Answer: C**



**Watch Video Solution**

12. If  $y(t)$  satisfies the differential equation

$y'(t) + 2y(t) = 2e^{-2t}$ ,  $y(0) = 2$  then  $y(1)$  equals

A.  $\frac{3}{e}$

B.  $\frac{3}{e^2}$

C.  $\frac{4}{e}$

D.  $\frac{4}{e^2}$

**Answer: D**



**Watch Video Solution**

13. If  $\frac{dy}{dx} - y \log_e 2 = 2^{\sin x} (\cos x - 1) \log_e 2$ , then  $y =$

A.  $2^{\sin x} + c2^x$

B.  $2^{\cos x} + c2^x$

C.  $2^{\sin x} + c2^{-x}$

D.  $2^{\cos x} + c2^{-x}$

**Answer: A**



**Watch Video Solution**

14. If  $ye^y dx = (y^3 + 2xe^y) dy$ ,  $y(0) = 1$ , then the value of  $x$  when  $y = 0$  is

A.  $-1$

B.  $0$

C.  $1$

D.  $2$



**Answer: B**

 [Watch Video Solution](#)

15. If  $y_1(x)$  is a solution of the differential equation  $\frac{dy}{dx} - f(x)y = 0$ , then a solution of the differential equation  $\frac{dy}{dx} + f(x)y = r(x)$  is

A.  $y = \frac{1}{y_1(x)} \int r(x)y_1(x)dx + \frac{c}{y_1(x)}$

B.  $y = y_1(x) \int \frac{r(x)}{y_1(x)} dx + c$

C.  $y = \int r(x)y_1(x)dx + c$

D. None of these

**Answer: A**

 [View Text Solution](#)

16. The general solution of  $x \left( \frac{dy}{dx} \right) + (\log x)y = x^{-\frac{1}{2} \log x}$  is

$$A. y = x^{1 - \frac{1}{2}\log x} + cx^{-\frac{1}{2}\log x}$$

$$B. y. x^{\frac{1}{2}\log x} = x^{\frac{1}{2}\log x} + c$$

$$C. y = e^{\frac{(\log x)^2}{2}}(x + c)$$

$$D. y = e^{\frac{1}{2}(\log x)^2} \left( x^{1 - \frac{1}{2}(\log x)} - x^{-\frac{1}{2}\log x} \right) + c$$

**Answer: A**



**Watch Video Solution**

17. Find the general solution of the differential equation

$$(1 + \tan y)(dx - dy) + 2xdy = 0$$

$$A. x(\sin y + \cos y) = \sin y + ce^y$$

$$B. x(\sin y + \cos y) = \sin y + ce^{-y}$$

$$C. y(\sin x + \cos x) = \sin x + ce^x$$

D. None of these

**Answer: B**

 [Watch Video Solution](#)

18. Solution of differential equation  $x^2y - x^3 \frac{dy}{dx} = y^4 \cos x$  is

A.  $x^2y^{-3} = 2 \sin x + c$

B.  $x^2y^{-3} = 3 \cos x + c$

C.  $x^3y^{-3} = 3 \sin x + c$

D.  $x^2y^3 = 3 \sin x + cx^2y$

**Answer: C**

 [Watch Video Solution](#)

19. Suppose a solutions of the differential equation

$(xy^3 + x^2y^7) \frac{dy}{dx} = 1$  satisfies the initial conditions  $y\left(\frac{1}{4}\right) = 1$ . Then

the value of  $\frac{dy}{dx}$  when  $y = -1$  is

A.  $-\frac{3}{20}$

B.  $-\frac{20}{3}$

C.  $-\frac{5}{16}$

D.  $-\frac{16}{5}$

**Answer: D**



**View Text Solution**

20. The general solution of the differential equation

$$\frac{dy}{dx} = y \tan x - y^2 \sec x \text{ is}$$

A.  $\tan x = (c + \sec x)y$

B.  $\sec y = (c + \tan y)x$

C.  $\sec x = (c + \tan x)y$

D. None of these

**Answer: C**



**Watch Video Solution**

21. The solution of differential equation  $x^2(xdy + ydx) = (xy - 1)^2 dx$  is (where  $c$  is an arbitrary constant)

A.  $xy - 1 = cx$

B.  $xy - 1 = cx^2$

C.  $\frac{1}{xy - 1} = \frac{1}{x} + c$

D. None of these

Answer: C



[View Text Solution](#)

22. Solution of the differential  $(x + 2y^3) = \frac{dx}{dy}y$  is

A.  $x = y^2(c + y^2)$

B.  $x = y(c - y^2)$

C.  $x = 2y(c - y^2)$

$$D. x = y(c + y^2)$$

Answer: D



View Text Solution

23. General solution of differential equation

$$x^2 \left( x + y \frac{dy}{dx} \right) + \left( x \frac{dy}{dx} - y \right) \sqrt{x^2 + y^2} = 0 \text{ is}$$

A.  $\frac{1}{\sqrt{x^2 + y^2}} + \frac{y}{x} = c$

B.  $\sqrt{x^2 + y^2} - \frac{y}{x} = c$

C.  $\sqrt{x^2 + y^2} + \frac{y}{x} = c$

D.  $2\sqrt{x^2 + y^2} + \frac{y}{x} = c$

Answer: C



View Text Solution

24. Solution of the differential  $y' = \frac{3yx^2}{x^3 + 2y^4}$  is

A.  $x^3y^{-1} = \frac{2}{3}y^3 + c$

B.  $x^2y^{-1} = \frac{2}{3}y^3 + c$

C.  $xy^{-1} = \frac{2}{3}y^3 + c$

D. None of these

Answer: A



Watch Video Solution

25. For  $y > 0$  and  $x \in R$ ,  $ydx + y^2dy = xdy$  where  $y = f(x)$ . If  $f(1)=1$ , then the value of  $f(-3)$  is

A. 1

B. 2

C. 3

D. 4

**Answer: C**

 [Watch Video Solution](#)

26. An equation of the curve satisfying  $xdy - ydx = \sqrt{x^2 - y^2}dx$  and  $y(1) = 0$  is

A.  $y = x^2 \log|\sin x|$

B.  $y = x \sin(\log|x|)$

C.  $y^2 = x(x - 1)^2$

D.  $y = 2x^2(x - 1)$

**Answer: B**

 [Watch Video Solution](#)



27. The solution of

$$(y(1 + x^{-1}) + \sin y)dx + (x + \log x + x \cos y)dy = 0 \text{ is}$$

A.  $(1 + y^{-1} \sin y) + x^{-1} \log x = c$

B.  $(y + \sin y) + xy \log x = C$

C.  $xy + y \log x + x \sin y = C$

D. None of these

**Answer: C**



[Watch Video Solution](#)

28. The solution of  $(1 + x) \frac{dy}{dx} + 1 = e^{x-y}$  is

A.  $e^y(x + 1) = c$

B.  $e^y(x + 1) = e^x + c$

C.  $e^y(x + 1) = ce^x$

$$D. (x + 1) = e^x + c$$

**Answer: B**



**Watch Video Solution**

29. The solution of differential equation

$$x \sec\left(\frac{y}{x}\right)(ydx + xdy) = y \operatorname{cosec}\left(\frac{y}{x}\right)(xdy - ydx) \text{ is}$$

A.  $xy = c \operatorname{cosec}\left(\frac{y}{x}\right)$

B.  $xy^2 \sin \frac{y}{x} = c$

C.  $xy \operatorname{cosec} \frac{y}{x} = c$

D.  $xy = c \sin\left(\frac{x}{y}\right)$

**Answer: C::D**



**Watch Video Solution**

30. The general solution of the differential equation

$$\sqrt{1 - x^2y^2}dx = ydx + xdy \text{ is}$$

A.  $\sin(xy) = x + c$

B.  $\sin^{-1}(xy) + x = c$

C.  $\sin(x + c) = xy$

D.  $\sin(xy) + x = c$

**Answer: C**



**Watch Video Solution**

31. Solve  $(1 + xy)ydx + (1 - xy)x dy = 0$

A.  $\frac{x}{y} + \frac{1}{xy} = k$

B.  $\log\left(\frac{x}{y}\right) = \frac{1}{xy} + k$

C.  $\frac{x}{y} - \frac{1}{xy} = k$

D.  $\log\left(\frac{y}{x}\right) = xy + k$

**Answer: B**



**Watch Video Solution**

**32.** Solution of the differential equation  $\frac{dy}{dx} = \frac{y^3}{e^{2x} + y^2}$ , is

A.  $e^{-2x}y^2 + 2\ln|y| = c$

B.  $e^{2x}y^2 = 2\ln|y| + c$

C.  $e^x + \ln|y| = c$

D. None of these

**Answer: A**



**Watch Video Solution**

**33.** A population grows at the rate of 10% of the population per year. How long does it take for the population to double ?

A.  $2 \log 10$  years

B.  $20 \log 2$  years

C.  $10 \log 2$  years

D.  $5 \log 2$  years

**Answer: C**



[View Text Solution](#)

**34.** A tangent drawn to the curve  $y = f(x)$  at  $P(x, y)$  cuts the  $x$  and  $y$  axes at  $A$  and  $B$ , respectively, such that  $AP : PB = 1 : 3$ . If  $f(1) = 1$  then the curve passes through  $\left(k, \frac{1}{8}\right)$  where  $k$  is

A. 1

B. 2

C. 3

D. 4

**Answer: B**



[View Text Solution](#)

35. A curve 'C' with negative slope through the point(0,1) lies in the I Quadrant. The tangent at any point 'P' on it meets the x-axis at 'Q'. Such that  $PQ = 1$ . Then

The curve in parametric form is

A.  $x = \cos \theta + \ln \tan(\theta/2), y = \sin \theta$

B.  $x = -\cos \theta + \ln \tan(\theta/2), y = \sin \theta$

C.  $x = -\cos \theta - \ln \tan(\theta/2), y = \sin \theta$

D. None of these

**Answer: C**



[Watch Video Solution](#)

36. Tangent to a curve intercepts the y-axis at a point  $P$ . A line perpendicular to this tangent through  $P$  passes through another point

(1,0). The differential equation of the curve is (a)

(b)  $y \frac{dy}{dx} - x \left( \frac{dy}{dx} \right)^2 = 1$  (c)  $x \frac{d^2y}{dx^2} + \left( \frac{dy}{dx} \right)^2 = 1$  (d)  $y \frac{dx}{dy} + x = 1$  (e) None of these

(y) (b) [Math Processing Error] (eee) (c)

(d)  $y \frac{dx}{dy} + x = 1$  (e) None of these

A.  $y \frac{dy}{dx} - x \left( \frac{dy}{dx} \right)^2 = 1$

B.  $x \frac{d^2y}{dx^2} + \left( \frac{dy}{dx} \right)^2 = 1$

C.  $y \frac{dx}{dy} + x = 1$

D. None of these

Answer: A

 Watch Video Solution

37. The orthogonal trajectories of the family of curves  $a^{n-1}y = x^n$  are given by (A)  $x^n + n^2y = \text{constant}$  (B)  $ny^2 + x^2 = \text{constant}$  (C)

$$n^2x + y^n = \text{constant} \quad (D) y = x$$

A.  $x^n + n^2y = \text{const}$

B.  $ny^2 + x^2 = \text{const}$

C.  $n^2x + y^n = \text{const}$

D.  $n^2x - y^n = \text{const}$

**Answer: B**



[View Text Solution](#)

## Multiple Correct Answer Type

1. The differential equation for the family of curves  $y = c \sin x$  can be given by

A.  $\left(\frac{dy}{dx}\right)^2 = y^2 \cot^2 x$

B.  $\left(\frac{dy}{dx}\right)^2 - \left(\sec x \frac{dy}{dx}\right)^2 + y^2 = 0$



$$C. \left(\frac{dy}{dx}\right)^2 = \tan^2 x$$

$$D. \frac{dy}{dx} = y \cot x$$

**Answer: A::B::D**



**Watch Video Solution**

2. If the solution of the equation  $\frac{d^2x}{dt^2} + 4\frac{dx}{dt} + 3x = 0$  given that for  $t = 0, x = 0$  and  $\frac{dx}{dt} = 12$  is in the form  $x = Ae^{-3t} + Be^{-t}$ , then

A.  $A + B = 0$

B.  $A + B = 12$

C.  $|AB| = 36$

D.  $|AB| = 49$

**Answer: A::C**



**Watch Video Solution**

3. The solution of  $\left(\frac{dy}{dx}\right)^2 - 2\left(x + \frac{1}{4x}\right)\frac{dy}{dx} + 1 = 0$

A.  $y = x^2 + c$

B.  $y = \frac{1}{2}\ln(x) + c, x > 0$

C.  $y = \frac{x}{2} + c$

D.  $y = \frac{x^2}{2} + c$

Answer: A:B

 [Watch Video Solution](#)

4.  $\left(\frac{dy}{dx}\right)^2 + 2y \cot x \frac{dy}{dx} = y^2$  has the solution

A.  $y + \frac{c}{1 + \cos x} = 0$

B.  $y = \frac{c}{1 - \cos x}$

C.  $x = \sin^{-1} \sqrt{\frac{c}{2y}}$

D.  $x = \cos^{-1} \left(\frac{c}{2y}\right)$

**Answer: A::B**



**Watch Video Solution**

5. Let  $\frac{dy}{dx} + y = f(x)$  where  $y$  is a continuous function of  $x$  with  $y(0) = 1$  and  $f(x) = \begin{cases} e^{-x} & \text{if } 0 \leq x \leq 2 \\ e^{-2} & \text{if } x > 2 \end{cases}$  Which of the following hold(s) good ?

A.  $y(1) = 2e^{-1}$

B.  $y'(1) = -e^{-1}$

C.  $y(3) = -2e^{-3}$

D.  $y'(3) = -2e^{-3}$

**Answer: A::B::D**



**View Text Solution**

6. A differentiable function satisfies

$$f(x) = \int_0^x (f(t)\cot t - \cos(t - x))dt.$$

Which of the following hold(s) good?

- A.  $f(x)$  has a minimum value  $1 - e$
- B.  $f(x)$  has a maximum value  $1 - e^{-1}$
- C.  $f''\left(\frac{\pi}{2}\right) = e$
- D.  $f'(0) = 1$

Answer: A::B::C



[View Text Solution](#)

7.  $y = f(x)$  which has differential equation  $y(2xy + e^x)dx - e^x dy = 0$  passing through the point  $(0, n 1)$ . Then which of the following is/are true about the function?

- A.  $x = 1 + \sqrt{2}$  is point of local maxima

B.  $x = 1 - \sqrt{2}$  is point of local minima

C.  $\lim_{x \rightarrow \infty} f(x) = -\infty$

D.  $\lim_{x \rightarrow -\infty} f(x) = 0$

**Answer: A::B::C::D**



[View Text Solution](#)

8. Suppose a curve whose sub tangent is  $n$  times the abscissa of the point of contact and passes through the point  $(2, 3)$ . Then

A. for  $n = 1$ , equation of the curve is  $2y = 3x$

B. for  $n = 1$ , equation of the curve is  $2y^2 = 9x$

C. for  $n = 2$ , equation of the curve is  $2y = 3x$

D. for  $n = 2$ , equation of the curve is  $2y^2 = 9x$

**Answer: A::D**



[View Text Solution](#)

9. Let  $C$  be a curve such that the normal at any point  $P$  on it meets  $x$ -axis and  $y$ -axis at  $A$  and  $B$  respectively. If  $BP : PA = 1 : 2$  (internally) and the curve passes through the point  $(0, 4)$ , then which of the following alternative(s) is/are correct?

- A. The curves passes through  $(\sqrt{10}, -6)$ .
- B. The equation of tangent at  $(4, 4\sqrt{3})$  is  $2x + \sqrt{3}y = 20$ .
- C. The differential equation for the curve is  $yy' + 2x = 0$ .
- D. The curve represents a hyperbola.

**Answer: A:D**

 [View Text Solution](#)

10. The normal at a general point  $(a, b)$  on curve makes an angle  $\theta$  with  $x$ -axis which satisfies  $b(a^2 \tan \theta - \cot \theta) = a(b^2 + 1)$ . The equation of curve can be

A.  $y = e^{x^2/2} + c$

B.  $\log ky^2 = x^2$

C.  $y = ke^{x^2/2}$

D.  $x^2 - y^2 = k$

**Answer: B::C::D**



**View Text Solution**

## Comprehension Type

1. Tangent is drawn at the point  $(x_i, y_i)$  on the curve  $y = f(x)$ , which intersects the x-axis at  $(x_{i+1}, 0)$ . Now, again a tangent is drawn at  $(x_{i+1}, y_{i+1})$  on the curve which intersect the x-axis at  $(x_{i+2}, 0)$  and the process is repeated  $n$  times, i.e.  $i = 1, 2, 3; n$ . If  $x_1, x_2, x_3, ; x_n$  form an arithmetic progression with common difference equal to  $(\log)_2 e$  and curve passes through  $(0, 2)$ . Now if curve passes through the point  $(-2, k)$ , then the value of  $k$  is\_\_\_\_\_

A. (1, 4)

B. (5, 1/16)

C. (2, 1/2)

D. None of these

**Answer: B**



[Watch Video Solution](#)

2. Tangent is drawn at the point  $(x_i, y_i)$  on the curve  $y = f(x)$ , which intersects the x-axis at  $(x_{i+1}, 0)$ . Now again tangent is drawn at  $(x_{i+1}, y_{i+1})$  on the curve which intersects the x-axis at  $(x_{i+2}, 0)$  and the process is repeated n times i.e.  $i = 1, 2, 3, \dots, n$ .

If  $x_1, x_2, x_3, \dots, x_n$  form a geometric progression with common ratio equal to 2 and the curve passes through (1, 2), then the curve is

A. circle

B. hyperbola



C. ellipse

D. parabola

**Answer: B**



[View Text Solution](#)

## Question Bank

1. Find the real value of  $m$  for which the substitution  $y = u^m$  will transform the differential equation  $2x^4y \frac{dy}{dx} + y^4 = 4x^6$  in to a homogenous equation.



[Watch Video Solution](#)

2. A function  $y = f(x)$  satisfies the differential equation  $\frac{dy}{dx} + x^2y = -2x$ ,  $f(1) = 1$ . The value of  $|f''(1)|$  is



[Watch Video Solution](#)

3. If the differential equation representing the family of curves  $y = C_1 \cos 2x + C_2 \cos^2 x + C_3 \sin^2 x + C_4$  is  $\lambda y' = y'' \tan 2x$ , then  $\lambda$  is

 [Watch Video Solution](#)

4. Given  $y(0) = 2000$  and  $\frac{dy}{dx} = 32000 - 20y^2$ , then find the value of  $\lim_{x \rightarrow \infty} y(x)$ .

 [Watch Video Solution](#)

5. Number of values of  $m \in \mathbb{N}$  for which  $y = e^{mx}$  is a solution of the differential equation  $\frac{d^3y}{dx^3} - 3\frac{d^2y}{dx^2} - 4\frac{dy}{dx} + 12y = 0$  (a) 0 (b) 1 (c) 2 (d)

More than 2

 [Watch Video Solution](#)

6. If  $x(t)$  is a solution of  $\frac{(1+t)dy}{dx} - ty = 1$  and  $y(0) = -1$  then  $y(1)$  is (a) (b)(c) - (d)  $\frac{1}{e}2(f)(g)(h)$  (i) (b) (j)(k)e + (l)  $\frac{1}{m}2(n)(o)(p)$  (q) (c) (d)(e)e - (f)  $\frac{1}{g}2(h)(i)(j)$  (k) (d) (l)(m)(n)  $\frac{1}{o}2(p)(q)(r)$  (s)



**Watch Video Solution**

7. Let  $y = f(x)$  be a function satisfying the differential equation  $\frac{xdy}{dx} + 2y = 4x^2$  and  $f(1) = 1$ . Then  $f(-3)$  is equal to



**Watch Video Solution**