



MATHS

BOOKS - CENGAGE

STRAIGHT LINES

Examples

1. Find the equation of the line through $(2, 3)$ which is (i) parallel to the x-axis and (ii) parallel to the y-axis.

[Watch Video Solution](#)

2. Find the equation of line passing through point $(2, -5)$ which is

(i) parallel to the line $3x + 2y - 4 = 0$

(ii) perpendicular to the line $3x + 2y - 4 = 0$



[Watch Video Solution](#)

3. Find the equation of the perpendicular bisector of the line segment joining the points $A(2, 3)$ and $B(6, -5)$.



[Watch Video Solution](#)

4. Find the locus of a point P which moves such that its distance from the line $y = \sqrt{3}x - 7$ is the same as its distance from $(2\sqrt{3}, -1)$



[Watch Video Solution](#)

5. Consider a triangle with vertices $A(1, 2)$, $B(3, 1)$, and $C(-3, 0)$. Find the equation of altitude through vertex A . the equation of median through vertex A . the equation of internal angle bisector of $\angle A$.



[View Text Solution](#)

6. Find the coordinates of the foot of the perpendicular drawn from the point P(1,-2) on the line $y = 2x + 1$. Also, find the image of P in the line.

 [Watch Video Solution](#)

7. If the line $\left(\frac{x}{a}\right) + \left(\frac{y}{b}\right) = 1$ moves in such a way that $\left(\frac{1}{a^2}\right) + \left(\frac{1}{b^2}\right) = \left(\frac{1}{c^2}\right)$, where c is a constant, prove that the foot of the perpendicular from the origin on the straight line describes the circle $x^2 + y^2 = c^2$.

 [View Text Solution](#)

8. In what ratio does the line joining the points (2, 3) and (4, 1) divide the segment joining the points (1, 2) and (4, 3)?

 [View Text Solution](#)

9. $ABCD$ is a square whose vertices are $A(0, 0)$, $B(2, 0)$, $C(2, 2)$, and $D(0, 2)$. The square is rotated in the XY - plane through an angle 30° in the anticlockwise sense about an axis passing through A perpendicular to the XY - plane. Find the equation of the diagonal BD of this rotated square.



[View Text Solution](#)

10. In a triangle ABC , side AB has equation $2x + 3y = 29$ and side AC has equation $x + 2y = 16$. If the midpoint of BC is $(5, 6)$, then find the equation of BC .



[View Text Solution](#)

11. Two consecutive sides of a parallelogram are $4x + 5y = 0$ and $7x + 2y = 0$. If the equation of one diagonal is $11x - 7y = 9$, find the equation of the other diagonal.



[View Text Solution](#)

[View Text Solution](#)

12. If one of the sides of a square is $3x - 4y - 12 = 0$ and the center is $(0, 0)$, then find the equations of the diagonals of the square.



[View Text Solution](#)

13. A vertex of an equilateral triangle is $2, 3$ and the opposite side is $x + y = 2$. Find the equations of other sides.



[View Text Solution](#)

14. A line $4x + y = 1$ passes through the point $A(2, -7)$ and meets line BC at B whose equation is $3x - 4y + 1 = 0$, the equation of line AC such that $AB = AC$ is (a) $52x + 89y + 519 = 0$ (b) $52x + 89y - 519 = 0$ (c) $82x + 52y + 519 = 0$ (d) $89x + 52y - 519 = 0$



[View Text Solution](#)

15. A ray of light is sent along the line $x - 2y - 3 = 0$ upon reaching the line $3x - 2y - 5 = 0$, the ray is reflected from it. Find the equation of the line containing the reflected ray.



[View Text Solution](#)

16. Find the equation of the line which intersects the y-axis at a distance of 2 units above the origin and makes an angle of 30° with the positive direction of the x-axis.



[View Text Solution](#)

17. Find the equation of a straight line cutting off an intercept -1 from y-axis and being equally inclined to the axes.



[View Text Solution](#)

18. Find the equation of a line that has $-y$ -intercept 4 and is a perpendicular to the line joining $(2, -3)$ and $(4, 2)$.



[View Text Solution](#)

19. Find equation of the line passing through the point $(2, 2)$ and cutting off intercepts on the axes whose sum is 9.



[View Text Solution](#)

20. Find the equation of the straight line that (i) makes equal intercepts on the axes and passes through the point $(2;3)$ (ii) passes through the point $(-5;4)$ and is such that the portion intercepted between the axes is divided by the point in the ratio $1 : 2$



[View Text Solution](#)

21. Line segment AB of fixed length c slides between coordinate axes such that its ends A and B lie on the axes. If O is origin and rectangle OAPB is completed, then show that the locus of the foot of the perpendicular drawn from P to AB is $x^{\frac{2}{3}} + y^{\frac{2}{3}} = c^{\frac{2}{3}}$.



[View Text Solution](#)

22. Reduce the line $2x - 3y + 5 = 0$ in slope-intercept, intercept, and normal forms.



[View Text Solution](#)

23. Find the equation of the line which satisfy the given conditions :
Perpendicular distance from the origin is 5 units and the angle made by the perpendicular with the positive axis is 30° .



[View Text Solution](#)

24. A straight line is drawn through the point $P(2;3)$ and is inclined at an angle of 30° with the x -axis . Find the coordinates of two points on it at a distance 4 from point P.



[View Text Solution](#)

25. The line joining two points $A(2,0)$ and $B(3,1)$ is rotated about A in anticlockwise direction through an angle of 15° . find the equation of line in the new position. If b goes to c in the new position what will be the coordinates of C.



[View Text Solution](#)

26. A line through point $A(1,3)$ and parallel to the line $x-y+1 = 0$ meets the line $2x-3y + 9 = 0$ at point P. Find distance AP without finding point P.



[View Text Solution](#)

27. Two adjacent vertices of a square are $(1, 2)$ and $(-2, 6)$ Find the other vertices.

 [View Text Solution](#)

28. A Line through the variable point $A(1 + k; 2k)$ meets the lines $7x + y - 16 = 0$; $5x - y - 8 = 0$ and $x - 5y + 8 = 0$ at B;C;D respectively.

Prove that AC;AB and AD are in HP.

 [View Text Solution](#)

29. if P is the length of perpendicular from origin to the line $\frac{x}{a} + \frac{y}{b} = 1$

then prove that $\frac{1}{a^2} + \frac{1}{b^2} = \frac{1}{p^2}$

 [View Text Solution](#)

30. Find the coordinates of a point on $x + y + 3 = 0$, whose distance from $x + 2y + 2 = 0$ is $\sqrt{5}$.

 [Watch Video Solution](#)

31. Find the least and greatest values of the distance of the point $(\cos \theta, \sin \theta)$, $\theta \in R$, from the line $3x - 4y + 10 = 0$.

 [View Text Solution](#)

32. Prove that the product of the lengths of the perpendiculars drawn from the points $(\sqrt{a^2 - b^2}, 0)$ and $(-\sqrt{a^2 - b^2}, 0)$ to the line $\frac{x}{a} \cos \theta + \frac{y}{b} \sin \theta = 1$ is b^2 .

 [View Text Solution](#)

33. Find the least value of $(x - 1)^2 + (y - 2)^2$ under the condition $3x + 4y - 2 = 0$.

 [View Text Solution](#)

34. ABC is an equilateral triangle with $A(0, 0)$ and $B(a, 0)$, ($a > 0$). L , M and N are the foot of the perpendiculars drawn from a point P to the side AB , BC , and CA , respectively. If P lies inside the triangle and satisfies the condition $PL^2 = PM^2 + PN^2$, then find the locus of P .

 [View Text Solution](#)

35. Line L has intercepts a and b on the coordinate axes. When the axes are rotated through a given angle keeping the origin fixed, the same line L has intercepts p and q . Then $a^2 + b^2 = p^2 + q^2$ $\frac{1}{a^2} + \frac{1}{b^2} = \frac{1}{p^2} + \frac{1}{q^2}$
 $a^2 + p^2 = b^2 + q^2$ (d) $\frac{1}{a^2} + \frac{1}{p^2} = \frac{1}{b^2} + \frac{1}{q^2}$

 [View Text Solution](#)

36. Two sides of a square lie on the lines $x + y = 1$ and $x + y + 2 = 0$.

What is its area?



[View Text Solution](#)

37. Find equation of the line which is equidistant from parallel lines

$$9x + 6y - 7 = 0 \text{ and } 3x + 2y + 6 = 0.$$



[View Text Solution](#)

38. If one side of the square is $2x - y + 6 = 0$, then one of the vertices is $(2, 1)$. Find the other sides of the square.



[View Text Solution](#)

39. Prove that the area of the parallelogram contained by the lines

$$4y - 3x - a = 0, 3y - 4x + a = 0, 4y - 3x - 3a = 0, \quad \text{and}$$

$$3y - 4x + 2a = 0 \text{ is } \left(\frac{2}{7}\right)a^2.$$

 [View Text Solution](#)

40. The equation of straight line passing through $(-2,-7)$ and having an intercept of length 3 between the straight lines : $4x + 3y = 12$, $4x + 3y = 3$ are : (A) $7x + 24y + 182 = 0$ (B) $7x + 24y + 18 = 0$ (C) $x + 2 = 0$ (D) $x - 2 = 0$

 [View Text Solution](#)

41. A line L is drawn from $P(4, 3)$ to meet the lines L_1 and L_2 given by $3x + 4y + 5 = 0$ and $3x + 4y + 15 = 0$ at points A and B , respectively. From A , a line perpendicular to L is drawn meeting the line L_2 at A_1 . Similarly, from point B , a line perpendicular to L is drawn meeting the line L_1 at B_1 . Thus, a parallelogram AA_1BB_1 is formed. Then the equation of L so that the area of the parallelogram AA_1BB_1 is the least is $x - 7y + 17 = 0$ $7x + y + 31 = 0$ $x - 7y - 17 = 0$ $x + 7y - 31 = 0$

 [View Text Solution](#)

42. Are the points $(3, 4)$ and $(2, -6)$ on the same or opposite sides of the line $3x - 4y = 8$?

 [View Text Solution](#)

43. Find the set of positive values of b for which the origin and the point $(1, 1)$ lie on the same side of the straight line, $a^2x + aby + 1 = 0, \forall a \in R$.

 [View Text Solution](#)

44. If the point $(a^2, a + 1)$ lies in the angle between the lines $3x - y + 1 = 0$ and $x + 2y - 5 = 0$ containing the origin, then find the value of a .

 [View Text Solution](#)

45. If the point (a, a) is placed in between the lines $|x + y| = 4$, then find the values of a .

 [View Text Solution](#)

46. The complete set of real values of 'a' such that the point lies triangle $p(a, \sin a)$ lies inside the triangle formed by the lines $x - 2y + 2 = 0$; $x + y = 0$ and $x - y - \pi = 0$

 [View Text Solution](#)

47. Determine all the values of α for which the point (α, α^2) lies inside the triangle formed by the lines. $2x + 3y - 1 = 0$ $x + 2y - 3 = 0$
 $5x - 6y - 1 = 0$

 [View Text Solution](#)

48. Sketch the origin in which the points satisfying the following inequality lie.

(i) $2x - 3y - 5 > 0$ (ii) $-3x + 4y + 7 > 0$

(iii) $x > 2$ (iv) $y > -3$

 [View Text Solution](#)

49. Sketch the origin in which the points satisfying the following inequalities lie.

(i) $|x + y| < 2$ (ii) $|2x - y| > 3$ (iii) $|x| > |y|$

 [View Text Solution](#)

50. Find the values of b for which the points $(2b + 3, b^2)$ lies above of the line $3x - 4y - a(a - 2) = 0 \quad \forall a \in R$.

 [Watch Video Solution](#)

51. Plot the region of the points P (x,y) satisfying $|x| + |y| < 1$.



[Watch Video Solution](#)

52. Plot the region of the points P(x,y) satisfying $2 > \max.$

$\{|x|, |y|\}$.



[View Text Solution](#)

53. IF one of the vertices of a square is (3,2) and one of the diagonals is along the line $3x+4y+8=0$, then find the centre of the square and other vertices.



[View Text Solution](#)

54. In $\triangle ABC$, vertex A is (1, 2). If the internal angle bisector of $\angle B$ is $2x - y + 10 = 0$ and the perpendicular bisector of AC is $y = x$, then find the equation of BC



[Watch Video Solution](#)

55. Find the locus of image of the variable point $(\lambda^2, 2\lambda)$ in the line mirror $x-y+1=0$, where λ is a parameter.



[Watch Video Solution](#)

56. Lines $L_1 \equiv ax + by + c = 0$ and $L_2 \equiv lx + my + n = 0$ intersect at the point P and make an angle θ with each other. Find the equation of a line different from L_2 which passes through P and makes the same angle θ with L_1 .



[Watch Video Solution](#)

57. For the straight lines $4x + 3y - 6 = 0$ and $5x + 12y + 9 = 0$, find the equation of the bisector of the obtuse angle between them, bisector of the acute angle between them, and bisector of the angle which contains $(1, 2)$



[Watch Video Solution](#)

58. The equations of bisectors of two lines L_1 & L_2 are $2x - 16y - 5 = 0$ and $64x + 8y + 35 = 0$. If the line L_1 passes through $(-11, 4)$, the equation of acute angle bisector of L_1 & L_2 is:



[Watch Video Solution](#)

59. If $x + y = 0$ is the angle bisector of the angle containing the point $(1, 0)$, for the line $3x + 4y + b = 0$; $4x + 3y + b = 0$, $4x + 3y - b = 0$ then



[Watch Video Solution](#)

60. Two equal sides of an isosceles triangle are given by $7x - y + 3 = 0$ and $x + y = 3$, and its third side passes through the point $(1, -10)$. Find the equation of the third side.



[Watch Video Solution](#)

61. The vertices B and C of a triangle ABC lie on the lines $3y = 4x$ and $y = 0$, respectively, and the side BC passes through the point $\left(\frac{2}{3}, \frac{2}{3}\right)$. If $ABOC$ is a rhombus lying in the first quadrant, O being the origin, find the equation of the line BC .

 View Text Solution

62. Two sides of a rhombus lying in the first quadrant are given by $3x - 4y = 0$ and $12x - 5y = 0$. If the length of the longer diagonal is 12, then find the equations of the other two sides of the rhombus.

 Watch Video Solution

63. If the line $ax + by = 1$ passes through the point of intersection of $y = x \tan \alpha + p \sec \alpha$, $y \sin(30^\circ - \alpha) - x \cos(30^\circ - \alpha) = p$, and is inclined at 30° with $y = \tan \alpha x$, then prove that $a^2 + b^2 = \frac{3}{4p^2}$.



Watch Video Solution

64. Find the value of λ , if the lines $3x - 4y - 13 = 0$, $8x - 11y - 33 = 0$ and $2x - 3y + \lambda = 0$ are concurrent.



Watch Video Solution

65. If the lines $a_1x + b_1y + 1 = 0$, $a_2x + b_2y + 1 = 0$ and $a_3x + b_3y + 1 = 0$ are concurrent, show that the points (a_1, b_1) , (a_2, b_2) and (a_3, b_3) are collinear.



Watch Video Solution

66. Show that the straight lines given by $x(a + 2b) + y(a + 3b) = a + b$ for different values of a and b pass through a fixed point.



Watch Video Solution

 Watch Video Solution

67. Let $ax + by + c = 0$ be a variable straight line, where a , b and c are the 1st, 3rd, and 7th terms of an increasing AP, respectively. Then prove that the variable straight line always passes through a fixed point. Find that point.

 Watch Video Solution

68. Prove that all the lines having the sum of the intercepts on the axes equal to half of the product of the intercepts pass through the point. Find the fixed point.

 Watch Video Solution

69. Find the straight line passing through the point of intersection of $2x + 3y + 5 = 0$, $5x - 2y - 16 = 0$, and through the point $(-1, 3)$.

 Watch Video Solution

70. Consider a family of straight lines $(x + y) + \lambda(2x - y + 1) = 0$.

Find the equation of the straight line belonging to this family that is farthest from $(1, -3)$.

 [Watch Video Solution](#)

71. Let the sides of a parallelogram be $U=a$, $U=b$, $V=a'$ and $V=b'$, where $U=lx+my+n$, $V=l'x+m'y+n'$. Show that the equation of the diagonal through the point of intersection of

$U = a$, $V = a'$ and $U = b$, $V = b'$ is given by
$$\begin{vmatrix} U & V & 1 \\ a & a' & 1 \\ b & b' & 1 \end{vmatrix} = 0.$$

 [Watch Video Solution](#)

72. A variable line passes through a fixed point P. The algebraic sum of the perpendiculars drawn from the points $(2,0)$, $(0,2)$ and $(1,1)$ on the line is zero. Find the coordinate of the point P.



Watch Video Solution

73. Show that the lines $4x + y - 9 = 0$, $x - 2y + 3 = 0$, $5x - y - 6 = 0$ make equal intercepts on any line of slope 2.



Watch Video Solution

74. The equations of two sides of a triangle are $3y - x - 2 = 0$ and $y + x - 2 = 0$. The third side, which is variable, always passes through the point $(5, -1)$. Find the range of the values of the slope of the third side, so that the origin is an interior point of the triangle.



Watch Video Solution

75. Find the locus of the circumcenter of a triangle whose two sides are along the coordinate axes and the third side passes through the point of

intersection of the line $ax + by + c = 0$ and $lx + my + n = 0$.

 [Watch Video Solution](#)

76. Let ABC be a triangle with $AB = AC$. If D is the midpoint of BC , E is the foot of the perpendicular drawn from D to AC , and F is the midpoint of DE , then prove that AF is perpendicular to BE .

 [Watch Video Solution](#)

77. A diagonal of rhombus $ABCD$ is member of both the families of lines $(x + y - 1) + \lambda_1(2x + 3y - 2) = 0$ and $(x - y + 2) + \lambda_2(2x - 3y + 5) = 0$ and rhombus is $(3, 2)$. If the area of the rhombus is $12\sqrt{5}$ sq. units, then find the remaining vertices of the rhombus.

 [Watch Video Solution](#)

78. Let ABC be a given isosceles triangle with $AB = AC$. Sides AB and AC are extended up to E and F , respectively, such that $BE \times CF = AB^2$. Prove that the line EF always passes through a fixed point.

 [Watch Video Solution](#)

79. Find the straight line passing through the point of intersection of lines $2x+3y+5=0$ and $5x-2y-16=0$ and through the point $(-1,3)$ using the concept of family of lines.

 [Watch Video Solution](#)

80. Find the normal to the curve $x = a(1 + \cos \theta)$, $y = a \sin \theta$. Prove that it always passes through a fixed point and find that fixed point.

 [Watch Video Solution](#)

81. Consider two lines L_1 and L_2 given by $x - y = 0$ and $x + y = 0$, respectively, and a moving point $P(x, y)$. Let $d(P, L_i), i = 1, 2$, represents the distance of point P from the line L_i . If point P moves in a certain region R in such a way that $2 \leq d(P, L_2) + d(P, L_1) \leq 4$, find the area of region R .

 [Watch Video Solution](#)

82. about to only mathematics

 [Watch Video Solution](#)

83. A line through $A(-5, -4)$ meets the lines $x + 3y + 2 = 0$, $2x + y + 4 = 0$ and $x - y - 5 = 0$ at the points B, C and D respectively, if $\left(\frac{15}{AB}\right)^2 + \left(\frac{10}{AC}\right)^2 = \left(\frac{6}{AD}\right)^2$ find the equation of the line.

 [Watch Video Solution](#)

84. A rectangle $PQRS$ has its side PQ parallel to the line $y = mx$ and vertices P, Q and S on the lines $y = a, x = b,$ and $x = -b,$ respectively. Find the locus of the vertex R .

 [Watch Video Solution](#)

Exercise 2 1

1. Find the equation of the right bisector of the line segment joining the points $(3, 4)$ and $(-1, 2)$.

 [Watch Video Solution](#)

2. If the coordinates of the points A, B, C and D be $(a, b), (a', b'), (-a, b)$ and $(a', -b')$ respectively, then the equation of the line bisecting the line segments AB and CD is

 [Watch Video Solution](#)

3. If the coordinates of the vertices of triangle ABC are $(-1, 6)$, $(-3, -9)$ and $(5, -8)$, respectively, then find the equation of the median through C .

 [Watch Video Solution](#)

4. Find the equation of the line perpendicular to the line $\frac{x}{a} - \frac{y}{b} = 1$ and passing through a point at which it cuts the x-axis.

 [Watch Video Solution](#)

5. If the middle points of the sides BC , CA , and AB of triangle ABC are $(1, 3)$, $(5, 7)$, and $(-5, 7)$, respectively, then find the equation of the side AB .

 [Watch Video Solution](#)

6. Find the equations of the lines which pass through the origin and are inclined at an angle $\tan^{-1} m$ to the line $y = mx + c$.

 [Watch Video Solution](#)

7. If $(-2,6)$ is the image of the point $(4,2)$ with respect to line $L=0$, then L is:

 [Watch Video Solution](#)

8. Find the area bounded by the curves $x + 2|y| = 1$ and $x = 0$.

 [Watch Video Solution](#)

9. Find the equation of the straight line passing through the intersection of the lines $x - 2y = 1$ and $x + 3y = 2$ and parallel to $3x + 4y = 0$.

 [Watch Video Solution](#)

10. If the foot of perpendicular from the origin to a straight line is at the point (3,-4). Then the equation of the line is

 [Watch Video Solution](#)

11. A straight line through the point (2, 2) intersects the lines $\sqrt{3}x + y = 0$ and $\sqrt{3}x - y = 0$ at the points A and B. The equation of AB so that the triangle OAB is equilateral, where O is the origin.

 [Watch Video Solution](#)

12. The equation of the straight line passing through the point (4, 3) and making intercepts on the coordinate axes whose sum is -1 is

 [Watch Video Solution](#)

13. A straight line through the point $A(3, 4)$ is such that its intercept between the axis is bisected at A . its equation is

 [Watch Video Solution](#)

14. A straight line L is perpendicular to the line $5x - y = 1$. The area of the triangle formed by line L , and the coordinate axes is 5. Find the equation of line L .

 [Watch Video Solution](#)

15. One side of a rectangle lies along the line $4x + 7y + 5 = 0$. Two of its vertices are $(-3, 1)$ and $(1, 1)$. Find the equations of the other three sides.

 [Watch Video Solution](#)

16. The point of intersection of the curves $y^2 = 4x$ and the line $y = x$ is

 [Watch Video Solution](#)

17. The diagonals AC and BD of a rhombus intersect at $(5, 6)$. If $A \equiv (3, 2)$, then find the equation of diagonal BD .

 [Watch Video Solution](#)

18. Find the equation of the straight line which passes through the origin and makes angle 60° with the line $x + \sqrt{3}y + \sqrt{3} = 0$.

 [Watch Video Solution](#)

19. If $A(-6, -6)$ and $B(-6, 4)$ be two points that a point P on the line AB satisfies $AP = \frac{2}{9}AB$. find the point P

 [Watch Video Solution](#)

20. In the adjoining figure , ΔPQR is an equilateral triangle. $QR = RN$.

Prove that $PN^2 = 3PR^2$



[Watch Video Solution](#)

21. Two fixed points A and B are taken on the coordinates axes such that $OA = a$ and $OB = b$. Two variable points A' and B' are taken on the same axes such that $OA' + OB' = OA + OB$. Find the locus of the point of intersection of AB' and $A'B$.



[Watch Video Solution](#)

22. A regular polygon has two of its consecutive diagonals as the lines $\sqrt{3}x + y - \sqrt{3}$ and $2y = \sqrt{3}$. Point $(1, c)$ is one of its vertices. Find the equation of the sides of the polygon and also find the coordinates of the vertices.



[Watch Video Solution](#)

23. Find the direction in which a straight line must be drawn through the point $(1, 2)$ so that its point of intersection with the line $x + y = 4$ may be at a distance of 3 units from this point.



[Watch Video Solution](#)

Exercise 2.2

1. Find the points on the line $x + y = 1$ that lie at a distance 3 units from the line $5x + 12y = 3$.



[Watch Video Solution](#)

2. The center of a square is at the origin and its one vertex is $A(2, 1)$. Find the coordinates of the other vertices of the square.



Watch Video Solution

3. The straight line passing through $P(x_1, y_1)$ and making an angle α with x-axis intersects $Ax + By + C = 0$ in Q then PQ=



Watch Video Solution

4. The centroid of an equilateral triangle is $(0,0)$. If two vertices of the triangle lie on $x+y = 2\sqrt{2}$, then find all the possible vertices fo triangle.



Watch Video Solution

Exercise 2 3

1. Find the points on $y - a\xi s$ whose perpendicular distance from the line $4x - 3y - 12 = 0$ is 3.



Watch Video Solution

2. If p and q are the lengths of perpendiculars from the origin to the lines $x \cos \theta - y \sin \theta = k \cos 2\theta$ and $x \sec \theta + y \csc \theta = k$, respectively, prove that $p^2 + 4q^2 = k^2$.



[Watch Video Solution](#)

3. Prove that the lengths of the perpendiculars from the points $(m^2, 2m)$, $(mm', m + m')$, and $(m'^2, 2m')$ to the line $x + y + 1 = 0$ are in GP.



[Watch Video Solution](#)

4. The ratio in which the line $3x+4y+2 = 0$ divides the distance between $3x + 4y + 5 = 0$ and $3x + 4y - 5 = 0$



[Watch Video Solution](#)

5. Find the rectangular form of the complex numbers.

$$\left(\cos \frac{\pi}{6} + i \sin \frac{\pi}{6}\right) \left(\cos \frac{\pi}{12} + i \sin \frac{\pi}{12}\right)$$



[Watch Video Solution](#)

6. Find the equations of lines parallel to $3x - 4y - 5 = 0$ at a unit distance from it.



[Watch Video Solution](#)

7. Find the equation of a straight line passing through the point $(-5, 4)$ and which cuts off an intercept of $\sqrt{2}$ units between the lines $x + y + 1 = 0$ and $x + y - 1 = 0$.



[Watch Video Solution](#)

1. The point $(8, -9)$ with respect to the lines $2x + 3y - 4 = 0$ and $6x + 9y + 8 = 0$ lies on the same side of the lines the different sides of the line one of the line none of these

 [Watch Video Solution](#)

2. How the following pairs of points are placed w.r.t the line $3x - 8y - 7 = 0$?

(i) $(-3, -4)$ and $(1, 2)$ (ii) $(-1, -1)$ and $(3, 7)$

 [Watch Video Solution](#)

3. Find the range of $(\alpha, 2 + \alpha)$ and $\left(\frac{3\alpha}{2}, a^2\right)$ lie on the opposite sides of the line $2x + 3y = 6$.

 [Watch Video Solution](#)

4. If the point $P(a^2, a)$ lies in the region corresponding to the acute angle between the lines $2y = x$ and $4y = x$, then find the values of a .



Watch Video Solution

5. If $(a, 3a)$ is a variable point lying above the straight line $2x+y+4=0$ and below the line $x+4y-8=0$, then find the values of a .



Watch Video Solution

6. Find the values of α such that the variable point $(\alpha, \tan\alpha)$ lies inside the triangle whose sides are

$$y = x + \sqrt{3} - \frac{\pi}{3}, x + y + \frac{1}{\sqrt{3}} + \frac{\pi}{6} = 0 \text{ and } x - \frac{\pi}{2} = 0$$



Watch Video Solution

7. Find the area of the region in which points satisfy

$$3 \leq |x| + |y| \leq 5.$$



Watch Video Solution

8. $x - y \leq 2$



Watch Video Solution

Exercise 2 5

1. Find the equation of the bisector of the obtuse angle between the lines $3x - 4y + 7 = 0$ and $12x + 5y - 2 = 0$.



Watch Video Solution

2. The incident ray is along the line $3x - 4y - 3 = 0$ and the reflected ray is along the line $24x + 7y + 5 = 0$. Find the equation of mirrors.



Watch Video Solution

3. If the two sides of rhombus are $x + 2y + 2 = 0$ and $2x + y - 3 = 0$, then find the slope of the longer diagonal.

 [Watch Video Solution](#)

4. In triangle ABC , the equation of the right bisectors of the sides AB and AC are $x + y = 0$ and $y - x = 0$, respectively. If $A \equiv (5, 7)$, then find the equation of side BC .

 [Watch Video Solution](#)

5. Show that the reflection of the line $ax + by + c = 0$ on the line $x + y + 1 = 0$ is the line $b + ay + (a + b - c) = 0$ where $a \neq b$.

 [Watch Video Solution](#)

6. The joint equation of two altitudes of an equilateral triangle is $(\sqrt{3}x - y + 8 - 4\sqrt{3})(-\sqrt{3}x - y + 12 + 4\sqrt{3}) = 0$ The third altitude has the equation

 [Watch Video Solution](#)

7. The equations of the perpendicular bisectors of the sides AB and AC of triangle ABC are $x - y + 5 = 0$ and $x + 2y = 0$, respectively. If the point A is $(1, -2)$, then find the equation of the line BC .

 [Watch Video Solution](#)

8. about to only mathematics

 [Watch Video Solution](#)

1. If a and b are two arbitrary constants, then prove that the straight line $(a - 2b)x + (a + 3b)y + 3a + 4b = 0$ will pass through a fixed point.

Find that point.



[Watch Video Solution](#)

2. If a, b, c are in harmonic progression, then the straight line $\left(\frac{x}{a}\right) + \left(\frac{y}{b}\right) + \left(\frac{1}{c}\right) = 0$ always passes through a fixed point. Find that point.



[Watch Video Solution](#)

3. A variable line passes through a fixed point P. The algebraic sum of the perpendiculars drawn from the points $(2,0)$, $(0,2)$ and $(1,1)$ on the line is zero. Find the coordinate of the point P.



[Watch Video Solution](#)

4. Consider the family of lines

$$5x + 3y - 2 + \lambda_1(3x - y - 4) = 0 \text{ and } x - y + 1 + \lambda_2(2x - y - 2) = 0$$

. Find the equation of a straight line that belongs to both the families.



[Watch Video Solution](#)

5. If the straight lines $x + y - 2 = 0$, $2x - y + 1 = 0$ and

$ax + by - c = 0$ are concurrent, then the family of lines

$2ax + 3by + c = 0$ (a, b, c are nonzero) is concurrent at (2, 3) (b)

$$\left(\frac{1}{2}, \frac{1}{3}\right) \left(-\frac{1}{6}, -\frac{5}{9}\right) \text{ (d) } \left(\frac{2}{3}, -\frac{7}{5}\right)$$



[Watch Video Solution](#)

Exercise Single

1. Find the equations of the diagonals of the square formed by the lines

$$x = 0, y = 0, x = 1 \text{ and } y = 1.$$

A. $y=x, y+x=1$

B. $y=x, x+y=2$

C. $2y = x, y+x = 1/3$

D. $y=2x, y+2x = 1$

Answer: A



Watch Video Solution

2. The coordinates of two consecutive vertices A and B of a regular hexagon $ABCDEF$ are $(1, 0)$ and $(2, 0)$, respectively. The equation of the diagonal CE is

A. $\sqrt{3}x + y = 4$

B. $x + \sqrt{3}y + 4 = 0$

C. $x + \sqrt{3}y = 4$

D. none of these

Answer: C



Watch Video Solution

3. If each of the points $(x, 4)$, $(-2, y)$ lie on the line joining the points $(2, -1)$ and $(5, -3)$ then the point $P(x_1, y_1)$ lies on the line

A. $6(x + y) - 25 = 0$

B. $2x + 6y + 1 = 0$

C. $2x + 3y - 6 = 0$

D. $6(x + y) + 25 = 0$

Answer: B



Watch Video Solution

4. The equation of a straight line which passes through the point $(a \cos^3 \theta, a \sin^3 \theta)$ and perpendicular to $x \sec \theta + y \operatorname{cosec} \theta = a$

A. $x \cos \theta - y \sin \theta = a \cos 2\theta$

B. $x \cos \theta + y \sin \theta = a \cos 2\theta$

C. $x \sin \theta + y \cos \theta = a \cos 2\theta$

D. none of these

Answer: A



Watch Video Solution

5. The line PQ whose equation is $x - y = 2$ cuts the x-axis at P , and Q is $(4,2)$. The line PQ is rotated about P through 45° in the anticlockwise direction. The equation of the line PQ in the new position is $y = -\sqrt{2}$

(b) $y = 2$ (c) $x = 2$ (d) $x = -2$

A. $y = -\sqrt{2}$

B. $y=2$

C. $x=2$

D. $x=-2$

Answer: C



Watch Video Solution

6. If $P(a,b)$ is the mid point of a line segment between the axes, then:

A. $x+y=2c$

B. $x+y=c$

C. $2(x+y)=c$

D. $2x+y=c$

Answer: C



Watch Video Solution

7. If the x intercept of the line $y = mx + 2$ is greater than $\frac{1}{2}$ then the gradient of the line lies in the interval

A. $(-1,0)$

B. $\left(\frac{-1}{4}, 0\right)$

C. $(-\infty, -4)$

D. $(-4, 0)$

Answer: D



Watch Video Solution

8. The equation of a straight line on which the length of perpendicular from the origin is four units and the line makes an angle of 120° with the x-axis is $x\sqrt{3} + y + 8 = 0$ $x\sqrt{3} - y = 8$ $x\sqrt{3} - y = 8$ (d)

$x - \sqrt{3}y + 8 = 0$

A. $x\sqrt{3} + y + 8 = 0$

B. $x\sqrt{3} - y = 8$

C. $x\sqrt{3} - y = 8$

D. $x - \sqrt{3}y + 8 = 0$

Answer: A



[Watch Video Solution](#)

9. $ABCD$ is a square $A \equiv (1, 2)$, $B \equiv (3, -4)$. If line CD passes through $(3, 8)$, then the midpoint of CD is

A. (2,6)

B. (6,2)

C. (2,5)

D. $(\frac{28}{5}, \frac{1}{5})$

Answer: D



[Watch Video Solution](#)

10. Find the equation of a line which passes through the point $(2, 3, 4)$ and which has equal intercepts on the axes.

A. $9x-20y+96=0$

B. $9x+20y=24$

C. $20x+9y+53=0$

D. none of these

Answer: A



Watch Video Solution

11. A square of side a lies above the x -axis and has one vertex at the origin.

The side passing through the origin makes an angle α ($0 < \alpha < \frac{\pi}{4}$)

with the positive direction of x -axis. The equation of its diagonal not passing through the origin is

A. $y(\cos\alpha + \sin\alpha) + x(\sin\alpha - \cos\alpha) = a$

B. $y(\cos\alpha + \sin\alpha) + x(\sin\alpha + \cos\alpha) = a$

C. $y(\cos\alpha + \sin\alpha) + x(\cos\alpha - \sin\alpha) = a$

D. $y(\cos\alpha - \sin\alpha) - x(\sin\alpha - \cos\alpha) = a$

Answer: C



Watch Video Solution

12. Let $P=(-1, 0)$, $Q=(0,0)$ and $R=(3, 3\sqrt{3})$ be three points. The equation of the bisector of the angle PQR is

A. $(\sqrt{3}/2)x + y = 0$

B. $x + \sqrt{3}y = 0$

C. $\sqrt{3}x + y = 0$

D. $x + (\sqrt{3}/2)y = 0$

Answer: C



Watch Video Solution

13. The equation of a line through the point $(1, 2)$ whose distance from the point $(3, 1)$ has the greatest value, is

A. $y=2x$

B. $y=x+1$

C. $x+2y=5$

D. $y=3x-1$

Answer: A



Watch Video Solution

14. If one of the sides of a square is $3x - 4y - 12 = 0$ and the center is $(0, 0)$, then find the equations of the diagonals of the square.

A. $7x-8y+9=0, 8x+7y-22=0$

B. $9x-8y+7=0, 8x+9y-26=0$

C. $23x-7y-9=0, 7x+23y-53=0$

D. none of these

Answer: C

 [Watch Video Solution](#)

15. The locus of the centers of the circles $(x - 1)^2 + y^2 = 10$ and $x^2 + (y - 2)^2 = 5$ intersect is $\frac{\pi}{6}$ (b) $\frac{\pi}{4}$ (c) $\frac{\pi}{3}$ (d) $\frac{\pi}{2}$

A. $\frac{\pi}{4}$

B. $\frac{\pi}{2}$

C. $\frac{\pi}{3}$

D. $\frac{\pi}{6}$

Answer: B

 [Watch Video Solution](#)

16. A line with positive rational slope, passes through the point A(6,0) and is at a distance of 5 units from B (1,3). The slope of line is

A. $\frac{15}{8}$

B. $\frac{8}{15}$

C. $\frac{5}{8}$

D. $\frac{8}{5}$

Answer: B



Watch Video Solution

17. The line $2x - y = 1$ bisects angle between two lines. If equation of one line is $y = x$, then the equation of the other line is

A. $3x+3y-1=0$

B. $x-3y+2=0$

C. $5x+5y-3=0$

D. none of these

Answer: C

 [Watch Video Solution](#)

18. If R is any point on the x-axis and Q is any point on the y-axis and P is a variable point on RQ with $RP=b$, $PQ=a$, then find the equation of locus of P.

A. $x+y=1$

B. $x+y=2$

C. $x+y=2xy$

D. $2x+2y=1$

Answer: A

 [Watch Video Solution](#)

19. The number of possible straight lines passing through point $(2,3)$ and forming a triangle with coordinate axes whose area is 12 sq. unit is: a. one b. two c. three d. four

A. one

B. two

C. three

D. four

Answer: C



Watch Video Solution

20. Two parallel lines lying in the same quadrant make intercepts a and b on x and y axes, respectively, between them. The distance between the lines is (a) $\frac{ab}{\sqrt{a^2 + b^2}}$ (b) $\sqrt{a^2 + b^2}$ (c) $\frac{1}{\sqrt{a^2 + b^2}}$ (d) $\frac{1}{a^2} + \frac{1}{b^2}$

A. $\sqrt{a^2 + b^2}$

B. $\frac{ab}{\sqrt{a^2 + b^2}}$

C. $\frac{1}{\sqrt{a^2 + b^2}}$

D. $\frac{1}{a^2} + \frac{1}{b^2}$

Answer: B



Watch Video Solution

21. The line $L_1 \equiv 4x + 3y - 12 = 0$ intersects the x-and y-axies at A and B , respectively. A variable line perpendicular to L_1 intersects the x- and the y-axis at P and Q , respectively. Then the locus of the circumcenter of triangle ABQ is $3x - 4y + 2 = 0$ $4x + 3y + 7 = 0$ $6x - 8y + 7 = 0$ (d) none of these

A. $3x - 4y + 2 = 0$

B. $4x + 3y + 7 = 0$

C. $6x - 8y + 7 = 0$

D. none of these

Answer: C



Watch Video Solution

22. A beam of light is sent along the line $x - y = 1$, which after refracting from the x-axis enters the opposite side by turning through 30° towards the normal at the point of incidence on the x-axis. Then the

equation of the refracted ray is $(2 - \sqrt{3})x - y = 2 + \sqrt{3}$

$$(2 + \sqrt{3})x - y = 2 + \sqrt{3}$$

$$(2 - \sqrt{3})x + y = (2 + \sqrt{3})$$

$$y = (2 - \sqrt{3})(x - 1)$$

A. $(2 - \sqrt{3})x - y = 2 + \sqrt{3}$

B. $(2 + \sqrt{3})x - y = 2 + \sqrt{3}$

C. $(2 - \sqrt{3})x + y = (2 + \sqrt{3})$

D. $y = (2 + \sqrt{3})(x - 1)$

Answer: D



Watch Video Solution

23. The number of integral values of m for which the x-coordinate of the point of intersection of the lines $3x + 4y = 9$ and $y = mx + 1$ is also an

integer is (a) 2 (b) 0 (c) 4 (d) 1

A. 2

B. 0

C. 4

D. 1

Answer: A



[Watch Video Solution](#)

24. If the sum of the distances of a point from two perpendicular lines in a plane is 1, then its locus is a (a) square (b) a circle (c) a straight line (d) two intersecting lines

A. a square

B. a circle

C. a straight line

D. two intersecting lines

Answer: A



[Watch Video Solution](#)

25. From the differential equation of family of lines situated at a constant distance p from the origin.

A. $x+y+2=0$

B. $x+y+4=0$

C. $x \cos \alpha + y \sin \alpha = 2$

D. $x \cos \alpha + y \sin \alpha = \frac{1}{2}$

Answer: C



[Watch Video Solution](#)

26. The lines $y = m_1x$, $y = m_2x$ and $y = m_3x$ make equal intercepts on the line $x + y = 1$. Then

$$2(1 + m_1)(1 + m_3) = (1 + m_2)(2 + m_1 + m_3)$$

$$(1 + m_1)(1 + m_3) = (1 + m_2)(1 + m_1 + m_3)$$

$$(1 + m_1)(1 + m_2) = (1 + m_3)(2 + m_1 + m_3)$$

$$2(1 + m_1)(1 + m_3) = (1 + m_2)(1 + m_1 + m_3)$$

A. $2(1 + m_1)(1 + m_3) = (1 + m_2)(2 + m_1 + m_3)$

B. $(1 + m_1)(1 + m_3) = (1 + m_2)(1 + m_1 + m_3)$

C. $(1 + m_1)(1 + m_2) = (1 + m_3)(2 + m_1 + m_3)$

D. $2(1 + m_1)(1 + m_3) = (1 + m_2)(1 + m_1 + m_3)$

Answer: A



Watch Video Solution

27. The condition on a and b , such that the portion of the line $ax + by - 1 = 0$ intercepted between the lines $ax + y = 0$ and

$x + by = 0$ subtends a right angle at the origin, is (a) $a = b$ (b) $a + b = 0$

(c) $a = 2b$ (d) $2a = b$

A. $a = b$

B. $a + b = 0$

C. $a = 2b$

D. $2a = b$

Answer: B



Watch Video Solution

28. The area of the triangle formed by the lines $y = ax$, $x + y - a = 0$, and y -axis is equal to

A. $\frac{1}{2|1+a|}$

B. $\frac{a^2}{|1+a|}$

C. $\frac{1}{2} \frac{a}{|1+a|}$

D. $\frac{a^2}{2|1+a|}$

Answer: D



Watch Video Solution

29. The line $\frac{x}{a} + \frac{y}{b} = 1$ meets the x-axis at A , the y-axis at B , and the line $y = x$ at C such, that the area of ΔAOC is twice the area of ΔBOC . Then the coordinates of C are $\left(\frac{b}{3}, \frac{b}{3}\right)$ (b) $\left(\frac{2a}{3}, \frac{2a}{3}\right)$ $\left(\frac{2b}{3}, \frac{2b}{3}\right)$ (d) none of these

A. $\left(\frac{b}{3}, \frac{b}{3}\right)$

B. $\left(\frac{2a}{3}, \frac{2a}{3}\right)$

C. $\left(\frac{2b}{3}, \frac{2b}{3}\right)$

D. none of these

Answer: C



Watch Video Solution

30. The line $\frac{x}{3} + \frac{y}{4} = 1$ meets the y - and x -axes at A and B , respectively. A square $ABCD$ is constructed on the line segment AB away from the origin. The coordinates of the vertex of the square farthest from the origin are (a) (7, 3) (b) (4, 7) (c) (6, 4) (d) (3, 8)

A. 7,3

B. 4,7

C. 6,4

D. 3,8

Answer: B



[Watch Video Solution](#)

31. The area of a parallelogram formed by the lines $ax \pm bx \pm c = 0$ is (a)

$\frac{c^2}{(ab)}$ (b) $\frac{sc^2}{(ab)}$ (c) $\frac{c^2}{2ab}$ (d) none of these

A. $c^2 / (ab)$

B. $sc^2 / (ab)$

C. $c^2 / 2ab$

D. none of these

Answer: D



[Watch Video Solution](#)

32. One diagonal of a square is $3x-4y+8=0$ and one vertex is $(-1,1)$, then the area of square is

A. $\frac{1}{50}$ sq.unit

B. $\frac{1}{25}$ sq.unit

C. $\frac{3}{50}$ sq.unit

D. $\frac{2}{25}$ sq.unit

Answer: D



[Watch Video Solution](#)

33. In an isosceles triangle OAB , O is the origin and $OA=OB=6$. The equation of the side AB is $x-y+1=0$ Then the area of the triangle is

A. $2\sqrt{21}$

B. $\sqrt{142}$

C. $\sqrt{\frac{142}{2}}$

D. $\sqrt{\frac{71}{2}}$

Answer: D



Watch Video Solution

34. A straight line through the origin 'O' meets the parallel lines $4x + 2y = 9$ and $2x + y = -6$ at points P and Q respectively. Then the point 'O' divides the segment PQ in the ratio

A. 1:2

B. 3: 4

C. 2: 01

D. 4: 3

Answer: B



Watch Video Solution

35. The coordinates of the foot of the perpendicular from the point $(2, 3)$ on the line $-y + 3x + 4 = 0$ are given by $\left(\frac{37}{10}, -\frac{1}{10}\right)$ (b) $\left(-\frac{1}{10}, \frac{37}{10}\right)$ $\left(\frac{10}{37}, -10\right)$ (d) $\left(\frac{2}{3}, -\frac{1}{3}\right)$

A. $(37/10, -1/10)$

B. $(-1/10, 37/10)$

C. $(10/37, -10)$

D. $(2/3, -1/3)$

Answer: B



[Watch Video Solution](#)

36. The straight lines $7x - 2y + 10 = 0$ and $7x + 2y - 10 = 0$ form an isosceles triangle with the line $y = 2$. The area of this triangle is equal to

(a) $\frac{15}{7}$ sq units (b) $\frac{10}{7}$ sq units (c) $\frac{18}{7}$ sq units (d) none of these

A. $15/7$ sq. units

B. $10/7$ sq. units

C. $18/7$ sq. units

D. none of these

Answer: C



[Watch Video Solution](#)

37. The equations of the sides of a triangle are $x+y-5=0$, $x-y+1=0$, and $y-1=0$.

Then the coordinates of the circumcenter are

A. 2,1

B. 1,2

C. 2,-2

D. 1,-2

Answer: A



Watch Video Solution

38. If the intercepts made by the line $y = mx$ by lines $x = 2$ and $x = 5$ is less than 5, then the range of values of m is a. $\left(-\infty, -\frac{4}{3}\right) \cup \left(\frac{4}{3}, \infty\right)$ b. $\left(-\frac{4}{3}, \frac{4}{3}\right)$ c. $\left(-\frac{3}{4}, \frac{4}{3}\right)$ d. none of these

A. $\left(-\infty, -\frac{4}{3}\right) \cup \left(\frac{4}{3}, +\infty\right)$

B. $\left(-\frac{4}{3}, \frac{4}{3}\right)$

C. $\left(-\frac{3}{4}, \frac{4}{3}\right)$

D. none of these

Answer: A



Watch Video Solution

39. The range of values of θ in the interval $(0, \pi)$ such that the points $(3, 2)$ and $(\sin \theta, \cos \theta)$ lie on the same side of the line $x + y - 1 = 0$, is

A. $0 < \theta < \frac{\pi}{4}$

B. $0 < \theta < \frac{\pi}{2}$

C. $0 < \theta < \pi$

D. $\frac{\pi}{4} < \theta < \frac{3\pi}{4}$

Answer: B



Watch Video Solution

40. about to only mathematics

A. $\frac{5}{\sqrt{2}}$

B. $5\sqrt{2}+k$

C. 10

D. 5

Answer: D

 [Watch Video Solution](#)

41. Consider the points $A(0, 1)$ and $B(2, 0)$, and P be a point on the line $4x + 3y + 9 = 0$. The coordinates of P such that $|PA - PB|$ is maximum are $\left(-\frac{12}{5}, \frac{17}{5}\right)$ (b) $\left(-\frac{84}{5}, \frac{13}{5}\right)$ $\left(\frac{31}{7}, \frac{31}{7}\right)$ (d) $(, 0)$

 [Watch Video Solution](#)

42. Consider the point $A = (3, 4)$, $B(7, 13)$. If 'P' be a point on the line $y = x$ such that $PA + PB$ is minimum then coordinates of P is (A) $\left(\frac{13}{7}, 13, 7\right)$ (B) $\left(\frac{23}{7}, \frac{23}{7}\right)$ (C) $\left(\frac{31}{7}, \frac{31}{7}\right)$ (D) $\left(\frac{33}{7}, \frac{33}{7}\right)$

A. $(12/7, 12/7)$

B. $(-24/5, 17/5)$

C. $(31/7, 31/7)$

D. $(0, 0)$

Answer: C



[Watch Video Solution](#)

43. The area enclosed by $2|x| + 3|y| \leq 6$ is 3 sq. units (b) 4 sq. units 12 sq. units (d) 24 sq. units

A. 3 sq. units

B. 4 sq. units

C. 12 sq. units

D. 24 sq. units

Answer: C

 [Watch Video Solution](#)

44. ABC is a variable triangle such that A is $(1,2)$ and B and C lie on line $y = x + \lambda$ (where λ is a variable). Then the locus of the orthocentre of triangle ABC is $(x - 1)^2 + y^2 = 4$ $x + y = 3$ $2x - y = 0$ (d) none of these

A. $x+y=0$

B. $x-y=0$

C. $x^2 + y^2 = 4$

D. $x+y=3$

Answer: D

 [Watch Video Solution](#)

45. In ABC , the coordinates of the vertex A are $(4, -1)$, and lines $x - y - 1 = 0$ and $2x - y = 3$ are the internal bisectors of angles

Band C. Then, the radius of the encircle of triangle ABC is $\frac{4}{\sqrt{5}}$ (b) $\frac{3}{\sqrt{5}}$

(c) $\frac{6}{\sqrt{5}}$ (d) $\frac{7}{\sqrt{5}}$

A. $4/\sqrt{5}$

B. $3/\sqrt{5}$

C. $6/\sqrt{5}$

D. $7/\sqrt{5}$

Answer: C



Watch Video Solution

46. P is a point on the line $y + 2x = 1$, and Q and R two points on the line $3y + 6x = 6$ such that triangle PQR is an equilateral triangle. The length of the side of the triangle is (a) $\frac{2}{\sqrt{5}}$ (b) $\frac{3}{\sqrt{5}}$ (c) $\frac{4}{\sqrt{5}}$ (d) none of these

A. $2/\sqrt{15}$

B. $3/\sqrt{5}$

C. $4/\sqrt{5}$

D. none of these

Answer: A



Watch Video Solution

47. If the equation of base of an equilateral triangle is $2x - y = 1$ and the vertex is $(-1, 2)$, then the length of the side of the triangle is

A. $\sqrt{20/3}$

B. $2/\sqrt{15}$

C. $\sqrt{8/15}$

D. $\sqrt{15/2}$

Answer: A



Watch Video Solution

48. The locus of a point that is equidistant from the lines $x + y - 2\sqrt{2} = 0$ and $x + y - \sqrt{2} = 0$ is (a) $x + y - 5\sqrt{2} = 0$ (b) $x + y - 3\sqrt{2} = 0$ (c) $2x + 2y - 3\sqrt{2} = 0$ (d) $2x + 2y - 5\sqrt{5} = 0$

A. $x + y - 5\sqrt{2} = 0$

B. $x + y - 3\sqrt{2} = 0$

C. $2x + 2y - 3\sqrt{2} = 0$

D. $2x + 2y - 5\sqrt{2} = 0$

Answer: C



Watch Video Solution

49. If the quadrilateral formed by the lines $ax + by + c = 0$, $a'x + b'y + c = 0$, $ax + by + c' = 0$, $a'x + b'y + c' = 0$ has perpendicular diagonals, then $b^2 + c^2 = b'^2 + c'^2$ (a) $c^2 + a^2 = c'^2 + a'^2$ (b) $a^2 + b^2 = a'^2 + b'^2$ (c) $c^2 + a^2 = c'^2 + a'^2$ (d) none of these

A. $b^2 + c^2 = b'^2 + c'^2$

B. $c^2 + a^2 = c'^2 + a'^2$

C. $a^2 + b^2 = a'^2 + b'^2$

D. none of these

Answer: C



Watch Video Solution

50. A line of fixed length 2 units moves so that its ends are on the positive x-axis and that part of the line $x + y = 0$ which lies in the second quadrant. Then the locus of the midpoint of the line has equation. a.

$x^2 + 5y^2 + 4xy - 1 = 0$ b. $x^2 + 5y^2 + 4xy + 1 = 0$ c.

$x^2 + 5y^2 - 4xy - 1 = 0$ d. $4x^2 + 5y^2 + 4xy + 1 = 0$

A. $x^2 + 5y^2 + 4xy - 1 = 0$

B. $x^2 + 5y^2 + 4xy + 1 = 0$

C. $x^2 + 5y^2 - 4xy - 1 = 0$

D. $x^2 + 5y^2 - 4xy - 1 = 0$

Answer: A



Watch Video Solution

51. If the extremities of the base of an isosceles triangle are the points $(2a, 0)$ and $(0, a)$, and the equation of one of the side is $x = 2a$, then the area of the triangle is (a) $5a^2$ sq units (b) $\frac{5a^2}{2}$ sq units (c) $\frac{25a^2}{2}$ sq units (d) none of these

A. $5a^2$ sq. units

B. $5a^2 / 2$ sq. units

C. $25a^2 / 2$ sq. units

D. none of these

Answer: B



Watch Video Solution

52. $A \equiv (-4, 0)$, $B \equiv (4, 0)$ and N are the variable points of the y -axis such that M lies below N and $MN = 4$. Lines AM and BN intersect at P . The locus of P is

a. $2xy - 16 - x^2 = 0$ b. $2xy + 16 - x^2 = 0$ c. $2xy + 16 + x^2 = 0$ d. $2xy - 16 + x^2 = 0$

A. $2xy - 16 - x^2 = 0$

B. $2xy + 16 - x^2 = 0$

C. $2xy + 16 + x^2 = 0$

D. $2xy - 16 + x^2 = 0$

Answer: D



Watch Video Solution

53. The number of triangles that the four lines $y = x + 3$, $y = 2x + 3$, $y = 3x + 2$, and $y + x = 3$ form is (a) 4 (b) 2 (c) 3 (d) 1

A. 4

B. 2

C. 3

D. 1

Answer: C



Watch Video Solution

54. A variable line $\frac{x}{a} + \frac{y}{b} = 1$ moves in such a way that the harmonic mean of a and b is 8. Then the least area of triangle made by the line with the coordinate axes is (1) 8 sq. unit (2) 16 sq. unit (3) 32 sq. unit (4) 64 sq. unit

A. 8 sq. unit

B. 16 sq. unit

C. 32 sq. unit

D. 64 sq. unit

Answer: C



Watch Video Solution

55. Two $A(0, 0)$ and $B(x, y)$ with $x \in (0, 1)$ and $y > 0$. Let the slope of line AB be m_1 . Point C lies on line $x = 1$ such that the slope of BC is equal to m_2 , where $\frac{m_1}{m_2} = 0$

A. 1

B. $1/2$

C. $1/4$

D. $1/8$

Answer: D



Watch Video Solution

56. A triangle is formed by the lines $x + y = 0$, $x - y = 0$, and $lx + my = 1$. If l and m vary subject to the condition $l^2 + m^2 = 1$, then the locus of its circumcenter is $(x^2 - y^2)^2 = x^2 + y^2$

$$(x^2 + y^2)^2 = (x^2 - y^2)(x^2 + y^2)^2 = 4x^2y^2(x^2 - y^2)^2 = (x^2 + y^2)^2$$

- A. $(x^2 - y^2)^2 = x^2 + y^2$
- B. $(x^2 - y^2)^2 = (x^2 - y^2)$
- C. $(x^2 - y^2) = 4x^2y^2$
- D. $(x^2 - y^2)^2 = (x^2 + y^2)^2$

Answer: A



Watch Video Solution

57. Let P be $(5, 3)$ and a point R on $y = x$ and Q on the x-axis be such that $PQ + QR + RP$ is minimum. Then the coordinates of Q are $\left(\frac{17}{4}, 0\right)$ (b) $(17, 0)$ $\left(\frac{17}{2}, 0\right)$ (d) none of these

A. $(17/4,0)$

B. $(17,0)$

C. $(17/2,0)$

D. none of these

Answer: A



Watch Video Solution

58. If a pair of perpendicular straight lines drawn through the origin forms an isosceles triangle with the line $2x + 3y = 6$, then area of the triangle so formed is $36/13$ (b) $12/17$ (c) $13/5$ (d) $17/14$

A. $\frac{36}{13}$ sq. unit

B. $\frac{12}{17}$ sq. unit

C. $\frac{13}{5}$ sq. unit

D. $\frac{17}{13}$ sq. unit

Answer: A



[Watch Video Solution](#)

59. A point $P(x, y)$ moves that the sum of its distance from the lines $2x - y - 3 = 0$ and $x + 3y + 4 = 0$ is 7. The area bounded by locus P is (in sq. unit)

A. 70

B. $70\sqrt{2}$

C. $35\sqrt{2}$

D. 140

Answer: B



[Watch Video Solution](#)

60. If AD, BE and CF are the altitudes of $\triangle ABC$ whose vertex A is (-4,5). The coordinates of points E and F are (4,1) and (-1,-4), respectively.

Equation of BC is

A. $3x-4y+28=0$

B. $4x+3y+28=0$

C. $3x-4y-28=0$

D. $x+2y+7=0$

Answer: C



[Watch Video Solution](#)

61. The vertex A of $\triangle ABC$ is (3,-1). The equation of median BE and angle bisector CF are $x-4y+10=0$ and $6x+10y-59=0$, respectively. Equation of AC is

A. $5x+18y=37$

B. $15x+8y=37$

C. $15x-8y=37$

D. $15x+8y+37=0$

Answer: B

 [Watch Video Solution](#)

62. Suppose A, B are two points on $2x - y + 3 = 0$ and $P(1, 2)$ is such that $PA=PB$. Then the mid point of AB is

A. $\left(\frac{-1}{5}, \frac{13}{5}\right)$

B. $\left(\frac{-7}{5}, \frac{9}{5}\right)$

C. $\left(\frac{7}{5}, \frac{-9}{5}\right)$

D. $\left(\frac{-7}{5}, \frac{-9}{5}\right)$

Answer: A

 [Watch Video Solution](#)

63. about to only mathematics

- A. equilateral
- B. right angled
- C. scalene
- D. none of these

Answer: D



Watch Video Solution

64. A light ray coming along the line $3x + 4y = 5$ gets reflected from the

line $ax + by = 1$ and goes along the line $5x - 12y = 10$. Then,

$$a = \frac{64}{115}, b = \frac{112}{15} \quad a = \frac{14}{15}, b = -\frac{8}{115} \quad a = \frac{64}{115}, b = -\frac{8}{115}$$

$$a = \frac{64}{15}, b = \frac{14}{15}$$

A. $a = \frac{64}{115}, b = \frac{112}{15}$

B. $a = \frac{14}{15}, b = -\frac{18}{115}$

$$C. a = \frac{64}{115}, b = -\frac{8}{115}$$

$$D. a = \frac{64}{15}, b = \frac{14}{15}$$

Answer: C



Watch Video Solution

65. The point $A(2, 1)$ is translated parallel to the line $x - y = 3$ by a distance of 4 units. If the new position A' is in the third quadrant, then the coordinates of A' are (A) $(2 + 2\sqrt{2}, 1 + 2\sqrt{2})$ (B) $(-2 + \sqrt{2}, -1 - 2\sqrt{2})$ (C) $(2 - 2\sqrt{2}, 1 - 2\sqrt{2})$ (D) none of these

A. $(2 + 2\sqrt{2}, 1 + 2\sqrt{2})$

B. $(-2 + \sqrt{2}, -1 - 2\sqrt{2})$

C. $(2 - 2\sqrt{2}, 1 - 2\sqrt{2})$

D. none of these

Answer: C



Watch Video Solution

66. One of the diagonals of a square is the portion of the line $\frac{x}{2} + \frac{y}{3} = 2$ intercepted between the axes. Then the extremities of the other diagonal are: (a) $(5, 5), (-1, 1)$ (b) $(0, 0), (4, 6)$ (c) $(0, 0), (-1, 1)$ (d) $(5, 5), (4, 6)$

A. $(5,5), (-1,1)$

B. $(0,0), (4,6)$

C. $(0,0), (-1,1)$

D. $(5,5), (4,6)$

Answer: A

[Watch Video Solution](#)

67. The point $P(2,1)$ is shifted through a distance $3\sqrt{2}$ units measured parallel to the line $x+y=1$ in the direction of decreasing ordinates, to reach

at Q . The image of Q with respect to given line is

A. (3,-4)

B. (-3,2)

C. (0,-1)

D. none of these

Answer: A



[Watch Video Solution](#)

68. Let O be the origin. If $A(1, 0)$ and $B(0, 1)$ and $P(x, y)$ are points such that $xy > 0$ and $x + y < 1$, then P lies either inside the triangle OAB or in the third quadrant. P cannot lie inside the triangle OAB P lies inside the triangle OAB P lies in the first quadrant only

A. P lies either inside the triangle OAB or in the third quadrant

B. P cannot lie inside the triangle OAB

C. P lies inside the triangle OAB

D. P lies in the first quadrant only

Answer: A



[Watch Video Solution](#)

69. In a triangle ABC , the bisectors of angles B and C lie along the lines $x = y$ and $y = 0$. If A is $(1, 2)$, then the equation of line BC is $2x + y = 1$ (b) $3x - y = 5$ (c) $x - 2y = 3$ (d) $x + 3y = 1$

A. $2x+y=1$

B. $3x-y=5$

C. $x-2y=3$

D. $x+3y=1$

Answer: B



[Watch Video Solution](#)

70. Line $ax + by + p = 0$ makes angle $\frac{\pi}{4}$ with $x \cos \alpha + y \sin \alpha = p, p \in R^+$. If these lines and the line $x \sin \alpha - y \cos \alpha = 0$ are concurrent, then (a) $a^2 + b^2 = 1$ (b) $a^2 + b^2 = 2$ (c) $2(a^2 + b^2) = 1$ (d) none of these

A. $a^2 + b^2 = 1$

B. $a^2 + b^2 = 2$

C. $2(a^2 + b^2) = 1$

D. none of these

Answer: B



[Watch Video Solution](#)

71. The equation of the line AB is $y = x$. If A and B lie on the same side of the line mirror $2x - y = 1$, then the equation of the image of AB is (a) $x + y - 2 = 0$ (b) $8x + y - 9 = 0$ (c) $7x - y - 6 = 0$ (d) None of these

A. $x+y=2$

B. $8x+y=9$

C. $7x-y=6$

D. none of these

Answer: C



Watch Video Solution

72. The equation of the bisector of the acute angle between the lines

$2x - y + 4 = 0$ and $x - 2y = 1$ is $x - y + 5 = 0$ $x - y + 1 = 0$

$x - y = 5$ (d) none of these

A. $x+y+5=0$

B. $x-y+1=0$

C. $x-y=5$

D. none of these

Answer: B



Watch Video Solution

73. The straight lines $4ax + 3by + c = 0$, where $a + b + c = 1$ (a) $(4, 3)$ (b) $\left(\frac{1}{4}, \frac{1}{3}\right)$ (c) $\left(\frac{1}{2}, \frac{1}{3}\right)$ (d) none of these

A. (4,3)

B. (1/4,1/3)

C. (1/2,1/3)

D. none of these

Answer: B



Watch Video Solution

74. If the lines $ax + y + 1 = 0$, $x + by + 1 = 0$ and $x + y + c = 0$ (a, b, c being distinct and different from 1) are concurrent, then prove that

$$\frac{1}{1-a} + \frac{1}{1-b} + \frac{1}{1-c} = 1.$$

A. 0

B. 1

C. $1/(a+b+c)$

D. none of these

Answer: B



Watch Video Solution

75. If lines $x + 2y - 1 = 0$, $ax + y + 3 = 0$, and $bx - y + 2 = 0$ are concurrent, and S is the curve denoting the locus of (a, b) , then the least distance of S from the origin is $\frac{5}{\sqrt{57}}$ (b) $5/\sqrt{51}$ $5/\sqrt{58}$ (d) $5/\sqrt{59}$

A. $5/\sqrt{57}$

B. $5/\sqrt{51}$

C. $5/\sqrt{58}$

D. $5/\sqrt{59}$

Answer: C



Watch Video Solution

76. The straight lines $x + 2y - 9 = 0$, $3x + 5y - 5 = 0$, and $ax + by - 1 = 0$ are concurrent, if the straight line $35x - 22y + 1 = 0$ passes through the point (a)(a, b) (b) (b, a) (c) ($-a, -b$) (d) none of these

A. (a, b)

B. (b, a)

C. ($-a, -b$)

D. none of these

Answer: A



Watch Video Solution

77. If the straight lines $2x + 3y - 1 = 0$, $x + 2y - 1 = 0$, and $ax + by - 1 = 0$ form a triangle with the origin as orthocentre, then (a, b) is given by

A. (6,4)

B. (-3,3)

C. (-8,8)

D. (0,7)

Answer: C



[Watch Video Solution](#)

78. If $\frac{a}{\sqrt{bc}} - 2 = \sqrt{\frac{b}{c}} + \sqrt{\frac{c}{b}}$, where $a, b, c > 0$, then the family of lines $\sqrt{ax} + \sqrt{by} + \sqrt{c} = 0$ passes through the fixed point given by (a) (1, 1) (b) (1, -2) (c) (-1, 2) (d) (-1, 1)

A. (1,1)

B. (1,-2)

C. (-1,2)

D. (-1,1)

Answer: D



Watch Video Solution

79. Distance possible to draw a line which belongs to all the given family of _____ lines

$$y - 2x + 1 + \lambda_1(2y - x - 1) = 0, 3y - x - 6 + \lambda_2(y - 3x + 6) = 0, ax +$$

, then $a = 4$ (b) $a = 3$ $a = -2$ (d) $a = 2$

A. $a=4$

B. $a=3$

C. $a=-2$

D. $a=2$

Answer: A



Watch Video Solution

80. If two members of family $(2 + \lambda)x + (1 + 2\lambda)y - 3(1 + \lambda) = 0$ and line $x+y=0$ make an equilateral triangle, the the incentre of triangle so formed is

A. $\left(\frac{1}{3}, \frac{1}{3}\right)$

B. $\left(\frac{7}{6}, -\frac{5}{6}\right)$

C. $\left(\frac{5}{6}, \frac{5}{6}\right)$

D. $\left(-\frac{3}{2}, -\frac{3}{2}\right)$

Answer: A



Watch Video Solution

81. The set of lines $x \tan^{-1} a + y \sin^{-1} \left(\frac{1}{\sqrt{1+a^2}} \right) + 2 = 0$ where $a \in (0, 1)$ are concurrent at

 [Watch Video Solution](#)

82. If $\sin(\alpha + \beta)\sin(\alpha - \beta) = \sin \gamma(2 \sin \beta + \sin \gamma)$, where $\alpha, \beta, \gamma \in (0, \pi)$

A. (1,1)

B. (-1,1)

C. (1,-1)

D. none of these

Answer: C

 [Watch Video Solution](#)

Exercise Multiple

1. If P is a point (x, y) on the line $y = -3x$ such that P and the point $(3, 4)$ are on the opposite sides of the line $3x - 4y = 8$, then $x > \frac{8}{15}$
(b) $x > \frac{8}{5}$ $y < -\frac{8}{5}$ (d) $y < -\frac{8}{15}$

A. $x > 8/15$

B. $x > 8/5$

C. $x < -8/5$

D. $y < -8/15$

Answer: A::C



Watch Video Solution

2. If (x, y) is a variable point on the line $y = 2x$ lying between the lines $2(x + 1) + y = 0$ and $x + 3(y - 1) = 0$, then $x \in \left(-\frac{1}{2}, \frac{6}{7}\right)$ (b)
 $x \in \left(-\frac{1}{2}, \frac{3}{7}\right)$ $y \in \left(-1, \frac{3}{7}\right)$ (d) $y \in \left(-1, \frac{6}{7}\right)$

A. $x \in (-1/2, 6/7)$

B. $x \in (-1/2, 3/7)$

C. $y \in (-1, 3/7)$

D. $y \in (-1, 6/7)$

Answer: B::D



Watch Video Solution

3. Let $P(\sin \theta, \cos \theta)$ ($0 \leq \theta \leq 2\pi$) be a point and let OAB be a triangle with vertices $(0, 0)$, $(\sqrt{\frac{3}{2}}, 0)$ and $(0, \sqrt{\frac{3}{2}})$ Find θ if P lies inside $\triangle OAB$

A. $0 < \theta < \pi/12$

B. $5\pi/2 < \theta < \pi/2$

C. $0 < \theta < 5\pi/2$

D. $5\pi/2 < \theta < \pi$

Answer: A::B



Watch Video Solution

4. The lines $x + 2y + 3 = 0$, $x + 2y - 7 = 0$, and $2x - y - 4 = 0$ are the sides of a square. The equation of the remaining side of the square can be $2x - y + 6 = 0$ (b) $2x - y + 8 = 0$ $2x - y - 10 = 0$ (b) $2x - y - 14 = 0$

A. $2x - y + 6 = 0$

B. $2x - y + 8 = 0$

C. $2x - y - 10 = 0$

D. $2x - y - 14 = 0$

Answer: A::D



Watch Video Solution

5. Angle made with the x-axis by a straight line drawn through (1, 2) so that it intersects $x + y = 4$ at a distance $\frac{\sqrt{6}}{3}$ from (1, 2) is 105° (b) 75°

(c) 60° (d) 15°

A. 105°

B. 75°

C. 60°

D. 15°

Answer: B::D



[Watch Video Solution](#)

6. Given three straight lines $2x + 11y - 5 = 0$, $24x + 7y - 20 = 0$, and $4x - 3y - 2 = 0$. Then, they form a triangle one line bisects the angle between the other two two of them are parallel

A. they form a triangle

B. they are concurrent

C. one line bisects the angle between the other two

D. two of them are parallel

Answer: C

 [Watch Video Solution](#)

7. A triangle is formed by the lines whose equations are AB: $x+y-5=0$, BC: $x+7y-7=0$ and CA: $7x+y+14=0$.

Then

A. angle at A is acute

B. angle at C is acute

C. internal angle bisector at angle B is $3x+6y-16=0$

D. external angle bisector at angle C is $8x+8y+7=0$

Answer: A::C::D

 [Watch Video Solution](#)

8. If the points $\left(\frac{a^3}{(a-1)}\right)$, $\left(\frac{(a^2-3)}{(a-1)}\right)$, $\left(\frac{b^3}{(b-1)}\right)$, $\left(\frac{(b^2-3)}{(b-1)}\right)$, and $\left(\frac{(c^2-3)}{(c-1)}\right)$, where a, b, c are different from 1, lie on the

$lx + my + n = 0$, then $a + b + c = -\frac{m}{l}$ $ab + bc + ca = \frac{n}{l}$

$$abc = \frac{(m+n)}{l} abc - (bc + ca + ab) + 3(a + b + c) = 0$$

A. $a + b + c = -\frac{m}{l}$

B. $ab + bc + ca = \frac{n}{l}$

C. $abc = \frac{(m+n)}{l}$

D. $abc - (bc + ca + ab) + 3(a + b + c) = 0$

Answer: A::B::D

 [Watch Video Solution](#)

9. Two sides of a rhombus OABC (lying entirely in first quadrant or fourth quadrant) of area equal to 2 sq. units, are $y = \frac{x}{\sqrt{3}}$, $y = \sqrt{3}x$. Then possible coordinates of B is / are ('O' being the origin)

A. $(1 + \sqrt{3}, 1 + \sqrt{3})$

B. $(-1 - \sqrt{3}, -1 - \sqrt{3})$

C. $(3 + \sqrt{3}, 3 + \sqrt{3})$

D. $(\sqrt{3} - 1, \sqrt{3} - 1)$

Answer: A:B



Watch Video Solution

10. If the ellipse $\frac{x^2}{4} + y^2 = 1$ meets the ellipse $x^2 + \frac{y^2}{a^2} = 1$ at four distinct points and $a = b^2 - 5b + 7$, then b does not lie in (a) $[4, 5]$ (b) $(-\infty, 2) \cup (3, \infty)$ (c) $(-\infty, 0)$ (d) $[2, 3]$

A. (1,1)

B. (1,-1)

C. (2,-2)

D. (3,3)

Answer: A::B::C::D



Watch Video Solution

$$11. \frac{d}{dx} \left[\tan^{-1} \left(\frac{\sqrt{x}(3-x)}{1-3x} \right) \right] = \frac{1}{2(1+x)\sqrt{x}} \quad (b) \quad \frac{3}{(1+x)\sqrt{x}}$$

$\frac{2}{(1+x)\sqrt{x}}$ (d) $\frac{3}{2(1+x)\sqrt{x}}$

A. $\left(2 \left(1 - \frac{3\sqrt{3}}{4} \right), \frac{3}{2} \left(1 - \frac{4}{\sqrt{3}} \right) \right)$

B. $\left(-2(1 + \sqrt{3}), \frac{3}{2}(1 - \sqrt{3}) \right)$

C. $\left(2(1 + \sqrt{3}), \frac{3}{2}(1 + \sqrt{3}) \right)$

D. $\left(2 \left(1 + \frac{3\sqrt{3}}{4} \right), \frac{3}{2} \left(1 + \frac{4}{\sqrt{3}} \right) \right)$

Answer: A::D



Watch Video Solution

12. The equation of the lines passing through the point $(1, 0)$ and at a distance $\frac{\sqrt{3}}{2}$ from the origin is (a) $\sqrt{3}x + y - \sqrt{3} = 0$ (b) $x + \sqrt{3}y - \sqrt{3} = 0$ (c) $\sqrt{3}x - y - \sqrt{3} = 0$ (d) $x - \sqrt{3}y - \sqrt{3} = 0$

A. $\sqrt{3}x + y - \sqrt{3} = 0$

B. $x + \sqrt{3}y - \sqrt{3} = 0$

C. $\sqrt{3}x - y - \sqrt{3} = 0$

D. $x - \sqrt{3}y - \sqrt{3} = 0$

Answer: A:C



Watch Video Solution

13. The sides of a triangle are the straight lines $x + y = 1$, $7y = x$, and $\sqrt{3}y + x = 0$. Then which of the following is an interior point of the triangle? (a) Circumcenter (b) Centroid (c) Incenter (d) Orthocenter

A. Circumcenter

B. Centroid

C. Incenter

D. Orthocenter

Answer: B::C



Watch Video Solution

14. If the straight line $ax + cy = 2b$, where $a, b, c > 0$, makes a triangle of area 2 sq. units with the coordinate axes, then a, b, c are in GP a, -b; c are in GP a, 2b, c are in GP (d) a, - 2b, c are in GP

A. a,b,c are in GP

B. a,-b, c are in GP

C. a,2b,c are in GP

D. a,-2b, c are in GP

Answer: A::B



[Watch Video Solution](#)

15. Consider the equation $y - y_1 = m(x - x_1)$. If m and x_1 are fixed and different lines are drawn for different values of y_1 , then (a) the lines will pass through a fixed point (b) there will be a set of parallel lines (c) all the lines intersect the line $x = x_1$ (d) all the lines will be parallel to the line $y = x_1$

- A. the lines will pass through a fixed point
- B. there will be a set of parallel lines
- C. all the lines intersect the line $x = x_1$
- D. all the lines will be parallel to the line $y = x_1$

Answer: B::C



[Watch Video Solution](#)

16. Equation(s) of the straight line(s), inclined at 30° to the x-axis such that the length of its (each of their) line segment(s) between the coordinate axes is 10 units, is (are) (a) $x + \sqrt{3}y + 5\sqrt{3} = 0$ (b) $x - \sqrt{3}y + 5\sqrt{3} = 0$ (c) $x + \sqrt{3}y - 5\sqrt{3} = 0$ (d) $x - \sqrt{3}y - 5\sqrt{3} = 0$

A. $x + \sqrt{3}y + 5\sqrt{3} = 0$

B. $x - \sqrt{3}y + 5\sqrt{3} = 0$

C. $x + \sqrt{3}y - 5\sqrt{3} = 0$

D. $x - \sqrt{3}y - 5\sqrt{3} = 0$

Answer: B::D



Watch Video Solution

17. The lines $x + y - 1 = 0$, $(m - 1)x + (m^2 - 7)y - 5 = 0$, and $(m - 2)x + (2m - 5)y = 0$ are concurrent for three values of m concurrent for no value of m parallel for one value of m parallel for two value of m

A. concurrent for three values of m

B. concurrent for one value of m

C. concurrent for no value of m

D. parallel for $m=3$

Answer: C::D



Watch Video Solution

18. The equation of a straight line passing through the point $(2, 3)$ and inclined at an angle of $\tan^{-1}\left(\frac{1}{2}\right)$ with the line $y + 2x = 5$ is (a) $y = 3$ (b) $x = 2$ (c) $3x + 4y - 18 = 0$ (d) $4x + 3y - 17 = 0$

A. $y=3$

B. $x=2$

C. $3x+4y-18=0$

D. $4x+3y-17=0$

Answer: B::C



Watch Video Solution

19. The equation of the line on which the perpendicular from the origin makes an angle of 30° with x - axis and which forms a triangle of area $\frac{50}{\sqrt{3}}$ with the axes is

A. $\sqrt{3}x + y - 10 = 0$

B. $\sqrt{3}x + y + 10 = 0$

C. $x + \sqrt{3}y - 10 = 0$

D. $x - \sqrt{3}y - 10 = 0$

Answer: A::B



Watch Video Solution

20. A line is drawn perpendicular to line $y = 5x$, meeting the coordinate axes at A and B . If the area of triangle OAB is 10 sq. units, where O is the origin, then the equation of drawn line is (a) $3x - y - 9 = 0$ (b) $5y + x = 10$ (c) $5y + x = -10$ (d) $x - 4y = 10$

A. 12

B. -12

C. 10

D. -10

Answer: A::B



Watch Video Solution

21. If $x - 2y + 4 = 0$ and $2x + y - 5 = 0$ are the sides of an isosceles triangle having area 10 sq units, the equation of the third side is (a) $3x - y = -9$ (b) $3x - y + 11 = 0$ (c) $x - 3y = 19$ (d) $3x - y + 15 = 0$

A. $x+3y=-1$

B. $x+3y=19$

C. $3x-y=-9$

D. $3x-y=11$

Answer: A::B::C::D



Watch Video Solution

22. The number of values of a for which the lines $2x + y - 1 = 0$,
 $ax + 3y - 3 = 0$, and $3x + 2y - 2 = 0$ are concurrent is 0 (b) 1 (c) 2 (d)
infinite

A. -3

B. -1

C. 1

D. 4

Answer: A::B::C::D



Watch Video Solution

23. Three lines $px + qy + r = 0$, $qx + ry + p = 0$ and $rx + py + q = 0$ are concurrent if

A. $p + q + r = 0$

B. $p^2 + q^2 + r^2 = pr + rp + pq$

C. $p^3 + q^3 + r^3 = 3pqr$

D. none of these

Answer: A::B::C



Watch Video Solution

24. The equation of bisector of two lines L_1 and L_2 are $2x - 16y - 5 = 0$ and $64x + 8y + 35 = 0$. If the line L_1 passes through $(-11, 4)$, then identify the

equation of acute angle bisector of L_1 and L_2 .

$$\text{A. } \frac{x - x_1}{\cos\left(\frac{\theta_1 + \theta_2}{2}\right)} = \frac{y - y_1}{\sin\left(\frac{\theta_1 + \theta_2}{2}\right)}$$

$$\text{B. } \frac{x - x_1}{-\sin\left(\frac{\theta_1 - \theta_2}{2}\right)} = \frac{y - y_1}{\cos\left(\frac{\theta_1 - \theta_2}{2}\right)}$$

$$\text{C. } \frac{x - x_1}{\sin\left(\frac{\theta_1 + \theta_2}{2}\right)} = \frac{y - y_1}{\cos\left(\frac{\theta_1 + \theta_2}{2}\right)}$$

$$\text{D. } \frac{x - x_1}{-\sin\left(\frac{\theta_1 + \theta_2}{2}\right)} = \frac{y - y_1}{\cos\left(\frac{\theta_1 + \theta_2}{2}\right)}$$

Answer: A:D



Watch Video Solution

25. Consider the lines $L_1 \equiv 3x - 4y + 2 = 0$ and $L_2 \equiv 3y - 4x - 5 = 0$. Now, choose the correct statement(s).

A. The line $x+y=0$ bisects the acute angle between L_1 and L_2 containing the origin.

B. The line $x-y+1=0$ bisects the obtuse angle between L_1 and L_2 not containing the origin.

C. The line $x+y+3=0$ bisects the obtuse angle between L_1 and L_2 containing the origin.

D. The line $x-y+1=0$ bisects the acute angle between L_1 and L_2 not containing the origin.

Answer: A::B



Watch Video Solution

26. The sides of a rhombus are parallel to the lines $x + y - 1 = 0$ and $7x - y - 5 = 0$. It is given that the diagonals of the rhombus intersect at $(1, 3)$ and one vertex, A of the rhombus lies on the line $y = 2x$. Then the coordinates of vertex A are $\left(\frac{8}{5}, \frac{16}{5}\right)$ (b) $\left(\frac{7}{15}, \frac{14}{15}\right)$ $\left(\frac{6}{5}, \frac{12}{5}\right)$ (d) $\left(\frac{4}{15}, \frac{8}{15}\right)$

A. $(8/5, 16/5)$

B. $(7/15, 14/15)$

C. $(6/5, 12/5)$

D. $(4/15, 8/15)$

Answer: A::C



Watch Video Solution

27. The system of equations $x + 2y + 3z = 1$, $x - y + 4z = 0$, $2x + y + 7z = 1$ has

A. $y + 3x = 0$ and $3y + 2x = 0$

B. $2y + 3x = 0$ and $3y + x = 0$

C. $2y = 3x$ and $3y = 0$

D. $y = 3x$ and $3y = 2x$

Answer: A::B::C::D



Watch Video Solution

28. Let $u \equiv ax + by + abz = 0$, $v \equiv bx - ay + baz = 0$, $a, b \in R$, be two straight lines. The equations of the bisectors of the angle formed by $k_1u - k_2v = 0$ and $k_1u + k_2v = 0$, for nonzero and real k_1 and k_2 are

(a) $u = 0$ (b) $k_2u + k_1v = 0$ (c) $k_2u - k_1v = 0$ (d) $v = 0$

A. $u=0$

B. $k_2u + k_1v = 0$

C. $k_2u - k_1v = 0$

D. $v=0$

Answer: A:D



Watch Video Solution

29. Two sides of a triangle are parallel to the coordinate axes. If the slopes of the medians through the acute angles of the triangle are 2 and m , the

$m = \frac{1}{2}$ (b) 2 (c) 4 (d) 8

A. $1/2$

B. 2

C. 4

D. 8

Answer: A::D

 [Watch Video Solution](#)

30. about to only mathematics

A. $PR = 3\sec\theta$

B. $PS = 4 \operatorname{cosec}\theta$

C. $PR + PS = \frac{2(3\sin\theta + 4\cos\theta)}{\sin 2\theta}$

D. $\frac{9}{(PR)^2} + \frac{16}{(PS)^2} = 1$

Answer: A::B::C::D

 [Watch Video Solution](#)

Exercise Comprehension

1. Let L be the line belonging to the family of straight lines $(a+2b)x + (a-3b)y + a - 8b = 0$, $a, b \in \mathbb{R}$, which is the farthest from the point $(2, 2)$.

Area enclosed by the line L and the coordinate axes is

A. $x+4y+7=0$

B. $2x+3y+4=0$

C. $4x-y-6=0$

D. none of these

Answer: A



[Watch Video Solution](#)

2. Let l be the line belonging to the family of straight lines $(a + 2b)x + (a - 3b)y + a - 8b = 0$, $a, b \in \mathbb{R}$, which is farthest from

the point $(2, 2)$, then area enclosed by the line L and the coordinate axes is

- A. $4/3$ sq. units
- B. $9/2$ sq. units
- C. $49/8$ sq. units
- D. none of these

Answer: C



[Watch Video Solution](#)

3. Let L be the line belonging to the family of straight lines $(a+2b)x + (a-3b)y + a - 8b = 0$, $a, b \in \mathbb{R}$, which is the farthest from the point $(2, 2)$.

If L is concurrent with the lines $x - 2y + 1 = 0$ and $3x - 4y + \lambda = 0$, then the value of λ is

- A. 2
- B. 1

C. -4

D. 5

Answer: D



Watch Video Solution

4. The perimeter of an equilateral triangle is 30 cm. The area is

A. 1

B. 2

C. 3

D. 4

Answer: B



Watch Video Solution

5. The perimeter of an equilateral triangle is 30 cm. The area is

A. 0,0

B. $0, 2\sqrt{3}$

C. $3, -\sqrt{3}$

D. none of these

Answer: D



[Watch Video Solution](#)

6. The perimeter of an equilateral triangle is 30 cm. The area is

A. $1, \sqrt{3}$

B. $0, \sqrt{3}$

C. 0, 2

D. none of these

Answer: A



Watch Video Solution

7. a variable line L is drawn through $O(0, 0)$ to meet the lines $L_1: y - x - 10 = 0$ and $L_2: y - x - 20 = 0$ at point A & B respectively

A point P is taken on line L the (1) if $\frac{2}{OP} = \frac{1}{OA} + \frac{1}{OB}$ then locus of

P is (2) if $(OP)^2 = (OA) \cdot (OB)$ then locus of P is (3) if

$\frac{1}{(OP)^2} = \frac{1}{(OA)^2} + \frac{1}{(OB)^2}$ then locus of point P is:

A. $3x+3y=40$

B. $3x+3y+40 = 0$

C. $3x-3y=40$

D. $3y-3x=40$

Answer: D



Watch Video Solution

8. A variable line L is drawn through $O(0,0)$ to meet the line L_1 and L_2 given by $y-x-10=0$ and $y-x-20=0$ at Points A and B , respectively.

Locus of P , if $OP^2 = OA \times OB$, is

A. $(y - x)^2 = 100$

B. $(y + x)^2 = 50$

C. $(y - x)^2 = 200$

D. none of these

Answer: C

 [Watch Video Solution](#)

9. a variable line L is drawn trough $O(0, 0)$ to meet the lines $L_1: y - x - 10 = 0$ and $L_2: y - x - 20 = 0$ at point $A \& B$ respectively

A point P is taken on line L the (1) if $\frac{2}{OP} = \frac{1}{OA} + \frac{1}{OB}$ then locus of

P is (2) if $(OP)^2 = (OA) \cdot (OB)$ then locus of P is (3) if

$\frac{1}{(OP)^2} = \frac{1}{(OA)^2} + \frac{1}{(OB)^2}$ then locus of point P is:

A. $(y - x)^2 = 80$

B. $(y - x)^2 = 100$

C. $(y - x)^2 = 64$

D. none of these

Answer: A



[Watch Video Solution](#)

10. The line $6x+8y=48$ intersects the coordinates axes at A and B, respectively. A line L bisects the area and the perimeter of triangle OAB, where O is the origin.

Slope of Line L is

A. 1

B. 2

C. 3

D. more than 3

Answer: A



[Watch Video Solution](#)

11. The line $6x+8y=48$ intersects the coordinates axes at A and B, respectively. A line L bisects the area and the perimeter of triangle OAB, where O is the origin.

The slope of line L can be

A. $(10 + 5\sqrt{6}) / 10$

B. $(10 - 5\sqrt{6}) / 10$

C. $(8 + 3\sqrt{6}) / 10$

D. none of these

Answer: B



[Watch Video Solution](#)

12. The line $6x+8y=48$ intersects the coordinates axes at A and B, respectively. A line L bisects the area and the perimeter of triangle OAB, where O is the origin.

Slope of Line L is

 [Watch Video Solution](#)

13. $A(1, 3)$ and $c\left(-\frac{2}{5}, -\frac{2}{5}\right)$ are the vertices of a ΔABC and the equation of the angle bisector of $\angle ABC$ is $x + y = 2$.

A. $7x+3y-4=0$

B. $7x+3y+4=0$

C. $7x-3y+4=0$

D. $7x-3y-4=0$

Answer: B

 [Watch Video Solution](#)

14. $A(1, 3)$ and $c\left(-\frac{2}{5}, -\frac{2}{5}\right)$ are the vertices of a ΔABC and the equation of the angle bisector of $\angle ABC$ is $x + y = 2$.

A. $(3/10, 17/10)$

B. $(17/10, 3/10)$

C. $(-5/2, 9/2)$

D. $(1,1)$

Answer: C



Watch Video Solution

15. $A(1, 3)$ and $c\left(-\frac{2}{5}, -\frac{2}{5}\right)$ are the vertices of a ΔABC and the equation of the angle bisector of $\angle ABC$ is $x + y = 2$.



Watch Video Solution

16. Let $ABCD$ be a parallelogram the equation of whose diagonals are $AC: x + 2y = 3$; $BD: 2x + y = 3$. If length of diagonal $AC = 4$ units and area of $ABCD = 8$ sq. units. (i) The length of the other diagonal is (ii) the length of side AB is equal to

A. $10/3$

B. 2

C. $20/3$

D. none of these

Answer: C



[Watch Video Solution](#)

17. Let $ABCD$ be a parallelogram whose equations for the diagonals AC and BD are $x+2y=3$ and $2x+y=3$, respectively.

The length of side AB is equal to

A. $2\sqrt{58}/3$

B. $4\sqrt{58}/9$

C. $3\sqrt{58}/9$

D. $4\sqrt{58}/9$

Answer: A

 [Watch Video Solution](#)

18. Let ABCD be parallelogram whose equations for the diagonals AC and BD are $x+2y=3$ and $2x+y=3$, respectively. If length of diagonal AC=4 units and area of parallelogram ABCD=8 sq. units then

(i) the length of other diagonal BD is

(a) $\frac{10}{3}$ (b) $\frac{20}{3}$ (c) 2 (d) 5

(ii) length of side AB equals to

(a) $\frac{2\sqrt{58}}{3}$ (b) $\frac{2\sqrt{58}}{9}$ (c) $\frac{3\sqrt{58}}{9}$ (d) $\frac{4\sqrt{58}}{9}$

 [Watch Video Solution](#)

19. Consider a triangle PQR with coordinates of its vertices as P(-8,5), Q(-15, -19), and R (1, -7). The bisector of the interior angle of P has the equation which can be written in the form $ax+2y+c=0$.

The distance between the orthocenter and the circumcenter of triangle PQR is

A. $25/2$

B. $29/2$

C. $37/2$

D. $51/2$

Answer: A



[Watch Video Solution](#)

20. Evaluate $\int_{-1}^1 (x - [x])dx$, where $[.]$ denotes the greatest integer function.



[Watch Video Solution](#)

21. Evaluate $\int_2^5 (x - [x])dx$, where $[.]$ denotes the greatest integer function.



Watch Video Solution

22. The base of an isosceles triangle measures 4 units base angle is equal to 45° . A straight line cuts the extension of the base at a point M at the angle θ and bisects the lateral side of the triangle which is nearest to M.

The area of quadrilateral which the straight line cuts off from the given triangle is

- A. $\frac{3 + \tan\theta}{1 + \tan\theta}$
- B. $\frac{3 + 5\tan\theta}{1 + \tan\theta}$
- C. $\frac{3 + \tan\theta}{1 - \tan\theta}$
- D. $\frac{3 + 2\tan\theta}{1 + \tan\theta}$

Answer: B



Watch Video Solution

23. The base of an isosceles triangle measures 4 units base angle is equal to 45° . A straight line cuts the extension of the base at a point M at the angle θ and bisects the lateral side of the triangle which is nearest to M. The length of portion of straight line inside the triangle may lie in the range



Watch Video Solution

24. The base of an isosceles triangle measures 4 units base angle is equal to 45° . A straight line cuts the extension of the base at a point M at the angle θ and bisects the lateral side of the triangle which is nearest to M. The length of portion of straight line inside the triangle may lie in the range

A. (2,4)

B. $\left(\frac{3}{2}, \sqrt{3}\right)$

C. $(\sqrt{2}, 2)$

D. $(\sqrt{2}, \sqrt{3})$

Answer: C



Watch Video Solution

25. Consider point $A(6, 30)$, point $B(24, 6)$ and line $AB: 4x+3y = 114$.

Point $P(0, \lambda)$ is a point on y-axis such that

$0 < \lambda < 38$ and point $Q(0, \lambda)$ is a point on y-axis such that $\lambda > 38$.

For all positions of point P, angle APB is maximum when point P is

A. $(0, 12)$

B. $(0, 15)$

C. $(0, 18)$

D. $(0, 21)$

Answer: C



Watch Video Solution

26. Consider point $A(6, 30)$, point $B(24, 6)$ and line $AB: 4x+3y = 114$.

Point $P(0, \lambda)$ is a point on y -axis such that

$0 < \lambda < 38$ and point $Q(0, \lambda)$ is a point on y -axis such that $\lambda > 38$.

The maximum value of angle APB is

A. $\frac{\pi}{3}$

B. $\frac{\pi}{2}$

C. $\frac{2\pi}{3}$

D. $\frac{3\pi}{3}$

Answer: B

[Watch Video Solution](#)

27. Consider point $A(6, 30)$, point $B(24, 6)$ and line $AB: 4x+3y = 114$.

Point $P(0, \lambda)$ is a point on y -axis such that

$0 < \lambda < 38$ and point $Q(0, \lambda)$ is a point on y-axis such that $\lambda > 38$.

For all positions of point Q, and $\angle AQB$ is maximum when point Q is

A. (0, 54)

B. (0, 58)

C. (0, 60)

D. (0, 1)

Answer: B



[Watch Video Solution](#)

Exercise Matrix

1. Consider the lines represented by equation $(x^2 + xy - x) \times (x - y) = 0$ forming a triangle. Then match the

following lists:

List I	List II
a. Orthocenter of triangle	p. $(1/6, 1/2)$
b. Circumcenter	q. $(1/(2 + 2\sqrt{2}), 1/2)$
c. Centroid	r. $(0, 1/2)$
d. Incenter	s. $(1/2, 1/2)$



Watch Video Solution

2. Consider the triangle formed by the lines

$$y+3x+2=0, 3y-2x-5=0, 4y+x-14=0$$

Match the following lists:

List I	List II
a. Values of α if $(0, \alpha)$ lies inside the triangle	p. $(-\infty, 7/3) \cup (13/4, \infty)$
b. Values of α if $(\alpha, 0)$ lies inside the triangle	q. $-4/3 < \alpha < 1/2$
c. Values of α if $(\alpha, 2)$ lies inside the triangle	r. No value of α
d. Value of α if $(1, \alpha)$ lies outside the triangle	s. $5/3 < \alpha < 7/2$



Watch Video Solution

3. A straight line with negative slope passing the point $(1, 4)$ meets the coordinate axes at A and B. The minimum value of $OA + OB =$

 [Watch Video Solution](#)

4. If the point P is symmetric to the point $Q(4,-1)$ with respect to the bisector of the first quadrant then the length of PQ is

 [Watch Video Solution](#)

5. The point $A(1, 2)$, $B(2, -3)$, $C(-1, -5)$ and $D(-2, 4)$ in order are the vertices of

 [Watch Video Solution](#)

6. Differentiate $y = \sin(x^2 + 3)$.



7. Consider the lines given by

$$L_1 : x + 3y - 5 = 0$$

$$L_2 : 3x - ky - 1 = 0$$

$$L_3 : 5x + 2y - 12 = 0$$

Match the following lists.

List I	List II
a. L_1, L_2, L_3 are concurrent if	p. $k = -9$
b. One of L_1, L_2, L_3 is parallel to at least one of the other two if	q. $k = -6/5$
c. L_1, L_2, L_3 form a triangle if	r. $k = 5/6$
d. L_1, L_2, L_3 do not form a triangle if	s. $k = 5$

8. Consider a ΔABC in which sides AB and AC are perpendicular to $x - y - 4 = 0$ and $2x - y - 5 = 0$, respectively. Vertex A is $(-2, 3)$ and the circumcenter of ΔABC is $(3/2, 5/2)$.

8. Consider a ΔABC in which sides AB and AC are perpendicular to $x - y - 4 = 0$ and $2x - y - 5 = 0$, respectively. Vertex A is $(-2, 3)$ and the circumcenter of ΔABC is $(3/2, 5/2)$.

The equation of the line in List I is of the form $ax + by + c = 0$, where

$a, b, c \in I$. Match it with the corresponding value of c in list II and then choose the correct code.

List I	List II
a. Equation of the perpendicular bisector of side AB	p. -1
b. Equation of the perpendicular bisector of side AC .	q. 1
c. Equation of side AC	r. -16
d. Equation of the median through A	s. -4

Codes :

$a \ b \ c \ d$

$r \ s \ p \ q$

$s \ r \ q \ p$

$q \ p \ s \ r$

$r \ p \ s \ q$



Watch Video Solution

Exercise Numerical

1. straight line L with negative slope passes through the point $(9,4)$ cuts the positive coordinate axes at the point P and W As L . Varies, find the minimum value of $|OP|+|OQ|$, where O is origin .

 [Watch Video Solution](#)

2. The number of values of k for which the lines $(k + 1)x + 8y = 4k$ and $kx + (k + 3)y = 3k - 1$ are coincident is _____

 [Watch Video Solution](#)

3. The sides of a triangle ABC lie on the lines $3x + 4y = 0$, $4x + 3y = 0$ and $x = 3$. Let (h, k) be the centre of the circle inscribed in $\triangle ABC$. The value of $(h + k)$ equals

 [Watch Video Solution](#)

4. The absolute value of the sum of the abscissas of all the points on the line $x + y = 4$ that lie at a unit distance from the line $4x + 3y - 10 = 0$ is _____

 [Watch Video Solution](#)

5. Two sides of a rectangle are $3x+4y+5=0$, $4x-3y+15=0$ and one of its vertices is $(0, 0)$. The area of rectangle is ___.

 [Watch Video Solution](#)

6. The line $x = c$ cuts the triangle with corners $(0, 0)$, $(1, 1)$ and $(9, 1)$ into two regions. Two regions to be the same c must be equal to (A) $\frac{5}{2}$ (B) 3 (C) $\frac{7}{2}$ (D) 5 or 15

 [Watch Video Solution](#)

7. For all real values of a and b , lines $(2a + b)x + (a + 3b)y + (b - 3a) = 0$ and $mx + 2y + 6 = 0$ are concurrent. Then m is equal to _____

 [Watch Video Solution](#)

8. The line $3x + 2y = 24$ meets the y-axis at A and the x-axis at B . The perpendicular bisector of AB meets the line through $(0, -1)$ parallel to the x-axis at C . If the area of triangle ABC is A , then the value of $\frac{A}{13}$ is _____

 [Watch Video Solution](#)

9. Consider a $\triangle ABC$ whose sides AB , BC and CA are represented by the straight lines $2x + y = 0$, $x + py = q$ and $x - y = 3$ respectively. The point P is $(2, 3)$. If P is orthocentre, then find the value of $(p+q)$ is

 [Watch Video Solution](#)

10. Triangle ABC with $AB = 13$, $BC = 5$, and $AC = 12$ slides on the coordinates axes with A and B on the positive x-axis and positive y-axis respectively. The locus of vertex C is a line $12x - ky = 0$. Then the value of k is _____

 [Watch Video Solution](#)

11. The line $y = \frac{3x}{4}$ meets the lines $x - y = 0$ and $2x - y = 0$ at points A and B , respectively. If P on the line $y = \frac{3x}{4}$ satisfies the condition $PA \cdot PB = 25$, then the number of possible coordinates of P is ____

 [Watch Video Solution](#)

12. In a plane there are two families of lines $y = x + r$, $y = -x + r$, where $r \in \{0, 1, 2, 3, 4\}$. The number of squares of diagonals of length 2 formed by the lines is:

 [Watch Video Solution](#)

13. If $5a + 5b + 20c = t$, then find the value of t for which the line $ax + by + c - 1 = 0$ always passes through a fixed point.

 [Watch Video Solution](#)

1. The line L given by $\frac{x}{5} + \frac{y}{b} = 1$ passes through the point (13,32).the line K is parallel to L and has the equation $\frac{x}{c} + \frac{y}{3} = 1$ then the distance between L and K is

A. $\frac{23}{\sqrt{17}}$

B. $\frac{23}{\sqrt{15}}$

C. $\sqrt{17}$

D. $\frac{17}{\sqrt{15}}$

Answer: A



Watch Video Solution

2. The line $L_1: y - x = 0$ and $L_2: 2x + y = 0$ intersect the line $L_3: y + 2 = 0$ at P and Q respectively. The bisector of the acute angle between L_1 and L_2 intersects L_3 at R. Statement-1 : The ratio $PR:RQ$

equals $2\sqrt{2} : \sqrt{5}$ Statement-2 : In any triangle, bisector of an angle divides the triangle into two similar triangles. Statement-1 is true, Statement-2 is true ; Statement-2 is correct explanation for Statement-1 Statement-1 is true, Statement-2 is true ; Statement-2 is not a correct explanation for Statement-1 Statement-1 is true, Statement-2 is false Statement-1 is false, Statement-2 is true

- A. Statement 1 is true, statement 2 is false.
- B. Statement 1 is true, statement 2 is true, statement 2 is the correct explanation of statement1.
- C. Statement 1 is true, statement 2 is true, statement 2 is not the correct explanation of statement 1.
- D. Statement 1 is false, statement 2 is true.

Answer: A



[Watch Video Solution](#)

3. A line is drawn through the point $(1, 2)$ to meet the coordinate axes at P and Q such that it forms a triangle OPQ, where O is the origin. If the area of the triangle OPQ is least, then the slope of the line PQ is

A. $-\frac{1}{4}$

B. -4

C. -2

D. $-\frac{1}{2}$

Answer: C



[Watch Video Solution](#)

4. The x-coordinate of the incentre of the triangle where the midpoints of the sides are $(0, 1)$, $(1, 1)$ and $(1, 0)$ is

A. $2 + \sqrt{2}$

B. $2 - \sqrt{2}$

C. $1 + \sqrt{2}$

D. $1 - \sqrt{2}$

Answer: B



Watch Video Solution

5. A ray of light along $x + \sqrt{3}y = \sqrt{3}$ gets reflected upon reaching x-axis, the equation of the reflected ray is

A. $y = x + \sqrt{3}$

B. $\sqrt{3}y = x - \sqrt{3}$

C. $y = \sqrt{3}x - \sqrt{3}$

D. $\sqrt{3}y = x - 1$

Answer: B



Watch Video Solution

6. Let a, b, c and d be non-zero numbers. If the point of intersection of the lines $4ax + 2ay + c = 0$ and $5bx + 2by + d = 0$ lies in the fourth quadrant and is equidistant from the two axes, then

A. $2bc - 3ad = 0$

B. $2bc + 3ad = 0$

C. $3bc - 2ad = 0$

D. $3bc + 2ad = 0$

Answer: C



[Watch Video Solution](#)

7. Let PS be the median of the triangle with vertices $P(2, 2)$, $Q(6, -1)$ and $R(7, 3)$. The equation of the line passing through $(1, -1)$ and parallel to PS is (1) $4x - 7y - 11 = 0$ (2) $2x + 9y + 7 = 0$ (3) $4x + 7y + 3 = 0$ (4) $2x - 9y - 11 = 0$

A. $4x - 7y - 11 = 0$

B. $2x+9y+7=0$

C. $4x+7y+3=0$

D. $2x-9y-11=0$

Answer: B



Watch Video Solution

8. Locus of the image of the point $(2, 3)$ in the line $(2x - 3y + 4) + k(x - 2y + 3) = 0, k \in \mathbb{R}$, is a : (1) straight line parallel to x-axis. (2) straight line parallel to y-axis (3) circle of radius $\sqrt{2}$ (4) circle of radius $\sqrt{3}$

A. Straight line parallel to x-axis

B. straight line parallel to y-axis

C. circle of radius $\sqrt{2}$

D. circle of radius 3

Answer: C



Watch Video Solution

9. Two sides of a rhombus are along the lines, $x - y + 1 = 0$ and $7x - y - 5 = 0$. If its diagonals intersect at $(-1, -2)$, then which one of the following is a vertex of this rhombus ? (1) $(-3, -9)$ (2) $(-3, -8)$ (3) $\left(\frac{1}{3}, -\frac{8}{3}\right)$ (4) $\left(-\frac{10}{3}, -\frac{7}{3}\right)$

A. $(-3, -8)$

B. $\left(\frac{1}{3}, -\frac{8}{3}\right)$

C. $\left(-\frac{10}{3}, -\frac{7}{3}\right)$

D. $(-3, -9)$

Answer: B



Watch Video Solution

1. about to only mathematics

A. a hyperbola

B. a parabola

C. an ellipse

D. a straight line

Answer: D



Watch Video Solution

2. about to only mathematics

A. $y + \sqrt{3}x + 2 - 3\sqrt{3} = 0$

B. $y - \sqrt{3}x + 2 + 3\sqrt{3} = 0$

C. $\sqrt{3}y - x + 3 + 2\sqrt{3} = 0$

D. $\sqrt{3}y + x - 3 + 2\sqrt{3} = 0$

Answer: B



Watch Video Solution

3. about to only mathematics

A. $a + b - c > 0$

B. $a - b + c < 0$

C. $a - b + c > 0$

D. $a + b - c < 0$

Answer: A



Watch Video Solution

4. For a point P in the plane, let $d_1(P)$ and $d_2(P)$ be the distances of the point P from the lines $x - y = 0$ and $x + y = 0$ respectively. The area of the region R consisting of all points P lying in the first quadrant of the plane and satisfying $2 \leq d_1(P) + d_2(P) \leq 4$, is



Watch Video Solution