



MATHS

BOOKS - KC SINHA ENGLISH

MATRICES - FOR BOARDS

Solved Examples

1. Find the number of matrices having 12 elements

[Watch Video Solution](#)

2. Construct a matrix $[a_{ij}]_{3 \times 3}$, where $a_{ij} = 2i - 3j$.

[Watch Video Solution](#)

3. Consider the following information regarding the number of men and women workers in three factories. I, II and III.

	Men workers	Women workers
I	30	25
II	25	31
III	27	26

Represent the above information in the form of 3×2 matrix. What does the entry in the third row and second column represent?



Watch Video Solution

4. Find the values of x, y, z and a which satisfy the matrix equation

$$\begin{bmatrix} x + 32y + xz - 14a - 6 \end{bmatrix} = \begin{bmatrix} 0 - 732a \end{bmatrix}$$


Watch Video Solution

5. If $\begin{bmatrix} x^2 - 4x & x^2 \\ x^2 & x^3 \end{bmatrix} = \begin{bmatrix} -3 & 1 \\ -x + 2 & 1 \end{bmatrix}$ (then find x)



Watch Video Solution

6.

If

$$[x + 3z + 42y - 7 - 6a - 10b - 3 - 210] = [063y - 2 - 6 - 32c + 22b + 4]$$

Find the values of a, b, c, x, y and z.



Watch Video Solution

7. If $A = \begin{bmatrix} a & b \\ -b & a \end{bmatrix}$, $B = \begin{bmatrix} -a & b \\ -b & -a \end{bmatrix}$ then find A+B



Watch Video Solution

8. Find matrix X if $X + \begin{bmatrix} 2 & 5 \\ 3 & -1 \end{bmatrix} = \begin{bmatrix} 3 & 4 \\ 2 & 0 \end{bmatrix}$



Watch Video Solution

9. Find X and Y if $X + Y = \begin{bmatrix} 7 & 0 \\ 2 & 5 \end{bmatrix}$ and $X - Y = \begin{bmatrix} 3 & 0 \\ 2 & 3 \end{bmatrix}$



Watch Video Solution

10. Find a matrix X such that $3A - 2B + X = 0$, where $A = \begin{bmatrix} 4 & 2 \\ 1 & 3 \end{bmatrix}$, $B = \begin{bmatrix} -2 & 1 \\ 3 & 2 \end{bmatrix}$



Watch Video Solution

11. If $A = \begin{bmatrix} -1 & 2 & 3 \\ -1 & 0 & 2 \\ 1 & -3 & 1 \end{bmatrix}$ and $B = \begin{bmatrix} 4 & 5 & 6 \\ -1 & 0 & 1 \\ 2 & 1 & 2 \end{bmatrix}$, find $4A - 3B$



Watch Video Solution

12. If $A = \begin{bmatrix} 3 & -1 & 4 \\ 2 & 3 & 1 \end{bmatrix}$, $B = \begin{bmatrix} 1 & 3 & 4 \\ 2 & 1 & 0 \\ -3 & 2 & 3 \end{bmatrix}$ find AB



Watch Video Solution

13. If $A = \begin{bmatrix} ab & b^2 \\ -a^2 & -ab \end{bmatrix}$, show that $A^2 = O$.



Watch Video Solution

14.

If

$$A = [1 \ -1 \ 2 \ 3], B = [2 \ 1 \ 1 \ 0], \text{ prove that } (A + B)^2 \neq A^2 + 2AB + B^2$$



Watch Video Solution

15. solve for x and y $\begin{bmatrix} 3 & -4 \\ 1 & 2 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 3 \\ 11 \end{bmatrix}$



Watch Video Solution

16.

Write

as

a

single

matrix:

$$\begin{pmatrix} -1 & 2 & 3 \end{pmatrix} \begin{pmatrix} -2 & -1 & 5 \\ 0 & -1 & 4 \\ 7 & 0 & 5 \end{pmatrix} - 2 \begin{pmatrix} 4 & -5 & -7 \end{pmatrix}$$



Watch Video Solution

17. Product of more than two Matrices :



Watch Video Solution

18. If $A = \begin{bmatrix} 1 & 2 & 3 \\ -1 & 0 & 2 \\ 1 & -3 & -1 \end{bmatrix}$, $B = \begin{bmatrix} 4 & 5 & 6 \\ -1 & 0 & 1 \\ 2 & 1 & 2 \end{bmatrix}$ and $C = \begin{bmatrix} -1 & -2 & 1 \\ -1 & 2 & 3 \\ -1 & -2 & 2 \end{bmatrix}$ verify that $A(B + C) = AB + AC$.



Watch Video Solution

19. about to only mathematics



Watch Video Solution

20. Find a 2×2 matrix B such that $B \begin{bmatrix} 1 & -2 & 14 \end{bmatrix} = \begin{bmatrix} 6 & 0 & 6 \end{bmatrix}$



Watch Video Solution

21. Find A , if $\begin{bmatrix} 4 \\ 1 \\ 3 \end{bmatrix} A = \begin{bmatrix} -4 & 8 & 4 \\ -1 & 2 & 1 \\ -3 & 6 & 3 \end{bmatrix}$



Watch Video Solution

22. Show that $A = \begin{bmatrix} -8 & 5 \\ 2 & 4 \end{bmatrix}$ satisfies the equation $A^2 + 4A - 42 = 0$



Watch Video Solution

23. If $A = \begin{bmatrix} -4 & 1 \\ 3 & 2 \end{bmatrix}$ find $f(A)$ if $f(x) = x^2 - 2x + 3$



Watch Video Solution

24. If $A = [122212221]$, then prove that $A^2 - 4A - 5I = O$.



Watch Video Solution

25. If $A^2 = 8A + kI$ where $A = \begin{bmatrix} 1 & 0 \\ -1 & 7 \end{bmatrix}$, then k is



Watch Video Solution

26. If $A = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$ then show that $A^n = \begin{bmatrix} \cos n\theta & -\sin n\theta \\ \sin n\theta & \cos n\theta \end{bmatrix}$



Watch Video Solution

27. Let $A = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix}$ show that $(aI + bA)^n = a^n I + na^{n-1}bA$, where I is the identity matrix of order 2 and $n \in \mathbb{N}$.



Watch Video Solution

28. If $A = \begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{bmatrix}$ then show that $A^n = \begin{bmatrix} 3^{n-1} & 3^{n-1} & 3^{n-1} \\ 3^{n-1} & 3^{n-1} & 3^{n-1} \\ 3^{n-1} & 3^{n-1} & 3^{n-1} \end{bmatrix}$.



Watch Video Solution

29. If A is any $m \times n$ such that AB and BA are both defined show that B is an $n \times m$ matrix.



Watch Video Solution

30. A, B are two matrices such that AB and $A + B$ are both defined; show that A, B are square matrices of the same order.



Watch Video Solution

31. A man buys 8 dozens of mangoes, 10 dozens of apples and 4 dozens of bananas. Mangoes cost Rs. 18 per dozen, apples Rs. 9 per dozen and bananas Rs 6 per dozen. Represent the quantities bought by a row matrix and the prices by a column matrix and hence obtain the total cost.



Watch Video Solution

32. If $A = \begin{bmatrix} 2 & 3 & 4 \\ 1 & 2 & 3 \\ 4 & -5 & 7 \end{bmatrix}$, $B = \begin{bmatrix} 2 & 5 & -1 \\ 3 & -1 & 2 \\ 7 & 2 & -3 \end{bmatrix}$, verify that

$$(A + B)' = A' + B'$$



Watch Video Solution

33. If $A = \begin{bmatrix} 2 & 4 & -1 \\ -1 & 0 & 2 \end{bmatrix}$, $B = \begin{bmatrix} 3 & 4 \\ -1 & 2 \\ 2 & 1 \end{bmatrix}$, Show that $(AB)' = B'A'$.



Watch Video Solution

34. If $A = \begin{bmatrix} 2 & 4 \\ 5 & 6 \end{bmatrix}$, show that $(A - A^T)$ is a skew symmetric matrix, where A^T is the transpose of matrix A.



Watch Video Solution

35. Express the following matrix as a sum of symmetric and skew-

symmetric matrices.
$$\begin{bmatrix} 1 & 3 & 5 \\ -6 & 8 & 3 \\ -4 & 6 & 5 \end{bmatrix}$$



Watch Video Solution

36. Find the inverse of the following matrices, if it exists, using elementary

operation:
$$\begin{bmatrix} 2 & 5 \\ 1 & 3 \end{bmatrix}$$



Watch Video Solution

37. Using elementary transformations, find the inverse of each of the

matrices $\begin{bmatrix} 2 & -6 & 1 & -2 \end{bmatrix}$



Watch Video Solution

38. find inverse using elementary operations
$$\begin{bmatrix} 6 & -3 \\ -2 & 1 \end{bmatrix}$$



[Watch Video Solution](#)

39. Find the inverse of the following matrix using elementary operation:

$$\begin{bmatrix} 1 & 3 & -2 \\ -3 & 0 & -5 \\ 2 & 5 & 0 \end{bmatrix}$$

[Watch Video Solution](#)

40. Obtain the inverse of the following matrix using elementary

operations $A = \begin{bmatrix} 0 & 1 & 2 \\ 1 & 2 & 3 \\ 3 & 1 & 1 \end{bmatrix}$.

[Watch Video Solution](#)

Exercise

1. The number of matrices having 7 elements is.....

[Watch Video Solution](#)

2. The number of matrices having 7 elements is.....



Watch Video Solution

3. If a matrix has 5 elements, write all possible orders it can have.



Watch Video Solution

4. If a matrix has 8 elements, what are the possible orders it can have?



Watch Video Solution

5. If a matrix has 24 elements, what are the possible orders it can have?

What, if it has 13 elements?



Watch Video Solution

6. What are the possible orders a matrix can have if it has 13 elements



Watch Video Solution

7. If a matrix has 18 elements, what are the possible orders it can have?

What, if it has 5 elements?



Watch Video Solution

8. construct 2×3 matrix whose elements are given by $a_{ij} = i + 2j$



Watch Video Solution

9. Construct a 3×2 matrix whose elements are given by $a_{ij} = j - i$.



Watch Video Solution

10. Construct 3×3 matrix whose elements are given by : $a_{ij} = i \times j$



Watch Video Solution

11. Construct 3×3 matrix whose elements are given by : $a_{ij} = i - j$



Watch Video Solution

12. Construct 3×3 matrix whose elements are given by : $a_{ij} = 2i - 3j$



Watch Video Solution

13. Construct 3×3 matrix whose elements are given by : $a_{ij} = i + j$



Watch Video Solution

14. Construct a 2×2 matrix $A = [a_{ij}]$ whose elements a_{ij} are given by:

$$a_{ij} = i + j$$



Watch Video Solution

15. Construct a 2×2 matrix $A = [a_{ij}]$ whose elements are given by

$$a_{ij} = (i + j)^2.$$



Watch Video Solution

16. Construct 3×4 matrix $A = [a_{ij}]$ whose elements are given by

$$a_{ij} = i + j$$



Watch Video Solution

17. Construct a 3×4 matrix $A = [a_{ij}]$ whose elements a_{ij} are given by:)

$$a_{ij} = 2i$$

[Watch Video Solution](#)

18. Construct 3×4 matrix $A = [a_{ij}]$ whose elements are: $a_{ij} = i \cdot j$

[Watch Video Solution](#)

19. Construct 3×4 matrix $A = [a_{ij}]$ whose elements are: $a_{ij} = \frac{i}{j}$

[Watch Video Solution](#)

20. Construct a 3×4 matrix, whose elements are given by: (i) $a_{ij} = 2i - j$

[Watch Video Solution](#)

21. Construct a 3×4 matrix $A = [a_{ij}]$ whose elements a_{ij} are given by:

(i) $a_{ij} = j$ (ii) $a_{ij} = \frac{1}{2}| - 3i + j|$

[Watch Video Solution](#)

22. In the matrix $\begin{bmatrix} 1 & 0 & 5 \\ 2 & -3 & 4 \end{bmatrix}$ i. number of rows is.....



Watch Video Solution

23. In the matrix $\begin{bmatrix} 1 & 0 & 5 \\ 2 & -3 & 4 \end{bmatrix}$ number of columns is.....



Watch Video Solution

24. In the matrix $\begin{bmatrix} 1 & 0 & 5 \\ 2 & -3 & 4 \end{bmatrix}$ order of matrix matrix is.....



Watch Video Solution

25. In the matrix $\begin{bmatrix} 1 & 0 & 5 \\ 2 & -3 & 4 \end{bmatrix}$ the number of entries is.....



Watch Video Solution

26. If A is a row matrix as well as a column matrix then what is the order of A?



Watch Video Solution

27. Let $A = [a_{ij}] = \begin{bmatrix} 1 & -2 & 5 \\ 3 & 4 & -6 \\ 9 & 15 & 13 \end{bmatrix}$ and

$B = b_{ij} = \begin{bmatrix} 2 & 5 & 19 & -7 \\ 35 & -2 & \frac{5}{2} & 12 \\ \sqrt{3} & 1 & -5 & 17 \end{bmatrix}$ Then: order of the matrix A is..



Watch Video Solution

28. Let $A = [a_{ij}] = \begin{bmatrix} 1 & -2 & 5 \\ 3 & 4 & -6 \\ 9 & 15 & 13 \end{bmatrix}$ and

$B = b_{ij} = \begin{bmatrix} 2 & 5 & 19 & -7 \\ 35 & -2 & \frac{5}{2} & 12 \\ \sqrt{3} & 1 & -5 & 17 \end{bmatrix}$ Then: $a_{23} \dots$



Watch Video Solution

29. Let $A = [a_{ij}] = \begin{bmatrix} 1 & -2 & 5 \\ 3 & 4 & -6 \\ 9 & 15 & 13 \end{bmatrix}$ and

$B = [b_{ij}] = \begin{bmatrix} 2 & 5 & 19 & -7 \\ 35 & -2 & \frac{5}{2} & 12 \\ \sqrt{3} & 1 & -5 & 17 \end{bmatrix}$ Then: $a_{31} = \dots$

 Watch Video Solution

30. Let $A = [a_{ij}] = \begin{bmatrix} 1 & -2 & 5 \\ 3 & 4 & -6 \\ 9 & 15 & 13 \end{bmatrix}$ and

$B = b_{ij} = \begin{bmatrix} 2 & 5 & 19 & -7 \\ 35 & -2 & \frac{5}{2} & 12 \\ \sqrt{3} & 1 & -5 & 17 \end{bmatrix}$ Then: order of matrix B is.

 Watch Video Solution

31. Let $A = [a_{ij}] = \begin{bmatrix} 1 & -2 & 5 \\ 3 & 4 & -6 \\ 9 & 15 & 13 \end{bmatrix}$ and

$B = b_{ij} = \begin{bmatrix} 2 & 5 & 19 & -7 \\ 35 & -2 & \frac{5}{2} & 12 \\ \sqrt{3} & 1 & -5 & 17 \end{bmatrix}$ Then: the number of elements in A

and B is.....

[Watch Video Solution](#)

32. Let $A = [a_{ij}] = \begin{bmatrix} 1 & -2 & 5 \\ 3 & 4 & -6 \\ 9 & 15 & 13 \end{bmatrix}$ and

$B = [b_{ij}] = \begin{bmatrix} 2 & 5 & 19 & -7 \\ 35 & -2 & \frac{5}{2} & 12 \\ \sqrt{3} & 1 & -5 & 17 \end{bmatrix}$ Then: write the elements

$b_{13}, b_{21}, b_{33}, b_{24}, b_{23}$

[Watch Video Solution](#)

33. If $\begin{bmatrix} x - y & 2 \\ 1 & x + y \end{bmatrix} = \begin{bmatrix} 3 & 2 \\ 1 & 7 \end{bmatrix}$ then $x = \dots$. $y =$

[Watch Video Solution](#)

34. If $\begin{bmatrix} x - y & 2x - x_1 \\ 2x - y & 3x + y_1 \end{bmatrix}, = \begin{bmatrix} -1 & 5 \\ 0 & 13 \end{bmatrix}$ and coordinats of points P and Q be (x, y) and (x_1, y_1) respectively then P, Q = ?

[Watch Video Solution](#)

35. If $\begin{bmatrix} x - y & 2x + z \\ 2x - y & 3z + \omega \end{bmatrix} = \begin{bmatrix} -1 & 5 \\ 0 & 13 \end{bmatrix}$ find $x + y + z + \omega$



Watch Video Solution

36. If $\begin{bmatrix} x + y & z \\ 9x + 3y & \omega \end{bmatrix} = \begin{bmatrix} -2 & 4 \\ 0 & 5 \end{bmatrix}$, find x, y, z, w



Watch Video Solution

37. If $\begin{bmatrix} x & 3x - y \\ 2x + z & 3y - w \end{bmatrix} = \begin{bmatrix} 3 & 2 \\ 4 & 7 \end{bmatrix}$, then find $x + y + z + w$.



Watch Video Solution

38. Find x, y, z from the following equation: $\begin{bmatrix} x + y & 2 \\ 5 + z & xy \end{bmatrix} = \begin{bmatrix} 6 & 2 \\ 5 & 8 \end{bmatrix}$



Watch Video Solution

39. Find x, y, z from the following equation: $\begin{bmatrix} 4 & 3 \\ x & 5 \end{bmatrix} = \begin{bmatrix} y & z \\ 1 & 5 \end{bmatrix}$



Watch Video Solution

40. Write the values of x, y and z from the following equation:

$$\begin{bmatrix} x + y + z \\ x + z \\ y + z \end{bmatrix} = \begin{bmatrix} 9 \\ 5 \\ 7 \end{bmatrix}$$



Watch Video Solution

41. Find the values of a, b, c and d from the following equations:

$$[2a + ba - 2b5c - d4c + 3d] = [4 - 31124]$$



Watch Video Solution

42. Let $A = \begin{bmatrix} 2 & 4 \\ 3 & 2 \end{bmatrix}$, $B = \begin{bmatrix} 1 & 3 \\ -2 & 5 \end{bmatrix}$, $C = \begin{bmatrix} -2 & 5 \\ 3 & 4 \end{bmatrix}$ find $A+B+C$



Watch Video Solution

43. Let $A = \begin{bmatrix} 2 & 4 \\ 3 & 2 \end{bmatrix}$, $B = \begin{bmatrix} 1 & 3 \\ -2 & 5 \end{bmatrix}$, $C = \begin{bmatrix} -2 & 5 \\ 3 & 4 \end{bmatrix}$ find $A-B$



Watch Video Solution

44. Let $A = \begin{bmatrix} 2 & 4 \\ 3 & 2 \end{bmatrix}$, $B = \begin{bmatrix} 1 & 3 \\ -2 & 5 \end{bmatrix}$, $C = \begin{bmatrix} -2 & 5 \\ 3 & 4 \end{bmatrix}$ find $3A-C$



Watch Video Solution

45. If $A = \begin{bmatrix} 1 & 4 & -1 \\ 2 & 6 & 5 \end{bmatrix}$ and $B = \begin{bmatrix} 3 & -2 & -6 \\ 2 & 0 & -7 \end{bmatrix}$, find $A+B$ and $A-B$



Watch Video Solution

46. If $A = [122331]$ and $B = [3 - 1 - 1032]$, then find $2A - B$.



Watch Video Solution

47. Given $A = \begin{bmatrix} \sqrt{3} & 2 & 1 \\ 3 & -1 & 0 \end{bmatrix}$ and $B = \begin{bmatrix} 2 & -2 & \sqrt{5} \\ 3 & 1 & \frac{1}{2} \end{bmatrix}$, find $A + B$.



Watch Video Solution

48. Compute the following: $\begin{bmatrix} 0 & 1 & 5 \\ -3 & 2 & 1 \end{bmatrix} + \begin{bmatrix} 6 & 2 & -3 \\ -1 & 4 & 2 \end{bmatrix}$



Watch Video Solution

49. Compute the following: $\begin{bmatrix} 2 & -1 \\ 3 & 5 \end{bmatrix} + \begin{bmatrix} 4 & 3 \\ 1 & -2 \end{bmatrix}$



Watch Video Solution

50. Compute the following: (i) $[ab - ba] + [aa]$ (ii)

$[a^2 + b^2b^2 + c^2a^2 + c^2a^2 + b^2] + [2ab2bc - 2ac - 2ab]$ (iii)

$[-14 - 68516285] + [1276805324]$ (iv)

$[\cos^2 x \sin^2 x \sin^2 x \cos^2 x] + [\sin^2 x \cos^2 x \cos^2 x \sin^2 x]$



Watch Video Solution

51. Compute the following: $\begin{bmatrix} \cos^2 x & \sin^2 x \\ \sin^2 x & \cos^2 x \end{bmatrix} + \begin{bmatrix} \sin^2 x & \cos^2 x \\ \cos^2 x & \sin^2 x \end{bmatrix}$



Watch Video Solution

52. Compute the following: $\begin{bmatrix} a^2 + b^2 & b^2 + c^2 \\ a^2 + c^2 & a^2 + b^2 \end{bmatrix} + \begin{bmatrix} 2ab & 2bc \\ -2ac & -2ab \end{bmatrix}$



Watch Video Solution

53. Compute the following: $\begin{bmatrix} -1 & 4 & -6 \\ 8 & 5 & 16 \\ 2 & 8 & 5 \end{bmatrix} + \begin{bmatrix} 12 & 7 & 6 \\ 8 & 0 & 5 \\ 3 & 2 & 4 \end{bmatrix}$



Watch Video Solution

54. Compute the following: $\begin{bmatrix} 2 & 3 & 1 \\ 5 & -1 & 2 \\ 0 & 3 & 5 \end{bmatrix} + \begin{bmatrix} 1 & -2 & 3 \\ -3 & 1 & 5 \\ 6 & 2 & 0 \end{bmatrix}$



Watch Video Solution

55. If $A = \begin{bmatrix} 1 & -2 & 3 \\ 4 & 5 & -1 \\ 5 & 0 & 2 \end{bmatrix}$ find $3A$



Watch Video Solution

56. If $A = \begin{bmatrix} 2 & -1 \\ 4 & 2 \end{bmatrix}$, $B = \begin{bmatrix} 4 & 3 \\ -2 & 1 \end{bmatrix}$, $C = \begin{bmatrix} -2 & -3 \\ -1 & 2 \end{bmatrix}$, compute the following: $A+(B+C)$



Watch Video Solution

57. If $A = \begin{bmatrix} 2 & -1 \\ 4 & 2 \end{bmatrix}$, $B = \begin{bmatrix} 4 & 3 \\ -2 & 1 \end{bmatrix}$, $C = \begin{bmatrix} -2 & -3 \\ -1 & 2 \end{bmatrix}$, compute the following: $(A+B)+C$



Watch Video Solution

58. If $A = \begin{bmatrix} 2 & -1 \\ 4 & 2 \end{bmatrix}$, $B = \begin{bmatrix} 4 & 3 \\ -2 & 1 \end{bmatrix}$, $C = \begin{bmatrix} -2 & -3 \\ -1 & 2 \end{bmatrix}$, compute the following: $-2A+(B+C)$



Watch Video Solution

59. If $A = \begin{bmatrix} 2 & -1 \\ 4 & 2 \end{bmatrix}$, $B = \begin{bmatrix} 4 & 3 \\ -2 & 1 \end{bmatrix}$, $C = \begin{bmatrix} -2 & -3 \\ -1 & 2 \end{bmatrix}$, compute the following: $A+(2B-C)$



Watch Video Solution

60.

If

$$A = \begin{bmatrix} 1 & 2 & 3 \\ -1 & 0 & 2 \\ 1 & -3 & 1 \end{bmatrix}, B = \begin{bmatrix} 4 & 5 & 6 \\ -1 & 0 & 1 \\ 2 & 1 & 2 \end{bmatrix} \text{ and } C = \begin{bmatrix} -1 & -2 & 1 \\ -1 & 2 & 3 \\ -1 & -2 & 2 \end{bmatrix}$$

verify that $A+(B+C)=(A+B)+C$



Watch Video Solution

61. Evaluate $\begin{bmatrix} \sin^2 \theta & 1 \\ \cos^2 \theta & 0 \end{bmatrix} + \begin{bmatrix} \cos^2 \theta & 0 \\ -\cos^2 \theta & 1 \end{bmatrix} + \begin{bmatrix} 0 & -1 \\ -1 & 0 \end{bmatrix}$



Watch Video Solution

62. From the following equation, find the values of x and y:

$$2 \begin{bmatrix} x & 5 \\ 7 & y-3 \end{bmatrix} + \begin{bmatrix} 3 & 4 \\ 1 & 2 \end{bmatrix} = \begin{bmatrix} 7 & 14 \\ 15 & 14 \end{bmatrix}$$



Watch Video Solution

63. Find the values of x and y satisfying the equation:

$$2 \begin{bmatrix} 1 & 3 \\ 0 & x \end{bmatrix} + \begin{bmatrix} y & 0 \\ 1 & 2 \end{bmatrix} = \begin{bmatrix} 5 & 6 \\ 1 & 8 \end{bmatrix}$$



Watch Video Solution

64. Solve the equation for x, y, z and t, if $2[xzyt] + 3[1 - 102] = 3[3546]$



Watch Video Solution

65. If $A = \text{diag}[1, 3, 2]$, $B = \text{diag}, [0, 2, 5]$, $C = \text{diag}[3, -2, 5]$. Find $4A - 3B$



Watch Video Solution

66. If $A = \text{diag}[1, 3, 2]$, $B = \text{diag}, [0, 2, 5]$, $C = \text{diag}[3, -2, 5]$. Find $A + B - 2C$



Watch Video Solution

67. Find matrix X, if $X + \begin{bmatrix} 2 & 5 \\ 3 & 2 \end{bmatrix} = \begin{bmatrix} 4 & 0 \\ -7 & 6 \end{bmatrix}$



Watch Video Solution

68. Find a matrix X such that $2A - B + X = 0$ where $A = \begin{bmatrix} 3 & 1 \\ 0 & 2 \end{bmatrix}$ and $B = \begin{bmatrix} -2 & 1 \\ 0 & 3 \end{bmatrix}$



Watch Video Solution

69. Find a matrix X such that $A + 2B + X = 0$ where

$$A = \begin{bmatrix} 2 & -1 \\ 3 & 5 \end{bmatrix}, B = \begin{bmatrix} -1 & 1 \\ 0 & 2 \end{bmatrix}$$



Watch Video Solution

70. Find a matrix X of order 3×2 such that $2A + 3X = 58B$, where

$$A = \begin{bmatrix} 8 & 0 \\ 4 & -2 \\ 3 & 6 \end{bmatrix}, B = \begin{bmatrix} 2 & -2 \\ 4 & 2 \\ -5 & 1 \end{bmatrix}$$



Watch Video Solution

71. Find matrices X and Y , if $X + Y = \begin{bmatrix} 5 & 2 & 0 & 9 \end{bmatrix}$ and $X - Y = \begin{bmatrix} 3 & 6 & 0 & -1 \end{bmatrix}$.



Watch Video Solution

72. Show that $\cos \theta \begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix} + \sin \theta \begin{bmatrix} \sin \theta & -\cos \theta \\ \cos \theta & \sin \theta \end{bmatrix} = I$

[Watch Video Solution](#)

73. If $2X + 3Y = \begin{bmatrix} 2 & 3 \\ 4 & 0 \end{bmatrix}$ and $3X + 2Y = \begin{bmatrix} -2 & 2 \\ 1 & -5 \end{bmatrix}$ find $X + Y$.

[Watch Video Solution](#)

74. Given $A = \begin{bmatrix} 1 & 2 & -3 \\ 5 & 0 & 2 \\ 1 & -1 & 1 \end{bmatrix}$ and $B = \begin{bmatrix} 3 & -1 & 2 \\ 4 & 2 & 5 \\ 2 & 0 & 3 \end{bmatrix}$, find the matrix C such that $A+C=B$.

[Watch Video Solution](#)

75. If $A = \begin{bmatrix} 2 & 3 & 4 \\ -3 & 0 & 2 \end{bmatrix}$, $B = \begin{bmatrix} 3 & -4 & -5 \\ 1 & 2 & 1 \end{bmatrix}$ and $C = \begin{bmatrix} 5 & -1 & 2 \\ 7 & 0 & 3 \end{bmatrix}$, find the matrix X such that $2A + 3B = X + C$

[Watch Video Solution](#)

76. If $A = \begin{bmatrix} 2 & 3 & -4 \\ 1 & 0 & 6 \\ -2 & 1 & 5 \end{bmatrix}$, $B = \begin{bmatrix} 5 & 1 & 2 \\ 6 & -1 & 4 \\ 5 & 3 & -4 \end{bmatrix}$, find $2A - 3B$.



Watch Video Solution

77. If $A = \begin{bmatrix} 1 & 2 & 3 \\ -1 & 0 & 2 \\ 1 & -3 & 1 \end{bmatrix}$, $B = \begin{bmatrix} 4 & 5 & 6 \\ -1 & 0 & 1 \\ 2 & 1 & 2 \end{bmatrix}$, $C = \begin{bmatrix} -1 & -2 & 1 \\ -1 & 2 & 3 \\ -1 & -2 & 2 \end{bmatrix}$

find $A - 2B + 3C$



Watch Video Solution

78. If $A = [2, 3, 5]$ and $B = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}$, find AB



Watch Video Solution

79. If $A = \begin{bmatrix} 2 & 1 & 3 \\ 4 & 1 & 0 \end{bmatrix}$ and $B = \begin{bmatrix} 1 & -1 \\ 0 & 2 \\ 5 & 0 \end{bmatrix}$ find AB and BA .



Watch Video Solution

[Watch Video Solution](#)

80. Evaluate the following : $\begin{bmatrix} 0 & 2 \\ 0 & 3 \end{bmatrix} \begin{bmatrix} 4 & 6 \\ 0 & 0 \end{bmatrix}$

[Watch Video Solution](#)

81. Evaluate the following : $\begin{bmatrix} 1 & 3 \\ 2 & 1 \end{bmatrix} \begin{bmatrix} 4 \\ -1 \end{bmatrix}$

[Watch Video Solution](#)

82. Evaluate the following : $\begin{bmatrix} 2 \\ 4 \\ 6 \end{bmatrix} \begin{bmatrix} 1 & 2 & 3 \end{bmatrix}$

[Watch Video Solution](#)

83. Evaluate the following : $\begin{bmatrix} 1 & 2 & 3 \end{bmatrix} \begin{bmatrix} 2 \\ 4 \\ 6 \end{bmatrix}$

[Watch Video Solution](#)

84. Evaluate the following : $\begin{bmatrix} 1 & 2 & -3 \\ -2 & 1 & 7 \end{bmatrix} \begin{bmatrix} 2 & 3 & 1 \\ 5 & 4 & 2 \\ 1 & 6 & 3 \end{bmatrix}$



Watch Video Solution

85. Evaluate the following : $\begin{bmatrix} 1 & 4 & 2 \\ 5 & -2 & 3 \end{bmatrix} \begin{bmatrix} 2 & -4 \\ 1 & -3 \\ 4 & 0 \end{bmatrix}$



Watch Video Solution

86. If $A = \begin{bmatrix} 2 & 9 \\ 4 & 3 \end{bmatrix}$ and $B = \begin{bmatrix} 1 & 5 \\ 7 & 2 \end{bmatrix}$ find $AB-BA$



Watch Video Solution

87. If $A = \begin{bmatrix} \cos \theta & \sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$, $B = \begin{bmatrix} \cos \phi & \sin \phi \\ \sin \phi & \cos \phi \end{bmatrix}$ show that $AB=BA$.



Watch Video Solution

88. If $A = \begin{bmatrix} 1 & 2 \\ 5 & 7 \end{bmatrix}$ and $B = \begin{bmatrix} 2 & 0 \\ 3 & -4 \end{bmatrix}$, show that $AB \neq BA$



Watch Video Solution

89. If $A = \begin{bmatrix} 1 & 2 \\ 3 & -4 \\ 5 & 6 \end{bmatrix}$ and $B = \begin{bmatrix} 4 & 5 & 6 \\ 7 & -8 & 2 \end{bmatrix}$, is $AB=BA$? Also find AB and BA .



Watch Video Solution

90. If $A = \begin{bmatrix} -1 & 2 \\ 3 & 4 \end{bmatrix}$ and $B = \begin{bmatrix} 2 & -3 \\ 5 & 1 \end{bmatrix}$ show that $AB \neq BA$



Watch Video Solution

91. If $A = \begin{bmatrix} 1 & 2 & 3 \\ 0 & 1 & 0 \\ 1 & 1 & 0 \end{bmatrix}$ and $B = \begin{bmatrix} -1 & 1 & 0 \\ 0 & -1 & 1 \\ 2 & 3 & 4 \end{bmatrix}$ show that $AB \neq BA$



Watch Video Solution

92. Evaluate the following: $\left(\begin{bmatrix} 1 & 3 \\ -1 & -4 \end{bmatrix} + \begin{bmatrix} 3 & -2 \\ -1 & 1 \end{bmatrix}\right) \begin{bmatrix} 1 & 3 & 5 \\ 2 & 4 & 6 \end{bmatrix}$



Watch Video Solution

93. Evaluate the following : Find $\begin{bmatrix} 1 & -1 \\ 0 & 2 \\ 2 & 3 \end{bmatrix} \left(\begin{bmatrix} 1 & 0 & 2 \\ 2 & 0 & 1 \end{bmatrix} - \begin{bmatrix} 0 & 1 & 2 \\ 1 & 0 & 2 \end{bmatrix} \right)$



Watch Video Solution

94. Evaluate the following : $\begin{bmatrix} 1 & 1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 4 \\ 4 \\ 4 \end{bmatrix}$



Watch Video Solution

95. Evaluate : $\begin{bmatrix} 1 & 3 & 5 \end{bmatrix} \begin{bmatrix} 1 & 0 & 3 \\ 2 & 0 & 1 \\ 0 & 1 & 2 \end{bmatrix} \begin{bmatrix} 1 \\ 4 \\ 6 \end{bmatrix}$



Watch Video Solution

96. Evaluate the following : $\begin{bmatrix} 1 & -1 \\ 0 & 2 \\ 2 & 3 \end{bmatrix} \left(\begin{bmatrix} 1 & 0 & 2 \\ 2 & 0 & 1 \end{bmatrix} - \begin{bmatrix} 0 & 1 & 3 \\ 1 & 0 & 2 \end{bmatrix} \right)$



Watch Video Solution

97. If $P(x) = [(\cos x, \sin x), (-\sin x, \cos x)]$, then show that $P(x) \cdot P(y) = P(x + y) = P(y) \cdot P(x)$.



Watch Video Solution

98. If $F(x) = [\cos x - \sin x, 0, \sin x, \cos x, 0, 0, 0, 1]$, show that $F(x)F(y) = F(x + y)$.



Watch Video Solution

99. If $A = \begin{bmatrix} 2 & 3 \\ -1 & 5 \end{bmatrix}$, $B = \begin{bmatrix} 3 & -1 \\ 4 & 7 \end{bmatrix}$ and $C = \begin{bmatrix} 5 & -1 \\ 0 & 3 \end{bmatrix}$, show that $A(B + C) = AB + AC$



Watch Video Solution

[Watch Video Solution](#)

100. If $A = \begin{bmatrix} 2 & -1 \\ -1 & 2 \end{bmatrix}$ and $B = \begin{bmatrix} 1 & 4 \\ -1 & 1 \end{bmatrix}$ then find $(A + B)^2$

[Watch Video Solution](#)

101. If $A = \begin{bmatrix} 2 & 3 \\ 4 & 5 \end{bmatrix}$, $B = \begin{bmatrix} 3 & 4 \\ 7 & 2 \end{bmatrix}$, $C = \begin{bmatrix} 1 & 0 \\ 0 & 7 \end{bmatrix}$ then verify that $(AB)C = A(BC)$

[Watch Video Solution](#)

102. If $A = \begin{bmatrix} \cos \alpha & \sin \alpha \\ -\sin \alpha & \cos \alpha \end{bmatrix}$ then show that $A^2 = \begin{bmatrix} \cos 2\alpha & \sin 2\alpha \\ -\sin 2\alpha & \cos 2\alpha \end{bmatrix}$

[Watch Video Solution](#)

103. If $A = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$, $B = \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix}$ and $C = \begin{bmatrix} i & 0 \\ 0 & -i \end{bmatrix}$, show that $A^2 = B^2 = -C^2 = I_2$ and $AB = -BA$, $AC = -CA$ and $BC = -CB$



Watch Video Solution

104. If $A = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix}$, $B = \begin{bmatrix} 0 & 5 & 7 \\ 0 & 0 & 6 \\ 0 & 0 & 0 \end{bmatrix}$ and $C = \begin{bmatrix} -1 & 3 & 5 \\ 1 & -3 & -5 \\ -1 & 3 & 5 \end{bmatrix}$,

show that $A^2 = I$



Watch Video Solution

105. If $A = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix}$, $B = \begin{bmatrix} 0 & 5 & 7 \\ 0 & 0 & 6 \\ 0 & 0 & 0 \end{bmatrix}$ and $C = \begin{bmatrix} -1 & 3 & 5 \\ 1 & -3 & -5 \\ -1 & 3 & 5 \end{bmatrix}$,

show that $C^2 = C$



Watch Video Solution

106. If $B = \begin{bmatrix} 0 & 5 & 7 \\ 0 & 0 & 6 \\ 0 & 0 & 0 \end{bmatrix}$, show that $B^4 = O$



Watch Video Solution

107. If $A = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 0 & -2 \end{bmatrix}$, $B = \begin{bmatrix} 1 & 1 & -1 \\ 2 & 0 & 3 \\ 3 & -1 & 2 \end{bmatrix}$ and $C = \begin{bmatrix} 1 & 3 \\ 0 & 2 \\ -1 & 4 \end{bmatrix}$

find $A(BC)$.



Watch Video Solution

108. Show that $\begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}^2 = \begin{bmatrix} 1 & 3 \\ 0 & 1 \end{bmatrix}$



Watch Video Solution

109. If $A = \begin{bmatrix} \cos 2\theta & \sin 2\theta \\ -\sin 2\theta & \cos 2\theta \end{bmatrix}$, find A^2



Watch Video Solution

110. If $A = \begin{bmatrix} 0.8 & 0.6 \\ -0.6 & 0.8 \end{bmatrix}$ then find A^3



Watch Video Solution

111. Two matrices A and B have in total 6 different elements (none repeated). How many different matrices A and B are possible such that products AB is defined.

$$A = \begin{bmatrix} \cos^2 \theta & \cos \theta \sin \theta \\ \cos \theta \sin \theta & \sin^2 \theta \end{bmatrix} \text{ and } B = \begin{bmatrix} \cos^2 \phi & \cos \phi \sin \phi \\ \cos \phi \sin \phi & \sin^2 \phi \end{bmatrix} \text{ then}$$

is

 [Watch Video Solution](#)

112. If $A = \begin{bmatrix} 0 & c & -b \\ -c & 0 & a \\ b & -a & 0 \end{bmatrix}$ and $B = \begin{bmatrix} a^2 & ab & ac \\ ab & b^2 & bc \\ ac & bc & c^2 \end{bmatrix}$, show that

$$AB = BA = O_3 \times 3.$$

 [Watch Video Solution](#)

113. If $A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ a & b & -1 \end{bmatrix}$, find A^2

 [Watch Video Solution](#)

114. Let $A = \begin{bmatrix} -1 & 1 & -1 \\ 3 & -3 & 3 \\ 5 & 5 & 5 \end{bmatrix}$ and $B = \begin{bmatrix} 0 & 4 & 3 \\ 1 & -3 & -3 \\ -1 & 4 & 4 \end{bmatrix}$, compute $A^2 - B^2$.



Watch Video Solution

115. If $A = \begin{bmatrix} 2 & 3 & 4 \\ 1 & 2 & 3 \\ -1 & 1 & 2 \end{bmatrix}$, $B = \begin{bmatrix} 1 & 3 & 0 \\ -1 & 2 & 1 \\ 0 & 0 & 2 \end{bmatrix}$, find AB and BA and show that $AB \neq BA$



Watch Video Solution

116. If ω is cube roots of unity, prove that

$$\left\{ \begin{bmatrix} 1 & \omega & \omega^2 \\ \omega & \omega^2 & 1 \\ \omega^2 & 1 & \omega \end{bmatrix} + \begin{bmatrix} \omega & \omega^2 & 1 \\ \omega^2 & 1 & \omega \\ \omega & \omega^2 & 1 \end{bmatrix} \right\} \begin{bmatrix} 1 \\ \omega \\ \omega^2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$



Watch Video Solution

117. If

$$A = \begin{bmatrix} 1 & 0 & -2 \\ 3 & -1 & 0 \\ -2 & 1 & 1 \end{bmatrix}, B = \begin{bmatrix} 0 & 5 & -4 \\ -2 & 1 & 3 \\ -1 & 0 & 2 \end{bmatrix} \text{ and } C = \begin{bmatrix} 1 & 5 & 2 \\ -1 & 1 & 0 \\ 0 & -1 & 1 \end{bmatrix}$$

verify that $A(B-C) = (AB-AC)$



Watch Video Solution

118. Find x if $\begin{bmatrix} x & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ -1 & -3 \end{bmatrix} \begin{bmatrix} x \\ 3 \end{bmatrix} = O$



Watch Video Solution

119. Solve the matrix equations: $\begin{bmatrix} 2x & 3 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ -3 & 0 \end{bmatrix} \begin{bmatrix} x \\ 8 \end{bmatrix} = 0$



Watch Video Solution

120. Find the values of a and b for which the following holds:

$$\begin{bmatrix} 4 & 2 \\ 3 & -1 \end{bmatrix} \begin{bmatrix} a \\ b \end{bmatrix} = \begin{bmatrix} -4 \\ 2 \end{bmatrix}$$



Watch Video Solution

121. Let $A = \begin{bmatrix} 2 & -13 & 4 \end{bmatrix}$, $B = \begin{bmatrix} 5 & 2 & 7 & 4 \end{bmatrix}$, $C = \begin{bmatrix} 2 & 5 & 3 & 8 \end{bmatrix}$. Find a matrix D such that

$$CD - AB = 0$$



Watch Video Solution

122. Find the matrix X so that

$$X \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix} = \begin{bmatrix} -7 & -8 & -9 \\ 2 & 4 & 6 \end{bmatrix}.$$



Watch Video Solution

123. Find the value of x , if $\begin{bmatrix} 1 & 3 & 2 \\ 2 & 5 & 1 \\ 15 & 3 & 2 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \\ x \end{bmatrix} = 0$



Watch Video Solution

124. Find the value of x such that $\begin{bmatrix} 1 & x & 1 \end{bmatrix} \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 3 & 2 & 5 \end{bmatrix} \begin{bmatrix} 1 \\ -2 \\ 3 \end{bmatrix} = O$



Watch Video Solution

125. Find x , if $\begin{bmatrix} x & -5 & -1 \end{bmatrix} \begin{bmatrix} 10 & 20 & 21 & 20 & 3 \end{bmatrix} \begin{bmatrix} x & 4 & 1 \end{bmatrix} = O$.



Watch Video Solution

126. For what values of x : $\begin{bmatrix} 1 & 2 & 1 \end{bmatrix} \begin{bmatrix} 12 & 20 & 11 & 10 & 2 \end{bmatrix} \begin{bmatrix} 0 & 2 & x \end{bmatrix} = O$?



Watch Video Solution

127. If $\begin{bmatrix} x & 4 & 1 \end{bmatrix} \begin{bmatrix} 2 & 1 & 2 \\ 1 & 0 & 2 \\ 0 & 2 & -4 \end{bmatrix} \begin{bmatrix} x \\ 4 \\ -1 \end{bmatrix} = 0$, find x .



Watch Video Solution

128. If $[1, 1, x] \begin{bmatrix} 1 & 0 & 2 \\ 0 & 2 & 1 \\ 2 & 1 & 0 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} = O$ find x



Watch Video Solution

129. Find a 2×2 matrix B such that $\begin{bmatrix} 6 & 5 \\ 5 & 6 \end{bmatrix} B = \begin{bmatrix} 11 & 0 \\ 0 & 11 \end{bmatrix}$



Watch Video Solution

130. Solve the matrix equation $\begin{bmatrix} 5 & 4 \\ 1 & 1 \end{bmatrix} X = \begin{bmatrix} 1 & -2 \\ 1 & 3 \end{bmatrix}$, where X is a 2×2 matrix.



Watch Video Solution

131. Without using the concept of inverse of a matrix, find the matrix $[xyzu]$ such that $[5 \ -7 \ -23][xyzu] = [-16 \ -672]$.



Watch Video Solution

132. Find the matrix A such that $\begin{bmatrix} 2 & -1 \\ 1 & 0 \\ -3 & 4 \end{bmatrix} A = \begin{bmatrix} -1 & -8 & -10 \\ 1 & -2 & -5 \\ 9 & 22 & 15 \end{bmatrix}$



Watch Video Solution

133. Find the matrix A such that $A \cdot \begin{bmatrix} 2 & 3 \\ 4 & 5 \end{bmatrix} = \begin{bmatrix} 0 & -4 \\ 10 & 3 \end{bmatrix}$.



Watch Video Solution

134. If $A = [3 \ -5 \ -42]$, find $A^2 - 5A - 14I$.



Watch Video Solution

135. Show that the matrix $A = \begin{bmatrix} 5 & 3 \\ 12 & 7 \end{bmatrix}$ then show $A^2 - 12A - I = O$.



Watch Video Solution

136. Show that the matrix $A = \begin{bmatrix} 2 & 3 & 1 & 2 \end{bmatrix}$ satisfies the equation $A^2 - 4A + I = 0$



Watch Video Solution

137. If $A^2 = 8A + kI$ where $A = \begin{bmatrix} 1 & 0 \\ -1 & 7 \end{bmatrix}$, then k is



Watch Video Solution

138. If $A = \begin{bmatrix} 3 & -2 \\ 4 & -2 \end{bmatrix}$, find k such that $A^2 - kA - 2I = 0$.



Watch Video Solution

139. If $A = \begin{bmatrix} 10 & -17 \end{bmatrix}$, find k such that $A^2 - 8A + kI = O$.



Watch Video Solution

140. If $f(x) = x^2 - 5x + 7$ and $A = \begin{bmatrix} 3 & 1 \\ -1 & 2 \end{bmatrix}$. Find $f(A)$



Watch Video Solution

141. If $A = \begin{bmatrix} 3 & 4 \\ -4 & -3 \end{bmatrix}$, find $f(A)$, where $f(x) = x^2 - 5x + 7$



Watch Video Solution

142. If $A = [1221]$ and $f(x) = x^2 - 2x - 3$, show that $f(A) = O$.



Watch Video Solution

143. If $A = \begin{bmatrix} 2 & -1 \\ 3 & 2 \end{bmatrix}$ and $B = \begin{bmatrix} 0 & 4 \\ -1 & 7 \end{bmatrix}$, find $3A^2 - 2B + I$.



Watch Video Solution

144. Questions of matrix polynomial Equation

[Watch Video Solution](#)

145. If $A = \begin{bmatrix} 1 & 2 & 3 \\ 3 & 2 & 1 \\ 4 & 2 & 1 \end{bmatrix}$ then show that $A^3 - 23A = 40I = 0$

[Watch Video Solution](#)

146. If $A = [102021203]$, prove that $A^3 - 6A^2 + 7A + 2I = 0$

[Watch Video Solution](#)

147. If $A = \begin{bmatrix} 3 & 1 \\ -1 & 2 \end{bmatrix}$, show that $A^2 - 5A + 7I = O$. Hence, find A^{-1} .

[Watch Video Solution](#)

148. If $\begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$, prove that $A^n = \begin{bmatrix} 1 & n \\ 0 & 1 \end{bmatrix}$ for all positive integers n .

[Watch Video Solution](#)

149. if $A = \begin{bmatrix} 3 & -4 \\ 1 & -1 \end{bmatrix}$, then prove that $A^n = \begin{bmatrix} 1 + 2n & -4n \\ n & 1 - 2n \end{bmatrix}$,

where n is any positive integer.



Watch Video Solution

150. If $A = \text{diag}(a \ b \ c)$, show that $A^n = \text{diag}(a^n \ b^n \ c^n)$ for all positive integer n .



Watch Video Solution

151. If $A = \begin{bmatrix} \cos \alpha & \sin \alpha \\ -\sin \alpha & \cos \alpha \end{bmatrix}$, prove by mathematical induction that,
 $A^n = \begin{bmatrix} \cos n\alpha & \sin n\alpha \\ -\sin n\alpha & \cos n\alpha \end{bmatrix}$ for every natural number n



Watch Video Solution

152. If $A = [\cos \theta \ i \sin \theta \ i \sin \theta \cos \theta]$, then prove by principle of mathematical induction that $A^n = [\cos n \theta \ i \sin n \theta \ i \sin n \theta \cos n \theta]$ for all $n \in N$.



Watch Video Solution

153. If A and B are square matrices of the same order such that $AB = BA$, then prove by induction that $AB^n = B^n A$. Further prove that $(AB)^n = A^n B^n$ for all $n \in N$.



Watch Video Solution

154. A trust fund has Rs 30,000 that must be invested in two different types of bonds. The first bond pays 5% interest per year, and the second bond pays 7% interest per year. Using matrix multiplication, determine how to divide Rs 30,000 among the tw



Watch Video Solution

155. A store has in stock 20 dozen shirts, 15 dozen trousers, and 25 dozen pair of socks. If the selling prices is Rs. 50 per shirt, Rs. 90 per trouser and Rs. 12 per pair of socks, then find the total amount the store owner will get after selling all the items in the stock.



Watch Video Solution

156. The cooperative stores of a particular school has 10 dozen physics books, 8 dozen chemistry books and 5 dozen mathematics books. Their selling prices are Rs. 8.30, Rs. 3.45 and Rs. 4.50 each respectively. Find the total amount the store will receive from selling all the items.



Watch Video Solution

157. The bookshop of a particular school has 10 dozen chemistry books, 8 dozen physics books, 10 dozen economics books. Their selling prices are Rs 80, Rs 60 and Rs 40 each respectively. Find the total amount the bookshop will receive from selling all

[Watch Video Solution](#)

158. Find the transpose of the matrix $\begin{bmatrix} 1 & 3 \\ 2 & 6 \\ 5 & -3 \end{bmatrix}$

[Watch Video Solution](#)

159. If $A = \begin{bmatrix} 3 \\ 1 \\ -2 \end{bmatrix}$, $B = \begin{bmatrix} 1 & 3 & -5 \end{bmatrix}$, verify $(AB)' = B'A'$

[Watch Video Solution](#)

160. If $A = \begin{bmatrix} -2 \\ 4 \\ 5 \end{bmatrix}$, $B = \begin{bmatrix} 1 & 3 & -6 \end{bmatrix}$, verify that $(AB)' = B'A'$.

[Watch Video Solution](#)

161. If $A = [-1/2/3]$ and $B = [-2 \ -1 \ -4]$, verify that $(AB)^T = B^T A^T$



Watch Video Solution

162. If $A = \begin{bmatrix} -2 \\ 4 \\ 5 \end{bmatrix}$, $B = [1 \ 3 \ -6]$, verify that $(AB)' = B' A'$.



Watch Video Solution

163. If $A = \begin{bmatrix} 10 & 3 \\ 1 & 2 \end{bmatrix}$ and $B = \begin{bmatrix} 1 & 0 \\ 1 & -1 \end{bmatrix}$, verify that $(AB)' = B' A'$



Watch Video Solution

164. If $A = \begin{bmatrix} 1 & 4 \\ 0 & 5 \\ 6 & 7 \end{bmatrix}$ and $B = \begin{bmatrix} 2 & 3 & -1 \\ 1 & 0 & -7 \end{bmatrix}$, verify that $(AB)' = B' A'$



Watch Video Solution

165. If $A = \begin{bmatrix} 2 & 1 & 3 \\ 4 & 1 & 0 \end{bmatrix}$, $B = \begin{bmatrix} 1 & -1 \\ 0 & 2 \\ 5 & 0 \end{bmatrix}$, verify that $(AB)' = B'A'$



Watch Video Solution

166. If $A = \begin{bmatrix} 5 & 1 \\ 6 & 7 \end{bmatrix}$, $B = \begin{bmatrix} 2 & 1 \\ 3 & 4 \end{bmatrix}$, $C = \begin{bmatrix} 1 & 3 \\ -1 & 4 \end{bmatrix}$, verify the following
 $(A + B)' = A' + B'$



Watch Video Solution

167. If $A = \begin{bmatrix} 5 & 1 \\ 6 & 7 \end{bmatrix}$, $B = \begin{bmatrix} 2 & 1 \\ 3 & 4 \end{bmatrix}$, $C = \begin{bmatrix} 1 & 3 \\ -1 & 4 \end{bmatrix}$, verify the following
 $(A')' = A$



Watch Video Solution

168. If $A = \begin{bmatrix} 5 & 1 \\ 6 & 7 \end{bmatrix}$, $B = \begin{bmatrix} 2 & 1 \\ 3 & 4 \end{bmatrix}$, $C = \begin{bmatrix} 1 & 3 \\ -1 & 4 \end{bmatrix}$, verify the following
 $(AC)' = C'A'$

[Watch Video Solution](#)

169. If $A = \begin{bmatrix} 5 & 1 \\ 6 & 7 \end{bmatrix}$, $B = \begin{bmatrix} 2 & 1 \\ 3 & 4 \end{bmatrix}$, $C = \begin{bmatrix} 1 & 3 \\ -1 & 4 \end{bmatrix}$, verify the following
 $(AB)' = B'A'$

[Watch Video Solution](#)

170. If (i) $A = [\cos \alpha \sin \alpha - \sin \alpha \cos \alpha]$, then verify that $A'A = I$
(ii) $A = [\sin \alpha \cos \alpha - \cos \alpha \sin \alpha]$, then verify that $A'A = I$.

[Watch Video Solution](#)

171. If $A = [\sin \alpha \cos \alpha - \cos \alpha \sin \alpha]$, verify that $A^T A = I_2$.

[Watch Video Solution](#)

172. Find x and y if the matrix $A = \frac{1}{3} \begin{bmatrix} 1 & 2 & 2 \\ 2 & 1 & -2 \\ x & 2 & y \end{bmatrix}$ satisfies the condition

$$AA' = A'A = I_3$$



Watch Video Solution

173. If $A = \begin{bmatrix} 0 & a & b \\ -a & 0 & c \\ -b & -c & 0 \end{bmatrix}$, find $\frac{1}{2}(A + A')$ and $\frac{1}{2}(A - A')$



Watch Video Solution

174. If $A = \begin{bmatrix} 3 & -4 \\ 1 & -1 \end{bmatrix}$, show that $A - A^T$ is a skew symmetric matrix.



Watch Video Solution

175. If $A = \begin{bmatrix} 3 & 4 \\ 5 & 1 \end{bmatrix}$, show that $A - A^T$ is a skew symmetric matrix, where A^T denotes the transpose of matrix A



Watch Video Solution

[Watch Video Solution](#)

176. If $A = \begin{bmatrix} 4 & 1 \\ 5 & 8 \end{bmatrix}$, show that $A + A^T$ is symmetric matrix, where A^T denotes the transpose of matrix A



[Watch Video Solution](#)

177. If $A = \begin{bmatrix} 2 & 3 \\ 4 & 5 \end{bmatrix}$, prove that $A - A^T$ is a skew-symmetric matrix.



[Watch Video Solution](#)

178. Expressing the following matrices as the sum of a symmetric and

skew symmetric matrix : $\begin{bmatrix} 1 & 2 & 4 \\ 6 & 8 & 1 \\ 3 & 5 & 7 \end{bmatrix}$



[Watch Video Solution](#)

179. Express the matrix $A = \begin{bmatrix} 2 & 4 & -6 \\ 7 & 3 & 5 \\ 1 & -2 & 4 \end{bmatrix}$ is the sum of a symmetric and skew symmetric matrix.



Watch Video Solution

180. Expressing the following matrices as the sum of a symmetric and skew symmetric matrix : $\begin{bmatrix} 2 & -2 & -4 \\ -1 & 3 & 4 \\ 1 & -2 & -3 \end{bmatrix}$



Watch Video Solution

181. Let $A = [325413067]$ Express A as sum of two matrices such that one is symmetric and the other is skew symmetric.



Watch Video Solution

182. Show that the following matrices are skew symmetric:

$$\begin{bmatrix} 0 & e & f \\ -e & 0 & g \\ -f & -g & 0 \end{bmatrix}$$



[Watch Video Solution](#)

183. Show that the following matrices are skew symmetric:

$$\begin{bmatrix} 0 & 1 & -1 \\ -1 & 0 & 1 \\ 1 & -1 & 0 \end{bmatrix}$$



[Watch Video Solution](#)

184. (i) Show that the matrix $A = \begin{bmatrix} 1 & -1 & 5 & -12 \\ 15 & 1 & -12 & 5 \\ 13 & 12 & 1 & -1 \\ 3 & 10 & 1 & 1 \end{bmatrix}$ is a symmetric matrix. (ii) Show that the matrix $A = \begin{bmatrix} 0 & 1 & -1 & -12 \\ 1 & 0 & 1 & -10 \\ 12 & 1 & 0 & 1 \\ 10 & 1 & 1 & 0 \end{bmatrix}$ is a skew symmetric matrix.



[Watch Video Solution](#)

185. Find the values of x, y, z if the matrix $A = \begin{bmatrix} 0 & 2y & z \\ x & y & -zx \\ y & -zx & yz \end{bmatrix}$ satisfy the equation $A' A = I$.



Watch Video Solution

186. If B is a square matrix and A is any square matrix of order equal to that of B , prove that $B' A B$ is symmetric or skew symmetric according as A is symmetric or skew symmetric.



Watch Video Solution

187. If A and B are symmetric matrices of the same order, show that $AB + BA$ is symmetric.



Watch Video Solution

188. Using elementary transformations, find the inverse of the matrix
 $[1327]$



Watch Video Solution

189. Using elementary transformations, find the inverse of the matrix
 $[2174]$



Watch Video Solution

190. Using elementary transformations, find the inverse of the matrix
 $[2111]$



Watch Video Solution

191. By using elementary operations, find the inverse of the matrix
$$A = \begin{bmatrix} 1 & 2 \\ 2 & -1 \end{bmatrix}.$$

[Watch Video Solution](#)

192. Using elementary transformations, find the inverse of the matrix
 $[2357]$

[Watch Video Solution](#)

193. Using elementary transformations, find the inverse of the matrix
 $[4534]$

[Watch Video Solution](#)

194. Using elementary transformations, find the inverse of the matrix
 $[3 - 1 - 42]$

[Watch Video Solution](#)

195. Using elementary transformations, find the inverse of the matrix

$$\begin{bmatrix} 1 & -123 \end{bmatrix}$$



Watch Video Solution

196. Using elementary transformations (operations), find the inverse of

the following matrices, if it exists: $\begin{bmatrix} 2 & -3 \\ -1 & 2 \end{bmatrix}$



Watch Video Solution

197. Using elementary transformation, find the inverse of the matrix

$$\begin{bmatrix} 10 & -2 \\ -5 & -1 \end{bmatrix}$$



Watch Video Solution

198. Using elementary transformations (operations), find the inverse of the

following matrices, if it exists: $\begin{bmatrix} 3 & 9 \\ 1 & 3 \end{bmatrix}$

[Watch Video Solution](#)

199. Using elementary transformations, find the inverse of the matrix :

$$\begin{bmatrix} 2 & 0 & -1 \\ 5 & 1 & 0 \\ 0 & 1 & 3 \end{bmatrix}$$

[Watch Video Solution](#)

200. Using elementary transformations, find the inverse of the matrix

$$\begin{bmatrix} 2 & -3 & 3 \\ 2 & 2 & 3 \\ 3 & -2 & 2 \end{bmatrix}$$

[Watch Video Solution](#)

201. Find the inverse the matrix (if it exists) given in $\begin{bmatrix} 2 & 1 & 3 & 4 \\ -1 & 0 & -7 & 2 \end{bmatrix}$

[Watch Video Solution](#)

202. Using elementary transformations (operations), find the inverse of

the following matrices, if it exists $\begin{bmatrix} 2 & -1 & 3 \\ -5 & 3 & 1 \\ -3 & 2 & 3 \end{bmatrix}$



Watch Video Solution