



BIOLOGY

AAKASH INSTITUTE ENGLISH

ANATOMY OF FLOWERING PLANTS

Examples

1. Name the primary meristems seen in flowering plants.



Watch Video Solution

2. What is the role of secondary meristems in the body of some plants?



Watch Video Solution

3. Give two characteristic features of parenchymatous cells.



Watch Video Solution

4. Collenchyma differs from sclerenchyma in

(i) Retaining protoplasm at maturity.

(ii) Having thick walls.

(iii) Being meristematic

(iv) Providing mechanical support to the plant.



Watch Video Solution

5. What is endarch condition? Where is it present?



Watch Video Solution

6. Name the vessel-less families of angiosperms.



Watch Video Solution

7. What are pit field? Where are they present?



Watch Video Solution

8. Which component of the phloem provides the mechanical support to the plants?



Watch Video Solution

9. Write two features of epidermal cells of plants.



Watch Video Solution

10. Define stomatal apparatus.



Watch Video Solution

11. What are the parts of cortex in a dicotyledon stem and a monocotyledon root?



Watch Video Solution

12. What is the role of vascular cambium in dicots?



Watch Video Solution

13. Name the part of general cortex in dicot root which is made up barrel shaped cells.



Watch Video Solution

14. Name one structure which is very distinct in roots but poorly developed or even absent in stems.



Watch Video Solution

15. What is the function/role of bulliform cells in grasses ?



Watch Video Solution

16. What is the difference in the appearance of spring and autumn wood?



Watch Video Solution

17. Discuss the differential activity of vascular cambium w.r.t secondary xylem and secondary pholem.



Watch Video Solution

18. What is the location of the oldest and youngest layers of secondary pholem in a dicot stem?



Watch Video Solution

19. What is the role of pericycle in secondary growth of dicot roots?



Watch Video Solution

Try Yourself

1. Why apical intercalary meristems are called primary meristems?



Watch Video Solution

2. The location of apical meristms is

(1) Lateral sides of roots and shoots

(2)At any part of plants

(3) At the tips of roots and shoots.

(4) Between apex and mature areas of roots and shoots.



Watch Video Solution

3. Give two exmaples of secondary meristems.



Watch Video Solution

4. Which of the following statements is incorrect for secondary meristems?

(1) They regenerate the parts of grasses damaged by grazers.

(2) They are the cylindrical meristems.

(3) They occur in the mature regions of roots and shoots.

(4) They appear later than primary meristems in the life of a plant.



Watch Video Solution

5. Why parenchyma is called a simple tissue?



Watch Video Solution

6. All of the following statements are correct for cells of parenchyma, except

A. They are generally isodiametric

B. They have lignified cell walls.

C. They may be spherical, oval, polygonal or elongated in shape.

D. They are the mature cells.

Answer: b



Watch Video Solution

7. Collenchyma tissue is a.....tissue.

- (1) Living and non-mechanical.
- (2) Dead and mechanical
- (3) Living and mechanical
- (4) Dead and non-mechanical



Watch Video Solution

8. Sclerenchyma tissue is atissue.

(1) Living and non-mechanical

(2) Dead and mechanical

(3) Living and mechanical

(4) Dead and non-mechanical.



Watch Video Solution

9. Lignin is deposited in a net-like manner in which of the following types of thickening in inner walls of tracheids?

(1) Annular

(2) Spiral

(3) Scalariform

(4) Reticulate



Watch Video Solution

10. Which of the following is a living component of xylem?

A. Xylem fibres

B. Tracheids

C. Vessels

D. Xylem parenchyma

Answer: d



Watch Video Solution

11. Xylem fibres posses obliterated central lumen due to very thick walls. This helps to perform the function of

A. Conduction

B. Storage of food

C. Mechanical support

D. Both (2) and (3)

Answer: c



Watch Video Solution

12. Tracheids are found in all the following plants, except

A. Pteridophytes

B. Gymnosperms

C. Bryophytes

D. Angiosperms

Answer: c



Watch Video Solution

13. The albuminous cells occur in the

A. Dicots

B. Monocots

C. Gymnosperms

D. All of these

Answer: c



Watch Video Solution

14. Which of the following component of phloem is made up of sclerenchymatous cells?

A. Phloem fibres

B. Sieve tubes

C. Companion cells

D. Pholem parenchyma

Answer: a



Watch Video Solution

15. Companion cells are seen associated with:

A. Vessels

B. Sieve tube elements

C. Tracheids

D. Pholem fibres

Answer: b



Watch Video Solution

16. Component of pholem which is absent most of the monocots is

A. Pholem parenchyma

B. Sieve pores

C. Sieve tube

D. Companion cell

Answer: a



Watch Video Solution

17. Which of the following statement is incorrect for the epidermis?

(1) It is the outermost layer of the primary plant body.

(2) It is often covered by cuticle externally

(3) It is usually multilayered.

(4) It is made up of completely arranged cells.



Watch Video Solution

18. Cuticle is absent in

A. Monocot leaf epidermis

B. Dicot leaf epidermis

C. Stem epidermis

D. Root epidermis

Answer: D



Watch Video Solution

19. Which of the following components of epidermal tissue system posses chloroplast?

- A. Trichomes
- B. Guard cells
- C. Subsidiary cells
- D. Cuticle

Answer: B



Watch Video Solution

20. All of the following statements are incorrect for guard cells, except

- A. They are bean-shaped in grasses
- B. Non green cells
- C. They do not enclose a stomatal pore

D. Their outer wall are thin and inner wall are thick

Answer: d



Watch Video Solution

21. Ground tissue system is not differential into which of the following components?

A. Epidermis

B. Hypodermis

C. Endodermis

D. Medulla

Answer: A



Watch Video Solution

22. Name the innermost layer of cortex that has barrel shaped cells.

A. Epidermis

B. Hypodermis

C. Endodermis

D. Pith

Answer: c



Watch Video Solution

23. Vascular bundles with cambium are called

A. Closed

B. Open

C. Exarch

D. Endarch

Answer: B



Watch Video Solution

24. In a stem, the vascular bundles and xylary elements areand respectively,

A. Radial, endarch

B. Radial, exarch

C. Conjoint, endarch

D. Conjoint, exarch

Answer: c



Watch Video Solution

25. Lateral roots arise from

A. cortex

B. Endodermis

C. Cork cambium

D. Pericycle

Answer: d



Watch Video Solution

26. Differentiate between the following

Dicot root and Monocot root

- A. Radial, exarch vascular bundles
- B. Unicelled root hairs
- C. Pericycle forms the lateral roots
- D. Well developed pith

Answer: D



Watch Video Solution

27. Vascular bundle is enclosed within a well developed sclerenchymatous sheath in



Watch Video Solution

28. Pith is well developed in

A. Dicot stem and monocot root

B. Monocot stem and dicot root

C. Monocot root and monocot stem

D. Dicot root and dicot stem

Answer: a



Watch Video Solution

29. The bulliform cells are present in the

A. Spongy parenchyma

B. Adaxial epidermis

C. Palisade parenchyma

D. Abaxial epidermis

Answer: b



Watch Video Solution

30. What is the true about a monocot leaf?

A. Presence of reticualte venation

B. Absence of bulliform cells.

C. Absence of vascular bundles

D. Absence of differential mesophyll

Answer: d



Watch Video Solution

31. The annual rings are bands of

A. Uniform environmental conditions

B. Non-uniform environmental conditions.

C. Absence of secondary growth

D. Absence of primary growth

Answer: B



Watch Video Solution

32. Which of the following is not true about early wood?

- A. a) It is formed during spring season
- B. b) It has a lower density than late wood
- C. c) Its vessels have wider lumen

D. d) It is produced when activity of cambium is lesser.

Answer: d



Watch Video Solution

33. Name the meristematic tissue present in the cortex of the stem. Which secondary tissues does it give rise to ?

A. Intercalary meristem

B. Vascular cambium

C. Interfascicular cambium

D. Cork cambium

Answer: a



Watch Video Solution

34. Vascular cambium ring is constituted by

A. Interfascicular and cork cambium

B. Intrafascicular and cork cambium

C. Intrafascicular and interfascicular
cambium

D. Only cork cambium

Answer: C



Watch Video Solution

35. In a dicot stem, youngest layer of
secondary xylem is situated

A. In between pith and primary xylem

B. Just outside the vascular cambium

C. Just on inner side of the vascular
cambium

D. Just inside the phellogen

Answer: C



Watch Video Solution

36. Which of the following is called early wood?

A. Spring wood

B. Autumn wood

C. Primary xylem

D. Secondary phloem

Answer: A



Watch Video Solution

37. In a dicotyledonous stem, the sequences of tissues from the outside to the inside is

A. Phellem, phellogen- endodermis-
secondary cortex.

B. Phellem-phellogen- phloem- xylem

C. Phellem-cork cambium -secondary
pholem- phelloderm

D. Pericycle-phellem- pholem-endodermis

Answer: b



Watch Video Solution

38. The cambium ring of roots resembles the cambium ring of stem in

- A. Its mode of function
- B. Its origin from ground tissue
- C. Its wavy outline
- D. All of these

Answer: a



Watch Video Solution

Exercise

1. Intercalary meristem is related to all, except
- A. Present between permanent cells
 - B. Primary meristem
 - C. Increasing the growth of axis
 - D. Regenerates part of grasses removed by grazing herbivores.

Answer: C



Watch Video Solution

2. Choose odd one out w.r.t. origin

A. Interfascicular cambium

B. Intercalary meristem

C. Apical meristem

D. Intrafascicular cambium

Answer: A



Watch Video Solution

3. Find odd one out w.r.t histogens found in shoot apex according to Hanstein

A. Tunica

B. Periblem

C. Plerome

D. Dermatogen

Answer: A



Watch Video Solution

4. Select an incorrect match

A. Tunica-Corpus theory- Schmidt

B. Histrogen theory- Hanstein

C. Quiescent center theory - Clowes

D. Korper -Kappe theory- Hofmeister

Answer: D



Watch Video Solution

5. Thickening material present in wall of collenchyma is

- A. Pectin, cellulose, hemicellulose
- B. Lignin, cellulose, hemicellulose
- C. Hemicellulose, suberin, cellulose
- D. Suberin, pectin, cellulose

Answer: A



Watch Video Solution

6. Sclereids possess

- A. Tapering ends
- B. Highly thickened wall
- C. Hemicellulosic wall
- D. Broad lumen

Answer: B



Watch Video Solution

7. Major xylary element in wood of a gymnospermic plant is

A. Vessel

B. Tracheid

C. Xylem fibre

D. Xylem parenchyma

Answer: B



Watch Video Solution

8. Which of the following cells helps in maintaining the pressure gradient in the sieve tubes?

A. Pholem parenchyma

B. Bast fibre

C. Companion cells

D. Wood fibre

Answer: C



Watch Video Solution

9. Which component is not found in phloem of angiosperms?

A. Albuminous cells

B. Sieve tube

C. Companion cells

D. Bast fibre

Answer: A



Watch Video Solution

10. Sclerenchyma fibres are absent in

A. Protoxylem

B. Protophloem

C. Metaxylem

D. More than one option is correct

Answer: D



Watch Video Solution

11. Green cells of epidermis are

- A. Bulliform cells
- B. Subsidiary cells
- C. motor cells
- D. guard cells

Answer: D



Watch Video Solution

12. Trichomes

- A. Are not hair
- B. May be branched
- C. Are usually single celled
- D. Are always non-secondary

Answer: B



Watch Video Solution

13. Multilayered epidermis is found in the leaves of

A. Nerium and Ficus

B. Ficus and Vanda

C. Equisetum and Grasses

D. Vanda and Nerium

Answer: A



Watch Video Solution

14. Dicot root is similar in all the given characters to monocot root, except.



Watch Video Solution

15. The casparian rings are found in

A. Stem endodermis

B. Root endodermis

C. Pericycle

D. Outer cortex

Answer: B



Watch Video Solution

16. Starch grains are abundant in the endodermal cells of

- A. Monocot root
- B. Dicot stem
- C. Monocot stem
- D. Dicot root

Answer: B



Watch Video Solution

17. Stem of barley shows

- A. Presence of collenchyma in hypodermis
- B. Scattered vascular bundles
- C. Presence of parenchymatous pericytes
- D. Presence of parenchymatous pericytes.

Answer: B



Watch Video Solution

18. (a) Palisade tissue is present towards upper epidermis in monocot leaves.

(b) Lower layer of mesophyll cells is loosely packed with few chloroplasts in dicot leaves.

(c) Dicot leaves have conjoint, collateral and closed vascular bundles.

A. All are incorrect

B. Only b is correct

C. Only a is incorrect

D. Only c is incorrect

Answer: C



Watch Video Solution

19. Conjoint, collateral, endarch and closed vascular bundles are found in

A. Monocot root

B. Monocot stem

C. Dicot root

D. Dicot stem

Answer: B



Watch Video Solution

20. Bulliform cells are

A. Empty

B. Colourless

C. Found in grass leaves

D. All of these

Answer: D



Watch Video Solution

21. Cambium ring in dicot system is

- A. Primary meristem in origin
- B. Secondary meristem in origin
- C. Promeristem in origin
- D. Both primary and secondary meristem in origin

Answer: D



Watch Video Solution

22. Dendrochronology deals with the study of

- A. Phylogeny
- B. Numerical taxonomy
- C. Age of trees
- D. Grasses

Answer: C



Watch Video Solution

23. Heartwood is characterised by all, except

- A. Presence of tyloses
- B. Presence of tannins, resins, oils, gums
etc.
- C. Its commercial importance for timber
- D. Active in water conduction

Answer: D



Watch Video Solution

24. Sapwood is

- A. Secondary xylem
- B. Secondary phloem
- C. Phellem
- D. Secondary cortex

Answer: A



Watch Video Solution

25. Peripheral region of secondary xylem is

- A. Light and functional
- B. Dark and non-functional
- C. Hard and durable
- D. Resistant and light

Answer: A



Watch Video Solution

26. Bark includes

- A. Periderm + Secondary phloem
- B. Periderm + primary xylem
- C. Secondary phloem + xylem
- D. Secondary xylem + Cork cambium

Answer: B



Watch Video Solution

27. Which of the following tissue makes phellogen during the secondary growth in dicot roots?

A. Endodermis

B. Hypodermis

C. Epidermis

D. Periderm

Answer: D



Watch Video Solution

28. Which of the following structures is not formed by activity of cork cambium?

A. Phellem

B. Phelloderm

C. Secondary xylem

D. Corky layer

Answer: C



Watch Video Solution

29. Choose correct option w.r.t. Lenticels

- A. Lens shaped
- B. Permit gaseous exchange
- C. Occur in most woody trees
- D. More than one option is correct

Answer: D



Watch Video Solution

30. Choose correct option w.r.t. origin of cork cambium in dicot stem and root

A. Completely primary in both

B. Completely secondary in both

C. Partly primary and partly secondary in
both

D. Primary in stem and secondary in root

Answer: B



Watch Video Solution

31. Stomata are the component of

A. Endodermis

B. Hypodermis

C. Epidermis

D. Periderm

Answer: D



Watch Video Solution

32. Specialised epidermal cells surrounding the guard cell are called

A. Phellem

B. Phelloderm

C. Secondary xylem

D. Corky layer

Answer: C



Watch Video Solution

Assignment Section A Objective Type Questions

1. Which of the following tissue makes phellogen during the secondary growth in dicot roots?

- A. Epidermal tissue system
- B. Ground tissue system
- C. Conducting tissue system
- D. Vascular tissue system

Answer: A



[Watch Video Solution](#)

2. Which of the following structures is not formed by activity of cork cambium?

A. Bulliform cells

B. Companion cells

C. Subsidiary cells

D. Endodermal cells

Answer: C



[Watch Video Solution](#)

3. Tissues are

- A. Groups of cells which are similar in origin and function
- B. Groups of organs which are similar in origin and function
- C. Cells which are similar in function but not in origin.

D. Groups of cells which are not similar in origin and function.

Answer: A



Watch Video Solution

4. Plant tissues are divided into meristematic and permanent tissues on which of the following basis?

A. Whether the plant is a dicot or a monocot

B. Whether the cells being formed are capable of dividing or not

C. Position

D. Origin

Answer: B



Watch Video Solution

5. The axillary bud is constituted by the cells 'left behind' from-

- A. Root apical meristem
- B. Shoot apical meristem
- C. Intercalary meristem
- D. Lateral meristem

Answer: B



Watch Video Solution

6. The woody axis of flowering plants is produced by

- A. Apical meristem
- B. Primary meristem
- C. Intercalary meristem
- D. Secondary meristem

Answer: D



Watch Video Solution

7. All of the following are lateral meristems, except

- A. Intercalary meristem
- B. Fascicular vascular cambium
- C. Interfascicular cambium
- D. Cork cambium

Answer: A



Watch Video Solution

8. What is the importance of lateral meristematic region?

- A. It gives rise to the lateral branches
- B. It increases girth of the plant axis
- C. It increases girth as well as length of the
plant axis
- D. It increases only length of the plant axis

Answer: B



Watch Video Solution

9. The parenchymatous cells are

A. Dead

B. Thick-walled

C. Thin-walled

D. Thick-walled and dead

Answer: C



Watch Video Solution

10. Collenchyma differs from parenchyma in having

(a) Living protoplasm

(b) Cellulose walls

(c) Vacuoles

(d) Pectin and cellulose deposits at corners

A. Possessing thick cell wall

B. Lacking protoplasm

C. Containing chloroplasts usually

D. Being meristematic

Answer: A



Watch Video Solution

11. Which among the following are absent in the collenchyma?

- A. Chloroplasts
- B. Vacuoles
- C. Intercellular spaces
- D. Pectin deposition

Answer: C



Watch Video Solution

12. The elongated, thick-walled and tapering cells are

- A. Parenchymatous
- B. Collenchymatous
- C. Chlorenchymatous
- D. Sclerenchymatous

Answer: D



Watch Video Solution

13. Sclereids are commonly found in

- A. Young stems and petioles of leaves
- B. Fruit walls of nuts
- C. Roots
- D. Fleshy stems

Answer: B



Watch Video Solution

14. What is the function of vessels in flowering plants?

- A. Transport of food
- B. To get rid of excess water
- C. Photosynthesis
- D. Transport of water and minerals

Answer: D



15. Which of the following statements is correct?

A. Angiosperms lack vessels in their xylem

B. The presence of vessels is a characteristic feature of angiosperms

C. The cells of vessels are living

D. Vessels is a long cylindrical tube-like cells made up of many vessel members

Answer: B



Watch Video Solution

16. The central lumens are obliterated in

- A. Xylem fibres
- B. Pholem parenchyma
- C. Xylem parenchyma
- D. Sieve tubes

Answer: A



Watch Video Solution

17. Which of the following is true for endarch type of primary xylem?

A. Protoxylem lies towards the periphery of the organ

B. Metaxylem lies towards the periphery of the organ

C. Metaxylem lies towards the pith of the organ

D. Protophloem lies towards the path of
the organ

Answer: B



Watch Video Solution

18. Which of the following is absent in most of
the monocotyledons?

A. a) Phloem parenchyma

B. b) Tracheids

C. c) Vessels

D. d) Xylem parenchyma

Answer: A



Watch Video Solution

19. Stomata are the component of

A. Epidermal tissue system

B. Ground tissue system

C. Conducting tissue system

D. Vascular tissue system

Answer: A



Watch Video Solution

20. Specialised epidermal cells surrounding the guard cell are called

A. Bulliform cells

B. Companion cells

C. Subsidiary cells

D. Endodermal cells

Answer: C



Watch Video Solution

21. Root hairs are the

A. Multicellular elongations of epidermal cells

B. Acellular elongations of epidermal cells

C. Unicellular elongations of epidermal cells

D. Multicellular elongations of endodermal cells.

Answer: C



Watch Video Solution

22. The distinguishing feature of open vascular bundles is presence of

A. a) Cambium

B. b) Xylem and phloem

C. c) Pericycle

D. d) Endodermis

Answer: A



Watch Video Solution

23. Which type of arrangement of vascular bundles occurs in the roots of monocots?

A. Conjoint open

B. Radial

C. Conjoint closed

D. Bicollateral

Answer: B



Watch Video Solution

24. The presence of cambium in the vascular bundles provides them the ability to

A. Radially transport the food

B. Form secondary tissues

C. Prevent water loss due to transpiration

D. Conduct photosynthesis

Answer: B



Watch Video Solution

25. Lateral roots arise from

A. Endodermis

B. Pericycle

C. Conjunctive tissue

D. Cambium ring

Answer: B



Watch Video Solution

26. Polyarch xylem bundles are found in

A. Monocot root

B. Dicot root

C. Monocot stem

D. Dicot stem

Answer: A



Watch Video Solution

27. The casparian strips are present on the plant cells of roots which are

A. Bean-shaped

B. Dumb-bell shaped

C. Barrel-shaped

D. Lens-shaped

Answer: C



Watch Video Solution

28. The conjunctive tissue lies between the

A. a) Xylem and phloem

B. b) Pericycle and endodermis

C. c) Epidermis and cortex

D. d) Epidermis and hypodermis

Answer: A



Watch Video Solution

29. In which of the following characters, a monocot root differs from a dicot root?

A. Radial vascular bundles

B. Large pith

C. Conjunctive tissue in between xylem and
pholem

D. Single layered endodermis

Answer: B



Watch Video Solution

30. Match the following

Column-I	Column-II
a. Hypodermis in dicot stem	(i) Absent
b. Pericycle in dicot stem	(ii) Parenchymatous
c. Ground tissue in monocot stem	(iii) Collenchymatous
d. Phloem parenchyma in monocot stem	(iv) Sclerenchymatous

A. a(iv), b(i), c(iii), d(ii)

B. a(i), b(ii), c(iv), d(iii)

C. a(iii), b(iv), c(ii), d(i)

D. a(ii), b(ii), c(i), d(iv)

Answer: C



[Watch Video Solution](#)

31. Vascular bundles surrounded by a sclerenchymatous bundle sheath is a feature of

- A. Dicot root
- B. Monocot root
- C. Dicot root
- D. Monocot stem

Answer: D



Watch Video Solution

32. The central most portion of stem of dicotyledonous plants is occupied by

A. Vascular bundles

B. Pericycle

C. Pith

D. Cortex

Answer: C



33. Which of the following is not true for the vascular bundles of monocotyledonous stems?

- A. Scattered in the ground tissue
- B. Possess water-containing cavities
- C. Ring' arrangement
- D. Conjoint and closed

Answer: C



34. The epidermis in a dorsiventral leaf

(a) Covers both adaxial and abaxial surfaces

(b) Is not covered by cuticle

(c) Bears more stomata on the upper side

(d) May even lack stomata on the upper side

Which of the above statements are correct?

A. (a) and (c)

B. (b) and (d)

C. (a) and (d)

D. (b) and (c)

Answer: C



Watch Video Solution

35. Choose correct option w.r.t. spongy mesophyll in dicot leaf.

A. Numerous large spaces and air cavities between its cells.

B. A large number of chloroplasts

C. Present on the adaxial surface

D. Vertical and parallel arrangemetn of cells

Answer: A



Watch Video Solution

36. The size of vascular bundles in a dorsiventral leaf is dependent on

A. Size of lamina

B. Size of veins

C. Number of stomata

D. Number of veins

Answer: B



Watch Video Solution

37. The stomata in an isobilateral leaf

A. Are present only on the adaxial epidermis

B. Are present only on the abaxial epidermis

C. Are absent on both the surfaces of the epidermis

D. Are present on both the surfaces of the epidermis

Answer: D



Watch Video Solution

38. During water stress, the bulliform cells

(a) Become turgid

(b) Become flaccid

(c) Make the leaves curl inwards

(d) Make the leaf surface exposed

The correct options are

A. (a) and (c)

B. (b) and (d)

C. (a) and (d)

D. (b) and (c)

Answer: D



Watch Video Solution

39. The tissue which participate in the secondary growth is

A. a) Lateral meristem

B. b) Apical meristem

C. c) Intercalary meristem

D. d) Primary meristem

Answer: A



Watch Video Solution

40. The infrascicular cambium

- A. Is a simple permanent tissues
- B. Is a meristematic tissue
- C. Is a complex permanent tissues
- D. Is secondary meristern

Answer: B



Watch Video Solution

41. The cambial ring is generally (Option 1)
More active on the inner side than on the
outer (Option 2) More active on the outer
side than the inner (Option 3) Equally active
towards both sides (Option 4) Equally inactive
towards both sides

A. a) More active on the inner side than on
the outer

B. b) More active on the outer side than
the inner

C. c) Equally active towards both sides

D. d) Equally inactive towards both sides

Answer: A



Watch Video Solution

42. In a dicot stem, the interfascicular cambium strip arises

A. a) Between xylem and phloem

B. b) From medullary rays

C. c) From pith

D. d) From pericycle

Answer: B



Watch Video Solution

43. Wood is

A. Primary xylem

B. Primary phloem

C. Secondary xylem

D. Secondary phloem

Answer: C



Watch Video Solution

44. An annual ring is formed by

A. Two consecutive rings of spring wood

B. Two alternate rings of spring wood and autumn wood.

C. Two consecutive rings of autumn wood

D. Two alternate rings of sapwood and heartwood

Answer: B



Watch Video Solution

45. Growth rings are well marked in trees growing in

A. Mumbai

B. Chennai

C. Shimla

D. Kolkata

Answer: C



Watch Video Solution

46. Which of the following is the function of heartwood?

A. a) Mechanical support

B. b) Radial conduction of water and minerals

C. c) Lateral conduction of food

D. d) Minimise water loss in water stress

Answer: A



Watch Video Solution

47. The heartwood cannot conduct water because of

- A. Peripheral locations in the stems
- B. Suberized cell walls
- C. Central location in the stems
- D. Deposition of organic compounds

Answer: D



Watch Video Solution

48. Besides phellem, phellogen, root periderm comprises of

A. Secondary cortex

B. Phelloderm

C. Bark

D. More than one option is correct

Answer: B



Watch Video Solution

49. The feature which is common to both heartwood and sapwood is

- A. Both are the regions of secondary xylem
- B. Both are involved in the conduction of water
- C. Both comprise dead elements with accumulation of aromatic compounds
- D. Both are located in the central layers of the stem.

Answer: A



Watch Video Solution

50. In a plant organ which is covered by periderm and in which the stomata are absent, some gaseous exchange still takes place through

- A. Bulliform cells
- B. Lenticles
- C. Pneumatophores
- D. Trichomes

Answer: B



Watch Video Solution

51. Bark refers to all tissues exterior to the

- A. cork cambium
- B. Pericycle
- C. Vascular cambium
- D. Periderm

Answer: C



Watch Video Solution

52. Which of the following is true for the dicot roots?

A. Both vascular and cork cambium are responsible for the secondary growth

B. Vascular cambium arises during the primary growth

C. They do not show secondary growth just like monocot roots

D. Cork cambium develops during the primary growth

Answer: A



Watch Video Solution

Assignment Section B Objective Type Questions

1. Meristem is characterized by

A. Isodiametric cells with cellulosic thin wall

B. Absence of intercellular space and vacuole

C. Absence of reserve food materials and plastids

D. All of these

Answer: D



Watch Video Solution

2. According to histogen theory, stem epidermis is derived from

A. Calyptrogen

B. Dermatogen

C. Protoderm

D. Periblem

Answer: B



Watch Video Solution

3. (A) According to Clowes roots apex consists of an inverted cup like structure.

(B) Low amount of RNA, DNA and protein is characteristic of waiting meristem.

(c) Cells of reserve meristem can divide only when the root apex gets injured.

A. All are correct

B. A and C are incorrect

C. A and B are incorrect

D. B and C are incorrect

Answer: A



Watch Video Solution

4. Collenchyma is a type of mechanical tissue but it is not as efficient as sclerenchyma.

However, it has certain advantages like

(a) It offers no resistance to the growing organs

(b) It has the power of growth

(c) It is flexible

(d) Through it has the power of growth, it

offers no resistance to the growing organs
and it is flexible

A. it offers no resistance to the growing
organs

B. it has no cellulose In the cell wall

C. it is flexible

D. It has the power of growth, it offers no
resistance to the growing organs and is
not flexible

Answer: C



Watch Video Solution

5. Isodiametric sclereids, found in hard endocarp of coconut and fleshy portion of some fruits are

A. Brachysclereids

B. Asterosclereids

C. Osteosclereids

D. Trichosclereids

Answer: A



Watch Video Solution

6. Main water-conducting element of xylem in homoxylous plants is

- (a) Trachea
- (b) Vessel
- (c) Tracheid
- (d) Xylem parenchyma

A. Tracheary element

B. Vessel

C. Tracheid

D. Xylem parenchyma

Answer: C



Watch Video Solution

7. In trees, the death of protoplasm is essential for a vital function such as

- A. Food transport
- B. Water transport
- C. Both (1) and (2)

D. Stomatal movements

Answer: B



Watch Video Solution

8. Find set of cells connected by pit fields between their common longitudinal walls

A. Companion cell and phloem fibres

B. Companion cell and sieve tube

C. Sieve cell and albuminous cell

D. Sieve tube and phloem fibre

Answer: B



Watch Video Solution

9. In old sieve tubes at the end of growing season, which of the following gets deposited over sieve plate to regulate sugar transport :

A. P-protein

B. Callose

C. Lignin

D. ABA

Answer: B



Watch Video Solution

10. Parenchymatous structure with intercellular spaces is

A. Epidermis

B. Endodermis

C. Cortex

D. Pericycle

Answer: C



Watch Video Solution

11. Select correct features w.r.t. trichomes in shoot system

(a) Usually unicelled

(b) Branched or unbranched

(c) May be secretory

(d) Soft or stiff

(e) Helps against transpiration

A. a,b,d and e

B. All except c and d

C. All except a

D. a, c and e

Answer: C



Watch Video Solution

12. The pericycle of roots is never sclerenchymatous because it

A. Does not act as a mechanical tissue in roots

B. Is the place of origin of root branches

C. Gives rise to root hairs

D. Gives rise to root hairs (when the root is young), and to root branches (at maturity)

Answer: B



Watch Video Solution

13. Tissue commonly known as the passport point or biological check post is characterized by

- A. Bulliform cells and raphides
- B. Cystolith and motor cells
- C. Casparian bands and passage cells
- D. Passage cells and fats

Answer: C



Watch Video Solution

14. Seat of origin of lateral root and formation of cork cambium are features related to



Watch Video Solution

15. Centripetal and centrifugal xylems are important features of

(a) Root and stem, respectively

(b) Exarch and endarch, respectively

(c) Endarch and exarch, respectively

(d) Both a and b

A. Root and stem respectively

B. Exarch and endarch respectively

C. Endarch and exarch respectively

D. Both (1) and (2)

Answer: D



Watch Video Solution

16. Memembers of Winteraceae, Tetracentraceae and trochodendraceae

- A. Do not have tracheids
- B. Do not have albuminous cells
- C. Do not have vessels
- D. More than one option is correct

Answer: D



Watch Video Solution

17. Which of the following vascular bundles are always open?

- A. Radial
- B. Collateral
- C. Bicollateral
- D. Concentric

Answer: C



Watch Video Solution

18. The vascular bundles in the stems of several dicots are conjoint, collateral, and open. In each of these bundles,

A. Xylem and phloem are on the same radius with phloem towards the pith and xylem towards the pericycle without a strip of cambium between them

B. Xylem and phloem are on the same radius with xylem situated towards the

pericycle and a strip of cambium separates the two

C. Xylem completely surrounds the phloem on all sides but the two are separated by the cambium.

D. Phloem completely surrounds the xylem and a strip of cambium separates the two

Answer: B



Watch Video Solution

19. Which is not true for monocot stem ?

A. Sclerenchymatous hypodermis

B. Presence of water canals in pitch

C. Conjoint, collateral closed vascular
bundles

D. Presence of bundle sheath

Answer: B



Watch Video Solution

20. Vascular bundles are conjoint, collateral, endarch and lack cambium between xylem and phloem in all, but not in

A. Maize

B. Barley

C. Wheat

D. Sunflower

Answer: D



Watch Video Solution

21. Select a set having correct match

Dicot stem	Monocot stem
(1) Sclerenchymatous hypodermis	Collenchymatous hypodermis
(2) Parenchymatous pericycle	Sclerenchymatous pericycle
(3) Epidermis with trichomes	Water containing cavities in vascular bundles
(4) Oval bundles	Wedge shaped bundles



Watch Video Solution

22. State true or false. Monocot stem differs from dicot stem.



Watch Video Solution

23. Vascular cambium is a meristematic layer that cuts off

- A. Primary xylem and primary phloem
- B. Xylem vessels and xylem tracheids
- C. Primary xylem and secondary xylem
- D. Secondary xylem, secondary phloem and medullary rays

Answer: D





[Watch Video Solution](#)

24. Derivatives of the secondary meristem in the steler region are

- A. Phellem and phelloderm
- B. Alburnum and primary pholem
- C. Duramen and alburnum
- D. Primary xylem and secondary pholem

Answer: C



[Watch Video Solution](#)

25. What is the position of oldest secondary phloem ?

- A. Just outside the pericycle
- B. Just outside the vascular cambium
- C. Just below the pericycle
- D. Below the vascular cambium

Answer: C



Watch Video Solution

26. In a mature dicot stem which has undergone secondary growth, youngest layer of secondary xylem is situated

- A. Between pith and primary xylem
- B. Just outside vascular cambium
- C. Just inside vascular cambium
- D. Just inside cork cambium

Answer: C



Watch Video Solution

27. (A) heart wood is durable, dark and central in position.

(B) Tyloses are balloon like structures of xylem parenchyma in vessel lumen.

(C) Late wood is formed during spring season.

A. All are correct

B. Only A is correct

C. Only B is incorrect

D. Only C is incorrect

Answer: D



Watch Video Solution

28. Secondary growth in extrasteler region is due to activity of

- A. Interfascicular cambium
- B. Intrafascicular cambium
- C. Cork cambium
- D. Intercalary meristem

Answer: C



29. Seasonal activity of vascular cambium is influenced by many factors, except

- A. Geographical location of plant
- B. Relative humidity and temperature
- C. Photoperiod and water supply
- D. Leaf orientation

Answer: D



30. When secondary growth is initiated in dicot stem, what will happen first?

A. The cells of cambium divide periclinally
to form xylem mother cells

B. Interfascicular cambium join with
intrafascicular cambium

C. Parenchymatous cells present between
vascular bundles become meristematic

D. Pith get obliterated

Answer: C



Watch Video Solution

31. All given tissues are formed as a result of redifferentiation process, except

A. Phellum

B. Phelloderm

C. Secondary xylem

D. Interfascicular cambium

Answer: D



Watch Video Solution

32. Vascular cambium of dicot root is purely secondary in origin and arise from

A. Cells of conjunctive parenchyma just below phloem

B. Cells of pericycle just outside protoxylem

C. Cells of parenchyma between xylem and phloem

D. More than one option is correct

Answer: D



Watch Video Solution

33. Select wrong statement regarding secondary phloem

A. Arise due to activity of procambium

B. Occurs during secondary growth

C. No distinction between protophloem
and metaphloem

D. Secondary permanent tissue

Answer: A



Watch Video Solution

34. Vascular bundles in dicot leaves are

A. Scattered, conjoint, collateral, open

B. Scattered , conjoint, collateral, closed

C. Scattered, conjoint, collateral, open

D. Ringed, conjoint, collateral, closed

Answer: B



Watch Video Solution

35. Lenticles are formed due to rupture in epidermis because of pressure exerted by

A. Epiderm cells

B. Tyloses

C. Complementary cells

D. Phellem

Answer: C



Watch Video Solution

Section C Previous Year Questions

1. Identify the wrong statement in context of heartwood.

A. Organic compounds are deposited in it

B. It is highly durable

C. It conducts water and minerals
efficiently

D. It comprises dead elements with highly
lignified walls

Answer: C



Watch Video Solution

2. Which of the following is made up of dead cells ?

A. Xylem parenchyma

B. Collenchyma

C. Phellem

D. Pholem

Answer: C



Watch Video Solution

3. The vascular cambium normally gives rise to

- A. Phelloderm
- B. Primary pholem
- C. Secondary xylem
- D. Periderm

Answer: C



Watch Video Solution

4. Cortex is the region found between

A. Epidermis and sale

B. Pericycle and endodermis

C. Endodermis and pith

D. Endodermis and vascular bundle

Answer: A



Watch Video Solution

5. The ballone-shaped structures called tyloses

A. Originate in the lumen of vessels

B. Characterize the sapwood

C. Are extensions of xylem parenchyma cells into vessels.

D. Are linked to the ascent of sap through xylem vessels.

Answer: C



Watch Video Solution

6. Specialised epidermal cells surrounding the guard cell are called

- A. Lenticels
- B. Complementary cells
- C. Subsidiary cells
- D. Bulliform cells

Answer: C



Watch Video Solution

7. Read the different components from (a) to (d) in the list given below and tell the correct order of the components with reference to their arrangement from outer side to inner side in a woody dicot stem:

(a) Secondary cortex

(b) Wood

(c) Secondary phloem

(d) Phellem The correct order is :

A. Secondary cortex

B. Wood

C. Secondary phloem + xylem

D. Phellem

Answer: D



Watch Video Solution

8. Vascular bundles in monocotyledons are considered closed because

A. Xylem is surrounded all around by phloem

B. A bundle sheath surrounds each bundle

C. Cambium is absent

D. There are no vessels with perforations

Answer: C



Watch Video Solution

9. A major characteristic of the monocot root is the presence of

- A. Cambium sandwiched between phloem and xylem along the radius
- B. Open vascular bundles
- C. Scattered vascular bundles
- D. Vasculature without cambium

Answer: D



Watch Video Solution

10. You are given a fairly old piece of dicot stem and root. Which of the following anatomical structures will you use to distinguish between the two?

- A. Secondary xylem
- B. Secondary phloem
- C. Protoxylem
- D. Cortical cells

Answer: C



Watch Video Solution

11. Tracheids differ from other tracheary elements in

- A. Having casparian strips
- B. Being imperforate
- C. Lacking nucleus
- D. Being lignified

Answer: B



Watch Video Solution

12. Age of tree can be estimated by

A. Biomass

B. Number of annual rings

C. Diameter of its heartwood

D. Its height and girth

Answer: B



Watch Video Solution

13. Interfascicular cambium develops from the cells of

A. Xylem parenchyma

B. Endodermis

C. Pericycle

D. Medullary rays

Answer: D



Watch Video Solution

14. Water containing cavities in vascular bundles are found in

A. Cycas

B. Pinus

C. Sunflower

D. Maize

Answer: D



Watch Video Solution

15. Closed vascular bundles lack

A. Cambium

B. Pith

C. Ground tissue

D. Conjunctive tissue

Answer: A



Watch Video Solution

16. Companion cells are closely associated with

- A. Trichomes
- B. Guard cells
- C. Sieve elements
- D. Vessel elements

Answer: C



Watch Video Solution

17. cork/bottle cork is formed from

A. Xylem

B. Vascular cambium

C. Dermatogen

D. Phellogen

Answer: D



Watch Video Solution

18. As compared to a dicot, root, a monocot root has

- A. More abundant secondary xylem
- B. Many xylem bundles
- C. Inconspicuous annual rings
- D. Relatively thicker periderm

Answer: B



Watch Video Solution

19. Function of companion cells is

- A. Loading of sucrose into sieve elements by passive transport
- B. Loading of sucrose into sieve elements
- C. Providing energy to sieve elements for active transport
- D. Providing water to phloem

Answer: B



Watch Video Solution

20. Ground tissue includes

- A. All tissues internal to endodermis
- B. All tissues external to endodermis
- C. All tissues except epidermis and vascular bundles
- D. Epidermis and cortex

Answer: C



Watch Video Solution

21. In land plants, the guard cells differ from other epidermal cells in having

A. Chloroplasts

B. Cytoskeleton

C. Mitochondria

D. Endoplasmic reticulum

Answer: A



Watch Video Solution

22. The cork cambium, cork and secondary cortex are collectively called

A. Phellem

B. Phelloderm

C. Phellogen

D. Peridem

Answer: D



Watch Video Solution

23. Which one of the following is wrongly matched

A. Cassia-Imbricate aestivation

B. Root pressure- Guttation

C. Puccinia-Smut

D. Root - Exarch protoxylem

Answer: C



Watch Video Solution

24. Some vascular bundles are described as open because these

- A. Possess conjunctive tissue between xylem and phloem
- B. Are not surrounded by pericycle
- C. Are surrounded by pericycle but no endodermis
- D. Are capable of producing secondary xylem and phloem

Answer: D



Watch Video Solution

25. The chief water conducting elements of xylem in gymnosperms are

- A. Tracheids
- B. Vessels
- C. Fibers
- D. Transfusion tissue

Answer: A



Watch Video Solution

26. Which one of the following is not a lateral meristem ?

- A. Intercalary meristem
- B. Intrafascicular cambium
- C. Interfascicular cambium
- D. Phellogen

Answer: A



Watch Video Solution

27. Heartwood differs form sapwood in

- A. Being susceptible to pests and pathogens
- B. Presence of rays and fibres
- C. Absence of vesseles and parenchyma

D. Having dead and non-conducting elements

Answer: D



Watch Video Solution

28. Transport of food material in higher plants takes place through

A. Companion cells

B. Transfusion tissue

C. Tracheids

D. Sieve elements

Answer: D



Watch Video Solution

29. Anatomically fairly old dicotyledonous root is distinguished from the dicotyledonous stem by

A. Absence of secondary phloem

B. Presence of cortex

C. Position of protoxylem

D. Absence of secondary xylem

Answer: C



Watch Video Solution

30. The annular and spirally thickened conducting elements generally develop in the protoxylem when the root or stem is

A. Elongating

B. Widening

C. Differentiating

D. Maturing

Answer: D



Watch Video Solution

31. Palisade parenchyma is absent in leaves of:

A. Mustard

B. Soybean

C. Grain

D. Sorghum

Answer: D



Watch Video Solution

32. In barely vascular bundles are

A. Closed and scattered

B. Open and in ring

C. Closed and radial

D. Open and scattered

Answer: A



Watch Video Solution

33. Reduction in vascular tissue, mechanical tissue and cuticle is characteristic of

A. Mesophytes

B. Epiphytes

C. Hydrophytes

D. Xerophytes

Answer: C



Watch Video Solution

34. Vascular tissues in flowering plants develop from

A. Dermatogen

B. Phellogen

C. Plermoe

D. Perjiblem

Answer: C



Watch Video Solution

35. The length of different internodes in a culm of sugarcane is variable because of

A. Intercalary meristem

B. Shoot apical meristem

C. Position of axillary buds

D. Size of leaf lamina at the node below
each internode

Answer: A



Watch Video Solution

36. For a critical study of secondary growth in plants, which one or the following pairs of plants is suitable ?

A. Wheat and maiden hair fern

B. Sugarcane and sunflower

C. Teak and pine

D. Deodar and fern

Answer: C



Watch Video Solution

37. Passage cells are walled cells found in

A. Central region of style through which the pollen tube grows towards the ovary.

B. Endodermis of roots facilitating rapid transport of water from cortex to pericycle .

C. Phloem elements that serve as entry points for substances for transport to other plant parts.

D. Testa of seed to enable emergence of growing embryonic axis during seed

germination.

Answer: B



Watch Video Solution

38. A common structural feature of vessel elements and sieve tube elements is

- A. Thick secondary walls
- B. Pores on lateral walls
- C. Presence of P-protein

D. Enucleate condition

Answer: D



Watch Video Solution

39. In a woody dicotyledonous tree, which of the following parts will mainly consist of primary tissue?

A. Stem and root

B. All parts

C. Shoot tips and root tips

D. Flowers, fruits and leaves

Answer: D



Watch Video Solution

40. Meristematic tissue responsible for increase in girth of tree trunk is

A. Apical meristem

B. Intercalary meristem

C. Lateral meristem

D. Phellogen

Answer: C



Watch Video Solution

41. Axillary bud an terminal bud are derived from the activity of

A. Lateral meristem

B. Intercalary meristem

C. Apical meristem

D. Parenchyma

Answer: C



Watch Video Solution

42. Procambium is situated just behind apical meristem. Procambium gives rise to

A. Primary vascular bundle

B. Fascicular cambium

C. Cork cambium

D. Both (1) and (2)

Answer: D



Watch Video Solution

43. The cells of the quiescent centre are characterised by

A. Having dense cytoplasm and prominent nuclei

- B. Having light cytoplasm and small nuclei
- C. Dividing regularly to add to the corpus
- D. Dividing regularly to add to tunica

Answer: B



Watch Video Solution

44. The apical meristem of the root is present

- A. Only in radicals
- B. Only in tap roots

C. Only in adventitious roots

D. In all the roots

Answer: D



Watch Video Solution

45. What is not true about sclereids ?

A. These are groups of living cells

B. These are found in nut shells, guava
pulp, pear

C. These are also called stone cells

D. These are form of sclerenchyma

Answer: A



Watch Video Solution

46. Chlorenchyma is known to develop in the

A. Cytoplasm of Chlorella

B. Mycelium of a green mould such as

Aspergillus

C. Capsule of a moss

D. Pollen tube of Pinus

Answer: C



Watch Video Solution

47. Vessels are found in

A. All angiosperms and some gymnosperm

B. Most of angiosperms and few
gymnosperms

C. All angiosperms, all gymnosperms and some pteridophyta

D. All pteridophyta

Answer: B



Watch Video Solution

48. Where are casparian rings found?

A. Endodermis

B. Exodermis

C. Pericycle

D. Epidermis

Answer: A



Watch Video Solution

49. Vascular bundles in monocotyledons are considered closed because

A. Xylem is surrounded all around by
phloem

B. There are no vessels with perforations

C. A bundle sheath surrounds each bundle

D. There is no secondary growth

Answer: D



Watch Video Solution

50. At maturity which of the following is enucleate ?

A. Palisade cell

B. Cortical cell

C. Sieve cell

D. Companion cell

Answer: C



Watch Video Solution

51. What happens during vascularisation in plants ?

A. Differentiation of procambium,
formation of primary phloem followed
by formation of primary xylem

B. Differentiation of procambium
followed by the formation of primary
phloem and xylem simultaneously

C. Formation of procambium, secondary
phloem and xylem simultaneously

D. Differentiation of procambium followed
by the formation of secondary xylem

Answer: B



Watch Video Solution

52. Which of the following statements is true?

A. Vessels are multicellular is true?

B. Tracheids are multicellular with narrow
lumen

C. Vessels are unicellular with narrow
lumen

D. Tracheids are unicellular with wide lumen

Answer: D



Watch Video Solution

53. Four radial vascular bundles are found in

A. Dicot root

B. Monocot root

C. Dicot root

D. Monocot stem

Answer: A



Watch Video Solution

54. Which one of the following have sunken stomata ?

A. Nerium

B. Hydrilla

C. Mango

D. Guava

Answer: A



Watch Video Solution

55. Gymnosperms are also called soft wood spermatophytes because they lack

A. Thick-walled tracheids

B. Xylem fibres

C. Cambium

D. Phloem fibres

Answer: B



Watch Video Solution

56. In an annual ring, the light coloured part is known as

A. Early wood

B. Late wood

C. Heartwood

D. Sapwood

Answer: D



Watch Video Solution

57. As a tree grows older, which increases rapidly in thickness

A. Heart wood

B. Sap wood

C. Phloem elements that serve as entry points for substances for transport to other plant parts.

D. Cortex

Answer: A



Watch Video Solution

58. Diffuse porous woods are characteristics of plants growing in

- A. Alpine region
- B. Cold winter regions
- C. Temperate climate
- D. Tropics

Answer: D



Watch Video Solution

59. Lenticles are involved in

- A. Gaseous exchange

B. Food transport

C. Photosynthesis

D. Transpiration

Answer: A



Watch Video Solution

60. The periderm includes

A. Secondary phloem

B. Cork

C. Cambium

D. All of these

Answer: B



Watch Video Solution

61. Which of the following meristems is responsible for extrastelar secondary growth in dicotyledonous stem

A. Interfascicular cambium

B. Intercalary meristem

C. Phellogen

D. Intrafascicular cambium

Answer: C



Watch Video Solution

Section D Assertion Reason Type Questions

1. Assertion: In maize stem, endodermis is present between general cortex and pericycle.

Reason: Eustele is present in maize Stem.

A. If both Assertion & Reason are true and the reason is the correct explanation of the assertion. Then mark (1).

B. If Assertion & Reason are true but the reason is not the correct explanation of the assertion, then mark (2)

C. If Assertion is true statement but Reason is false, then mark (3).

D. If both Assertion and Reason are false statements, then mark (4)

Answer: D



Watch Video Solution

2. A: Bast fibres are collenchymatous fibres.

R: Fibres are absent in secondary phloem.

A. Both Assertion & Reason are true and the reason is the correct explanation of

the assertion.

B. Assertion & Reason are true but the reason is not the correct explanation of the assertion.

C. Assertion is true statement but Reason is false.

D. Both Assertion and Reason are false statements.

Answer: D



Watch Video Solution

3. A: Oldest layer of sapwood lies just outside vascular cambium.

R: Sapwood contains actively conducting vessels and occupies central part of stem.

A. If both Assertion & Reason are true and the reason is the correct explanation of the assertion. Then mark (1).

B. If Assertion & Reason are true but the reason is not the correct explanation of

the assertion, then mark (2)

C. If Assertion is true statement but

Reason is false, then mark (3).

D. If both Assertion and Reason are false

statements, then mark (4)

Answer: D



Watch Video Solution

4. A: Bark is the tissue outside steler cambium.

R: Bark is formed due to the activity of phellogen only.

A. If both Assertion & Reason are true and the reason is the correct expalanation of the assertion. Then mark (1).

B. If Assertion & Reason are true but the reason is not the correct explanation of the assertion, then mark (2)

C. If Assertion is true statement but Reason is false, then mark (3).

D. If both Assertion and Reason are false statements, then mark (4)

Answer: C



Watch Video Solution

5. A: According to Clowes, there are three histogens in monocot root.

R: In dicot roots, innermost groups of initials form root cap.

A. If both Assertion & Reason are true and the reason is the correct explanation of the assertion. Then mark (1).

B. If Assertion & Reason are true but the reason is not the correct explanation of the assertion, then mark (2)

C. If Assertion is true statement but Reason is false, then mark (3).

D. If both Assertion and Reason are false statements, then mark (4)

Answer: D



Watch Video Solution

6. A: Intercalary meristems are commonly located at the base of leaves, above the nodes or below the nodes.

R: Vacuoles are large sized in the cells of intercalary meristem.

A. If both Assertion & Reason are true and the reason is the correct explanation of the assertion. Then mark (1).

B. If Assertion & Reason are true but the reason is not the correct explanation of the assertion, then mark (2)

C. If Assertion is true statement but Reason is false, then mark (3).

D. If both Assertion and Reason are false statements, then mark (4)

Answer: C



Watch Video Solution

7. A: Cells of sclerenchyma have thickened secondary walls.

R: Cells have deposition of lignin.

A. If both Assertion & Reason are true and the reason is the correct explanation of the assertion. Then mark (1).

B. If Assertion & Reason are true but the reason is not the correct explanation of the assertion, then mark (2)

C. If Assertion is true statement but Reason is false, then mark (3).

D. If both Assertion and Reason are false statements, then mark (4)

Answer: A



Watch Video Solution

8. A: Sieve tubes and companion cells are related ontogenetically.

R: Both develop from same mother cell.

A. If both Assertion & Reason are true and the reason is the correct explanation of the assertion. Then mark (1).

B. If Assertion & Reason are true but the reason is not the correct explanation of the assertion, then mark (2)

C. If Assertion is true statement but Reason is false, then mark (3).

D. If both Assertion and Reason are false statements, then mark (4)

Answer: A



Watch Video Solution

9. A: In monocot roots, pericycle gives rise to lateral roots and cork cambium.

R: It has tetrach vascular bundles usually.

A. If both Assertion & Reason are true and the reason is the correct explanation of the assertion. Then mark (1).

B. If Assertion & Reason are true but the reason is not the correct explanation of the assertion, then mark (2)

C. If Assertion is true statement but Reason is false, then mark (3).

D. If both Assertion and Reason are false statements, then mark (4)

Answer: D



Watch Video Solution

10. A: Complementary cells are cut-off by phellogen towards outside.

R: Phellogen is partially secondary in origin.

A. If both Assertion & Reason are true and the reason is the correct explanation of the assertion. Then mark (1).

B. If Assertion & Reason are true but the reason is not the correct explanation of the assertion, then mark (2)

C. If Assertion is true statement but Reason is false, then mark (3).

D. If both Assertion and Reason are false statements, then mark (4)

Answer: C



Watch Video Solution

