



## CHEMISTRY

### AAKASH INSTITUTE ENGLISH

### CHEMICAL KINETICS

#### Example

1. For a reaction,  $2A + D \rightarrow 3C$ , the rate of appearance of C at time t is  $1.2 \times 10^{-4} \text{ mol L}^{-1} \text{ s}^{-1}$ . Find the rate of reaction.



[Watch Video Solution](#)

2. The decomposition of  $NH_3$  on platinum surface is a zero order reaction. What would be the rate of production of  $N_2$  and  $H_2$  if  $k = 2.5 \times 10^{-4} \text{ mol L}^{-1} \text{ s}^{-1}$ ?





 [Watch Video Solution](#)

3. The form of rate law for a reaction is expressed as,  $\text{rate} = k[\text{Cl}_2][\text{NO}]^2$

Find out the order of the reaction with respect to  $\text{Cl}_2$ , with respect to NO and also the overall order of the reaction.

 [Watch Video Solution](#)

4. From the rate expression for the following reactions, determine their order of reaction and the dimensions of the rate constants.



 [Watch Video Solution](#)

5. The half life period of a first order reaction is 60 min. What percentage will be left after 240 min.

 [Watch Video Solution](#)

6. A first order reaction has specific reaction rate of  $10^{-3} s^{-1}$  . How much time will it takes for 10 gm of the reactant to reduce to 2.5 gm ?

 [Watch Video Solution](#)

7. A first order reaction is found to have a rate constant  $k = 7.39 \times 10^{-5} s^{-1}$  . Find the half life of this reaction.

 [Watch Video Solution](#)

8. Half life for a first order reaction is 20 minutes . What is its rate constant ?



 [Watch Video Solution](#)

9. Half life time for first order reaction is 25 minutes . What time will be required for 99% completion of reaction ?

 [Watch Video Solution](#)

10. A first order reaction has a specific rate constant of  $2 \times 10^{-3} \text{ s}^{-1}$  . How much time it will take for 20 gm to reduce to 5 g ?

 [Watch Video Solution](#)

11. The rate constant  $k$  for the order gas phase decomposition of ethyl iodide,  $C_2H_5I \rightarrow C_2H_4 + HI$  is  $1.60 \times 10^{-5} \text{ s}^{-1}$  at 600 K and  $6.36 \times 10^{-3} \text{ s}^{-1}$  at 700 K. Calculate the energy of activation for this reaction.

 [Watch Video Solution](#)

12. What would be the energy of activation for a reaction when a change of temperature from  $20^{\circ}\text{C}$  to  $30^{\circ}\text{C}$  exactly triples the reaction rate constant ?

 [Watch Video Solution](#)

13. The rate constant of a reaction is  $1.5 \times 10^7 \text{ s}^{-1}$  at  $50^{\circ}\text{C}$  and  $4.5 \times 10^7 \text{ s}^{-1}$  at  $100^{\circ}\text{C}$ . Calculate the value of activation energy for the reaction ( $R = 8.314 \text{ JK}^{-1} \text{ mol}^{-1}$ )

 [Watch Video Solution](#)

14. Calculate the activation energy of a reaction whose reaction rate at  $310 \text{ K}$  gets doubled for  $10 \text{ K}$  rise in temperature.

 [Watch Video Solution](#)

15. Half life of a first order chemical reaction is 69 hr at 300 K. Also , rate of this reaction is doubled as temperature is increased from 300 K to 310 K. Determine activation energy and pre - exponential factor for this reaction.

 [Watch Video Solution](#)

16. If the rate of a reaction gets doubled as the temperature is increased from  $27^{\circ}$  to  $37^{\circ}C$ . Find the activation energy of reaction?

 [Watch Video Solution](#)

17. Why reactions having higher molecularity are less observed ?

 [Watch Video Solution](#)

18. Rate constant  $k$  of a reaction varies with temperature according to the equation

$$\log k = \text{constant} - \frac{E_a}{2.303R} \left( \frac{1}{T} \right)$$

where  $E_a$  is the energy of activation for the reaction . When a graph is plotted for  $\log k$  versus  $\frac{1}{T}$  , a straight line with a slope - 6670 K is obtained. Calculate the energy of activation for this reaction

 [Watch Video Solution](#)

19. Why do reaction rates depend on temperature ? Explain.

 [Watch Video Solution](#)

20. What is the effect of catalyst on:

(i) Gibbs energy ( $\Delta G$ ) and


(ii) activation energy of a reaction ?

 [Watch Video Solution](#)

## Try Yourself

1. The rate of formation of nitric oxide (NO) in the following reaction is

$$3.6 \times 10^{-3} \text{ mol L}^{-1} \text{ s}^{-1} .$$



Find the rate of disappearance of oxygen.



[Watch Video Solution](#)

2. For the reaction  $R \rightarrow P$ , the concentration of a reactant changes from 0.03M to 0.02M in 25 minutes. Calculate the average rate of reaction using units of time both in minutes and seconds.



[Watch Video Solution](#)

3. For the reaction  $A + B \rightarrow C + D$ , doubling the concentration of both the reactants increases the reaction rate by 8 times and doubling the



initial concentration of only  $B$  is simply doubles the reaction rate. What is the rate law for the reaction ?

 [Watch Video Solution](#)

4. For a reaction  $X \rightarrow Y$ , the rate of reaction becomes twenty seven times when the concentration of  $X$  is increased three times. What is the order of the reaction?

 [Watch Video Solution](#)

5. The rate constant for first reaction is  $0.0005 \text{ min}^{-1}$ . Calculate its half life.

 [Watch Video Solution](#)

6. A first order reaction has a rate constant  $1.15 \times 10^{-3} \text{ s}^{-1}$ . How long will 5 g of this reactant take to reduce to 3 g?

 [Watch Video Solution](#)

7. The half life time ,  $t_{1/2}$  for the first order decomposition of  $NH_4NO_2$  is 123 minutes at  $15^\circ C$  .

(a) What is the value of  $k$  ?

(b) How long will it take for 2 g of  $NH_4NO_2$  to decompose until 0.2 remains ?

 [Watch Video Solution](#)

8. The rate of reaction triples when temperature changes form  $20^\circ C$  to  $50^\circ C$ . Calculate the energy of activation for the reaction ( $R = 8.314JK^{-1}mol^{-1}$ ).

 [Watch Video Solution](#)

9. The rate constant of a reaction is  $1.2 \times 10^{-3}s^{-1}$  at  $30^\circ C$  and  $2.1 \times 10^{-3}s^{-1}$  at  $40^\circ C$ . Calculate the energy of activation of the

reaction.


 [Watch Video Solution](#)

10. The activation energy of a reaction is  $94.14 \text{ kJ mol}^{-1}$  and the value of rate constant at  $313 \text{ K}$  is  $18 \times 10^{-5} \text{ s}^{-1}$ . Calculate the frequency factor or pre-exponential factor,  $A$ .

 [Watch Video Solution](#)

## Exercise

1. In a reaction,



rate =  $-\frac{1}{a} \frac{d[A]}{dt}$ , the -ve sign in it represents

- A. The rate of reaction increases with time
- B. The rate is negative for the reaction

C. The concentration of the reactant 'A' decreases during the reaction

D. The concentration of reactant 'A' increases in the reaction

**Answer: C**

 [Watch Video Solution](#)

2. The unit of rate and rate constant are same for a

A. mol/Ls

B. Same for instantaneous rate and average rate


C. Same for all reactions and not depend upon the nature of reaction

D. All of these

**Answer: D**

 [Watch Video Solution](#)

3. For the reaction,



Select the correct statement

- A. Rate of formation of  $O_2$  is same as rate of formation of  $NO_2$
- B. Rate of disappearance of  $N_2O_5$  is two times the rate of formation of  $NO_2$
- C. Rate of formation of  $O_2$  is 0.5 time rate disappearance of  $N_2O_5$
- D. Rate of formation of  $NO_2$  is equal to rate of disappearance of  $N_2O_5$

Answer: C



Watch Video Solution

4. The rate constant of  $n^{th}$  order has units :

A.  $(mol/L)^n s^{-1}$

B.  $(mol / L s^{-1})^{n-1}$

C.  $\left(\frac{L}{mol} s^{-1}\right)^{n-1}$

D.  $(L / mol)^{n-1} s^{-1}$

**Answer: D**



**Watch Video Solution**

5.  $\frac{dx}{dt} = K[A]^{0.5}[B]^{0.5}[C]^{0.5}$ . What will be the order of the reaction ?

A. 1

B. 1.5

C. Zero

D. 2

**Answer: B**



**Watch Video Solution**

6.  $H_2 + I_2 \rightarrow 2HI$  (An elementary reaction) of second order

If the volume of the container containing the gaseous mixture is increased to two times, then final rate of the reaction.

A. Become four times

B. Become  $\frac{1}{4}$  th of the original rate

C. Become 2 times

D. Become  $\frac{1}{2}$  of the original rate

**Answer: B**



**Watch Video Solution**

7. In a certain reaction shown below,



If rate of formation of C is  $9.6 \times 10^{-2}$  mol/Ls . What will be rate of reaction ?

A.  $9.6 \times 10^{-2} \text{ mol/Ls}$

B.  $3.2 \times 10^{-2} \text{ mol/Ls}$

C.  $2.4 \times 10^{-2} \text{ mol/Ls}$

D.  $4.8 \times 10^{-2} \text{ mol/Ls}$

**Answer: B**

 [Watch Video Solution](#)

8. if the rate of a reaction is equal to the rate constant , the order of the reaction is

A. First order

B. Second order

C. Half order


D. Zero order

**Answer: D**



 Watch Video Solution

9. For the reaction,



select the correct statement

- A. Rate of disappearance of  $H_2O_2$  will be three times the rate of disappearance of  $I^-$
- B. Rate of disappearance of  $H_2O_2$  is  $\frac{1}{3}$  of rate of formation of  $I_3^-$
- C. Rate of disappearance of  $I^-$  ions will be three times the rate of formation  $I_3^-$  ions
- D. Rate of formation of  $H_2O$  is  $\frac{1}{2}$  of rate formation of  $I_3^-$

**Answer: C**

 Watch Video Solution

10. The rate constant depends on

- A. Temperature
- B. Concentration
- C. Catalyst
- D. Both (1) & (3)

**Answer: D**



[Watch Video Solution](#)

11.  $2A \rightarrow C + D$

In this reaction, if we double the concentration of A, reaction rate becomes two times. What is the order of this reaction?

- A. 1
- B. 3
- C. 2

D. 1.5

**Answer: A**



[Watch Video Solution](#)

**12.** In a zero order reaction , 20% of the reaction complete in 10 s. How much time it will take to complete 50% of the reaction ?

A. 20 s

B. 25 s

C. 30 s

D. 40 s

**Answer: B**



[Watch Video Solution](#)

13. If the half - life of the first order reaction is 50 s, what be the value of its rate constant ?

A.  $1.38 \times 10^{-2} s^{-1}$

B.  $25 s^{-1}$

C.  $34.66 s^{-1}$

D.  $1.38 \times 10^{-4} s^{-1}$

**Answer: A**



[Watch Video Solution](#)

14. In a first order reaction , on plotting a graph between  $t_{1/2}$  and concentration of reactant

A. A curve is obtained

B. A straight line is obtained having slope  $45^\circ$

C. A straight line is parallel to time axis

D. A straight line is parallel to the concentration axis

**Answer: D**

 [Watch Video Solution](#)

15. In data given below.

	[A]	[B]	rate
	<i>mol / L</i>	<i>mol / L</i>	<i>molL<sup>-1</sup>s<sup>-1</sup></i>
a.	0.02	1.2	$3.0 \times 10^{-3}$
b.	0.04	2.4	$6.0 \times 10^{-3}$

The reaction may be

- A. First order
- B. Second order
- C. Pseudo second order
- D. Second order w.r.t. A

**Answer: A**

 [Watch Video Solution](#)

16. For certain first order reaction, 75% of the reaction complete in 30 min. How much time it require to complete 99.9% of the reaction ?

- A. 150 min
- B. 100 min
- C. 90 min
- D. 300 min

**Answer: A**



[Watch Video Solution](#)

17. For zero order reaction ,  $t_{1/2}$  will be ( $A_0$  is the initial concentration ,  $K$  is rate constant)

- A.  $0.693/K$
- B.  $2.303/K$

C.  $\frac{A_0}{2K}$

D.  $\frac{\ln A_0}{2}$

**Answer: C**

 [Watch Video Solution](#)



**18.** In pseudo - order reactions

- A. The actual order of reaction is different from that expected using rate law expression
- B. The concentration of at least one reactant is taken in large excess
- C. The concentration of reactant taken in excess may be taken as constant
- D. All of these

**Answer: D**

 [Watch Video Solution](#)

19. The depletion of ozone involves the following steps :



The predicted order of the reaction will be

A. First

B. 1.5

C. - 1.5

D. Zero

**Answer: A**



[Watch Video Solution](#)

20. The half life for the zero order reaction will be

A.  $t_{1/2} \propto (\text{conc.})^0$



B.  $t_{1/2} \propto (\text{conc.})^1$

C.  $t_{1/2} \propto (\text{conc.})^{-1}$

D.  $t_{1/2} \propto (\text{conc.})^2$

**Answer: B**

 [Watch Video Solution](#)

**21. A catalyst cannot change**

A. Rate constant of the reaction

B. Equilibrium constant of the reaction

C. Activation energy of reaction

D. All of these

**Answer: B**

 [Watch Video Solution](#)

22. Select the incorrect statement

- A. The minimum amount of energy required by reactant molecules to give product is threshold energy
- B. Activation energy is the sum of threshold energy and average kinetic energy
- C. Threshold energy is the sum of initial potential energy of reactants and activation energy
- D. Lower is the activation energy faster is the reaction

**Answer: B**



[Watch Video Solution](#)

23. If the rate of reaction increases by two times with every  $10^{\circ}C$  rise in temperature, then if we raise the temperature by  $40^{\circ}C$ , the new rate of the reaction will be :-

- A. 4 times
- B. 8 times
- C. 16 times
- D. 32 times

**Answer: C**

 [Watch Video Solution](#)

**24.** According to collision theory, rate of the reaction increases , because -

- (a) Number of collision decreases
- (b) Number of molecules crossing energy barrier increases
- (c) Kinetic energy of molecules increases
- (d) Value of rate constant increases

- A. Number of collision decreases
- B. Number of molecules crossing energy barrier increases
- C. Kinetic energy of molecules increases

D. Value of rate constant increases

**Answer: B**

 [Watch Video Solution](#)

25.  $A + B \rightarrow C$ ,  $\Delta H = + 60 \text{ kJ/mol}$

$E_{a,f}$  is 150 kJ. What is the activation energy for the backward reaction ?

A. 210 kJ


B. 105 kJ

C. 90 kJ

D. 145 kJ

**Answer: C**

 [Watch Video Solution](#)



26.

Reaction progress →

- A. It is a 2 steps reaction , step 1 is slower than step 2
- B. It is a 2 steps reaction , step 2 is slower than step 1
- C. Single step reaction where B is a activated complex
- D. Single step reaction where B is a reaction intermediate

**Answer: A**

 [Watch Video Solution](#)

27. The rate constant of a reaction depends upon

- A. Nature of reactants
- B. Physical state of the reactants
- C. Temperature
- D. All of these

**Answer: D**

 [Watch Video Solution](#)

28. The correct expression for Arrhenius equation is (where K is rate constant)

- A.  $K = e^{-\Delta H / RT} A$
- B.  $\ln K = \frac{\Delta E_a}{RT} + \ln A$
- C.  $\ln K = \ln A + \frac{E_a}{RT}$
- D.  $\ln \left( \frac{A}{K} \right) = \frac{E_a}{RT}$

**Answer: D**

 [Watch Video Solution](#)

**29.** The temperature coefficient of a certain reaction is found to be 3. If temperature changes from  $25^{\circ}$  to  $55^{\circ}$ , the new rate of reaction will be

A. 8 times

B. 9 times

C. 16 times

D. 27 times

**Answer: D**

 [Watch Video Solution](#)

**30.** In Arrhenius equation,  $K = Ae^{-E_a/RT}$ . The A is

- A. Activation energy
- B. Enthalpy of reaction
- C. Free energy of reaction
- D. Rate constant of the reaction at infinite temperature

**Answer: D**

 [Watch Video Solution](#)

## Assignment Section A Objective Type Questions

1. The unit of rate and rate constant are same for a

- A. First order
- B. Zero order
- C. Second order
- D. Third order



**Answer: B**

 [Watch Video Solution](#)

2. For a reaction of type

$aA + bB \rightarrow \text{products}$ ,  $\frac{-d[A]}{dt}$  is equal to

A.  $\frac{-d[B]}{dt}$

B.  $-\frac{b}{a} \frac{d[B]}{dt}$

C.  $-\frac{a}{b} \times \frac{d[B]}{dt}$

D.  $-\frac{b}{a} \times \frac{d[B]}{dt}$

**Answer: C**

 [Watch Video Solution](#)

3. For a gaseous reaction, the rate of reaction may be expressed in the units

A. atm

B. atm s

C. atm/s

D.  $\text{atm} / \text{s}^2$

**Answer: C**



**Watch Video Solution**

4. A gaseous reaction  $A_2(g) \rightarrow B(g) + \frac{1}{2}C(g)$  shows increase in pressure from  $100\text{mm}$  to  $120\text{mm}$  in 5 min . What is the rate of disappearance of  $A_2$  ?

A.  $4\text{min min}^{-1}$

B.  $8\text{min min}^{-1}$

C.  $16\text{min min}^{-1}$

D.  $2\text{min min}^{-1}$

**Answer: B**

 [Watch Video Solution](#)

**5. Which of the following will react at the highest rate ?**

- A. 1 mol of A and 1 mol of B in a 1 L vessel
- B. 2 mol of A and 2 mol of B in a 2 L vessel
- C. 3 mol of A and 3 mol of B in a 3 L vessel
- D. All would react at the same rate

**Answer: D**

 [Watch Video Solution](#)

**6. Which of the following does not affect the rate of reaction?**

- A. Amount of the reactant taken

B. Physical state of the reactant

C.  $\Delta H$  of reaction

D. Size of vessel

**Answer: C**

 [Watch Video Solution](#)

7. In a reaction  $2X + Y \rightarrow X_2Y$ ,  $Y$  the reactant  $X$  will disappear at

A. Half the rate as that of disappearance of  $Y$

B. The same rate as that of disappearance of  $Y$

C. The same rate as that of appearance of  $X_2Y$

D. Twice the rate as that of appearance of  $X_2Y$

**Answer: D**

 [Watch Video Solution](#)

8. During the course of a chemical reaction , the rate constant \_\_\_.

- A. Remains constant throughout
- B. Increases as the reaction proceeds
- C. Decreases as the reaction proceeds
- D. First increases followed by a decreases

**Answer: C**

 [Watch Video Solution](#)

9. A reaction involving two different reactants can never be:

- A. (a) Second order reaction
- B. (b) Bimolecular reaction
- C. (c) Unimolecular reaction
- D. (d) First order reaction

Answer: C

 Watch Video Solution

10. For the hypothetical reaction  $2A \rightarrow 3C$ , the reaction rate  $r$  in terms of the rate of change of the concentration is given by

A.  $r = - \frac{d[A]}{dt}$

B.  $r = - \frac{1}{2} \frac{d[A]}{dt}$

C.  $r = \frac{1}{3} \frac{d[A]}{dt}$

D.  $r = \frac{d[A]}{dt}$

Answer: B

 Watch Video Solution

11. In the reaction,  $A + 2B \rightarrow 6C + 2D$ , if the initial rate  $-\frac{d[A]}{dt}$  at  $t=0$  is  $2.6 \times 10^{-2} M \text{ sec}^{-1}$ , what will be the value of  $-\frac{d[B]}{dt}$  at  $t=0$ ?

A.  $2.6 \times 10^{-2}$

B.  $5.2 \times 10^{-2}$

C.  $1.3 \times 10^{-2}$

D.  $1.0 \times 10^{-1}$

**Answer: B**

 [Watch Video Solution](#)

**12.** For the reaction

$N_2 + 3H_2 \rightarrow 2NH_3$  The rate of change of concentration for hydrogen is  $-0.3 \times 10^{-4} Ms^{-1}$  The rate of change of concentration of ammonia is :

A.  $-0.2 \times 10^4$

B.  $0.2 \times 10^{-4}$


C.  $0.1 \times 10^{-4}$

D.  $0.3 \times 10^{-4}$

Answer: B

 Watch Video Solution

13. The graph plotted between concentration versus time



A. It gives rate of disappearance of reactant

B. Rate =  $(d[C_2 - C_1]) \frac{)}{t_2 - t_1}$

C. Both (1) & (2)

D. It predicts the order of reaction

Answer: C





Watch Video Solution

14. For reaction,  $2B + A \rightarrow 2C$

Which of the following is correct ?

A. Rate of disappearance of B is twice that of rate of disappearance of

C

B. Rate of disappearance of A is twice that of rate of disappearance of

C

C. Rate of appearance of C is twice that of rate of disappearance of A

D. All of these

Answer: C



Watch Video Solution

15. For the homogeneous elementary reaction,  $A + B \rightarrow C$ , the unit of rate constant is

A.  $s^{-1}$

B.  $s^{-1} \text{mol L}^{-1}$

C.  $s^{-1} \text{mol}^{-1} \text{L}$

D.  $s$

**Answer: C**



[Watch Video Solution](#)

16. The rate law for a reaction between  $A$  and  $B$  is given by rate  $= k[A]^n[B]^m$ . On doubling the concentration of  $A$  and halving the concentration of  $B$ , the ratio of the new rate to the earlier rate of the reaction becomes

A.  $\frac{1}{2^{n+m}}$

B.  $m + n$

C.  $2^{n+m}$

D.  $2^{n-m}$

**Answer: C**

 [Watch Video Solution](#)

17. Consider the reaction,

$2A + B \rightarrow C + D$  , If the rate expression is  $r = K[A]^2[B]^1$  and if volume is reduced to  $\frac{1}{3}$  , the rate of reaction will increase by

A. 27 times

B. 9 times

C. 8 time

D. Rate will not get affected

**Answer: A**

[Watch Video Solution](#)

18. The rate constant is numerically the same for three reaction of first , second and third order respectively. Which one is true for the of three reaction, if concentration of reactant is greater than 1M ?

A.  $r_1 = r_2 = r_3$

B.  $r_1 > r_2 > r_3$

C.  $r_1 < r_2 < r_3$

D. All of these

Answer: C

[Watch Video Solution](#)

19. Assuring an elementary reaction

$H_2O_2 + 3I^- + 2H^+ \rightarrow 2H_2O + I_3^-$  . The effect on the rate of this

reaction brought about by doubling the concentration of  $I^-$  without changing the order

- A. The rate would increase by a factor of 3
- B. The rate would increase by a factor of 8
- C. The rate would decrease by a factor of  $1/3$
- D. The rate would increase by a factor of 9

**Answer: B**



[Watch Video Solution](#)

20. For a reaction  $A + B \rightarrow$  products, the rate of reaction was doubled when concentration of A was doubled. When concentration of A and B both was double, the rate was again doubled, order of reaction w.r.t. A and B are

- A. 1,1
- B. 2,0

C. 1,0

D. 0,1

**Answer: C**

 [Watch Video Solution](#)

21. The overall order of reaction between X & Y is 3. Which of the following rate equation must be correct, if on doubling the concentration of X, the rate of reaction gets doubled ?

A.  $r = K[X]^2[Y]^0$

B.  $r = K[X]^1[Y]^2$

C.  $r = K[X]^1[Y]^3$

D.  $r = K[X]^2[Y]^1$

**Answer: B**

 [Watch Video Solution](#)

22. For a zero order reaction,

$$K = 1 \times 10^{-3} \text{mol L}^{-1} \text{s}^{-1}$$

If initial concentration of the reactant is  $1.0 \text{mol L}^{-1}$ , the concentration after 10 minutes would be

A.  $1 \times 10^{-2} \text{mol L}^{-1}$

B.  $0.6 \text{mol L}^{-1}$

C.  $0.4 \text{mol L}^{-1}$

D.  $1.0 \text{mol L}^{-1}$

**Answer: C**



[Watch Video Solution](#)

23. Which of the following statement is not correct ?

A. Molecularity of a reaction cannot be fractional

B. Molecularity of a reaction cannot be more than three

C. Molecularity of a reaction may or may not be equal to the order of reaction

D. Molecularity of a reaction is obtained from balanced chemical equation

**Answer: D**

 [Watch Video Solution](#)

**24.** Consider the following in respect of zero order reaction.

I.  $t_{1/2}$  is directly proportional to the initial concentration

II. Time taken for the completion of the reaction is twice its  $t_{1/2}$

III Concentration of the reactant decreases linearly with time

Which of the statements given above are correct ?

A. I & II only

B. I & III only



C. II & III only

D. I, II & III

**Answer: D**

 [Watch Video Solution](#)

25. If initial concentration is reduced to  $1/4^{th}$  in a zero order reaction ,  
the time taken for half the reaction to complete

A. Remains same

B. Becomes 4 time

C. Become one fourth

D. Doubles

**Answer: C**

 [Watch Video Solution](#)

26. If initial concentration is doubled, the time for half reaction is also doubled, The order of reaction is


- A. Zero
- B. One
- C. Two
- D. Three

**Answer: A**



[Watch Video Solution](#)

27. The inversion of cane sugar is represented by



It is a reaction of

- A. (a) Unimolecular
- B. (b) Second order

C. (c) Zero order

D. (d) First order

**Answer: D**

 [Watch Video Solution](#)

**28.** For a 1st order reaction , a straight line is obtained if you plot

A. (a) log conc. vs time

B. (b) conc. vs time

C. (c) 1/conc. vs time

D. (d) log conc. vs 1/time

**Answer: A**

 [Watch Video Solution](#)

29. Which order reaction obeys the expression  $t_{1/2} \propto 1[A]$  ?

A. (a) First

B. (b) Second

C. (c) Third


D. (d) Zero

**Answer: B**



**Watch Video Solution**

30. The graph of  $t_{1/2}$  versus initial concentration 'a' is for



- A. (a) First order
- B. (b) Second order
- C. (c) Zero order
- D. (d) Can't predict

**Answer: A**



**Watch Video Solution**

31. Reactant 'A' (initial concentration, a) reacts according to zero order kinetics, the time taken for the completion of the reaction is

A.  $\frac{a}{k}$

B.  $\frac{a}{2k}$

C.  $\frac{k}{a}$

D.  $\frac{2k}{a}$

**Answer: A**



[Watch Video Solution](#)

32. A first order reaction completes 60% in 20 minutes. The time required for the completion of 90% of the reaction is approx.....

A. 30 minutes

B. 40 minutes

C. 50 minutes

D. 60 minutes

**Answer: C**

 [Watch Video Solution](#)

**33.** The half-life period of a substance is 50 min at a certain initial concentration. When the concentration is reduced to one-half of its initial concentration, the half-life period is found to be 25 min . Calculate the order of reaction.

A. Zero

B. First

C. Second

D. Third

**Answer: A**

 [Watch Video Solution](#)

34. The half life of a second order process,  $2A \rightarrow \text{products}$ , is

- A. (a) Independent of initial concentration
- B. (b) Directly proportional to initial concentration of A
- C. (c) Inversely proportional to initial concentration of A
- D. (d) Inversely proportional to square of initial concentration

**Answer: C**



[Watch Video Solution](#)

35. The rate law for the reaction

$RCl + NaOH(aq) \rightarrow ROH + NaCl$  is given by

Rate =  $k[RCl]$ . The rate of the reaction will be

- A. Doubled on doubling the concentration of NaOH
- B. Halved on reducing the concentration of alkyl halide to one half
- C. Decreases on increasing the temperature



D. Unaffected by increasing the temperature

**Answer: B**

 [Watch Video Solution](#)

**36.** The rate constant for forward and backward reactions of hydrolysis of ester are  $1.1 \times 10^{-2}$  and  $1.5 \times 10^{-3}$  per minute respectively. Equilibrium constant for the reaction is

A. 7.33

B. 0.733

C. 73.3

D. 733

**Answer: A**

 [Watch Video Solution](#)

37. For a reaction ,  $A \rightarrow B$  , it has been found that the order of the reaction , is zero with respect to A. Which of the following expressions correctly describes the reaction ?

A. (a)  $K = \frac{2.303}{t} \log. \frac{[A]_0}{[A]}$

B. (b)  $[A]_0 - [A] = Kt$

C. (c)  $t_{1/2} = \frac{0.693}{K}$

D. (d)  $t_{1/2} \propto \frac{1}{[A]_0}$

**Answer: B**



**Watch Video Solution**

38. A sample of a radioactive substance undergoes 80% decomposition in 345 minutes. Its half life is .....minutes

A. (a)  $\frac{\ln 2}{\ln 5} \times 345$

B. (b)  $\frac{\ln 5}{\ln 2} \times 345$

C. (c)  $\frac{\ln 5}{\ln 4} \times 345$

D. (d)  $\frac{\ln 4}{\ln 5} \times 345$

**Answer: A**



**Watch Video Solution**

39. 99 % at a first order reaction was completed in 32 min . When will 99.9 % of the reaction complete.

A. (a) 50 minute

B. (b) 46 minutes

C. (c) 49 minute

D. (d) 48 minute

**Answer: D**



**Watch Video Solution**

40. Check, which of the following statements is false ?

- A. A catalyst does not differentiate between forward and backward reaction
- B. Large activation energy is associated with low reaction rate
- C. Maxwell's distribution of velocities remains- unaltered under all conditions of temperature and pressure
- D. A catalyst does not affect the equilibrium state of reaction

**Answer: C**



[Watch Video Solution](#)

41. Arrhenius parameter (A) depends on

- A. Steric factor
- B. Collision frequency
- C. Both (1) & (2)

D. Neither (1) nor (2)

**Answer: C**

 [Watch Video Solution](#)

**42.** The chemical reactions in which the reactants require high amount of activation energy are generally

A. Slow

B. Fast

C. Instantaneous

D. Spontaneous


**Answer: A**

 [Watch Video Solution](#)


43. For an exothermic chemical process occurring in two steps occurring in two steps as follows




The progress of reaction can be best described by :



A.



B.



C.


D. None of these

Answer: B




Watch Video Solution


44. A chemical process occurring in two steps, is plotted as




Endothermic




Exothermic



Exothermic




Endothermic



Answer: B

 Watch Video Solution

45. For the chemical process energies are plotted in graph.



Which of the following is correct ?

A. It is the exothermic reaction ,  $\Delta H = b - a$

B. Threshold energy ,  $e = a + c$



C.  $(E_a)_f < (E_a)_b$

D. All of these

**Answer: D**

 [Watch Video Solution](#)

**46.** The rate of a reaction becomes 2 times for every  $10^\circ C$  rise in temperature . How many times rate of reaction will be increased when temperature is increased from  $30^\circ C$  to  $80^\circ C$  ?

A. 16

B. 32

C. 64

D. 128

**Answer: B**

 [Watch Video Solution](#)

47. The rate of a reaction increases by 2.5 times when the temperature is raised from 300K to 310K. If  $K$  is the rate constant at 310 K will be equal to

- A. (a)  $K$
- B. (b)  $2K$
- C. (c)  $2.5K$
- D. (d)  $3K$

**Answer: C**



[Watch Video Solution](#)

48. The minimum amount of energy that the reacting molecules must possess at the time of collisions in order to produce effective collision is called

- A. Activation energy

B. Threshold energy

C. Internal energy

D. Free energy

**Answer: B**



**Watch Video Solution**

**49.** The rate constant, the activation energy and Arrhenius parameter of a chemical reaction at  $25^\circ C$  are  $x$ ,  $10x$  kJ/mol and  $2xs^{-1}$ . Value of rate constant as  $T \rightarrow \infty$  is

A.  $xs^{-1}$

B.  $2xs^{-1}$

C.  $\infty$

D.  $10xs^{-1}$

**Answer: B**



[Watch Video Solution](#)

50. At particular concentration , the half life of the reaction is 100 minutes. When the concentration of the reactant become double half life becomes , 25 minutes , then what will be the order of the reaction ?

A. 1

B. 2

C. 0

D. 3

**Answer: D**



[Watch Video Solution](#)

**Assignment Section B Objective Type Questions**

1. Consider the data given below for hypothetical reaction  $A \rightarrow X$

Time (s)	Rate of the reaction ( $\text{molL}^{-1}\text{s}^{-1}$ )
0	$1.60 \times 10^{-2}$
10	$1.60 \times 10^{-2}$
20	$1.60 \times 10^{-2}$
30	$1.59 \times 10^{-2}$

From the above data, the order of reaction is:

A. Zero

B. 1

C. 2

D. 3

**Answer: A**



**Watch Video Solution**

2. The graph between  $\log k$  versus  $1/T$  is a straight line.

A.  $-\frac{2.303R}{E_a}$

B.  $-\frac{E_a}{2.303R}$

C.  $\frac{2.303R}{E_a}$

D.  $\frac{E_a}{2.303R}$

**Answer: B**



**Watch Video Solution**

3. The temperature coefficient of most of the reactions lies between

A. 1 & 3

B. 2 & 3

C. 1 & 4

D. 2 & 4

**Answer: B**



**Watch Video Solution**

4. If the volume of the vessel in which the reaction  $2NO + O_2 \rightarrow 2NO_2$  is occurring is diminished to  $1/3$  rd of its initial volume . The rate of the reaction will be increased by

- A. One third
- B. Three times
- C. Nine times
- D. Twenty seven times

**Answer: D**

 [Watch Video Solution](#)

5. For the raction  $A + B \rightarrow$  products, it is found that order of A is 2 and the order of B is 3. In the rate expression when the concentration of both A and B are doubled the rate will increases by a factor

- A. 10

B. 16

C. 32


D. 28

**Answer: C**





**Watch Video Solution**

**6. Nitric oxide (NO) reacts with oxygen to produce nitrogen dioxide**



If the mechanism of reaction is



then rate law is

A. Rate =  $K'[NO][O_2]$

B. Rate =  $K'[NO][O_2]^2$

C. Rate =  $K'[NO]^2[O_2]$



$$\text{D. Rate} = K'[\text{NO}]^3[\text{O}_2]$$

**Answer: C**



[Watch Video Solution](#)

7. For the first order reaction, the taken to reduce the initial concentration to a factor of  $\frac{1}{4}$  is 10 minutes if the reduction in concentration is carried out to a factor of  $\frac{1}{16}$ , then time required will be


- A. 10 minutes
- B. 20 minutes
- C. 40 minutes
- D. 60 minutes

**Answer: B**



[Watch Video Solution](#)

8. The rate of the reaction



can be written in three ways:

$$\frac{-d[N_2O_5]}{dt} = k[N_2O_5]$$

$$\frac{d[NO_2]}{dt} = k'[N_2O_5]$$

$$\frac{d[O_2]}{dt} = k''[N_2O_5]$$

The relationship between  $k$  and  $k'$  and between  $k$  and  $k''$  are-

A.  $2K_1 = K_2 = 4K_3$

B.  $K_1 = K_2 = K_3$

C.  $2K_1 = 4K_2 = K_3$

D.  $K_1 = 2K_2 = 3K_3$

**Answer: A**



**Watch Video Solution**

9. For the reaction ,  $N_2O_4(g) \xrightleftharpoons[K_2]{K_1} 2NO_2(g)$ , the rate of disappearance of  $NO_2$  will be

A.  $K_1[N_2O_4] - K_2[NO_2]^2$

B.  $2K_1[N_2O_4] - 2K_2[NO_2]^2$

C.  $K_2[NO_2]^2 - K_1[N_2O_4]$

D.  $2K_2[NO_2]^2 - 2K_1[N_2O_4]$

**Answer: D**

 [Watch Video Solution](#)

10. For a homogeneous gaseous reaction  $A \rightarrow B + C + D$  , the initial pressure was  $P_0$  white pressure after time 't' was P. if  $(P > P_0)$  The expression for the constant K is

A.  $K = \frac{2.303}{t} \log. \left( \frac{2P_0}{3P_0 - P} \right)$

B.  $K = \frac{2.303}{t} \log. \left( \frac{3P_0}{2P_0 - P} \right)$

$$C. K = \frac{2.303}{t} \log. \left( \frac{P_0}{P_0 - P} \right)$$

$$D. K = \frac{2.303}{t} \log. \left( \frac{P_0}{4P_0 - P} \right)$$

**Answer: A**

 **Watch Video Solution**

**11. form the gaseous reaction**

$2A + B_2 \rightarrow 2AB$ , the following rate data were obtained at  $300K$ .

	Rate of disappearance of $B_2$ ( $\text{mol L}^{-1} \text{min}^{-1}$ )	Concentration	
		[A] M	[ $B_2$ ] M
<b>i</b>	$1.8 \times 10^{-3}$	<b>0.015</b>	<b>0.15</b>
<b>ii</b>	$1.08 \times 10^{-2}$	<b>0.090</b>	<b>0.15</b>
<b>iii</b>	$5.4 \times 10^{-3}$	<b>0.015</b>	<b>0.45</b>

Calculate the rate constant for the reaction and the rate of formation of

$AB$  when  $[A]$  is  $0.02$  and  $[B_2]$  is  $0.04 \text{mol L}^{-1}$  at  $300K$ .

A.  $0.5 \text{mol}^{-1} \text{min}^{-1} \text{litre}$

B.  $0.8 \text{mol}^{-1} \text{min}^{-1} \text{litre}$

C.  $1.5 \text{mol}^{-1} \text{min}^{-1} \text{litre}$

D.  $2\text{mol}^{-1}\text{min}^{-1}\text{litre}$

**Answer: B**

 [Watch Video Solution](#)

12. The inversion of a sugar follows first order rate equation which can be followed by noting the change in the rotation of the plane of polarization of light in the polarimeter. If  $r_\infty$ ,  $r_t$  and  $r_0$  are the rotations at  $t = \infty$ ,  $t = t$ , and  $t = 0$ , then the first order reaction can be written as

A.  $K = \frac{1}{t} \log_{10} \frac{r_1 - r_\infty}{r_0 - r_\infty}$

B.  $K = \frac{1}{t} \ln \frac{r_0 - r_\infty}{r_t - r_\infty}$

C.  $K = \frac{1}{t} \ln \frac{r_\infty - r_0}{r_t - r_\infty}$

D.  $K = \frac{1}{t} \ln \frac{r_\infty - r_t}{r_\infty - r_0}$

**Answer: B**

 [Watch Video Solution](#)

13. Which of the following is correct

A.  $\log. \frac{K_2}{K_1} = \frac{E_a}{2.303} \left[ \frac{\Delta T}{T_1 T_2} \right]$

B. For zero order  $t_{1/2}$  is inversely proportional to initial concentration

C. Catalyst decreases the activation energy

D. All of these

Answer: C

 [Watch Video Solution](#)

14. The rate constant for a reaction is  $1.5 \times 10^{-7}$  at  $50^\circ\text{C}$  and  $4.5 \times 10^7 \text{ s}^{-1}$  at  $100^\circ\text{C}$ . What is the value of activation energy?

A.  $2.2 \times 10^3 \text{ J mol}^{-1}$

B.  $2300 \text{ J mol}^{-1}$

C.  $2.2 \times 10^4 \text{ J mol}^{-1}$

D.  $220 \text{ J mol}^{-1}$

**Answer: C**



**Watch Video Solution**

15. In Arrhenius equation ,  $k = Ae^{\frac{E_a}{RT}}$  , A may be termed as rate constant .

- A. When 100% reactant will convert into the product
- B. When the temperature becomes high
- C. When the fraction of molecule crossing over the energy barrier becomes unity
- D. At very low temperature

**Answer: D**



**Watch Video Solution**

16. The rate constant of the production of 2B (g) by the reaction ,  $A(g) \xrightarrow{\Delta} 2B(g)$  is  $2.48 \times 10^{-4} s^{-1}$  A 1 : 1 molar ratio of A to B in the

reaction mixture is attained after

A. 26.25 minute

B. 27.25 minute

C. 28.25 minute

D. 0 minute

**Answer: B**



[Watch Video Solution](#)

17. Two substances A and B are present such that  $[A_0] = 4[B_0]$  and half-life of A is 5 minutes and that of B is 15 minutes. If they start decaying at the same time following first order kinetics after how much time the concentration of both of them would be same ?

A. 15 minute

B. 10 minute

C. 5 minute



D. 12 minute

**Answer: A**



[Watch Video Solution](#)

**18.** If the rate of reaction increases by 27 times , when temperature is increased by 30 K, then temperature coefficient of the reaction is

A. 3

B. 2

C. 1

D. 2.5

**Answer: A**




[Watch Video Solution](#)

19. the reaction  $A \rightarrow B$  follows first order Kinetics the time taken for 0.8 mole of A to produce 0.6 mole of B is 1 H , what is the time taken for the conversion of 0.9 mole of A to 0.675mole of B?

- A. 1 hour
- B. 30 min
- C. 15 min
- D. 5 min

Answer: A

 Watch Video Solution



20. If

find B%

A. 25 %

B. 50 %

C. 75 %

D. 80 %

**Answer: A**



[Watch Video Solution](#)

### Assignment Section C Previous Year Questions

1. A first order reaction has a specific reaction rate of  $10^{-2} s^{-1}$ . How much time will it take for 20 g of the reactant to reduce to 5 g ?

A. 2.38.6 second

B. 138.6 second



C. 346.5 second

D. 693.0 second

**Answer: B**

 [Watch Video Solution](#)

2. A hypothetical reaction  $A_2 + B_2 \rightarrow 2AB$  follows the mechanism as given below:



The order of the overall reaction is

A. 1

B. 2

C. 0

D. 1.5

**Answer: D**

 [Watch Video Solution](#)

3. The decomposition of phosphine ( $PH_3$ ) on tungsten at low pressure is a first order reaction it is because the

- A. Rate is proportional to the surface coverage
- B. Rate is inversely proportional to the surface coverage
- C. Rate is independent of the surface coverage
- D. Rate of decomposition is very slow

**Answer: A**

 [Watch Video Solution](#)

4. The addition of a catalyst during a chemical reaction alters which of the following quantities ?

- A. Activation energy
- B. Entropy

C. Internal energy

D. Enthalpy

**Answer: A**

 [Watch Video Solution](#)

5. The rate of a first-order reaction is  $0.04 \text{ mol L}^{-1} \text{ s}^{-1}$  at 10 seconds and  $0.03 \text{ mol L}^{-1} \text{ s}^{-1}$  at 20 seconds after initiation of the reaction. The half-life period of the reaction is :

A. 54.1 s

B. 24.1 s

C. 34.1 s

D. 44.1 s

**Answer: B**

 [Watch Video Solution](#)

6. The rate constant of the reaction  $A \rightarrow B$  is  $0.6 \times 10^{-3}$  mole per second. If the concentration of  $A$  is  $5M$ , then concentration of  $B$  after 20 minutes is:

A. 0.36 M

B. 0.72 M

C. 1.08 M

D. 3.60 M

**Answer: B**



[Watch Video Solution](#)

7. the activation energy of a reaction can be determined from the slope of which of the following graphs?

A.  $\frac{T}{\ln K} - vs \frac{1}{T}$

B.  $\ln K$  vs  $T$

C.  $\frac{\ln K}{T}$  vs  $T$

D.  $\ln K$  vs  $\frac{1}{T}$

**Answer: D**

 [Watch Video Solution](#)

8. When initial concentration of a reactant is doubled in a reaction, its half-life period is not affected. The order of the reaction is

A. More than zero but less than first

B. Zero

C. First

D. Second

**Answer: C**

 [Watch Video Solution](#)



9. A reaction having equal energies of activation for forward and reverse reactions has

A.  $\Delta G = 0$

B.  $\Delta H = 0$

C.  $\Delta H = \Delta G = \Delta S = 0$

D.  $\Delta S = 0$

**Answer: B**



**Watch Video Solution**

10. what is the activation energy for a reaction if its rate doubles when the temperature is raised from  $20^{\circ}\text{C}$  to  $35^{\circ}\text{C}$ ?

A.  $269\text{kJ mol}^{-1}$

B.  $34.7\text{ kJ mol}^{-1}$

C.  $15.1 \text{ kJ mol}^{-1}$

D.  $342 \text{ kJ mol}^{-1}$

**Answer: B**



[Watch Video Solution](#)

11. in a zero order reaction for every  $10^\circ \text{ C}$  rise of temperature , the rate is Doubled if the temperature is increased from  $10^\circ \text{ C}$  to  $100^\circ \text{ C}$  the rate of the reaction will become

A. 64 times

B. 128 times

C. 256 times

D. 512 times

**Answer: D**



[Watch Video Solution](#)

12. in a reaction  $A + B \rightarrow$  product , rate is doubled when the concentration of B is doubled and rate increases by a factor of 8 when the concentration of both the Reactants (A and B) are doubled ,Rate law for the reactions can be written as

A. Rate =  $K[A][B]$

B. Rate =  $k[A]^2[B]$

C. Rate =  $k[A][B]^2$

D. Rate =  $k[A]^2[B]^2$

**Answer: B**



[Watch Video Solution](#)

13. which one of the following statements for the order of a reaction is incorrect ?

A. Order of reaction is always whole number

B. Order can be determined only experimentally

C. Order is not influenced by stoichiometric coefficient of the reactants

D. Order of reaction is sum of power to the concentration terms of reactants to express the rate reaction

**Answer: A**



[Watch Video Solution](#)

**14.** The unit of rate constant for a first order reaction

A.  $L^2 \text{mol}^{-2} s^{-1}$

B.  $s^{-1}$

C.  $\text{mol}L^{-1} s^{-1}$

D.  $L\text{mol}^{-1} s^{-1}$

**Answer: B**



[Watch Video Solution](#)

15. The half life of a substance in a certain enzymecatalysed reaction is 138 s. The time required for the concentration of the substance to fall from  $1.28\text{mgL}^{-1}$  to  $0.04\text{mgL}^{-1}$ , is -

A. 690 s

B. 276 s

C. 414 s


D. 552 s

**Answer: A**



[Watch Video Solution](#)

16. for the reaction



the value of rate of disappearance of  $N_2O_5$  is given as

$6.25 \times 10^{-3} \text{ Mol L}^{-1} \text{ S}^{-1}$  the rate of formation of  $\text{NO}_2$  and  $\text{O}_2$  is given respectively as

A.  $6.25 \times 10^{-3} \text{ mol L}^{-1}$  &  $6.25 \times 10^{-3} \text{ mol L}^{-1} \text{ s}^{-1}$

B.  $1.25 \times 10^{-2} \text{ mol L}^{-1}$  &  $3.125 \times 10^{-3} \text{ mol L}^{-1} \text{ s}^{-1}$

C.  $6.25 \times 10^{-3} \text{ mol L}^{-1}$  &  $3.125 \times 10^{-3} \text{ mol L}^{-1} \text{ s}^{-1}$

D.  $1.25 \times 10^{-2} \text{ mol L}^{-1}$  &  $6.25 \times 10^{-3} \text{ mol L}^{-1} \text{ s}^{-1}$

**Answer: B**



**Watch Video Solution**

17. for an endothermic reaction , energy of activation is  $E_\alpha$  and Enthalpy of reaction is  $\Delta H$  ( both of these in  $\text{kJ / mol}$  ) . Minimum value of  $E_\alpha$  will be

A. Less than  $\Delta H$

B. Equal to  $\Delta H$

C. More than  $\Delta H$

D. Equal to zero

Answer: C

 Watch Video Solution

18. During the kinetic study of the reaction,  $2A + B \rightarrow C + D$ , following results were obtained

Run	$[A] / \text{molL}^{-1}$	$[B] \text{molL}^{-1}$	Initial rate formation of $D / \text{molL}^{-1} \text{mi}$
<i>I</i>	0.1	0.1	$6.0 \times 10^{-3}$
<i>II</i>	0.3	0.2	$7.2 \times 10^{-2}$
<i>III</i>	0.3	0.4	$2.88 \times 10^{-1}$
<i>IV</i>	0.4	0.1	$2.40 \times 10^{-2}$

Based on the above data which one of the following is correct ?

A. Rate =  $K[A]^2[B]$

B. Rate =  $K[A][B]$

C. Rate =  $K[A]^2[B]^2$


D. Rate =  $K[A][B]^2$

Answer: D



Watch Video Solution

19. The rate of the reaction



is given by the rate equation

$$Rate = k[NO]^2[Cl_2]$$

The value of the rate constant can be increased by

- A. Increasing the temperature
- B. Increasing the concentration of NO
- C. Increasing the concentration of the  $Cl_2$
- D. Doing all of these

**Answer: A**



Watch Video Solution



20. For the reaction ,  $N_2 + 3H_2 \rightarrow 2NH_3$  if  $\frac{d[NH_3]}{dt} = 2 \times 10^{-4} \text{ mol L}^{-1} \text{ s}^{-1}$ , the value of  $\frac{-d[H_2]}{dt}$  would be

A.  $4 \times 10^{-4} \text{ mol L}^{-1} \text{ s}^{-1}$

B.  $6 \times 10^{-4} \text{ mol L}^{-1} \text{ s}^{-1}$

C.  $1 \times 10^{-4} \text{ mol L}^{-1} \text{ s}^{-1}$


D.  $3 \times 10^{-4} \text{ mol L}^{-1} \text{ s}^{-1}$

Answer: D



Watch Video Solution

21. in the reaction ,



the rate of appearance of bromine ( $Br_2$ ) is related to rate of Disappearance of bromide ions as following .

A.  $\frac{d(Br_2)}{dt} = -\frac{5}{3} \frac{d(Br^-)}{dt}$

$$\text{B. } \frac{d(\text{Br}_2)}{dt} = \frac{5}{3} \frac{d(\text{Br}^-)}{dt}$$

$$\text{C. } \frac{d(\text{Br}_2)}{dt} = \frac{3}{5} \frac{d(\text{Br}^-)}{dt}$$

$$\text{D. } \frac{d(\text{Br}_2)}{dt} = -\frac{3}{5} \frac{d(\text{Br}^-)}{dt}$$

**Answer: D**

 [Watch Video Solution](#)

**22.** Half life period of a first - order reaction is 1396 seconds. The specific rate constant of the reaction is

A.  $0.5 \times 10^{-2} \text{ s}^{-1}$

B.  $4.9 \times 10^{-4} \text{ s}^{-1}$

C.  $5.0 \times 10^{-2} \text{ s}^{-1}$

D.  $5.0 \times 10^{-3} \text{ s}^{-1}$

**Answer: B**

 [Watch Video Solution](#)

23. for the reaction  $A + B \rightarrow$  products , it is observed that

I. On doubling the initial concentration of A only the rate of reaction is also doubled and .

II. On doubling the initial concentration of both A and B there is a change by a factor of 8 in the rate of the reaction .

the rate of this reaction is , given by

A. Rate =  $K[A][B]^2$

B. Rate =  $K[A]^2[B]^2$

C. Rate =  $k[A][B]$

D. Rate =  $K[A]^2[B]$

**Answer: A**



**Watch Video Solution**

24. the rate constants  $k_1$  and  $k_2$  for two different reactions are  $10^{16} \cdot e^{-2000/T}$  and  $10^{15} \cdot e^{-1000/T}$ , respectively the temperature at which

$k_1 = k_2$  is

A.  $\frac{1000}{2.303} K$

B. 1000 K

C.  $\frac{2000}{2.303} K$


D. 2000 K

**Answer: A**



**Watch Video Solution**

25. The bromination of acetone that occurs in acid solution is represented by this equation



These kinetic data were obtained from given reaction concentrations.

Initial concentrations, (M)

$[CH_3COCH_3]$	$[Br_2]$	$[H^+]$
0.30	0.05	0.05
0.30	0.10	0.05
0.30	0.10	0.10
0.40	0.05	0.20

Initial rate, disappearance of  $Br_2$ ,  $M_s^{-1}$

$$5.7 \times 10^{-5}$$

$$5.7 \times 10^{-5}$$

$$1.2 \times 10^{-4}$$

$$3.1 \times 10^{-4}$$

Based on these data, the rate equation is :

A. Rate =  $k[CH_3COCH_3][Br_2][H^+]$

B. Rate  $k = [CH_3COCH_3][H^+]$

C. Rate  $\frac{O}{[CH_3 - C - CH_3][Br_2]}$

D. Rate =  $k[CH_3COCH_3][Br_2][H^+]^2$

Answer: B



Watch Video Solution

26. In a first-order reaction  $A \rightarrow B$ , if  $k$  is rate constant and initial concentration of the reactant  $A$  is  $0.5\text{ M}$  then the half-life is:

A.  $\frac{\ln 2}{K}$

B.  $\frac{0.693}{0.5K}$

C.  $\frac{\log 2}{K}$


D.  $\frac{\log 2}{K\sqrt{0.5}}$

**Answer: A**




[Watch Video Solution](#)

27. the reaction of hydrogen and iodine monochloride is given as



this reaction is of first order with respect to  $H_2(g)$  and  $ICl(g)$ , following mechanisms mechanism A



Mechanism B





which of the above mechanism (S) can be consistent with the given information about the reaction ?

A. A only

B. B only

C. Both (1) and (2)

D. Neither (1) nor (2)

**Answer: B**



[Watch Video Solution](#)

**28.** If 60% of a first order reaction was completed in 60 minutes, 50% of the same reaction would be completed in approximately:

( $\log 4 = 0.60$ ,  $\log 5 = 0.69$ )

A. 40 minutes

B. 50 minutes

C. 45 minutes

D. 60 minutes

**Answer: C**

 [Watch Video Solution](#)

29. For the reaction,  $2A + B \rightarrow 3C + D$ , which of the following does not express the reaction rate

A.  $-\frac{d[C]}{3dt}$

B.  $-\frac{d[B]}{dt}$

C.  $\frac{d[D]}{dt}$

D.  $-\frac{d[A]}{2dt}$

**Answer: A**

 [Watch Video Solution](#)



30. For the reaction  $N_2 \rightarrow 3H_2 \rightarrow 2NH_3$ , how are the rate of reaction expressions inter-related  $\frac{d[H_2]}{dt}$  and  $\frac{d[NH_3]}{dt}$ ?

A.  $\frac{d[NH_3]}{dt} = -\frac{1}{3} \frac{d[H_2]}{dt}$

B.  $+\frac{d[NH_3]}{dt} = -\frac{2}{3} \frac{d[H_2]}{dt}$

C.  $+\frac{d[NH_3]}{dt} = -\frac{3}{2} \frac{d[H_2]}{dt}$

D.  $\frac{d[NH_3]}{dt} = \frac{d[H_2]}{dt}$

**Answer: B**

 [Watch Video Solution](#)

31. For a first-order reaction  $A \rightarrow B$  the reaction rate at reactant concentration of  $0.01M$  is found to be  $2.0 \times 10^{-5} \text{ mol L}^{-1} \text{ s}^{-1}$ . The half-life period of the reaction is

A. 220 s

B. 30 s

C. 300 s

D. 347 s

**Answer: D**

 [Watch Video Solution](#)

**32.** A nuclide of an alkaine earth metal undergoes radioactive decay by emission of three  $\alpha$  – particles in sucesion. The group of the periodic tablle to which the resulting daughter element would belong to:

A. Group 14

B. Group 16

C. Group 4

D. Group 6

**Answer: A**

 [Watch Video Solution](#)

33. The rate of reaction between two reactants  $A$  and  $B$  decreases by factor 4 if the concentration of reactant  $B$  is doubled. The order of this reaction with respect to  $B$  is

A.  $-1$

B.  $-2$

C.  $1$

D.  $2$

**Answer: B**



[Watch Video Solution](#)

34. A reaction is 50 % complete in 2 hours and 75 % complete in 4 hours the order of reaction is

A.  $0$

B. 1

C. 2

D. 3

**Answer: B**

 [Watch Video Solution](#)

35. Activation energy ( $E_a$ ) and rate constants ( $k_1$  and  $k_2$ ) of a chemical reaction at two different temperature ( $T_1$  and  $T_2$ ) are related by

A.  $\ln \frac{k_2}{k_1} = -\frac{E_a}{R} \left( \frac{1}{T_1} - \frac{1}{T_2} \right)$

B.  $\ln \frac{k_2}{k_1} = -\frac{E_a}{R} \left( \frac{1}{T_2} - \frac{1}{T_1} \right)$

C.  $\ln \frac{k_2}{k_1} = -\frac{E_a}{R} \left( \frac{1}{T_2} + \frac{1}{T_1} \right)$

D.  $\ln \frac{k_2}{k_1} = \frac{E_a}{R} \left( \frac{1}{T_1} - \frac{1}{T_2} \right)$

**Answer: B::D**

 [Watch Video Solution](#)


36. The half life of 2g sample of radioactive nuclide 'X' is 15 min. The half life time of 1g sample of X is

- A. 7.5 min
- B. 15 min
- C. 22.5 min
- D. 30 min

**Answer: B**

 [Watch Video Solution](#)

37. The rate of the reaction



can be written in three ways:

$$\frac{-d[N_2O_5]}{dt} = k[N_2O_5]$$
$$\frac{d[NO_2]}{dt} = k'[N_2O_5]$$

$$\frac{d[O_2]}{dt} = k''[N_2O_5]$$

The relationship between  $k$  and  $k'$  and between  $k$  and  $k''$  are-

A.  $k' = 2k, k'' = 2k$

B.  $k' = k, k'' = k$

C.  $k' = 2k, k'' = k$

D.  $k' = 2k, k'' = \frac{k}{2}$

**Answer: D**



**Watch Video Solution**

**38.** For the following reaction  $C_6H_{12}(aq) + H_2(g) \rightleftharpoons C_6H_{14}O_6(aq)$

Which one of the following is not affected by the addition of catalyst ?

A. Rate of forward reaction

B. Rate of backward reaction

C. Time required to reach the equilibrium

## D. Spontaneity

**Answer: D**

 [Watch Video Solution](#)

**39.** A chemical reaction proceeds into the following steps

Step I,  $2A \rightleftharpoons X$  fast

Step II,  $X + B \rightarrow Y$  slow

Step III,  $Y + B \rightleftharpoons$  product fast

The rate law for the overall reaction is

A. Rate =  $k[A]^2$

B. Rate =  $k[B]^2$

C. Rate =  $k[A][B]$

D. Rate =  $k[A]^2[B]$

**Answer: D**

 [Watch Video Solution](#)

40. The data for the reaction:  $A + B \xrightarrow{k} C$ .

Experiment	$[A]_0$	$[B]_0$	Initial rate
1	0.012	0.035	0.10
2	0.024	0.070	0.80
3	0.024	0.035	0.10
4	0.012	0.070	0.80

The rate law corresponding to the above data is

(a) Rate =  $k[B]^3$ , (b) Rate =  $k[B]^4$

(c) Rate =  $k[A][B]^3$ , (d) Rate =  $k[A]^2[B]^2$

A. Rate =  $k[A][B]^3$

B. Rate =  $k[A]^2[B]^2$

C. Rate =  $k[B]^3$

D. Rate =  $k[B]^4$

**Answer: C**



**Watch Video Solution**



41. Half - life for radioactive  $^{14}\text{C}$  is 5760 years. In how many years 200 mg of  $^{14}\text{C}$  will be reduced to 25 mg ?

- A. 17280 years
- B. 23040 years
- C. 5760 years
- D. 11524 years

**Answer: D**



[Watch Video Solution](#)

42. A chemical reaction is catalysed by a catalyst X. hence X.

- A. Reduces enthalpy of the reaction
- B. Does not affect equilibrium constant of reaction
- C. Decreases rate constant of the reaction
- D. Increases activation energy of the reaction

**Answer: B**

 [Watch Video Solution](#)

**43.** The given elementary reaction  $2FeCl_3 + SnCl_2 \rightarrow 2FeCl_2 + SnCl_4$

is an example of

- A. Third order reaction
- B. First order reaction
- C. Second order reaction
- D. None of these

**Answer: A**

 [Watch Video Solution](#)

**44.** Carbon 14 dating method is based on the fact that

- A. Ratio of carbon - 14 and carbon - 12 is constant
- B. Carbon - 14 is the same in all objects
- C. Carbon - 14 is highly insoluble
- D. All of these

**Answer: A**

 [Watch Video Solution](#)

**45.** For the reaction  $H_2(g) + I_2(g) \rightleftharpoons 2HI(g)$ , the rate of reaction is expressed as

- A.  $\frac{\Delta[H_2]}{\Delta t} = \frac{1}{2} \frac{\Delta[I_2]}{\Delta t} = -\frac{\Delta[HI]}{\Delta t}$
- B.  $-\frac{\Delta[I_2]}{\Delta t} = \frac{\Delta[H_2]}{\Delta t} = \frac{1}{2} \frac{\Delta[HI]}{\Delta t}$
- C.  $-\frac{\Delta[I_2]}{\Delta t} = \frac{\Delta[H_2]}{\Delta t} = \frac{\Delta[HI]}{2\Delta t}$

D. None of these

**Answer: D**

 Watch Video Solution

46. The experiment data for the reaction  $2A + B_2 \rightarrow 2AB$  is

Experiment	$[A]M$	$[B_2]M$	Initial rate( $molL^{-1}s^{-1}$ )
<i>I</i>	0.50	0.5	$1.6 \times 10^{-4}$
<i>II</i>	0.50	1.0	$3.2 \times 10^{-4}$
<i>III</i>	1.00	1.0	$3.2 \times 10^{-4}$

Write the most probable rate equation for the reaction giving reason for your answer.

A. rate =  $K[A]^2[B_2]^2$

B. rate =  $K[A]^2[B_2]$

C. rate =  $k[B_2]$

D. rate =  $k[B_2]^2$

**Answer: C**

 Watch Video Solution

47. Activation energy of chemical reaction can be determined by \_\_\_\_\_

- A. Evaluating rate constants at two different temperatures
- B. Evaluating velocities of reaction at two different temperature
- C. Evaluating rate constant at standard
- D. Changing concentration of reactants

**Answer: A**



[Watch Video Solution](#)

48. for a first -order reaction , the half - life period in independent of

- A. First power of final concentration
- B. Cube root of initial concentration
- C. Initial concentration
- D. Square root of final concentration

**Answer: C**



**Watch Video Solution**

**49.** The half - life of  ${}_{6}C^{14}$ , if its  $\lambda$  is  $2.31 \times 10^{-4} \text{ year}^{-1}$  is

A.  $3.5 \times 10^4$  years

B.  $3 \times 10^3$  years

C.  $2 \times 10^2$  years

D.  $4 \times 10^3$  years

**Answer: B**



**Watch Video Solution**

**50.** A 300 gram radioactive sample has life of 3 hour's. After 18 hours, remaining quantity will be:

A. 4.68 gram

B. 2.34 gram

C. 3.34 gram

D. 9.37 gram

**Answer: A**

 [Watch Video Solution](#)

**51.** Enzymes enhance the rate of reaction by

A. By lowering activation energy

B. By changing activation energy

C. By changing equilibrium constant

D. By forming enzyme substrate complex

**Answer: A**

 [Watch Video Solution](#)

52. For the reaction,  $2N_2O_5 \rightarrow 4NO_2 + O_2$  rate and rate constant are  $1.02 \times 10^{-4} \text{ mol L}^{-1} \text{ s}^{-1}$  and  $3.4 \times 10^{-5} \text{ s}^{-1}$  respectively. The concentration of  $N_2O_5$  in  $\text{mol L}^{-1}$  will be

A. 1.732

B. 3

C.  $1.02 \times 10^{-4}$

D.  $3.4 \times 10^5$

**Answer: B**



[Watch Video Solution](#)

53. A human body required the 0.01 M activity of radioactive substance after 24 h. Half life of radioactive substance is 6h. Then injection of maximum activity of radioactive substance that can be injected will be



A. 0.08

B. 0.04

C. 0.16

D. 0.32

**Answer: C**



**Watch Video Solution**

54. When a biochemical reaction is carried out in laboratory from outside of human body in the absence of enzyme, the rate of reaction obtained is  $10^{-6}$  times, then activation energy of the reaction in the presence of enzyme is

(a)  $\frac{6}{RT}$

(b) P is required

(c) Different forms  $E_a$  obtained in laboratory

(d) cannot say any things

A.  $\frac{6}{RT}$

B. P is required

C. Different from  $E_a$  obtained in laboratory

D. Can't say anything

**Answer: C**

 [Watch Video Solution](#)

55.  $2A + 2C \rightarrow 2B$ , rate of reaction  $\frac{+d(B)}{dt}$  is equal to

A.  $-\frac{3}{2} \frac{d(A)}{dt}$


B.  $-\frac{d(A)}{dt}$

C.  $-\frac{1}{3} \frac{d(A)}{dt}$

D.  $+2 \frac{d(A)}{dt}$

**Answer: B**

 [Watch Video Solution](#)



It would be a zero order reaction when

- A. The rate of reaction is proportional to square of conc of A
- B. The rate of reaction remains same at any conc of A
- C. The rate remains unchanged at any conc of B and C
- D. The rate of reaction doubles if conc of B is increased to double

**Answer: B**



[Watch Video Solution](#)

57. the activation energy for a simple chemical reaction  $A \rightarrow B$  is  $E_a$  in forward direction .

the activation energy for reverse reaction

- A. Is negative of  $E_a$
- B. Is always less than  $E_a$

C. Can be less than or more than  $E_a$

D. Is always double of  $E_a$

**Answer: C**

 [Watch Video Solution](#)

58. the reaction  $A \rightarrow B$  follows first order Kinetics the time taken for 0.8 mole of A to produce 0.6 mole of B is 1 H , what is the time taken for the conversion of 0.9 mole of A to 0.675mole of B?

A. 1 hour

B. 0.5 hour

C. 0.25 hour

D. 2 hour

**Answer: A**

 [Watch Video Solution](#)

59. if the rate of a reaction is equal to the rate constant , the order of the reaction is

- A. 1
- B. 2
- C. 0
- D. 3

**Answer: A**



[Watch Video Solution](#)

60. the temperature dependance of rate constant (k) of a chemical reaction is written in terms of Arrhenius equation ,  $K = Ae^{-E^*/RT}$ .  
Activation energy ( $E^*$ ) of the reaction can be calculated by plotting

- A. k Vs T

B.  $k$  Vs  $\frac{1}{\log T}$

C.  $\log k$  Vs  $\frac{1}{T}$

D.  $\log k$  Vs  $T$

**Answer: C**



**Watch Video Solution**

61. The radio isotope, tritium ( $H^3$ ) has a half life of 12.3 yr. If the initial amount of tritium is 32 mg, how many milligrams of it would remain after 49.2 yr?

A. 1 mg

B. 2 mg

C. 4 mg

D. 8 mg

**Answer: B**



Watch Video Solution

62. The rate of a first order reaction is  $1.5 \times 10^{-2} \text{ mol L}^{-1} \text{ min}^{-1}$  at 0.8 M concentration of the reactant. The half - life of the reaction is

- A. 0.383 min
- B. 36.97 min
- C. 8.73 min
- D. 7.53 min

**Answer: B**



Watch Video Solution

### Assignment Section D Assertion Reason Type Questions

1. A : Rate of reaction depends upon the concentration of the reactants.


R : The order of reaction can be negative with respect to substance

present in the reaction.

- A. If both Assertion & Reason are true and the reason is the correct explanation of the assertion , then mark (1).
- B. If both Assertion & Reason are true but the reason is not the correct explanation of the assertion , then mark (2).
- C. If Assertion is true statement but Reason is false, then mark (3).
- D. If both Assertion and Reason are false statements , then mark (4)

**Answer: B**

 [Watch Video Solution](#)



$$\text{Rate} = K[NO_2]^2$$

R : Rate does not depend upon [CO] because it is involved in the first step.



- A. If both Assertion & Reason are true and the reason is the correct explanation of the assertion , then mark (1).
- B. If both Assertion & Reason are true but the reason is not the correct explanation of the assertion , then mark (2).
- C. If Assertion is true statement but Reason is false, then mark (3).
- D. If both Assertion and Reason are false statements , then mark (4)

**Answer: D**



**Watch Video Solution**

3. A : If temperature does not affect the rate of reaction ,  $E_a = 0$

R : Lesser the activation energy, slower will be the reaction.

- A. If both Assertion & Reason are true and the reason is the correct explanation of the assertion , then mark (1).

B. If both Assertion & Reason are true but the reason is not the correct explanation of the assertion , then mark (2).

C. If Assertion is true statement but Reason is false, then mark (3).

D. If both Assertion and Reason are false statements , then mark (4)

**Answer: C**

 [Watch Video Solution](#)

4. A : The rate constant of first order reaction is used to calculate population if growth rate is given.

R : The rate constant is independent of concentration for first order reaction.

A. If both Assertion & Reason are true and the reason is the correct explanation of the assertion , then mark (1).

B. If both Assertion & Reason are true but the reason is not the correct explanation of the assertion , then mark (2).

C. If Assertion is true statement but Reason is false, then mark (3).

D. If both Assertion and Reason are false statements , then mark (4)

**Answer: B**

 [Watch Video Solution](#)

5. A : A positive catalyst increases the rate of reaction .

R : A positive catalyst alters reaction mechanism and decreases activation energy.

A. If both Assertion & Reason are true and the reason is the correct explanation of the assertion , then mark (1).

B. If both Assertion & Reason are true but the reason is not the correct explanation of the assertion , then mark (2).

C. If Assertion is true statement but Reason is false, then mark (3).

D. If both Assertion and Reason are false statements , then mark (4)

**Answer: A**

 [Watch Video Solution](#)

**6. A :** The molecularity of reaction can never be fractional.

**R :** Molecularity is the number of molecules needed to form activated complex, which will never be fractional.

- A. If both Assertion & Reason are true and the reason is the correct explanation of the assertion , then mark (1).
- B. If both Assertion & Reason are true but the reason is not the correct explanation of the assertion , then mark (2).
- C. If Assertion is true statement but Reason is false, then mark (3).
- D. If both Assertion and Reason are false statements , then mark (4)

**Answer: A**

 [Watch Video Solution](#)

7. A : The decomposition of gaseous  $N_2O_5$  follows first order kinetics.

R : The plot of log of its partial pressure versus time is linear with slope ,  
 $-\frac{k}{2.303}$  and having intercept equal to log P.

A. If both Assertion & Reason are true and the reason is the correct explanation of the assertion , then mark (1).

B. If both Assertion & Reason are true but the reason is not the correct explanation of the assertion , then mark (2).

C. If Assertion is true statement but Reason is false, then mark (3).

D. If both Assertion and Reason are false statements , then mark (4)

**Answer: A**



[Watch Video Solution](#)

8. Assertion : Complex reaction takes place in different steps and the slowest step determines the rate of reaction.

Reason : Order and molecularity of a reaction are always equal.

- A. If both Assertion & Reason are true and the reason is the correct explanation of the assertion , then mark (1).
- B. If both Assertion & Reason are true but the reason is not the correct explanation of the assertion , then mark (2).
- C. If Assertion is true statement but Reason is false, then mark (3).
- D. If both Assertion and Reason are false statements , then mark (4)

**Answer: D**

 [Watch Video Solution](#)

9. A : Rate constant increases with temperature.

R : Rate of exothermic reaction increases with temperature.

- A. If both Assertion & Reason are true and the reason is the correct explanation of the assertion , then mark (1).

B. If both Assertion & Reason are true but the reason is not the correct explanation of the assertion , then mark (2).

C. If Assertion is true statement but Reason is false, then mark (3).

D. If both Assertion and Reason are false statements , then mark (4)

**Answer: B**

 [Watch Video Solution](#)

**10. A :** Hydrolysis of ester in acidic medium follows first order kinetics.

**R :** Hydrolysis of ester is independent of the concentration of acid used.

A. If both Assertion & Reason are true and the reason is the correct explanation of the assertion , then mark (1).

B. If both Assertion & Reason are true but the reason is not the correct explanation of the assertion , then mark (2).

C. If Assertion is true statement but Reason is false, then mark (3).

D. If both Assertion and Reason are false statements , then mark (4)

**Answer: A**

 [Watch Video Solution](#)

**11. A :** The rate constant of zero order reaction is equal to rate of reaction.

**R :**  $t_{1/2}$  for zero order reaction is directly proportional to initial concentration.

- A. Both Assertion & Reason are true and the reason is the correct explanation of the assertion
- B. Both Assertion & Reason are true but the reason is not the correct explanation of the assertion
- C. Assertion is true statement but Reason is false
- D. Both Assertion and Reason are false statements

**Answer: B**



12. A : For exothermic reaction,

$$\Delta H = E_a \text{ (forward)} - E_a \text{ (backward)}$$

R : The value of activation energy for forward direction is less than activation energy for backward reaction.

A. If both Assertion & Reason are true and the reason is the correct explanation of the assertion , then mark (1).

B. If both Assertion & Reason are true but the reason is not the correct explanation of the assertion , then mark (2).

C. If Assertion is true statement but Reason is false, then mark (3).

D. If both Assertion and Reason are false statements , then mark (4)

**Answer: A**

**13. A :** Arrhenius parameter

$(A) = P$  (steric factor)  $\times z$  (collision frequency)

**R :** On increasing temperature , the value of A increases.

A. If both Assertion & Reason are true and the reason is the correct explanation of the assertion , then mark (1).

B. If both Assertion & Reason are true but the reason is not the correct explanation of the assertion , then mark (2).

C. If Assertion is true statement but Reason is false, then mark (3).

D. If both Assertion and Reason are false statements , then mark (4)

**Answer: C**



[Watch Video Solution](#)

**14. A :** In presence of +ve catalyst, activation energy & threshold energy decreases.

R : Minimum energy required to permit a reaction is known as threshold energy.

- A. If both Assertion & Reason are true and the reason is the correct explanation of the assertion , then mark (1).
- B. If both Assertion & Reason are true but the reason is not the correct explanation of the assertion , then mark (2).
- C. If Assertion is true statement but Reason is false, then mark (3).
- D. If both Assertion and Reason are false statements , then mark (4)

**Answer: B**

 [Watch Video Solution](#)

15. A : when temperature becomes  $\infty$  then the value of rate constant is maximum.

R :  $k_{\infty}$  is also known as maximum rate constant.

- A. If both Assertion & Reason are true and the reason is the correct explanation of the assertion , then mark (1).
- B. If both Assertion & Reason are true but the reason is not the correct explanation of the assertion , then mark (2).
- C. If Assertion is true statement but Reason is false, then mark (3).
- D. If both Assertion and Reason are false statements , then mark (4)

**Answer: B**



[Watch Video Solution](#)