



CHEMISTRY

AAKASH INSTITUTE ENGLISH

MOCK TEST 1

Example

1. Life Saving drug(s) used in cancer therapy are-

(a) Cisplatin ,

(b) AZT ,

(c) Taxol

A. (a) & (c)

B. (a) & (b)

C. (a), (b) & (c)

D. (b) & (c)

Answer: A



[Watch Video Solution](#)

2. Which of the following given ratio of units of length gives the highest value?

A. $\frac{1Pm}{1fm}$

B. $\frac{1hm}{1Gm}$

C. $\frac{1dm}{1\mu m}$

D. $\frac{1Mm}{1nm}$

Answer: D



3. Given the density of the chloroform is 1.510gcm^{-3} , then the volume occupied by 20.050 g of chloroform (upto correct significant figures) is

A. 13.3cm^3





B. 13.278cm^3

C. 13.28cm^3

D. 13.2780cm^3

Answer: C

4. Which of the following pair of compounds illustrate the law of multiple proportions?




Answer: A



[Watch Video Solution](#)

5. Out of the given values of temperature, which one is the highest?



B. 130°C

C. 310°F

D. 310 K

Answer: C



Watch Video Solution

6. Which of the following is/are pure substance(s)?

A. $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$

B. Brass

C. Diamond

D. All of these

Answer: C

 [Watch Video Solution](#)

7. If the masses of Cr and O are in the ratio 13 : 12 in CrO_3 , then the ratio of O that combines with the same mass of Cr in Cr_2O_3 and Cr_2O_7 is

A. 1 : 2

B. 1 : 4

C. 3 : 7

D. 2 : 5

Answer: C

 [Watch Video Solution](#)

8. CaCO_3 decomposes to give CaO and CO_2 , if the masses of CaO and CO_2 produced are 5.6 g and 4.4 g respectively by heating 12 g of an impure CaCO_3 sample then the % impurity of the sample will be

A. 33.33 %

B. 16.67 %

C. 83.33 %

D. 20 %

Answer: B

 [Watch Video Solution](#)

9. Which of the following statement is/are correct?

(i) An element of a substance contains only one kind of atoms.

- (ii) In CO_2 , carbon and oxygen chemically combined in a fixed proportion of 3 : 8 by mass.
- (iii) The constituents of the pure substances can be separated by simple physical methods.
- (iv) Milk is a homogeneous mixture.

- A. (i) and (ii)
- B. (i) , (iii) and (iv)
- C. Only (i)
- D. (i) , (ii) , (iii) and (iv)

Answer: A

 [Watch Video Solution](#)

10. The value of 3.00 km in yards upto correct significant figures and the scientific notation is, (1m = 1.094 yards)

A. 3282

B. 32.8×10^2

C. 3.28×10^3

D. 328.2×10^1

Answer: C



Watch Video Solution

11. An analytical balance has uncertainty in measurement equal to ± 1 mg. Then the result in terms of percentage would be if the weight of a compound is 10 g

A. $10 \pm 0.1 \%$

B. $10 \pm 0.01 \%$

C. $10 \pm 1 \%$

D. $10 \pm 0.001 \%$

Answer: B

 [Watch Video Solution](#)

12. A metal forms two oxides. The higher oxide contains 80% metal. 0.72g of the lower oxide gave 0.8g of higher oxide when oxidised. Calculate the ratio of weight of oxygen that combines with the fixed weight of metal in the two oxides, and show that the data supports the law of multiple proportions.

A. 2 : 3

B. 1 : 2

C. 4 : 5

D. 3 : 2

Answer: B



Watch Video Solution

13. The result of the following addition, $3.0560 + 0.0348 + 5.340$ upto correct significant figures is

A. 8.4308

B. 8.43

C. 8.431

D. 8.4

Answer: C



Watch Video Solution

14. The answer to the correct scientific notation of the following given expression $\frac{2.320}{3.694} \times 0.050$ is,

A. 0.03

B. 3.1×10^{-2}

C. 0.031

D. 0.31×10^{-1}

Answer: B



[Watch Video Solution](#)

15. Which of the following postulate is incorrect regarding Dalton's Atomic Theory?

A. Atoms are minute, indivisible and indestructible

B. All the atoms of an element are not identical to each other.

They have different masses and sizes

C. The relative numbers and kind of atoms are always the same in a given compound

D. Chemical reactions only rearrange the way in which atoms are combined , the atoms themselves are not changed

Answer: B



[Watch Video Solution](#)

16. Vapour density of a gas having formula $[CO]_x$ is 70. Find out X

A. 3.5

B. 4.5

C. 5

D. 6

Answer: C



Watch Video Solution

17. Mass of one atom of an element is 4.0×10^{-24} g. This is equal to

A. 2.4 u

B. 24 u

C. 0.024 u

D. 0.24 u

Answer: A



18. Which of the following combination of volumes (mL) of C_2H_4 (g) and O_2 (g) respectively is required to obtain 400 mL of CO_2 (g)? All volumes are measured at STP.

A. 100 , 300

B. 200 , 600

C. 200 , 200

D. 300 , 300

Answer: B

19. An element A exist in two isotopic forms A^{15} and A^{16} . If the average atomic mass of A was found to be 15.24, then the % relative abundance of A^{15} will be

A. 0.1

B. 0.2

C. 0.4

D. 0.76

Answer: D

 [Watch Video Solution](#)

20. What will be the mass of O_2 molecule in grams?

A. 5.31×10^{-23} g

B. 32 g

C. 16 g

D. 2.6×10^{-23} g

Answer: A



[Watch Video Solution](#)

21. Gram molecular mass of CH_4 is

A. 16 g

B. 16 u

C. 32 g

D. 32 u

Answer: A

22. Four one litre flasks are separately filled with the gases CO_2 , F_2 , NH_3 and He at same room temperature and pressure. The ratio of total number of atoms of these gases present in the different flasks would be

A. 1 : 1 : 1 : 1

B. 1 : 2 : 2 : 3

C. 3 : 2 : 4 : 1

D. 2 : 1 : 3 : 2

Answer: C

23. Two flasks A and B of equal capacity of volume contain SO_3 and CO gas respectively under similar conditions of temperature and pressure. Then, which of the following statement is true?

- A. A has twice the number of moles as that of B
- B. B has twice the number of moles as that of A
- C. A and B have equal number of moles
- D. B has twice the number of atoms as that of A

Answer: C

 [Watch Video Solution](#)

24. "Equal volumes of all gases at the same temperature and pressure contain equal number of particles." This statement is a direct consequence of :

- A. In accordance with Dalton's atomic theory and is known as Berzelius Hypothesis
- B. Not in accordance with Dalton's atomic theory and is known as Avogadro's Hypothesis
- C. Not in accordance with Dalton's atomic theory and is known as Berzelius Hypothesis
- D. In accordance with Dalton's atomic theory and is known as Avogadro's Hypothesis

Answer: C



Watch Video Solution

25. The vapour densities of two gases are in the ratio of 2 : 5
Their molecular masses are in the ratio of

A. 5 : 2

B. 1 : 3

C. 2 : 5

D. 3 : 1

Answer: C



Watch Video Solution

26. Atomic mass of an element is

A. Actual mass of one atom of the element

B. Relative mass of an atom of the element

C. Average relative mass of different atoms of the element

D. Always a whole number

Answer: C

 [Watch Video Solution](#)

27. A compound was found to contain 14.34% oxygen. The minimum molecular weight of the compound is

A. 111.5 g

B. 223.15 g

C. 97.62 g

D. 195.26 g

Answer: A

 [Watch Video Solution](#)

28. Loschmidt number is the number of

- A. Atoms present in 1 gram mole of a gas at STP
- B. Atoms present in 1 mL of a gas at STP
- C. Molecules present in 1 gram mole of a gas at STP
- D. Molecules present in 1 mL of a gas at STP

Answer: D



[Watch Video Solution](#)

29. 1 amu is

A. 1.66×10^{-24} kg

B. 1.66×10^{-27} kg

C. $\frac{1}{N_A}$

D. Both (2) & (3)

Answer: D

 [Watch Video Solution](#)

30. A well stoppered thermos flask contains hot water .This is an example of

- A. closed system
- B. open system
- C. isolated system
- D. non thermodynamic system

Answer: C

 [Watch Video Solution](#)

31. Incorrect relation is

A. for isothermal reversible change $W = -P_{ext}(V_f - V_i)$

B. for isothermal reversible change $Q = 2.303nRt \frac{\log V_f}{V_i}$

C. for isothermal reversible change

$$W = -2.303nRT \frac{\log P_f}{P_i}$$

D. for adiabatic change $\Delta U = W$

Answer: C

 [Watch Video Solution](#)

32. Which of the following is zero for an isochoric process?

A. ΔP

B. ΔV

C. ΔT

D. ΔE

Answer: B

 [Watch Video Solution](#)

33. Identify an intensive property among the following

A. Gibbs free energy

B. volume

C. internal energy

D. temperature

Answer: D

 [Watch Video Solution](#)

34. An adiabatic process occurs in

- A. open system
- B. closed system
- C. isolated system
- D. all of the three systems

Answer: C



Watch Video Solution

35. Which of the following is not a state function?

- A. internal energy
- B. temperature

C. work

D. enthalpy

Answer: C



Watch Video Solution

36. Which of the following is a feature of adiabatic expansion?

A. $\Delta V < 0$

B. $\Delta U < 0$

C. $\Delta U > 0$

D. $\Delta T = 0$

Answer: B



Watch Video Solution

37. Find out the correct match. a.First law of thermodynamics
b.isothermal change c.state function d.adiabatic change i.

$$\Delta U = W \text{ ii. } q = -W \text{ iii. } q = W \text{ iv. } \Delta U = q + W$$

A. a(iv),b(ii),c(iii),d(iv)

B. a(ii),b(iii),c(iv),d(i)

C. a(iv),b(i),c(ii),d(iii)

D. a(i),b(iv),c(ii),d(iii)

Answer: A



Watch Video Solution

38. In an isothermal process for an ideal gas

A. $q = 0$ and $\Delta U = 0$

B. $q \neq 0$ and $\Delta U = 0$

C. $q = 0$ and $\Delta U \neq 0$

D. $q \neq 0$ and $\Delta U \neq 0$

Answer: B



Watch Video Solution

39. The work done by a system is 8 Joule, when 40 joule heat is supplied to it .What is the increase in internal energy of the system?

A. 25J

B. 30J

C. 32J

D. 28J

Answer: C



Watch Video Solution

40. Work done during isothermal expansion of one mole of an ideal gas from 10 atm to 1atm at 300 k is

A. -4938.8J

B. 4138.8J

C. -5744.1J

D. 6257.2J

Answer: C



Watch Video Solution

41. Out of the following, choose the one which is not a part of internal energy?

- A. kinetic energy
- B. potential energy
- C. chemical bond energy
- D. gravitational energy

Answer: D



[Watch Video Solution](#)

42. Pressure volume (PV) work done by an ideal gas system at constant volume is

A. $-\frac{\Delta P}{P}$

B. Zero

C. $-\frac{V}{\Delta P}$

D. $-\Delta V$

Answer: B



Watch Video Solution

43. 6mole of an ideal gas expand isothermally and reversibly from a volume of 1 litre to a volume of 10 litre at 27° C .The maximum work is done

A. 47kj

B. 100kj

C. 0

D. 34.46kj

Answer: D



Watch Video Solution

44. An ideal gas expands against a constant external pressure of 2.0 atmosphere from 20 litre to 40 litre and absorbs 10 kJ of heat from surrounding. What is the change in internal energy of the system? (Given : 1 atm-litre = 101.3 J)

A. 4053J

B. 5948J

C. 14052J

D. 9940J

Answer: B



Watch Video Solution

45. The equation $\frac{1}{2}H_2(g) + \frac{1}{2}Cl_2 \rightarrow HCl, \Delta H^\circ = -24080 \text{ cal}$


means

- A. the heat absorbed when one gram molecule of HCL is formed from its element at 25°C is 24080 kcal
- B. the heat given out when one gram molecule of HCL is formed from its element at 298K is 24 080 kcal
- C. the heat observed when one atom of hydrogen reacts with one atom of chlorine to form one molecule of at 25°C and atmospheric pressure is 24.0 80 kcal
- D. the intrinsic heat of one molecule of HCL is 24.080kcal more than the intrinsic heat of one atom of hydrogen and one atom of chlorine

Answer: B

 [Watch Video Solution](#)

46. when 50cm^3 of $0.2\text{ N } H_3SO_4$ is mixed with 50 cm^3 of 1N KOH , the heat liberated is (Given



A. 11.46kJ

B. 57.3kJ

C. 0.563kJ

D. 0.573kJ

Answer: D

 [Watch Video Solution](#)

47. molar heat capacity of Aluminium is $25JK^{-1}mol^{-1}$ the heat necessary to raise the temperature of 54 gram of aluminium (atomic mass $27gmol^{-1}$) from $30^{\circ}C$ to $50^{\circ}C$ is

A. 1.5kj

B. 0.5kj

C. 1.0kj

D. 2.5kj

Answer: C

 [Watch Video Solution](#)

48. What will be the heat of formation of ethane, if the heat of combustion of carbon is $-xkJ$, heat of formation of water is $-ykJ$ and heat liberated during complete combustion of ethane is zkJ

A. $(-2x - 2y + z)kJ$

B. $(-2x - 3y + z)kJ$

C. $(-2x + 3y + z)kJ$

D. $(-2x - 3y - z)kJ$

Answer: B

 [Watch Video Solution](#)

49. If for H_2 gas, $C_p - C_v = a$ and for O_2 gas, $C_p - C_v = b$, where C_p and C_v is heat capacity in cal/g-k, then select the correct relation

A. $b=8a$

B. $a=b$

C. $a=16b$

D. $a=4b$

Answer: C

 [Watch Video Solution](#)

50. In a constant volume calorimeter 5g of a gas with molecular weight 40 was burnt in excess of oxygen at 298 K. the temperature of the calorimeter was found to increase from 298 K to 298.75 K due to combustion process. Given that the heat capacity of the calorimeter is 2.5 kJ K^{-1} , a numerical value for the ΔU of combustion of the gas in kJ mol^{-1} is

A. 15

B. 12


C. 90

D. 8

Answer: A

 Watch Video Solution

51. When 1 mole of oxalic acid is treated with excess of NaOH in dilute aqueous solution 108kJ of heat is liberated ,then the enthalpy of ionization of the oxalic acid is(Given



A. $4.6kJmol^{-1}$

B. $-4.6kJmol^{-1}$

C. $-6.6kJmol^{-1}$

D. $6.6kJmol^{-1}$

Answer: D



Watch Video Solution

52. The bond dissociation energies of X_2 , Y_2 and XY are in the ratio of 1:0.5:1. ΔH for the formation of XY is -200 kJ mol^{-1}

. The bond dissociation energy of X_2 will be

A. 100 kJ mol^{-1}

B. 800 kJ mol^{-1}

C. 300 kJ mol^{-1}




D. 400 kJ mol^{-1}

Answer: B



Watch Video Solution

53. The species which by definition has zero standard molar enthalpy of formation at $298K$ is



Answer: D



[Watch Video Solution](#)

54. One mole of anhydrous salt AB dissolves in water and liberates $15Jmol^{-1}$ of heat. The value of $\Delta H_{hydration}^\circ$ of AB is $-20.05 J mol^{-1}$. Hence the enthalpy of dissolution of hydrated salt $AB \cdot 3H_2O(s)$ is

A. -5.5 J mol^{-1}

B. 5.5 J mol^{-1}





C. 35.5 J mol^{-1}

D. -35.5 J mol^{-1}

Answer: B

 [Watch Video Solution](#)

55. choose the reaction in which ΔH is not equal to ΔU



Answer: B

 [Watch Video Solution](#)

56. Hess's law is application for the determination of heat of

- A. reaction
- B. formation
- C. transition
- D. All of these

Answer: D

 [Watch Video Solution](#)

57. Enthalpy (H) is equal to

- A. Internal energy (E)
- B. Product of pressure (P) and volume (V) of gas
- C. Internal energy (E)+PV(work)
- D. work (W) done by a system

Answer: C

 [Watch Video Solution](#)

58. If $\Delta_f H^\circ(C_2H_4)$ and $\Delta_f H^\circ(C_2H_6)$ are x_1 and x_2 kcal mol⁻¹, then heat of hydrogenation of C_2H_4 is :

- A. $x_1 + x_2$
- B. $x_1 - x_2$
- C. $x_2 - x_1$
- D. $x_1 + 2x_2$

Answer: C

 [Watch Video Solution](#)

59. heat of combustion of gaseous compounds A(molar mass =16) B(molar mass=28) C(molar mass=30) and D(molar mass=40) are -890,-1411,-1560 and -1900kJ/mole respectively. Which has the highest calorific fuel(J/g)?

A. A

B. B

C. C

D. D

Answer: A

 [Watch Video Solution](#)

60. which of the following case entropy increases

- A. boiling of an egg
- B. crystallization of sugar from solution
- C. freezing of water
- D. stretching of rubber

Answer: A



[Watch Video Solution](#)

61. if the reaction is reversible all is at equilibrium ($\Delta S_{universe} = 0$), then the entropy of the system

- A. will change abruptly

B. is greater than 1

C. will remain constant

D. is equal to 0

Answer: C

 [Watch Video Solution](#)

62. State and explain the second law of thermodynamics.

Mention the essential conditions for the spontaneity of a chemical reaction.

A. in any spontaneous process entropy of the universe always

increases

B. energy can neither be created nor be destroyed

C. energy of the universe remains constant

D. $\Delta S_{\text{universe}} > 0$ for a spontaneous reaction

Answer: A

 [Watch Video Solution](#)

63. ΔH and ΔS are positive for a chemical reaction. Under what conditions is the reaction expected to occur spontaneously?

A. $\Delta H < T\Delta S$

B. $T\Delta S > \Delta H$

C. $\Delta H = T\Delta S$

D. $\Delta G > 0$

Answer: B

 [Watch Video Solution](#)

64. the incorrect expression among the following is

A. $K = e^{-\Delta \frac{G^\circ}{R} T}$

B. $\ln K = \frac{\Delta H^\circ - T\Delta S^\circ}{R} T$

C. $\Delta S_\Sigma = -\Delta H_s y \frac{s}{T}$

D. $\Delta S_s y s = q_r e \frac{v}{T}$

Answer: B



Watch Video Solution

65. ΔG° for the following reaction:



Given that $\Delta_f G^\circ HI(g) = 1.8 \text{ kJ mol}^{-1}$. $\Delta_f G^\circ H_2S(g) = 33.8 \text{ kJ mol}^{-1}$.

A. 30200 kJ

B. -30.2 kJ

C. -30200 J

D. -302 J

Answer: C



[Watch Video Solution](#)

66. A process A \rightarrow D is difficult to occur directly instead it takes place in three successive steps, $\Delta S(A \rightarrow B) = 40$ e.u., $\Delta S(B \rightarrow C) = 30$ e.u., $\Delta S(D \rightarrow C) = 20$ e.u. where e.u. is entropy unit then the entropy change ΔS for the process (A \rightarrow D) is

A. + 90 e.u.

B. + 50 e.u.

C. -90 e.u.

D. - 50 e.u.

Answer: B



Watch Video Solution

67. the entropy possessed by certain substances at absolute zero is known as

A. Residual entropy

B. positive entropy

C. negative entropy




D. excess entropy

Answer: A




Watch Video Solution

68. The following equilibrium are given below,



The equilibrium constant of the reaction



A. $K_1 \frac{K_2}{K_3}$

B. $K_1 \frac{K_3^2}{K_2}$

C. $K_2 \frac{K_3^3}{K_1}$

D. $K_1 K_2 K_3$

Answer: C

 [Watch Video Solution](#)

69. consider the given reaction, $3A(g) + B(g) \rightleftharpoons 2C(g)$ at a given temperature if a mixture of 2 mol each of A, B and C exist at equilibrium and $K_c = 9$ then volume of the flask will be

- A. 3L
- B. 6L
- C. 9L
- D. 36L

Answer: B

 [Watch Video Solution](#)

70. In a chemical reaction, equilibrium is said to have been established when the

- A. backward and forward reaction ceases
- B. concentration of reactants and products are equal
- C. rate of backward reaction is equal to the rate of forward reaction
- D. reaction ceases to generate heat

Answer: C

 [Watch Video Solution](#)

71. consider the reaction $2A_g + B_g \rightleftharpoons 2C_g$ for which $K_c = 350$.
if 0.001` mole of each of the reactant and product are mix in a 2.0

L flux in the reaction quotient and spontaneous direction of the system will be

- A. $Q_c = 0.002$, the equilibrium shifts to the left
- B. $Q_c = 2000$, the equilibrium shifts to the left
- C. $Q_c = 0.002$, the equilibrium shifts to right
- D. $Q_c = 2000$, the equilibrium shifts to right

Answer: B



[Watch Video Solution](#)

72. for the reaction $P + Q \rightleftharpoons R + 2S$, initially the concentration of P is equal to that of Q (1 molar) but at equilibrium the concentration of R will be twice of that of P, then the equilibrium constant of the reaction is

A. $\frac{4}{3}$

B. $\frac{32}{3}$

C. $\frac{3}{10}$

D. $\frac{1}{10}$

Answer: B



Watch Video Solution

73. stage comes when no more sugar dissolves, instead it settles down at the bottom of the solution is now said to be

A. condensed

B. in a state of equilibrium

C. saturated

D. both (2) & (3)

Answer: D

 [Watch Video Solution](#)

74. Which of the following is non-polar molecular solid?

A. SiC

B. Naphthlene

C. HCl

D. AlN

Answer: B

 [Watch Video Solution](#)

75. Solids for which physical properties like electric resistance or refractive index show different values when measured along different directions are called

- A. Pseudo solids
- B. Isotropic solids
- C. Polymorphic solids
- D. Anisotropic solids

Answer: D

 [Watch Video Solution](#)

76. Which of the following given list has molecular solids only?

- A. Carborundum, dry ice, diamond, solid H₂, benzene

B. Naphthalene, Na_2SO_4 copper, corundum, CI_4

C. Corundum, camphor, silicon carbide, ice, solid CS_2

D. Solid H_2 , camphor, dry ice, *solid* CS_2 , naphthalene

Answer: D



Watch Video Solution

77. Match the crystal system/unit cells mentioned in Column I with their characteristic features mentioned in Column II.

Column I		Column II	
(A)	simple cubic and face-centred cubic	(p)	have these cell parameters $a = b = c$ and $\alpha = \beta = \gamma$
(B)	cubic and rhombohedral	(q)	are two crystal systems
(C)	cubic and tetragonal	(r)	have only two crystallographic angles of 90°
(D)	hexagonal and monoclinic	(s)	belong to same crystal system

A. a(iii), b(iv), c(ii), d(i)

B. a(ii) b(iv), c(i), d(i)

C. a(i), b(v), c(iv), d(ii)

D. a(v), b(iii), c(ii), d(i)

Answer: B



Watch Video Solution

78. If all the three interaxial angles defining the unit cell are equal in magnitude, the crystal cannot belong to

(I) Orthorhombic system

(II) Monoclinic system

(III) Hexagonal system

(IV) Tetragonal system

A. II, III

B. I, IV

C. III, IV

D. I, II

Answer: A

 [Watch Video Solution](#)

79. Which of the following sets of axial angles and axial lengths represent maximum number in Bravais lattices?

A. $\alpha = \beta = \gamma = 90^\circ$ and $a = b \neq c$

B. $\alpha = \beta = \gamma = 90^\circ$ and $a \neq b \neq c$

C. $\alpha = \beta = \gamma \neq 90^\circ$ and $a = b = c$

D. $\alpha = \beta = \gamma = 90^\circ$ and $a = b = c$

Answer: B



 [Watch Video Solution](#)

80. Which of the following given crystal system is the most symmetrical and the most unsymmetrical system respectively?

- A. Cubic, Hexagonal
- B. Orthorhombic, Monoclinic
- C. Cubic, Tridinic
- D. Rhombohedral, Tetragonal

Answer: C

 [Watch Video Solution](#)

81. Sodium crystallizes in a face centred cubic lattice. The approximate number of unit cells in 5.0 g of sodium is (Atomic

mass of sodium = 23 mu)

A. $32.7 \times 10_{22}$

B. $3.27 \times 10_{(22)}$

C. $6.54 \times 10_{(22)}$

D. $65.4 \times 10_{22}$




Answer: B



Watch Video Solution

82. Three atoms A, B and C crystallize in a cubic solid lattice where A atoms are present at the body centre, B atoms are present at the edge centre as well as at the corners of the cube and C atoms are present at the face centres of the cube. Now if all the atoms are removed from the two 4-fold axis and the one





2-fold axis passing through the cube, then the formula of the compound is



Answer: A

 [Watch Video Solution](#)


83. Three atoms P, Q and R crystallize in a cubic solid lattice where P atoms are at the alternate faces, R atoms are at the centre of edges and Q atoms are at the $\frac{2}{3}$ rd of the total corners present, hence the formula of the compound is




Answer: B

 [Watch Video Solution](#)

84. A compound formed by elements X, Y and Z has a cubic structure in which X atoms are at the corner of the cube and also at alternate face centres. Y atoms are present at the body centre and Z atoms are present at the alternate edge centre. Then the molecular formula of the compound is









Answer: D



Watch Video Solution

85. An ionic compound is made up of A & B only. Ions A occupy all the corners and alternate edge centers while atoms B occupy all the face centers. The formula of compound will be



Answer: B



Watch Video Solution

86. An ionic compound is made up of A & B only. Ions of A occupy all the corners and alternate face centers while that of B occupy body center and edge centers. If B contains -1 charge then charge on atom A will be

A. +1

B. +2

C. +3

D. +4

Answer: B



Watch Video Solution

87. Which one of the following schemes of ordering closed packed sheets of equal sized spheres do not generate closet packed lattice?

A. ABCABC

B. ABACABAC

C. ABBAABBA

D. ABCBCABCBC





Answer: C



[Watch Video Solution](#)

88. If the anions (X) form hexagonal closed packing and cations (Y) occupy only $\frac{3}{8}$ th of octahedral voids in it, then the general


formula of the compound is






Answer: C

 [Watch Video Solution](#)

89. A solid is formed and it has three types of atoms X, Y, Z. X forms an FCC lattice with Y atoms occupying one-fourth of tetrahedral voids and Z atoms occupying half of the octahedral voids. The formula of the solid is





Answer: D



Watch Video Solution

90. The two ions A^+ and B^- have radii 40 pm and 120 pm respectively. In the closed packed crystal of compound AB, the coordination number of A^+ would be

A. 6

B. 8

C. 4




D. 12

Answer: C



Watch Video Solution

91. A crystal is made of particles X, Y and Z. X forms fcc packing. Y occupies all the octahedral voids of X and Z occupies all the tetrahedral voids of X. If all the particles along one body diagonal are removed then the formula of the crystal would be:



Answer: D



Watch Video Solution

92. The number of nearest neighbours of each atom in cubic close packing (ccp) and body-centred cubic arrangement (bcc) is respectively

A. 12, 12

B. 12, 8

C. 8, 6

D. 8, 8

Answer: B



[Watch Video Solution](#)

93. Minimum distance between two tetrahedral voids if a is the edge length of the cube is

A. $\frac{a}{4}$

B. $\frac{a}{2\sqrt{2}}$

C. $\frac{a}{2}$

D. $\frac{\sqrt{3}a}{4}$

Answer: C



Watch Video Solution

94. The distance between an octahedral and tetrahedral void in fcc lattice would be:

A. $a\sqrt{3}$

B. $\frac{a\sqrt{3}}{2}$

C. $\frac{a\sqrt{3}}{3}$

D. $\frac{a\sqrt{3}}{4}$

Answer: D

 [Watch Video Solution](#)

95. You are given 6 identical balls . The maximum number of square voids and triangular voids (in separate arrangements) that can be created respectively are

A. 2, 4

B. 4, 2

C. 4, 3

D. 3, 4

Answer: A

 [Watch Video Solution](#)

96. The number of octahedral voids in case of hcp unit cell is

A. 6

B. 12

C. 4

D. 8

Answer: A



Watch Video Solution

97. The number of nearest neighbours of each sphere in hexagonal closed packing pattern in its own layer will be

A. 4

B. 6

C. 12

D. 8

Answer: B



Watch Video Solution

98. In an arrangement of type ABABA identical atoms of first layer (A) and third layer (A) are joined by a line passing through their centers . Identify the correct statement .

A. No void is found on the line

B. Only tetrahedral voids are found on the line

C. Only octahedral voids are found on the line

D. Equal number of tetrahedral and octahedral voids are found on the line

Answer: B

 [Watch Video Solution](#)

99. Given an alloy of Cu, Ag and Au in which Cu atoms constitute the ccp arrangement . If the hypothetical formula of the alloy is Cu_4Ag_3Au , the probable locations of Ag and Au atoms are

- A. Ag- all tetrahedral voids , Au - all octahedral voids
- B. Ag- $\frac{3}{8}$ th tetrahedral voids , Au - $\frac{1}{4}$ th octahedral voids
- C. Ag- 1/2 octahedral voids , Au - 1/2 tetrahedral voids
- D. Ag- all octahedral voids , Au - all tetrahedral voids

Answer: B

 [Watch Video Solution](#)

100. Which of the following statement is false ?

- A. Two tetrahedral voids are formed on each of the four body diagonals of the cube
- B. When body centre of the cube is surrounded by six atoms of face centres , an octahedral voids is formed .
- C. Tetrahedral void is present at the centre of each of the 12 edges
- D. The shortest distance between two octahedral voids is $\frac{a}{\sqrt{2}}$ (a is the edge length of the unit cell).

Answer: C



Watch Video Solution

101. A TV in fcc is formed by atoms at

- A. 3 corners and 1 face centre
- B. 3 face centres and 1 corner
- C. 2 face centres and 2 corners
- D. 2 face centres , 1 corner and 1 body centre

Answer: B



[Watch Video Solution](#)

102. Relationship between atomic radius and the edge length a of a body-centred cubic unit cell is

A. $r = \frac{a}{2}$

B. $r = \frac{a}{2\sqrt{2}}$

$$C. r = \frac{a\sqrt{3}}{4}$$

$$D. \frac{\sqrt{2}a}{2}$$

Answer: B



Watch Video Solution

103. The fraction of the total volume occupied by the atoms present in a simple cube is

$$A. \frac{\pi}{2}$$

$$B. \frac{\sqrt{3}\pi}{8}$$





$$C. (\sqrt{2}\pi)6$$

$$D. \frac{\pi}{6}$$

Answer: D



104. In a close packed structure of mixed oxides , the lattice is composed of oxide ions, one-eighth of tetrahedral voids are occupied by divalent cations while one-half of octahedral voids are occupied by trivalent cations. The formula of the oxide is



Answer: C

105. An ionic solid $A^{\oplus}B^{\ominus}$ crystallizes as an bcc structure. The distance between cation and anion in the lattice is $338 \pm$. The edge length of cell is

A. 195.15 pm

B. 97.58 pm

C. 390.3 pm

D. 780.6 pm

Answer: C

 [Watch Video Solution](#)

106. In a metal M having bcc arrangement edge length of the unit cell is 400 pm . The atomic radius of the metal is

A. 173 pm

B. 100 pm

C. 141 pm

D. 200 pm

Answer: A



Watch Video Solution

107. A compound XY crystallizes in BCC lattice with unit cell - edge length of 480 pm , if the radius of Y is 225 pm , then the radius of X is

A. 95.34 pm

B. 225 pm

C. 127.5 pm

D. 190.7 pm

Answer: D



Watch Video Solution

108. What are the number of atoms per unit cell and the number of nearest neighbours in a body centered cubic structure?

A. 2, 12

B. 4, 12

C. 2, 8

D. 2, 6

Answer: C



Watch Video Solution

109. An element crystallizes in a face centered cubic lattice and the edge of the unit cell is 0.559nm. The density is $3.19\text{g}/\text{cm}^3$.

What is the atomic mass?

A. 100.6

B. 75.9

C. 95.8

D. 83.9

Answer: D

 [Watch Video Solution](#)

110. An element X (At. mass = $80\text{g}/\text{mol}$) has fcc structure.

Calculate no. of unit cell in 8gm of X:

A. $0.4 \cdot N_A$

B. $0.1 \cdot N_A$

C. $4 \cdot N_A$

D. $0.025 \cdot N_A$

Answer: D



Watch Video Solution

111. Molybdenum (At. mass = 96 g/mol^{-1}) crystallizes as bcc crystal. If density of crystal is 10.3 g/cm^3 , then radius of Mo atoms (use $N_A = 6 \times 10^{23}$):

A. 111 pm

B. 314 pm

C. 138.56 pm

D. 314 pm

Answer: C

 [Watch Video Solution](#)

112. The coordination number of metal crystallising in a hexagonal close packing is

A. 12

B. 8

C. 6

D. 4

Answer: A

 [Watch Video Solution](#)

113. The unit cell present in ABCABC, closet packing of atoms is:

- A. Hexagonal
- B. Tetragonal
- C. Face centred cube
- D. Simple cube

Answer: C



[Watch Video Solution](#)

114. The atomic radius of strontium (Sr) is 215pm and it crystallizes with a cubic close packing . Edge length of the cube is :

- A. 430 pm

B. 608.2 pm

C. 496.53 pm

D. 304.1 pm

Answer: B



[Watch Video Solution](#)

115. Crystalline CsCl has density 3.988 g cm^{-3} . The volume occupied by single CsCl ion pair in the crystal will be

A. $7.01 \cdot 10^{-23} \text{ cm}^3$

B. $6.02 \cdot 10^{-24} \text{ cm}^3$

C. 1 cm^3

D. $3.5 \cdot 10^{-23} \text{ cm}^3$

Answer: A

 [Watch Video Solution](#)

116. A binary solid (AB) has a rock salt structure . If the edge length is 500 pm , and radius of cation is 80 pm , find the radius of anion

- A. 100 pm
- B. 120 pm
- C. 250 pm
- D. 170 pm

Answer: D

 [Watch Video Solution](#)

117. The two ions A^{+} and B^{-} have radii 85 and 200 pm respectively. In the closed packed crystal of compound AB, the coordination number of A^{+} ion is

A. 3

B. 4

C. 6

D. 8

Answer: C



[Watch Video Solution](#)

118. Caesium and chloride ions are in contact along the body diagonal in a body-centred cubic lattice. The edge length of the

unit cell is 350 pm and Cs^+ has a radius of 133pm. Hence, the radius of Cl^- ion is approximately

A. 170

B. 133

C. 180

D. 150

Answer: A



[Watch Video Solution](#)

119. Which of the following given statement(s) is/are correct for both fluorite and antiferite structures? (i) coordination number of cation is 8 (ii) Number of formula unit is one unit cell is 4 (iii) 100% tetrahedral voids are occupied (iv) Radius ratio of cation and anion is 0.20

A. (i) and (ii)

B. (i), (ii) & (iii)

C. (ii) and (iii)

D. (i), (ii), (iii) & (iv)

Answer: C



[Watch Video Solution](#)

120. A solid AB has ZnS-type structure. The edge length of unit cell is 400 pm and the radius of B^{-} ion is 0.130 nm. Then the radius of A^{+} ion is

A. 35.8 pm

B. 43.2 pm

C. 60.5 pm

D. 53.2 pm

Answer: B



[Watch Video Solution](#)

121. Which of the following is correct?

- A. AgBr shows both Schottky and Frankel defect
- B. Frenkel defect is shown by ionic solids where there is large difference in size of anion and cation
- C. In Frankel defect, dielectric constant of crystal increases
- D. All are correct

Answer: D



[Watch Video Solution](#)

122. Antiferromagnetic substance possess:

- A. Low magnetic moment
- B. Large magnetic moment
- C. Zero magnetic moment
- D. Non-zero value of magnetic moment

Answer: C



[Watch Video Solution](#)

123. $MgFe_2O_4$ has spinel structure , then the percentge of tetrahedral voids and the octahedral voids occupied are respectively

- A. 25% & 37.5%

B. 12.5% & 50%

C. 25% & 25%

D. 37.5% & 25%

Answer: B



Watch Video Solution

124. If an ionic solid XY (X & Y are monovalent ions) is doped with 10^{-2} moles % of another ionic solid AY_3 , then the concentration of the cation vacancies created is

A. $6.023 \times 10^{19} \text{ mol}^{-1}$

B. '60.23xx10^18 mol^-1'

C. '12.05xx10^21 mol^-1'

D. $1.205 \times 10^{21} \text{ mol}^{-1}$

Answer: D

 [Watch Video Solution](#)

125. Which of the following given statements is incorrect?

- A. F-centres generation is responsible factor for imparting colour to the crystal
- B. Frenkel defect is usually shown by ionic compounds having low coordination number
- C. Stoichiometry of crystal remains unaffected due to Schottky defect
- D. Density of crystal always increases due to substitutional impurity defect

Answer: D



Watch Video Solution

126. Which of the following given statement for semiconductor is correct?





- A. p-type semiconductor is formed by doping Si with B
- B. p-type semiconductor is formed by doping Si with P
- C. n-type semiconductor is formed by doping Si with Al
- D. n-type semiconductor is formed by doping Ge with B

Answer: A



Watch Video Solution

127. Which of the following given molecules in a pair are paramagnetic and ferrimagnetic substance respectively?



Answer: C

 [Watch Video Solution](#)

128. In Na_2O structure

A. O^{2-} ions constitute CCP and Na^+ ions occupy all the octahedral holes

B. O^{2-} ions constitute CCP and Na^+ ions occupy all the tetrahedral holes

C. O^{2-} ions constitute CCP and Na^+ ions occupy 50% of tetrahedral holes and 100% octahedral holes

D. Na^+ ions constitute CCP and O^{2-} ions occupy half of octahedral holes

Answer: B



[Watch Video Solution](#)

129. When a crystal structure of NaCl type is pressurised

A. The coordination number is decreased to 8 and converted to CsCl type crystal structure

B. The coordination number is remains same

C. The coordination number is increased to 8 and converted to CsCl type crystal structure

D. The coordination number is increased to 4 and converted to ZnS type crystal structure

Answer: C

 [Watch Video Solution](#)

130. Ferromagnetism arises because of the spontaneous alignment of the magnetic moments due to unpaired electrons

as

A. $\uparrow \uparrow \uparrow \uparrow \uparrow$

B. $\uparrow \uparrow \uparrow \downarrow \downarrow$

C. $\uparrow \downarrow \uparrow \downarrow$

D. $\uparrow \uparrow \downarrow \uparrow \uparrow$

Answer: A



Watch Video Solution

131. Minimum distance between two tetrahedral voids if a is the edge length of the cube is

A. $\frac{\sqrt{3}a}{4}$

B. $\frac{a}{2}$

C. $\frac{a}{2 \times \sqrt{2}}$

D. $\frac{\sqrt{2}a}{2}$

Answer: B



 Watch Video Solution

132. 2 M of 100 mL Na_2SO_4 is mixed with 3 M of 100 mL NaCl solution and 1 M of 200 mL $CaCl_2$ solution . Then the ratio of the concentration of cation and anion is

A. 1 : 1

B. 2:1

C. 2:3

D. 1 : 2

Answer: A

 Watch Video Solution

133. Among the following , select the pair that does not form an ideal solution


- A. Carbon tetrachloride and Silicon tetrachloride
- B. Chlorobenzene and Bromobenzene
- C. Chloroform and Carbon tetrachloride
- D. Benzene and toluene

Answer: C




Watch Video Solution


134. Which of the following graphs is correct for solubility of O_2 and N_2 in water at 298K.




A.



B.



C.



D.

Answer: A

 [Watch Video Solution](#)

135. State True or False Schottky defects lower the density of related solids

 [Watch Video Solution](#)

136. The volume of water added to 500 mL , 0.5 M NaOH so that its strength becomes 10mg NaOH per mL is

A. 250 mL

B. 500 mL

C. 750 mL

D. 1000 mL

Answer: B



Watch Video Solution

137. Amount (in g) of sample containing 80% NaOH required to prepare 60 litre of 0.5 M solution is

A. 1000

B. 1200

C. 1500

D. 1600

Answer: C

 [Watch Video Solution](#)

138. Henry's law is not valid when

- A. Temperature is high
- B. Pressure is low
- C. The gas is not highly soluble
- D. The gas neither reacts chemically with solvent nor dissociates or associates in the solvent

Answer: A

 [Watch Video Solution](#)

139. Number of Faraday's required to generate one gram atom of calcium from molten $CaCl_2$ is

A. 1

B. 2

C. 3

D. 4

Answer: B



[Watch Video Solution](#)

140. For the production of X L H_2 at STP at cathode, cost of electricity is x then cost of production of X L O_2 at STP at anode from water will be

A. x

B. $\frac{x}{2}$

C. $2x$

D. $4x$

Answer: C



Watch Video Solution

141. When one coulomb of electricity is passed through an electrolytic solution, the mass of the element deposited on the electrode is equal to

A. Molecular weight

B. equivalent weight

C. one gram

D. Electrochemical equivalent

Answer: D

 [Watch Video Solution](#)



142. When an aqueous solution of $AgNO_3$ is electrolysed between platinum electrodes, the substances liberated at anode and cathode are

- A. Cu at anode and Ag at cathode
- B. O_2 at anode and Cu at cathode
- C. O_2 at anode and Ag at cathode
- D. NO_2 at anode and Ag at cathode



Answer: C

 [Watch Video Solution](#)

143. Which of the following reaction is not involved in corrosion of iron?



C.



Answer: B



Watch Video Solution

144. During electrolysis of $H_2SO_4(aq)$ with high charge density, $H_2S_2O_8$ is formed as by product. In such electrolysis 44.8 L $H_2(g)$ and 15 L $O_2(g)$ are liberated at STP. Hence, the moles of $H_2S_2O_8$ formed is approximately equal to

A. 0.25

B. 0.66

C. 2

D. 2.68

Answer: B



Watch Video Solution

145. When a lead storage battery is discharged then incorrect option(s) is/are

- A. only 1
- B. only 1 & 2
- C. 1, 2, & 3
- D. Only 4

Answer: C

 [Watch Video Solution](#)

146. Three moles of electrons are passed through three solutions in succession containing $AgNO_3$, $CuSO_4$ and $AuCl_3$ respectively the molar ratio of amounts of cations reduced at cathode will be

- A. 1 : 2 : 3
- B. 3 : 2 : 1

C. 2: 1: 3

D. 6: 3: 2

Answer: D



[Watch Video Solution](#)

147. The Zn acts as sacrificial or cathodic protection to prevent rusting of iron because

A. E_{op} of Zn = E_{op} of Fe`

B. E_{op} of Zn > E_{op} of Fe


C. E_{op} of Zn < E_{op} of Fe

D. Zn dose mot react with water

Answer: B



148. An aqueous solution of $NaCl$ on electrolysis gives $H_2(g)$, $Cl_2(g)$, and $NaOH$ according to the reaction :



A direct current of $25A$ with a current efficiency of 62% is passed through $20L$ of $NaCl$ solution (20% by weight). Write down the reactions taking place at the anode and cathode. How long will it take to produce $1kg$ of Cl_2 ? (Assume no loss due to evaporation .)

A. 48.71 hr, 1041M

B. 2880 min, 1041M

C. 17.54 hr, 2M

D. 170.54 min, 2M

Answer: A



[Watch Video Solution](#)

149. When an electric current is passed through a cell having an electrolyte, then the cations and anions move to their respective electrodes if the cathode is pulled out of the solution then

- A. Both cations and anions will move towards anode
- B. cations will start moving towards anode while anions will stop moving
- C. Anions will continue to move towards anode while cations will stop moving
- D. Both cations and anions will starts moving randomly

Answer: D



[Watch Video Solution](#)

150. Which of the following given batteries are rechargeable?

1. Dry-cell battery

2. Nickel-cadmium battery

3. Lithium battery

4. Fuel cell

5. Lead storage battery

A. 1, 2 & 4

B. 2, 3 & 5

C. 1, 2, 4 & 5

D. 2, 4 & 5

Answer: B



Watch Video Solution



151. Define Fuel cell

 Watch Video Solution

152. A first order reaction is found to have a rate constant $k = 11 \times 10^{-14} \text{ s}^{-1}$. Find the half life of the reaction.

 Watch Video Solution

153. When an acidified solution of Na_2MoO_n (atomic mass of Mo=36) is electrolyzed, O_2 gas is liberated corresponding to a volume of 0.112 L at STP and mass of MO deposited is 0.32 g. Then the formula of the salt and oxidation state of Mo is



C. Na_2MoO_2 , +2

D. Na_2MoO_3 , +4

Answer: B

 [Watch Video Solution](#)

Exercise

1. For an equilibrium reaction, if the value of standard Gibb's free energy, AG° is zero, then the value of equilibrium constant, K will be equal to

A. Zero

B. 2

C. 1

D. 10

Answer: C

 Watch Video Solution

2. The equilibrium constant k_p for the reaction
 $2NOCl(g) \rightleftharpoons 2NO(g) + Cl_2(g)$ is $6.95 \cdot 10^{-8}$ at 298K . The standard gibb free energy change ΔG^0 at 298 K will be ($R = 2calK^{-1}mol^{-1}$) (log 6.95 = 0.8420)

A. 4.912 kcal

B. 14.74 kcal

C. 7.3 kcal


D. 9.825kcal

Answer: D



[Watch Video Solution](#)

3. The yield of product in the reaction,



would be higher at:

- A. High temperature and high pressure
- B. High temperature and low pressure
- C. Low temperature and high pressure
- D. Low temperature and low pressure

Answer: C



[Watch Video Solution](#)

4. A chemical reaction is catalysed by a catalyst X. hence X.

- A. Changes the equilibrium constant of the reaction
- B. Changes the enthalpy of reaction (ΔH)
- C. Alters the concentration of both reactants and products in a state of equilibrium
- D. Increases the speed of both the forward and backward reactions to same extent in a reversible

Answer: D

 [Watch Video Solution](#)

5. Which one of the equation is correct ?

A. $\Delta G = \Delta G^0 + nRT \log Q$

B. $\Delta G^0 = \Delta G + nRT \log Q$


$$C. \Delta G = \Delta G^0 + nRT \ln Q$$

$$D. \Delta G^0 = \Delta G + nRT \ln Q$$

Answer: C

 [Watch Video Solution](#)

6. The yield of product in the reaction,



would be higher at:

A. 4.24

B. 2.12




C. 42.4

D. 8.48

Answer: A

 [Watch Video Solution](#)

7. In which of the following reaction, the formation of product is favoured by increase in pressure?



Answer: B

 [Watch Video Solution](#)

8. Consider the following reaction at equilibrium: $\text{NH}_4\text{HS}(s)$


- A. Equilibrium shifts in the backward direction
- B. Equilibrium shifts in the forward direction
- C. Equilibrium remains unaffected
- D. The value of K is increased

Answer: C



[Watch Video Solution](#)

9. Ammonia is a weak base that reacts with water according to the equation



Select the correct option (s) that can increase the moles of ammonium ion in water:

- A. Addition of HCl
- B. Addition of H₂O
- C. Addition of NaOH
- D. Addition of NH₄Cl

Answer: C



[Watch Video Solution](#)





10. The equilibrium, $BaCO_3(s) \rightleftharpoons BaO(s) + CO_2(g)$ will shift in left hand direction by

- A. Addition of BaO(s)
- B. Removal of CO₂(g)
- C. Removal of BaO(s)
- D. Decreasing the volume of the vessel

Answer: D

 [Watch Video Solution](#)




11. Le - Chatelier principle is not applicable to :



Answer: C

 [Watch Video Solution](#)

12. In which of the following reactions, increase in the pressure at constant temperature does not affect the moles at equilibrium?



Answer: B

 [Watch Video Solution](#)

13. For the gas phase exothermic reaction, $A(g) + 2B(g)$

A. Decreasing the temperature

B. Increasing the pressure


C. Adding inert gas at constant pressure

D. Removing C(g) at equilibrium

Answer: C

 [Watch Video Solution](#)

14. For the reaction



at a given temperature, the equilibrium amount of $CO_2(g)$ can be increased by

A. Increasing the pressure

B. Adding an inert gas at constant pressure

C. Increasing the volume of the container

D. Increasing the amount of CO(g)

Answer: D

 [Watch Video Solution](#)


15. Which among the following is a Lewis acid?



Answer: B

 [Watch Video Solution](#)

16. The species which can act both as Bronsted acid and base is



Answer: C



Watch Video Solution

17. An example of a strong electrolyte is

A. Glucose

B. Urea

C. Ammonium hydroxide

D. Sodium formate

Answer: D

 [Watch Video Solution](#)

18. For a weak acid HA of concentration C (moll^{-1}) and degree of dissociation (α), Ostwald's dilution law is represented by the equation

A. $K_a = \frac{C^2\alpha}{1 - \alpha}$

B. $K_a = \frac{\alpha^2 C}{1 - \alpha}$

C. $K_a = C\alpha$

D. $K_a = \frac{C\alpha^2}{1 - \alpha^2}$

Answer: B

 [Watch Video Solution](#)

19. In the given irreversible reaction,
 $H_2O + HCl \rightarrow H_3O^+ + Cl^-$ the species that acts as
Bronsted base is

- A. H_2O
- B. HCl
- C. H_3O^+
- D. Cl^-

Answer: A

 [Watch Video Solution](#)

20. pH of a 0.001 M NaOH solution will be

- A. 9

B. 3

C. 11

D. 12

Answer: C



Watch Video Solution

21. pH of a solution is 5. Thus. the concentration of hydroxyl ion in the solution is

A. 9molL^{-1}

B. 5molL^{-1}

C. 10^{-5}molL^{-1}

D. 10^{-9}molL^{-1}

Answer: D

 [Watch Video Solution](#)

22. The dissociation constant of an acid, HA is 1×10^{-5} The pH of 0.1 M solution of the acid will be

A. 3

B. 5

C. 4

D. 2

Answer: A

 [Watch Video Solution](#)

23. 100 mL of 0.01 M solution of NaOH is diluted to 1 litre. The pH of resultant solution will be

A. 3

B. 12

C. 11

D. 8

Answer: C



[Watch Video Solution](#)

24. At $80^{\circ}C$, pure distilled water has $[H_3O^+] = 1 \times 10^{-6} \text{ mol } L^{-1}$. The value of K_w at this temperature will be

A. $1 \cdot 10^{-8}$

B. $1 \cdot 10^{-14}$

C. $1 \cdot 10^{-12}$

D. $1 \cdot 10^{-7}$

Answer: C



Watch Video Solution

25. The pH of a solution obtained by mixing 50 mL of 2N HCl and 50 mL of 1 N NaOH is [$\log 5 = 0.7$]

A. 1.7

B. 1.3

C. 0.7

D. 0.3

Answer: D

 [Watch Video Solution](#)

26. The pH of a solution increased from 3 to 6. Its $[H^{\oplus}]$ will be

- A. Increased by 1000 times
- B. Reduced to half
- C. Reduced by 100 times
- D. Reduced by 1000 times

Answer: D

 [Watch Video Solution](#)

27. Ionic product of water increases, if

- A. H^+ ions are added
- B. OH ions are added
- C. Temperature decreases
- D. Temperature increases

Answer: D



Watch Video Solution

28. The hydrogen ion concentration of 0.1 M solution of acetic acid, which is 20% dissociated, is

- A. 0.02 M
- B. 2 M
- C. 0.2 M
- D. 0.002 M

Answer: A

 [Watch Video Solution](#)

29. A monobasic weak acid solution which is 0.002 M has pH value equal to 5, The percentage ionization value of the acid in the solution will be

- A. 0.5
- B. 0.005
- C. 5
- D. 0.05

Answer: A

 [Watch Video Solution](#)

30. In which of the following the solubility of AgCl will be minimum ?



C. Pure water



Answer: A



[Watch Video Solution](#)

31. If the solubility of $\text{Mg}(\text{OH})_2$ in water is $S\text{molL}^{-1}$ then its K_{sp} will be



B. $4S^3$

C. $27S^3$

D. $8S^3$

Answer: B



Watch Video Solution

32. Aqueous solution of sodium acetate is

A. Alkaline

B. Neutral

C. Weakly acidic

D. Strongly acidic

Answer: A

 [Watch Video Solution](#)

33. An acidic buffer solution can be prepared by mixing the solutions of

- A. Sodium chloride and sodium hydroxide
- B. Nitric acid and sodium nitrate
- C. Ammonium chloride and ammonium hydroxide
- D. Sodium acetate and acetic acid

Answer: D

 [Watch Video Solution](#)

34. Degree hydrolysis (h) of a salt of weak acid and a strong base is given by

A. $\frac{K_w}{K_b}$

B. $K_w/(K_a \cdot K_b)$

C. $\frac{K_w}{K_a}$

D. $\frac{K_a \cdot K_b}{K_w}$

Answer: C

 [Watch Video Solution](#)

35. The pH of a solution at 25°C containing 0.20 M sodium acetate and 0.06 M acetic acid is

(pK_a of $\text{CH}_3\text{COOH} = 4.74$ and $\log 3 = 0.477$)

A. 4.36

B. 5.26

C. 5.84

D. 6.32

Answer: B

 [Watch Video Solution](#)

36. The pH of 0.2 M aqueous solution of NH_4Cl will be
(pK_b of $NH_4OH = 4.74$, $\log 2 = 0.3$)

A. 4.98

B. 5.42

C. 4.76

D. 4.32

Answer: A

 [Watch Video Solution](#)

37. On adding ammonium chloride to a solution ammonium hydroxide

- A. Dissociation of NH_4OH increases
- B. Concentration of OH increases
- C. Concentration of OH decreases
- D. Concentration of OH remains unchanged

Answer: C

 [Watch Video Solution](#)

38. Aqueous solution of which salt will not be hydrolysed?

- A. Potassium nitrate
- B. Potassium cyanide

C. Potassium formate

D. Potassium acetate

Answer: A

 [Watch Video Solution](#)

39. If K_{sp} for $HgSO_4$ is 6.4×10^{-5} , then solubility of this substance in mole per m^3 is :

A. $5.4 \cdot 10^{-5} M$

B. $8 \cdot 10^{-3} M$

C. $8 \cdot 10^{-4} M$

D. $6.4 \cdot 10^{-3} M$

Answer: B



 Watch Video Solution

40. pH of 0.5 M aqueous NaCN solution is
(pK_a of HCN = 9.3, $\log 5 = 0.7$)

A. 10.3

B. 9.5

C. 10.6

D. 11.5

Answer: D

 Watch Video Solution

41. Calculate the pH of an aqueous solution of 1.0M ammonium formate assuming complete dissociation. (pK_a of formic acid is

3.8 and pK_a of ammonia is 4.8).

A. 5.5

B. 7.5

C. 6.1

D. 6.5

Answer: D



Watch Video Solution

42. How many grams of calcium oxalate should be dissolved in water to make one litre of saturated solution? (K_{sp} of CaC_2O_4 is 2.5×10^{-9} and its molecular weight is 128 u)

A. $6.4 \cdot 10^{-3} g$

B. $8.0 \cdot 10^{-3} g$

C. $1.28 \cdot 10^{-3} g$

D. $6.4 \cdot 10^3 g$

Answer: A

 [Watch Video Solution](#)

43. What is the pH at which $Mg(OH)_2$ begins to precipitate from a solution containing 0.1 M Mg^{2+} ions ?

$[K_{sp} \text{ for } Mg(OH)_2 = 1.0 \times 10^{-11}]$

 [Watch Video Solution](#)

44. A certain buffer solution contains equal concentration of X^{\ominus} and HX . The K_b for X^{\ominus} is 10^{-10} . The pH of the buffer is

A. 10

B. 4

C. 5

D. 11

Answer: B



Watch Video Solution

45. When ClO_3^- , changes to Cl^-

A. It gains six electrons

B. It gains four electrons

C. It loses six electrons

D. It gains three electrons

Answer: A

 [Watch Video Solution](#)


46. A reducing agent is a substance which can

- A. Accept electron
- B. Donate electron
- C. Reduce itself
- D. Oxidises another species

Answer: B

 [Watch Video Solution](#)

47. For the redox reaction:





The correct coefficients of the reactants for the balanced reaction are



Watch Video Solution

48. Which among the following will not act as a reducing agent?







Answer: C



Watch Video Solution

49. In which of the following reactions, H_2O_2 is acting as a reducing agent?




Answer: B



Watch Video Solution

50. In the given reaction:



A. P is reduced only

B. P undergoes disproportionation reaction

C. P is oxidised only

D. O is reduced

Answer: B

 [Watch Video Solution](#)

51. In the conversion of $I_2 \rightarrow IO_3^-$ the oxidation state of iodine changes from

A. 0 to +6

B. 0 to +5

C. 0 to -1

D. 0 to -3

Answer: B

 [Watch Video Solution](#)

52. The oxidation state of Mn in MnO_4^{2-} is

A. +8

B. +6

C. +7

D. +5

Answer: B



[Watch Video Solution](#)

53. The oxidation state of phosphorus varies from

A. (-3) to +5

B. (-1) to +3

C. (-3) to +3

D. (-5) to +5

Answer: A

 [Watch Video Solution](#)

54. The average oxidation state of sulphur atom in $S_4O_6^{2-}$ ion is

A. +2

B. +5




C. 0

D. +2.5


Answer: D

 [Watch Video Solution](#)

55. Which of the following is not an example of disproportionation reaction ?




D.



Answer: C

 [Watch Video Solution](#)

56. For the redox reaction,



coefficients of the reactants for the balanced reaction are respectively MnO_4^- , $C_2O_4^{2-}$, H^+ :

A. $\frac{1}{5}$

B. $\frac{2}{5}$

C. $\frac{5}{2}$



D. $\frac{3}{5}$



Answer: B



Watch Video Solution

57. The equivalent weight of $MnCl_2$, is half of its molecular weight when it is converted to






Answer: C

 [Watch Video Solution](#)

58. The value of n in the following half equation is



A. 3

B. 5

C. 4

D. 6

Answer: A



59. The oxidation state of Cr in CrO_5 is

- A. +4
- B. +6
- C. +8
- D. +10

Answer: B

60. For the galvanic cell:

$Zn(s) | Zn^{2+}(aq)(1.0M) || Ni^{2+}(aq)(1.0M) | Ni(s)$, E_{cell}^o will be

[Given $E_{\frac{Zn^{2+}}{Zn}}^0 = -0.76V$, $E_{\frac{Ni^{2+}}{Ni}}^0 = -0.25V$]

A. $-0.51V$

B. $-1.01V$

C. $0.51V$

D. $1.01V$

Answer: C



Watch Video Solution

61. Which of the following metals will not react dilute hydrochloric acid?

A. Cu

B. Zn

C. Fe





D. Ca

Answer: A



Watch Video Solution

62. Syngas is a mixture of







Answer: D



Watch Video Solution

63. $Zn(s) + Cu^{2+}(aq) \rightarrow Zn^{2+}(aq) + Cu(s)$ The cell representation for the above redox reaction is



Answer: B

 [Watch Video Solution](#)

64. In which of the compounds the oxidation state of hydrogen is -1?



B. CaH_2

C. HBr

D. H_2S

Answer: B



Watch Video Solution

65. In Iodometric titration which indicator is used to detect end point of titration reaction?

A. Diphenylamine

B. Starch

C. MnO_4^\ominus

D. Methyl orange

Answer: B

 [Watch Video Solution](#)

66. If a small Cu rod is placed in an aqueous solution of ferrous salt, then which of the following will be observed? (





$$E^0_{\frac{Cu^{2+}}{Cu}} = -0.34V, E^0_{\frac{Fe^{2+}}{Fe}} = -0.44V)$$

- A. Copper will be oxidised
- B. Fe^{2+} will be reduced
- C. No reaction will take place
- D. Fe^{2+} will be oxidised

Answer: C

 [Watch Video Solution](#)

67. The standard reduction potential of zinc and silver at 298 K are $E_{\frac{Zn^{2+}}{Zn}}^0 = -0.76V$, $E_{\frac{Ag^+}{Ag}}^0 = 0.80V$ Which of the following reactions actually takes place in a cell reaction?



Answer: B



Watch Video Solution

68. The standard electrode potentials of four elements A,B,C,D are - 3.05,- 1.66,-0.40 and 0.80 volts respectively. The highest chemical activity will be shown by:

A. D

B. A

C. C

D. B

Answer: B



Watch Video Solution

69. In faintly alkaline solution, 4 moles of permanganate anion quantitatively oxidize thiosulphate anions to produce X moles of sulphate anion. The value of X

A. 8

B. 6

C. 4

D. 3

Answer: D



Watch Video Solution