



CHEMISTRY

BOOKS - CENGAGE CHEMISTRY (ENGLISH)

D AND F BLOCK ELEMENTS



1. On what ground can you say that scandium (Z = 21) is a transition

element but zinc (Z = 30) is not?

Watch Video Solution

2. Why do the transition elements exhibit higher enthalpies of atomisation?

Watch Video Solution

3. (a) Phosphourous (P_4) and vanadium (V) both exhibit variable oxidation states but due to different reason.

(b) Name the transition metal which does not show variable O.S.

Watch Video Solution

4. (a). Why IE_1 of Zn, Cd and Hg are very high?

(b). The relative difference in IE of any two consecutive d-bloch elements


is smaller than those of s and p-block elements.

(c). IE_2 is usually very high for Cr and Cu. Why?

Watch Video Solution

5. IE_2 for Zn is comparatively lower as compared to other transition elements of same series.

6. Why is Cr^{2+} reducing and Mn^{3+} oxidising when both have d^4 configuration?



8. for the first series of transition metal the $E^0(\Theta)$ Value are $E^{\Theta}, V, Cr, Mn, Fe, Co, Ni, Cu$ $(M^{2+}/M^{+1}), -1.18, -0.91, -1.18, -0.44, -0.28, -0.25, +0.34$ `Explain the irregularity in the above Values. **9.** Why is the E^{Θ} value for the Mn^{3+}/Mn^{2+} couple much more positive than that for Cr^{3+}/Cr^{2+} or Fe^{3+}/Fe^{2+} ? Exaplain.

Watch Video Solution

10. How would you account for the increasing oxidising power in the Example series $VO_2^+ < Cr_2O_7^{2-} < MnO_4^-$

Watch Video Solution

11. The maggnetic moment of divalent ion in aqueous solution with atomic number 25 is

Watch Video Solution

12. Name a member of the lanthanoid series which is well known to exhibit +4 oxidation state.



13. (a). A strip of Cu dissolved in $FeCl_3$, however Cu cannot displace Fe.

(b). $\left[CuCl_4
ight]^{2-}$ exists but $\left[CuI_4
ight]^{2-}$ does not. Why?

Watch Video Solution

14. (a). Why Mn^{2+} compounds are more stable than Fe^{2+} towards oxidation to their +3 state? (b). Calculate the magnetic moment of V^{3+} ion. (c). $[Ti(H_2O)_{6-}^{3+}$ gives violet coloured aqueous solution but $[Mg(H_2O)_6]^{2+}$ solution is colourless. (d). $[Ti(H_2O)_6]^{3+}$ is coloured whereas $[Sc(H_2O)_6]^{3+}$ is colourless. Why? 15. Why Eu^{2+} and Yb^{2+} ions in solutions are good reducing agents but an aqueous solution of Ce^{4+} is a good oxidising agent?

Watch Video Solution

16. A hydrated metallic salt A, light green in colour, gives a white anhydrous residue B after being heated gradually. B is soluble in water and its aqueous solution reacts with NO to give a dark brown compound C. B on strong heating gives a brown residue D and a mixture of two gases E and F. The gaseous mixture, when passed through acidified $KMnO_4$ discharges the pink colour, when passed through acidified $BaCl_2$ solution, gives a white precipitate. Identifty A, B, C, D, E and F

Watch Video Solution

17. Which of the following statements is correct?

- (a). Nickel is a silver white metal.
- (b). Nickel dimethylglyoxime is square planer.
- (c). Among complexes of cobalt the most stable are those of Co^{2+} ,

(d). Among simple compounds of cobalt the most stable are those of CO^{2+} .



18. Which of the following statements (s) is (are) correct with reference

to the ferrous and ferric ions?

- (a). Fe^{2+} gives brown colour with potassium ferricyanide.
- (b). Fe^{2+} gives blue precipitate with potassium ferricyanide.
- (c). Fe^{3+} gives red colour with potassium thiocyanate.
- (d). Fe^{3+} gives brown colour with potassium thiocyanate.



19. Hydrazine reduce Fehling's solution to form:

(a). Cu(OH)

- (b) Cu_2O
- (c). CuO
- (d). $Cu+O_2$

Watch Video Solution

20. In the following reaction Y and Z are respectively:

- $4AgNO_3 + 2Cl_2(Dry)
 ightarrow 4X + 2Y + Z$
- (a). $AgCl, O_2$
- (b). NO_2, N_2O_4
- (c). $N_2O_5O_2$
- (d). None



21. Explain the following facts:

(a). Copper hydroxide is soluble in ammonium hydroxide but not in sodium hydeoxide.

(b). Addition of an alkali to cuprous chloride solution gives a yellow precipitate which gradually changes to a red colour.

Watch Video Solution

22. Write balanced equations for the reaction of alkaline perbromate with zinc giving tetrahydraoxozincate anion.

Watch Video Solution

23. Excess of dilute sodium hydroxide solution is gradually added with shaking to an aqueous solution of zinc sulphate. What would you observe?

24. (a). The addition of NaOH solution to a solution of zinc chloride produces a white precipitate which dissolves on further addition of NaOH.

(b). The addition of NH_4OH to $ZnSO_4$ solution produces white precipitate but no precipitate is formed if it contains NH_4Cl .



25. Identify A, B, C, D and E.

A while substance A reacts with dilute H_2SO_4 to produce a colourless gas B and a colourless solution C. The reaction between B and acidified $K_2Cr_2O_7$ solution produces a green solution and slightly coloured precipitate D. The substance D burns in air to produce a gas E which reacts with B to yield D and a colourless liquid. Anhydrous copper sulphate is turned blue on adiition of this colourless liquid. Addition of aqueous NH_3 or NaOH to C produces first a precipitate **26.** Mercurous ion is written as Hg_2^{2+} whereas cuprous ion is written as Cu^+ . Explain.

27. Give reasons for the following :


Variation in the radii of transition elements are not as pronounced as

those of representative elements.




28. (a). The addition of NaOH solution to a solution of zinc chloride produces a white precipitate which dissolves on further addition of NaOH.

(b). The addition of NH_4OH to $ZnSO_4$ solution produces white precipitate but no precipitate is formed if it contains NH_4Cl .



is greater than Cu2+. Give suitable explanation.

Watch Video Solution



colourless?



6. Describe giving reason which one of the following pairs has the property indicated?

(a). Fe or Cu has higher melting point.

(b). Co^{2+} or Ni^{2+} has lower magnetic moment.

Watch Video Solution

7. Give reason for the following features of transition metal chemistry:

(i). The lowerst oxide of a transition metal (say, chromium, atomic number 24) is basic whereas the highest oxide is usually acidic.

(ii) . Transitio metals sometimes oxhibit very low oxidation states such as +1 and 0.

Watch Video Solution

8. The 4d and 5d series of transition metals have more frequency metal-

metal bonding in their compounds than do the 3d metals. Explain.

Watch Video Solution

9. Most of the transition metals do not displace hydrogen from dilute

acids. Why?



10. (a). Of the lanthanides, cerium (Z = 58) forms a tetrapositive ion, Ce^{4+} in aqueous solution. Why? (b), The +3 oxidation states of lanthanum (Z = 57), gadolinium (Z = 64) and lutetium (Z = 71) are especially stable. Why?



11. (a). Why second asnd third transition series elements show similar size?

(b). Why electronic configuration of lanthanoids not known with certainty?

12. Chemistry of all the lanthanoids is quite similar.

Watch Video Solution

13. Identify A to E. Pyrolusite on heating with KOH in the presence of air gives a dark green compound (A). The solution of (A) on treatment with H_2SO_4 gives a purple coloured compound (B), which gives the following reactions:

(a). KI on reaction with alkaline solution of (B) changes into a compound (C).

(b). The colour of the compoud (B) disappears on treatment with the acidic solution of $FeSO_4$.

(c). With conc. H_2SO_4 compound (B) gives (D) which can compose to yield (E) and oxygen.



14. (a).
$$CuSO_{4.5}H_2O \xrightarrow{100^{\circ}} (A) \xrightarrow{230^{\circ}} (B) \xrightarrow{800^{\circ}} (C) + (D)$$

(b). $AgNO_3 \xrightarrow{redhot} (E) + (F) + O_2$

Watch Video Solution

15. (a). A powdered substance (A) on treatment with fusion mixture gives a green coloured compound (B).

(b). The solution of (B) The solution of (B) in boiling water on acidification with dilute H_2SO_4 gives a pink coloured compound (C).

Watch Video Solution

16. (a). An ore (A) on roasting with sodium carbonate and lime in the presence of air gives two compounds (B) and (C).

(b). The solution of (B) in conc. HCl on treatment with potassiu ferroyanide gives a blue colour or precipitate of compound (D).

17. Identify A to C.

(a). An aqueous solution of a compoud (A) is acidic towards litmus and (A) is sublimed at about $300^{\circ}C$.

(b). (A) on treatment with an excess of NH_4CNC gives a red coloured compound (B) and on treatment with a solution of $K_4Fe(CN)_6$ gives a blue coloured compound (C).

Watch Video Solution

18. How the following conversions are made?

(a). Copper chloride from copper nitrate.

(b). Cuprous oxide from copper sulphate.

(c). Cuprous oxide from copper.

Watch Video Solution

Ex 61 Subjective

1. (a). Explain why Mn^{3+} is less stable than Mn^{22+} and M^{4+} ions?

(b). Cu(I) compounds are not known in aqueous solution.



2. Write down the names of crystal structures in which the following elements are presents:

Sc, Ti, V, Cr, Mn, Fe, Co, Ni, Cu, Zn

Watch Video Solution

3. Write down the name of catalyst for the following:

Adams catalyst in reduction.



4. Arrange the following complexes in decreasing orger of magnetic

moment:

$$\left[Ni(H_2O)_4\right]^{2+}, \left[Ni(CN)_4\right]^{2-}, \left[Fe(CN)_6\right]^{3-}, \left[Fe(CN)_6\right]^{4-}$$



5. What happen when:

(a). CO_2 gas is passed through aqueous solution of Na_2CrO_4 .

(b). Ammonium dichromate is heated strongly.



6. Explain by giving suitable reason.

(a). Yellow coloured aqueous solution of sodium chromate changes to orange-red when CO_2 under pressure is passed.

(b). Green solution of potassium manganate, K_2MnO_4 , turns purple when CO_2 is circulated.

(c). $Hg^{2\,+}$ and $Hg^{2\,+}_2$ salts are colourless.

(d). Cu^{2+} salts are paramagnetic while Cu^{\oplus} salts are diamagnetic in nature.



7. Explain:

(a). Why is $K - 2Cr_2O_7$ generally preferred to $Na_2Cr_2O_7$ in volumetric

analysis?

(b). Which divalent metal ion has maximum paramagnetic character amont the first transition metals? Why?

(c). How the colour of $K_2 C r_2 O_7$ solution does depend on pH of the solution?

Watch Video Solution

8. (a). Why do second and third rows of transition elements resemble each other more closely than the first row?

Give reasons:

(b). Most transition metals show paramagnetic behaviours.

(c). Transition metal and their many compounds are coloured.

Watch Video Solution

9. Acidified $K_2Cr_2O_7$ solution turns green when Na_2SO_3 is added to it.

This is due to the formation of

Watch Video Solution

10. Identify D

When a white crystalline compound X is heated with $K_2Cr_2O_7$ and concentrated H_2SO_4 , a reddish brown gas A is evolved. On passing A into caustic soda solution a yellow coloured solution of B is obtained Neutralising the solution of B with acetic acid and on subsequent addition of lead acetate, a yellow precipitate C is obtained. When X is heated with NaOH solution a colourless gas is evolved and on passing this gas into K_2Hgl_4 solution a reddish brown precipitate D is formed.

Watch Video Solution

11. Complete and balance the following equation:

(c).
$$KMnO_4 \xrightarrow{\bigtriangleup}_{200\ ^\circ C}$$

(iv). $MnO_4^{\Theta} + AsO_3^{\Theta} + H^{\oplus} \rightarrow$
(v). $K_2Cr_2O_7 + \xrightarrow{NH_4Cl}$

Watch Video Solution

12. Wxplain the following giving suitable reson.

Transitin metals of 3d-series doe not react readily with dilute acids to liberate hydrogen although they have high negative reduction potential values. Explain. . 1. Complete and balance the following equation:

(i). $FeSO_4 \xrightarrow[2.K_4[Fe(CN)_6]]{} \xrightarrow{1.K_2Cr_2O_7} \xrightarrow{2.K_4[Fe(CN)_6]}$ (ii). $FeSO_4 \xrightarrow[tored \neq ss]{} \xrightarrow{heated} \xrightarrow{tored \neq ss}$

Watch Video Solution

2. Explain the following :

(i). $FeCl_3$ and $FeBr_3$ are well known, but FeI_3 has doubtful existence.

(ii). Anhydrous $FeCl_3$ cannot be obtained by heating hydrated ferric chloride.



3. Draw shape of $FeCl_3$ in :

(i). Water

(ii). Ether
(iii). Gaseous state.
Watch Video Solution
4. What happen when iron is treated with:
(i). Steam
(ii). Dilute HNO_3
Watch Video Solution
5. Colourless salt (A) decolourise I_2 solution and gives white precipitate (change to black) with $AgNO_3$ solution. (A) also produces pink colour

with $FeCl_3$ solution. Identify (A) and explain reactions.

Watch Video Solution

6. Iron forms iron (II) chloride, $FeCl_2$ and iron (III) chloride $FeCl_3$. One of these chlorides is a dark brown solid, melting point $306^{\circ}C$. The other is a white crystalline solid with greenish tint, melts at $674^{\circ}C$ Which description best fits iron (II) chloride Explain.

Watch Video Solution

7. Account for the following observation and write balanced chemical equations for them.

(i). When ferrous sulphate crystals are exposed to air, it turns brown.

(ii). A solution of ferric chloride in water gives a brown precipitate on standing.



Ex 6 2 Subjective Compound Of Cu

1. When copper is exposed to atmosphere for a longer time, a green mass deposits on its surface. Discuss the chemical reaction involved.

Watch	Video	So	lution

2. Complete and balance the following equations:

 $CuS+2H_2SO_4+\ldots \
ightarrow 2CuSO_4+2H_2O+S$

Watch Video Solution

3. Explain the following:

(i). Copper hydroxide is soluble in ammonium hydroxide but not in sodium hydroxide.

(ii). A dark blue precipitate is formed when sodium hydroxide solution is

added to copper sulphate solution. The precipitate darkens on heating.



1. (i). ZnO is used i as a white paint inspite of the fact that it has less convering power than white lead.

(ii). Hydrated $ZnCl_2$ cannot be dehydrated on heating?

(iii). Zn is used in galvanisation of Fe.

Watch Video Solution

Ex 6 2 Subjective Compound Of Ag

1. Preparation of looking mirrors involves the use of ammonical $AgNO_3$

+red lead+ HCHO. Explain the function of each.



2. Give reason for the following:

(i). Silver ornament gets tarnished when exposed to atmosphere for a

long time.

(ii) Silver nitrate solution is kept in dark coloured bottles.

(iii). Why does $AgNO_3$ produce a black stain on the skin.

Watch Video Solution

Ex 6 2 Subjective Compound Of Hg

1. An aqueous solution containing one mole of HgI_2 and two moles of Nal is orange in colour. On addition of excess Nal the solution becomes colourless. The orange colour reappears on subsequent addition of NaOCl. Explain with equation.



2. $HgCl_2$ and $SnCl_2$ cannot exist together in an aqeous solution . Explain.

Watch Video Solution

3. Compelete and balance the following equations:

(i). $Cu(OH)_2 + NH_4NO_3 + NH_4OH
ightarrow \ldots + H_2O$

(ii). $Au + HCl + HNO_3
ightarrow \ldots + \ldots + H_2O$

(iii). $Ag_2S + 2CuCl_2 + 2\mathrm{Hg}
ightarrow \ldots + \ldots S + 2Ag$

Watch Video Solution

Exercises Linked Comprehension

1. The colours of the transition metal are due to d-d excitation. The energy required for d-d electron axcitation is available in the visible range.

Transition metal ions have the tendency to absorb certain rediations from the visible region and exhibit the complementary colour.

The transition metal ions which have completely filled d-orbitals are colourless as the excitation of electron or electrons is not possible within d-orbitals. The transition metal ions which have completely empthy d-orbitals are also colourless. In $KMnO_4$ and $K_2Cr_2O_7$, there are no unpaired electrons at the central atom but they are dep in colour. The colour of these compounds is due to charge transfer spectrum. for example in MnO_4 electron is momentrily transferred from O to the metal and thuys oxygen changes from O^{2-} and O^{Θ} maganese from Mn^{7+} to Mn^{6+} .

Q. Which of the following is paramagnetic as well as coloured ion?

A. Cu^\oplus B. Cu^{2+} C. Sc^{3+}

D. Zn^{2+}

Answer: b



2. The colours of the transition metal are due to d-d excitation. The energy required for d-d electron axcitation is available in the visible

range.

Transition metal ions have the tendency to absorb certain rediations from the visible region and exhibit the complementary colour.

The transition metal ions which have completely filled d-orbitals are colourless as the excitation of electron or electrons is not possible within d-orbitals. The transition metal ions which have completely empthy d-orbitals are also colourless. In $KMnO_4$ and $K_2Cr_2O_7$, there are no unpaired electrons at the central atom but they are dep in colour. The colour of these compounds is due to charge transfer spectrum. for example in MnO_4 electron is momentrily transferred from O to the metal and thuys oxygen changes from O^{2-} and O^{Θ} maganese from Mn^{7+} to Mn^{6+} .

- Q. Which is a coloured ion?
 - A. $\left[Cr(H_2O)_6
 ight]^{3+}$
 - B. $\left[Cu(CN)_4\right]^{3-}$
 - $\mathsf{C.}\left[Ti(H_2O)_6\right]^{4\,+}$
 - D. $\left[Sc(H_2O)_6
 ight]^{3+}$

Answer: a

Watch Video Solution

3. The colours of the transition metal are due to d-d excitation. The energy required for d-d electron axcitation is available in the visible range.

Transition metal ions have the tendency to absorb certain rediations from the visible region and exhibit the complementary colour.

The transition metal ions which have completely filled d-orbitals are colourless as the excitation of electron or electrons is not possible within d-orbitals. The transition metal ions which have completely empthy d-orbitals are also colourless. In $KMnO_4$ and $K_2Cr_2O_7$, there are no unpaired electrons at the central atom but they are dep in colour. The colour of these compounds is due to charge transfer spectrum. for example in MnO_4 electron is momentrily transferred from O to the metal and thuys oxygen changes from O^{2-} and O^{Θ} maganese from ${Mn^{7\,+}}$ to ${Mn^{6\,+}}.$

Q. Select the correct statement:

A. colour of the transition metal ion arises due to d-d transition.

B. colour of certain oxysalts of transition metals is due to charge

tranfer.

C. Both are correct.

D. None is correct.

Answer: c

Watch Video Solution

4. The colours of the transition metal are due to d-d excitation. The energy required for d-d electron axcitation is available in the visible range.

Transition metal ions have the tendency to absorb certain rediations

from the visible region and exhibit the complementary colour.

The transition metal ions which have completely filled d-orbitals are colourless as the excitation of electron or electrons is not possible within d-orbitals. The transition metal ions which have completely empthy d-orbitals are also colourless. In $KMnO_4$ and $K_2Cr_2O_7$, there are no unpaired electrons at the central atom but they are dep in colour. The colour of these compounds is due to charge transfer spectrum. for example in MnO_4 electron is momentrily transferred from O to the metal and thuys oxygen changes from O^{2-} and O^{Θ} maganese from Mn^{7+} to Mn^{6+} .

Q. Which of the following compounds is (are) coloured due to charge transfer spectra and not due to d-d transition?

A. $KMnO_4$

B. K_2CrO_4

 $C. CrO_3$

D. all of these

Answer: d

5. The colours of the transition metal are due to d-d excitation. The energy required for d-d electron axcitation is available in the visible range.

Transition metal ions have the tendency to absorb certain rediations from the visible region and exhibit the complementary colour.

The transition metal ions which have completely filled d-orbitals are colourless as the excitation of electron or electrons is not possible within d-orbitals. The transition metal ions which have completely empthy d-orbitals are also colourless. In $KMnO_4$ and $K_2Cr_2O_7$, there are no unpaired electrons at the central atom but they are dep in colour. The colour of these compounds is due to charge transfer spectrum. for example in MnO_4 electron is momentrily transferred from O to the metal and thuys oxygen changes from O^{2-} and O^{Θ} maganese from Mn^{7+} to Mn^{6+} .

Q. $K_2Cr_2O_7$ gives coloured solution in water. The colour is due to

A. d-d transition in Cr-atoms

B. Presence of unpaired electron in d-orbital of oxygen

C. charge transfer from 0 to Cr

D. none of the above

Answer: c

Watch Video Solution

6. $KMnO_4$ is prepared from the mineral pyrolusite, MnO_2 (deep purple colour). It acts as an oxidising agent in the neutral, alkaline as well as acidic medium in acidic medium it is used in volumetric analysis for estimation of Fe^{2+} , $Cr_2O_4^{2-}$ salts etc. The titrations are carried out in presence of H_2SO_4 . However, before using it as a titrant, it is first standardised with standard oxalic acid solution or Mohr's salt solution . In one of the experiments on titration 26.8g of dry pure sodium oxalate $(Mw = 123gmol^{-1})$ was dissolved in 1L of distilled water and then 100 mL of $2MH_2SO_4$ were added. The solution was cooled. Now to this solution $0.1MKMnO_4$ solution was added till a very faint pink colour

persisted.

Q. The purple colour of $KMnO_4$ is due to

A. incomplete d-subshell

B. ionic nature of $KMnO_4$

C. Charge transfer

D. Resonance in MnO_4^{Θ} ion

Answer: c

Watch Video Solution

7. $KMnO_4$ is prepared from the mineral pyrolusite, MnO_2 (deep purple colour). It acts as an oxidising agent in the neutral, alkaline as well as acidic medium in acidic medium it is used in volumetric analysis for estimation of Fe^{2+} , $Cr_2O_4^{2-}$ salts etc. The titrations are carried out in presence of H_2SO_4 . However, before using it as a titrant, it is first standardised with standard oxalic acid solution or Mohr's salt solution . In one of the experiments on titration 26.8g of dry pure sodium oxalate

 $(Mw = 123 gmol^{-1})$ was dissolved in 1L of distilled water and then 100 mL of $2MH_2SO_4$ were added. The solution was cooled. Now to this solution $0.1MKMnO_4$ solution was added till a very faint pink colour persisted.

Q. Mohr's salt, $FeSO_4(NH_4)_2SO_{46}H_2O$, is preffered over $FeSO_{4.7}H_2O$ for standardisation of $KMnO_4$ solution because.

A. Mohr's salt is a double salt while ferrous sulphate is a single salt.

B. Mohr's salt is not hygroscopic but $FeSO_{4.7}H_2O$ is hygroscopic.

C. Mohr's salt contains only ferrous ions whereas ferrous sulphate

contains some ferric ions.

D. Mohr's salt solution can be titrated even in the absence of H_2SO_4 .

Answer: c



8. $KMnO_4$ is prepared from the mineral pyrolusite, MnO_2 (deep purple colour). It acts as an oxidising agent in the neutral, alkaline as well as acidic medium in acidic medium it is used in volumetric analysis for estimation of $Fe^{2\,+}$, $Cr_2O_4^{2\,-}\,$ salts etc. The titrations are carried out in presence of H_2SO_4 . However, before using it as a titrant, it is first standardised with standard oxalic acid solution or Mohr's salt solution . In one of the experiments on titration 26.8g of dry pure sodium oxalate $\left(Mw=123gmol^{-1}
ight)$ was dissolved in 1L of distilled water and then 100 mL of $2MH_2SO_4$ were added. The solution was cooled. Now to this solution $0.1MKMnO_4$ solution was added till a very faint pink colour persisted.

Q. When pyrolusite is fused with KOH and $KClO_3$, we get

A. $KMnO_4$

B. $K_2 MnO_4$

C. both $KMnO_4$ and K_2MnO_4

D. None of these

Answer: b

Watch Video Solution

9. $KMnO_4$ is prepared from the mineral pyrolusite, MnO_2 (deep purple colour). It acts as an oxidising agent in the neutral, alkaline as well as acidic medium in acidic medium it is used in volumetric analysis for estimation of $Fe^{2\,+}$, $Cr_2O_4^{2\,-}$ salts etc. The titrations are carried out in presence of H_2SO_4 . However, before using it as a titrant, it is first standardised with standard oxalic acid solution or Mohr's salt solution . In one of the experiments on titration 26.8g of dry pure sodium oxalate $(Mw = 123 gmol^{-1})$ was dissolved in 1L of distilled water and then 100 mL of $2MH_2SO_4$ were added. The solution was cooled. Now to this solution $0.1MKMnO_4$ solution was added till a very faint pink colour persisted.

Q. If $0.1MK_2Cr_2O_7$ solution were used in place of 0.1 $MKMnO_4$ solution used would be

B. 400 mL

C. 333.3 mL

D. 666.6 mL

Answer: d

Watch Video Solution

10. If instead of H_2SO_4 , HCl or HNO_3 of suitable concentration were

used, the volume of $KMnO_4$ solution used would have been

A. less in case of HCl but more in case of HNO_3

B. More in case of HCl but less in case of HNO_3

C. more in both cases

D. Less in both cases

Answer: b

11. $KMnO_4$ is prepared from the mineral pyrolusite, MnO_2 (deep purple colour). It acts as an oxidising agent in the neutral, alkaline as well as acidic medium in acidic medium it is used in volumetric analysis for estimation of Fe^{2+} , $Cr_2O_4^{2-}$ salts etc. The titrations are carried out in presence of H_2SO_4 . However, before using it as a titrant, it is first standardised with standard oxalic acid solution or Mohr's salt solution . In one of the experiments on titration 26.8g of dry pure sodium oxalate $(Mw = 123 gmol^{-1})$ was dissolved in 1L of distilled water and then 100 mL of $2MH_2SO_4$ were added. The solution was cooled. Now to this solution $0.1MKMnO_4$ solution was added till a very faint pink colour persisted.

Q. The volume of $KMnO_4$ solution that must have been added to obtain the faint pink colour at the end point must be

A. 100 mL

B. 200 mL

C. 400 mL

D. 800 mL

Answer: d



12. Transition metals and many of their compounds show paramagnetic behaviour where there are unpaired electron or electrons. The magnetic moment arises from the spin and orbital motions in ions or molecule. Magnetic moment of n unpaired electrons is given as

$$\mu=\sqrt{n(n+2)}$$
 Bohr magneton

Magnetic moment increases as the number of unpaired electrons increases.

Q. Which among the following ions has maximum value of magnetic moment

A. Cu^{2+}

B. Mn^{2+}

C. Cr^{2+}

D. Ti^{2+}

Answer: b



13. Arrange the following in increasing value of magnetic moments. (i) $[Fe(Cn)_6]^{4-}$ (ii) $[Fe(CN)_6]^{3-}$ (iii) $[Cr(NH_3)_6]^{3+}$ (iv) $[Ni(H_2O)_4]^{2+}$ A. I < II < III < IV

 $\mathsf{B}.\,IV < III < II < I$

 $\mathsf{C}.\,II < III < I < IV$

D. I < II < IV < III

Answer: d

14. Transition metals and many of their compounds show paramagnetic behaviour where there are unpaired electron or electrons. The magnetic moment arises from the spin and orbital motions in ions or molecule. Magnetic moment of n unpaired electrons is given as

$$\mu=\sqrt{n(n+2)}$$
 Bohr magneton

Magnetic moment increases as the number of unpaired electrons increases.

Q. In 3d series the maximum magnetic moment is shown by

- A. Sc
- B.V
- C. Cr

D. Fe

Answer: c

15. Transition metals and many of their compounds show paramagnetic behaviour where there are unpaired electron or electrons. The magnetic moment arises from the spin and orbital motions in ions or molecule. Magnetic moment of n unpaired electrons is given as

 $\mu=\sqrt{n(n+2)}$ Bohr magneton

Magnetic moment increases as the number of unpaired electrons increases.

Q. Magnetic moments of $Cr(Z=24), Mn^{\oplus}(Z=35)$ and $Fe^{2+}(Z=36)$ are x,y and z respectively. Which of the following order is correct?

A. x < y < zB. x = y < zC. z < x = yD. x = y = z

Answer: c

16. Transition metals and many of their compounds show paramagnetic behaviour where there are unpaired electron or electrons. The magnetic moment arises from the spin and orbital motions in ions or molecule. Magnetic moment of n unpaired electrons is given as

 $\mu=\sqrt{n(n+2)}$ Bohr magneton

Magnetic moment increases as the number of unpaired electrons increases.

Q. There are three unpaired electrons in $[Co(H_2O)_6]^{2+}$ and calculated value of magnetic moment on the basis of $\sqrt{n(n+2)}$ formula is 3.87 BM. which is lower than the experimental value of 4.40 BM. The reason for this difference is due to

A. increase in number of unpaired electrons during determination

B. Some contribution of the orbital motion of the electrons to the magnetic moment

C. d-d transition

D. Experimental error

Answer: b

Watch Video Solution

17. Transition metals and many of their compounds show paramagnetic behaviour where there are unpaired electron or electrons. The magnetic moment arises from the spin and orbital motions in ions or molecule. Magnetic moment of n unpaired electrons is given as

$$\mu=\sqrt{n(n+2)}$$
 Bohr magneton

Magnetic moment increases as the number of unpaired electrons increases.

Q. Magnetic moment of $\left[Ni(CN)_4\right]^{2-}$ is zero but that of $\left[Ni(H_2O)_4\right]^{2+}$ is 2.83*BM*. is because of :

A. CN^{θ} is a strong ligand making two unpaired electrons in Ni^{2+} to pair up, while in $[Ni(H_2O)_4]^{2+}$ two electrons remain unpaired as H_2O is a weak ligand.

B. Different oxidation state of Ni in two complexes

C. Both (a) and (b)

D. none of the above

Answer: a

Watch Video Solution

18. Photography is based on the nature of silver halides. Except AgF, the silver halides are photosensitive. These undergo decomposition in light and turn black due to formation of free silver.

 $2AgBr \xrightarrow{light} 2Ag + Br_2$

The photography films are prepared by adding 20% aqueous solution of $AgNO_3$ to NH_4Br solution containing gelatic. When such a film is exposed, emulsion gets affected and a latent image is formed on the film.

When this exposed film or plate is dipped ina developer which contains a reducing agent, the part affected most during exposure are reduced to the maximum. The image becomes visible. It is called a negative. The remaining sensitive emulsion on the negative is removed by dissolving it in hypo solution (fixer). Fihnally, a positive of the nagative already prepared is made on silver bromide paper.

Q. The compound formed on the unexposed photographic film or plate is

A. silver nitrate

B. ammonium bromide

C. diammine silver bromide

D. silver bromide

Answer: d



19. Photography is based on the nature of silver halides. Except AgF, the silver halides are photosensitive. These undergo decomposition in light and turn black due to formation of free silver.

 $2AgBr \stackrel{light}{\longrightarrow} 2Ag + Br_2$

The photography films are prepared by adding $20\,\%\,$ aqueous solution

of $AgNO_3$ to NH_4Br solution containing gelatic. When such a film is exposed, emulsion gets affected and a latent image is formed on the film.

When this exposed film or plate is dipped ina developer which contains a reducing agent, the part affected most during exposure are reduced to the maximum. The image becomes visible. It is called a negative. The remaining sensitive emulsion on the negative is removed by dissolving it in hypo solution (fixer). Fihnally, a positive of the nagative already prepared is made on silver bromide paper.

Q. The exposed part of the film or plate after developing contains

A. Silver metal

B. Silver oxide

C. Silver bromide

D. Silver nitrate

Answer: a

20. Photography is based on the nature of silver halides. Except AgF, the silver halides are photosensitive. These undergo decomposition in light and turn black due to formation of free silver.

$$2AgBr \stackrel{light}{\longrightarrow} 2Ag + Br_2$$

The photography films are prepared by adding 20% aqueous solution of $AgNO_3$ to NH_4Br solution containing gelatic. When such a film is exposed, emulsion gets affected and a latent image is formed on the film.

When this exposed film or plate is dipped ina developer which contains a reducing agent, the part affected most during exposure are reduced to the maximum. The image becomes visible. It is called a negative. The remaining sensitive emulsion on the negative is removed by dissolving it in hypo solution (fixer). Fihnally, a positive of the nagative already prepared is made on silver bromide paper.

Q. Silver halides are used in photographgy because these compounds

A. are insoluble in water

B. are affected by light

C. are soluble in ammonia solution

D. easily stick on the surface of photographic plate or film

Answer: b

Watch Video Solution

21. Photography is based on the nature of silver halides. Except AgF, the silver halides are photosensitive. These undergo decomposition in light and turn black due to formation of free silver.

 $2AgBr \stackrel{light}{\longrightarrow} 2Ag + Br_2$

The photography films are prepared by adding 20% aqueous solution of $AgNO_3$ to NH_4Br solution containing gelatic. When such a film is exposed, emulsion gets affected and a latent image is formed on the film.

When this exposed film or plate is dipped ina developer which contains a reducing agent, the part affected most during exposure are reduced to the maximum. The image becomes visible. It is called a negative. The remaining sensitive emulsion on the negative is removed by dissolving it in hypo solution (fixer). Fihnally, a positive of the nagative already prepared is made on silver bromide paper.

Q. The solution of the developer consists of

A. Alkaline solution of pyrogallol

B. Alkaline solution of quinil

C. Either (a) or (b)

D. Neither (a) nor (b)

Answer: c



22. Photography is based on the nature of silver halides. Except AgF, the silver halides are photosensitive. These undergo decomposition in light and turn black due to formation of free silver.

 $2AgBr \stackrel{light}{\longrightarrow} 2Ag + Br_2$

The photography films are prepared by adding $20\,\%\,$ aqueous solution

of $AgNO_3$ to NH_4Br solution containing gelatic. When such a film is exposed, emulsion gets affected and a latent image is formed on the film.

When this exposed film or plate is dipped ina developer which contains a reducing agent, the part affected most during exposure are reduced to the maximum. The image becomes visible. It is called a negative. The remaining sensitive emulsion on the negative is removed by dissolving it in hypo solution (fixer). Fihnally, a positive of the nagative already prepared is made on silver bromide paper.

Q. Silver bromide dissolves in hypo solution forming:

A. $Ag_2S_2O_3$

 $\mathsf{B.}\, Ag_2S$

 $\mathsf{C}.\,Na_3\big[Ag(SO_3)_2\big)\big]$

D. $NaAgS_2O_3$

Answer: c

23. In any transition series, from left to right, the d-orbitals are progressively filled and their properties vary accordingly.

Q. In the second transition series, the largest number of oxidation states are shown by

A. Tc

B. Ru

C. Rh

D. Pd

Answer: c

Watch Video Solution

24. In any transition series, from left to right, the d-orbitals are progressively filled and their properties vary accordingly.

Q. Which element do you expect to have the smallest atomic radius?

A. La

B.Zn

C. Sc

D. Hg

Answer: b

Watch Video Solution

25. In any transition series, from left to right, the d-orbitals are progressively filled and their properties vary accordingly.

Q. Which element do you expect to have the highest melting point?

A. La

B.W

C. Pt

D. Os

Answer: b



26. In any transition series, from left to right, the d-orbitals are progressively filled and their properties vary accordingly.

Q. Which of the following pair of compounds is expected to exhibit same colour in aqueous solution?

- A. $FeCl_2, CuCl_2$
- B. $VOCl_2, CuCl_2$
- $C.VOCl_2, FeCl_2$
- D. $FeCl_2$, $MnCl_2$

Answer: b

27. In any transition series, from left to right, the d-orbitals are progressively filled and their properties vary accordingly.

Q. Which of the following is the correct order of second ionisation energy?

- A. V>Cr>Mn
- $\operatorname{B.} V < Cr < Mn$
- $\mathsf{C}.\, V < Cr > Mn$
- $\mathsf{D.}\, V > Cr < Mn$

Answer: c

Watch Video Solution

28. $KMnO_4$ reacts with $Na_2S_2O_3$ in acidic, strongly basic and aqueous (neutral) media. 100mL of $LMnO_4$ reacts with 100 mL of 0.1 M $Na_2S_2O_3$ in acidic, basic and neutral media.

Q. The molarity (M) of $KMnO_4$ solution in the acidic medium is

A. 0.2 M

B. 0.02M

C. 0.4 M

D. 0.04 M

Answer: b

> Watch Video Solution

29. $KMnO_4$ reacts with $Na_2S_2O_3$ in acidic, strongly basic and aqueous (neutral) media. 100mL of $LMnO_4$ reacts with 100 mL of 0.1 M $Na_2S_2O_3$ in acidic, basic and neutral media.

Q. The molarity (M) of $KMnO_4$ solution in basic medium is:

A. 0.8 M

B. 0.08 M

C. 0.26 M

D. 0.026M

Answer: a



30. $KMnO_4$ reacts with $Na_2S_2O_3$ in acidic, strongly basic and aqueous (neutral) media. 100mL of $LMnO_4$ reacts with 100 mL of 0.1 M $Na_2S_2O_3$ in acidic, basic and neutral media.

Q. The molarity (M) of $KMnO_4$ in aqueous medium is

A. 0.8M

B. 0.08M

C. 0.26M

D. 0.026M

Answer: c

31. $KMnO_4$ reacts with $Na_2S_2O_3$ in acidic, strongly basic and aqueous (neutral) media. 100mL of $LMnO_4$ reacts with 100 mL of 0.1 M $Na_2S_2O_3$ in acidic, basic and neutral media.

Q. The molality (m) of $KMnO_4$ in the acidic medium is (Density of $KMnO_4$) solution $= 1.58^{-1}~Mw(KMnO_4) = 158gmol^{-1}$)

A.0.025

 $\mathsf{B}.\,0.25$

 $\mathsf{C}.\,0.12$

 $\mathsf{D}.\,0.012$

Answer: d



Exercises Multiple Correct

1. dz^2 orbital is involvedr in which of the following hybridisastion?

A. sp^3d

 $\mathsf{B}.\,dsp^3$

 $\mathsf{C.}\, sp^3d^2$

D. $d^2 s p^3$

Answer: a,b

Watch Video Solution

2. d_{xy}, d_{yz} and d_{xz} orbitals is involved in which of the following hybridisation?

A. dsp^2

 $\mathsf{B.}\, sp^3d$

C. sp^3d^2

 $\mathsf{D}.\,d^3sp^3$

Answer: c,d

3. $d_{x^2-y^2}$ and d_{z^2} orbitals is involved in which of the foollowing hybridisation?

A. sp^3d^2 B. d^2sp^3 C. sp^3d^3

D. d^3sp^3

Answer: a,b

Watch Video Solution

4. $d_x 2 -_y 2$ orbital is involved in which of the following hybridisation ?

A. sp^3d

 $\mathsf{B.}\, sp^3d^2$

 $\mathsf{C.}\, sp^3d^3$

D. None of these

Answer: b,c

Watch Video Solution

5. Which of following is/are correctly matched ?

A. d-block element: Electronic configuration is $ns^{0-2}(n-1)d^{1-10}$

B. p-block element:electronic configuration is $ns^{1-2}np^{1-6}$

C. s-block element:electronic configuration is ns^{1-2}

D. Ce: F-blocks's first member

Answer: a,c,d

6. Electrons are filled into atomic orbitals in the increasing order of orbital energy level according to ?



7. Which of the following represents the incoorect order of the properties indicated?

A.
$$Ni^{2+} > Cr^{2+} > Fe^{2+} > Mn^{2+}$$
 (size)

B. Sc > tI > Cr > Mn (size)

C. $Ni^{2+} < Co^{2+} < Fe^{2+} < Mn^{2+}$ (unpaired electron)

D. $H_3AsO_4 > H_3PO_4$ (acidic strength order)

Answer: a,d

8. What will be the correct representation of quatum numbers f the last

electron entered into Ce?

A.
$$n = 4, l = 3, m = -3, s = \frac{-1}{2}$$

B. $n = 4, l = 3, m = 0, s = \frac{-1}{2}$
C. $n = 4, l = 2, m = -3, s \frac{+1}{2}$
D. $n = 4, l = 3, m = +2, s = \frac{-1}{2}$

Answer: a,b,d



9. Which is true statement about $KMnO_4$?

A. Its solution is unstable in acidic medium.

B. It has purple colour.

C. MnO_4^{θ} changes to Mn^{2+} in basic solution.

D. It is self indicator in Fe&(2+) or $C_2O_4^{2-}$ titration.

Answer: a,b,d



10. Out of
$$[Fe(CN)_6]^{4-}$$
, $[Ni(CN)_4]^{2-}$, and $[Ni(CO)_4]$: select the

incorrect Statement(s):

A. All have identical geometry

B. All are paramagnetic

C. all are diamagnetic

D. $\left[Fe(CN)_6\right]^{4-}$ is dimagnetic but $\left[Ni(CN)_4\right]^{2-}$ and $\left[Ni(CO)_4\right]$

are paramagnetic

Answer: a,b,d

11. The ability of d-block elements to form complexes is due to:

A. Small and highly charged ions

B. Vacant low energy orbitals to accept lone pair of electrons from

ligands

C. Low polarising power of cation

D. None is correct

Answer: a,b

Watch Video Solution

12. Which one of the following ionic species will impact colour to an aqueous solutions?

A. $Ti^{4\,+}$

 $\mathsf{B.}\,Cu^{\,\oplus}$

C. Zn^{2+}

D. Cr^{3+}

Answer: a,b,c



13. A transition element X has a configuration $3d^4$ in its +3 oxidation state. Its atomic number is not

A. 25

B.26

C. 22

D. 19

Answer: b,c,d

14. The transition metals which do not form amalgams are

A. Zn

B. Fe

C. Cd

D. Pt

Answer: b,c

Watch Video Solution

15. The colour of the transition metal ions is due to

A. d-d transition

B. Charger transfer

C. Change in the geometry

D. None

Answer: a,b



16. Transition elements have greater tendency to form complexes because

- A. They have vacant d-orbitals
- B. They have large size
- C. They show variable oxidation state
- D. They have two electrons in their outermost shells.

Answer: a,c,d



17. Which out of the following belong to 3d-series?

A. Copper

B. Cobalt

C. Gold

D. Silver

Answer: a,b

Watch Video Solution

18. The elements which exist in the liquid state at rooom temperature are.

A. Na

 $\mathsf{B.}\,Br_2$

 $\mathsf{C}.Hg$

 $\mathsf{D.}\,Ga$

Answer: b,c,d

19. Which of the following statements (s) is (are) correct with reference to the ferrous and ferric ions?

(a). Fe^{2+} gives brown colour with potassium ferricyanide.

(b). Fe^{2+} gives blue precipitate with potassium ferricyanide.

(c). Fe^{3+} gives red colour with potassium thiocyanate.

(d). Fe^{3+} gives brown colour with potassium thiocyanate.

A. Fe^{3+} gives brown colour with potassium ferricyanide

- B. Fe^{2+} gives blue precipitate with potassium ferricyanide.
- C. Fe^{3+} gives red colour with potassium thiocyanate.
- D. Fe^{2+} gives brown colour with ammonium thiocyanate.

Answer: a,b,c



20. Which of the following represents the correct order of the properties inidicated?

A.
$$Ni^{2+}>Cr^{2+}>Fe^{2+}>Mn^{2+}$$
 (size)
B. $Sc>Ti>Cr>Mn$ (size)
C. $Ni^{2+}< Co^{2+}< Fe^{2+}< Mn^{2+}$ (unpaired electron)

D. $Cr^{3\,+}\,>\,Cr^{2\,+}$ (magnetic moment)

Answer: b,c

Watch Video Solution

21. The correct statement for d-block element is

A. it shows magnetic property

B. It has variable valency

C. It has tendency of formation of coloured ions.

D. it has complete d-orbitals.

Answer: a,b,c



22. The aqueous solution of the salt will be coloured in the case of

A. $Zn(CO_3)_2$

B. $LiNO_3$

 $C. Cl(NO_3)_2$

D. $CrCl_3$

Answer: c,d

Watch Video Solution

23. Potassium manganate $(K_2 M n O_4)$ is formed when

A. Chlorine is passed through aqueous $KMnO_4$ solution

B. Magnanese dioxide is fused with potassium hydroxide in air

C. Formaldehyde reacts with potassium

D. Potassium permaganate reacts with H_2SO_4

Answer: b,c

Watch Video Solution

24. Cl_2 gas is obtained by various reactions but not by:

A.
$$KMnO_4+conc.$$
 $HCl \stackrel{\Delta}{\longrightarrow}$

B.
$$KCl + K_2 Cr_2 O_7 + conc. \ H_2 SO_4 \overset{\Delta}{\longrightarrow}$$

C.
$$MnO_4 + conc. \ HCl \stackrel{\Delta}{\longrightarrow}$$

D.
$$KCl + F_2 \stackrel{\Delta}{\longrightarrow}$$

Answer: a,c,d

25. Which is /are true statement?

A. lons of d-block elements are coloured due to d-d transition.

B. lons of f-block elements are coloured due to f-f transition.

C. $\left[Sc(H_2O)_6\right]^{3+}$, $\left[Ti(H_2O)_6\right]^{4+}$, are coloured complexes.

D. Cu^{\oplus} is colourless ion.

Answer: a,b,d

Watch Video Solution

26. In $ig[Fe(CN)_5(NO)ig]^{2-}$, Fe has +2 state. It cannot be dicided by

- A. Magnetic measurement
- B. Colligative property
- C. Colour Hybridisation

Answer: b,c,d



27. Which one of the following statement is/are correct?

A. Zinc dissolves in sodium hydroxide solution.

B. Carbon monoxide reduces iron (III) oxide to iron.

C. Mercury (II) iodide dissolves in excess of potassium isodide

solution

D. Tin (IV) chloride is made by dissolving tin solution in concentrated

hydrochloric acid.

Answer: a,b,c



28. In the iodometric estimation in the laboratory which process is involved?

$$\begin{array}{l} \text{A. } Cr_2 O_7^{2^-} + H^{\oplus} + I^{\oplus} \rightarrow 2 C r^{3^+} I_2 \\ \\ I_2 + S_2 O_3^{2^-} \rightarrow S_4 O_6^{4^-} + I^{\oplus} \\ \text{B. } Mn O_4^{\oplus} + H^{\oplus} + I^{\oplus} \rightarrow Mn^{2^+} + I_2 \\ \\ I_2 + S_2 O_3^{\oplus} \rightarrow S_4 O_6^{2^-} I^{\oplus} \\ \text{C. } Mn O_4^{\oplus} + O H^{\oplus} + I^{\oplus} \rightarrow Mn O_2 + I_2 \\ \\ I_2 + S_2 O_3^{2^-} \rightarrow S_4 O_6^{2^-} + I^{\oplus} \\ \text{D. } Cr_2 O_7^{2^-} + O H^{\oplus} + I^{\oplus} \rightarrow 2 C r^{3^+} + I_2 \\ \\ I_2 + S_2 O_3^{2^-} \rightarrow S_4 O_6^{2^-} + I^{\oplus} \end{array}$$

Answer: a,b

1. In general the melting and boiling points of transition metals

A. increases gradually across the period from left to right

B. decreases gradually across the period from left to right

C. first increases till the middle of the period and then decreases

towards the end

D. First decreases regulary till the middle of the period and then increases towards the end.

Answer: c



2. Which metal has the highest melting point?

A. Pt

B. W

C. Pd

D. Au

Answer: b

Watch Video Solution

3. Which metal has the lowerst melting point?

A. Cs

B. Na

C. Hg

D. Sn

Answer: c

4. Which of the following electronic structures refers to transition elements?

A. 2, 8, 18, 5

B. 2, 8, 14, 2

C. 2, 8, 18, 32, 18, 8, 1

D.2, 8, 5

Answer: b

Watch Video Solution

5. The inner transition elements are the elements which the added electrons go to

A. (n-1) d-orbitals

B. (n-2) f-orbitals

C. (n-1) d-orbitals and (n-1) f-orbitals

D. (n-1) d-orbitals and ns orbitals.

Answer: b

Watch Video Solution

6. The electronic configuration of actinides Cannot be assigned with degree of certainty because of

A. Samll energy difference between 5f and 6d levels

B. overlapping of inner orbitals

C. Free movement of electrons over all the orbitals

D. None of above

Answer: a

7. The second series of transition element start with

A. Ytterium

B. Yttribium

C. Scandium

D. Rhodium

Answer: a

Watch Video Solution

8. Which of the following does not belong to 3d series of transition

elements?

A. Titanium

B. Iron

C. Palladium

D. Vanadium

Answer: c



9. Among the following series of transition metal ions the one where all meal ions have $3d^2$ electronic configuration is

A.
$$Ti^{3+}, V^{2+}, Cr^{3+}, Mn^{4+}$$

B. $Ti^{\oplus}, V^{4+}, Cr^{6+}, Mn^{7+}$
C. $Ti^{4+}, V^{3+}, Cr^{2+}, Mn^{3+}$
D. $Ti^{2+}, V^{3+}, Cr^{4+}, Mn^{5+}$

Answer: d



10. The first ionisation energies of the elements of the first transition

series (Ti to Cu)

A. Increases as the atomic number increases

B. Decreases as the atomic number increases

C. Do not show any change as the addition of electrons takes place in

the inner (n - i) d-orbitals.

D. Increases from Ti to Mn and then decreases from Mn to Cu.

Answer: a

Watch Video Solution

11. Which has the largest radius?

A. Co^{3+}

B. Mn^{3+}

C. Fe^{3+}

 $\mathsf{D.}\,Fe$

Answer: d

12. The atomic numbers of vanadium (V), chromium (Cr), manganese (Mn) and iron (Fe) are respectively 23, 24, 25 and 26. Which one of these may be expected to have the highest second ionisation enthalpy ?

V

Cr

Mn

Fe

A. V

B. Cr

C. Mn

D. Fe

Answer: b

13. The first ionisation energies of the elements of the transition series.

A. Increases as the atomic number increases

B. Decreases as the atomic number increases

C. Do not show any change as the addition of electrons takes place in

the inner (n-1) d-orbitals.

D. None of these

Answer: a

Watch Video Solution

14. Among 3d transition series the IE

A. Increases regularly in moving from left to right

B. Decrease regularly in moving from left to right

C. Remains constant within the period

D. increases gradually within the period but the relativ increase is not

sharp.

Answer: d

Watch Video Solution

15. Which of the following is the most suitable description of transition

elements?

A. Low melting points

B. No catalytic activity

C. Show variable oxidatio states

D. Exhibit inert pair effect

Answer: c

1. The tendency of the transition elements to form coloured compounds is attributed to

A. Transition of electrons from one atom to the other

B. Transition of electrons from s-orbitals of then outer shells to p-

orbitals.

C. d-d-transition in last but one shell.

D. None of the reason is correct.

Answer: c

Watch Video Solution

2. Which of the following compound is not cooured?

A. Copper (II) sulphat

B. Zinc(II) chloride

C. Chromium(II) sulphate

D. Manganese(II oxalate)

Answer: b

Watch Video Solution

3. Anhydrous cobalt(II) chloride is blue in blue in colour but on dissolving

in water it changes to pink in colour because

A. its oxidation state changes

B. Its magnetic character changes

C. its coordination number changes

D. in water it shows fluorescence

Answer: c

Exercises Single Correct D Block Elements Magnetic Properties

1. Which of the following contains the maximum number of unpaired electrons?

A. $TiCl_3$

B. $MnCl_3$

 $C. FeSO_4$

D. $CuSO_4$

Answer: b



2. Among the following pairs of ions, the lower oxidation state in aqueous solution is more stable than the other,in:

A. Ti^{\oplus}, Ti^{3+} B. Cu^{\oplus}, Cu^{2+} C. Cr^{2+}, Cr^{3+} D. V^{2+}, VO^{2+}

Answer: d

Watch Video Solution

3. In general, the Transition elements exhibit their highest oxidation states in their compounds with elements like:


A. C

B.S

C. S and P

D. F and O

Answer: d



Answer: b



5. In which of the following complexes the metal ion is in zero oxidation state?

A.
$$\left[Cu(NH_3)_4
ight] Cl_2$$


 $\mathrm{B.}\,Zn_2\big[Fe(CN)_6\big]$

 $\mathsf{C.}\,Mn_2(CO)_{10}$

D. $\left[Ag(NH_3)_2\right]Cl$

Answer: c





Answer: a

1. In general the order of reactivity of halogens with transition metals is

- A. $F_2 > Cl_2 > Br_2 > I_2$
- B. $F_2 < Cl_2 < Br_2 < I_2$
- C. $F_2 < Cl_2 > Br_2 < I_2$
- D. $F_2>I_2>Br_2>Cl_2$

Answer: a

Watch Video Solution

2. Which of the following cannot reduce the acidified solution of permanganate?

A. $(COOH)_2$

 $\mathsf{B}.\,H_2$

C. Nascent hydrogen

D. Fe^{2+} ions

Answer: b

Watch Video Solution

3. Which of the following can be employed for the conversion of potassium manganate to potassium permanganate?

A. O_3

 $\mathsf{B.}\,Cl_2$

C. Electrolysis

D. All

Answer: d

4. The blue colour produced on adding H_2O_2 , to acidified K_2Cr_2, O_7 is

due to the formation of

A. CrO_5

B. Cr_2O_3

 $C. CrO_4$

D. CrO_3

Answer: a

Watch Video Solution

5. The equilibrium
$$Cr_2O_7^{2\,-} \, \Leftrightarrow 2CrO_4^{2\,-}$$

A. exist in acidic medium

B. exists in basic medium

C. Exists in neutral medium

D. It does not exist

Answer: b



6. Which of the following compounds is used as the starting material for the preparation of potassium dichromate ?

- A. K_2SO_4 . $Cr_2(SO_4)$
- B. $PbCrO_4$ (chrome yellow)
- C. $FeCr_2O_4$ (chromite)
- D. $PbCrO_4 \ _ \ 4PbO$ (chrome red)

Answer: c

7. The image on an exposed and developed photographic film is due to

A. AgBr

 $\mathsf{B.}\left[Ag(C_2O_3)_2\right]^{3+}$

 $\mathsf{C}.Ag$

D. Ag_2O

Answer: c

Watch Video Solution

8. During titration H_2SO_4 is preferably used over HCl and HNO_3 to make the solution acidic because

A. H_2SO_4 is a strong oxidising agent and it reacts with $KMnO_4$

during titration

B. Some $KMnO_4$ is consumed during the reaction with H_2SO_4

C. H_2SO_4 does not react with $KMnO_4$ or the reducing agent used

D. H_2SO_4 can turn colourless $KMnO_4$ to pink at the end point

Answer: c

Watch Video Solution

9. When $KMnO_4$ solution is added to oxalic acid solution, the decolourisation is slow in the beginning but becomes instantaneous after some time because

A. $Mn^{2\,+}$ acts as autocatalyst

B. CO_2 is formed as the oproduct

C. Reaction is exothermic

D. MnO_4^{θ} catalyses the reaction.

Answer: a

10. Which of the following of manganese is amphoteric?

A. MnO_2

B. Mn_2O_3

 $\mathsf{C}. Mn_2O_7$


D. MnO

Answer: a

Watch Video Solution

11. The number of moles of $K_2 C r_2 O_7$ reduced by one mole of $S n^{2+}$ ions

is



Answer: a



12. CrO_3 dissolves in aqueous NaOh to give

A. $CrO_4^{2\,-}$

- $\mathsf{B.}\, Cr(OH)_3$
- C. $Cr_2O_7^{2\,-}$

 $\mathsf{D.}\, Cr(OH)_2$

Answer: a

13. An explosion takes place when conc. H_2SO_4 is added to $KMnO_4$.

Which of the following is formed?

A. Mn^2O_7

 $\mathsf{B.}\,MnO_2$

C. $MnSO_4$

D. Mn_2O_3

Answer: a



14. Formation of interstitial compound makes the transition metal

A. More soft

B. More ductile

C. More metallic

D. More hard

Answer: d



15. The lanthanide contraction relates to

A. Atomic radii

B. Atomic as well as M^{3+} radii

C. Valence electrons

D. Oxidation states

Answer:

Watch Video Solution

16. The properties of Zr and Hf are similar because

A. Both belong to d-block

B. Both belong to same group of the periodic table

C. both have simular radii

D. both have same number of electrons

Answer: c

Watch Video Solution

17. Although +3 oxidation states is the characteristic oxidation state of lanthanoids but cerium shows +4 oxidation state also. Why?

A. It resembles alkali metals

B. It has very low IE

C. it has tendency to attain noble has configuration

D. It has tendency to attain f^0 configuration

Answer: d

18. Which of the two have almost similar size

- A. $_{-}(22)Ti$ and $_{-}(40)Zr$
- B. $_(41)Nb$ and $_(73)Ta$
- C. $_{-}(39)Y$ and $_{-}(57)La$
- D. $_{-}(20)Ca$ and $_{-}(31)Ir$

Answer: b

Watch Video Solution

19. State True/False

 $La(OH)_3$ is less basic than $Lu(OH)_3$

20. The reason for the stability of Gd^{3+} ion is

A. 4f-subshell-half filled

B. 4f subshell-completely filled

C. Possesses the general electronic configuration of noble gases

D. 4f-subshell empty.

Answer: a

Watch Video Solution

21. Arrange
$$(I)Ce^{3+}, (II)La^{3+}, (III)Pm^{3+}$$
 and $(IV)Yb^{3+}$ in

increasing order of their ionic radii.

A. IV < III < I < II

 $\mathsf{B}.\, I < IV < III < II$

 $\mathsf{C}.\,IV < III < II < I$

D. III < II < I < IV

Answer: a



22. The radius of $La^{3+}(Z=57)$ is 106 pm. Which one of the following

given values will be closest to the radius of Lu^{3+} (Z=71)?

A. 160 pm

B. 140 pm

C. 106 pm

D. 85 pm

Answer: d



23. Lanthanides and actinides resemble in

- A. Electron configuration
- B. Oxidation state
- C. lonisation energy
- **D.** Formation Complexes

Answer: d

> Watch Video Solution

24. Misch metal is

A. An alloy of aluminium

B. A mixture of chromium and lead chromate

C. An alloy of lanthanoid metals

D. An alloy of copper

Answer: c

25. In aqueous solutions Eu^{2+} acts as

A. An oxidising agent

B. A reducing agent

C. Can act either of these

D. Can act as redox agent

Answer: b

Watch Video Solution

26. Which transition metal has lowest density?

A. Sc

B. Ti

C. Zn

Answer: a



27. Among the transition elements the element with lowest melting point belongs to

A. Group 3

B. group 11

C. group 6

D. group 12

Answer: d

28. The stability of particular oxidation state of a metal in aqueous solution is determined by

A. Enthalpy of sublimation of the metal

B. Ionisation energy

C. Enthalpy of hydration of the metal ion

D. All of these

Answer: d

Watch Video Solution

29. Which of the following is not a condition for complex formation?

A. Small atomic size

B. High nuclear charge

C. Variable oxidation states

D. Availability of vacant d orbitals

Answer: c



30. Which is not amphoteric?

A. Al^{3+}

- B. Cr^{3+}
- C. Fe^{3+}
- D. Zn^{2+}

Answer: c

Watch Video Solution

31. Both acid and base may react with which of the following oxides?

A. CaO

 $\mathsf{B.}\,Na_2O_2$

C. ZnO

D. Mn_3O_4

Answer: c

Watch Video Solution

32. Coinage metals show the properties of

A. Typical elements

B. Normal elements

C. Transitional elements

D. Inert elements

Answer: c

33. When $(NH_4)_2 CrO_2O_7$ is heated, the gas evolved is

A. N_2

 $\mathsf{B.}\,NO_2$

 $\mathsf{C}.O_2$

D. Na_2O

Answer: a

Watch Video Solution

34. Which of the following oxides of Cr is amphoteric.

A. CrO_2

 $\mathsf{B.}\, Cr_2O_3$

 $C. CrO_5$

D. CrO_3

Answer: a



Exercises Single Correct Compounds Of Transition Elements Iron

1. Important ore of iron is

A. Garnierite

B. Pentalandite

C. Haematite

D. Smaltite

Answer: c

2. A compound of iron exists as a dimer in vapour state. It is hygroscopic in nature and disolves in water giving brown acidic solution. The compound is

A. Fe_3O_4

B. $FeSO_4$

 $C. FeCl_3$

D. FeO_3

Answer: c

Watch Video Solution

3. Which among FeO and Fe_2O_3 is more basic?

A. FeO

B. Fe_2O_3

C. both have same basic length

D. None of them is basic

Answer: a



4. The colour of FeF_3 is

A. Brown

B. Red Brown

C. Light green

D. White

Answer: d

Watch Video Solution

5. Iron is obtained on a large scale from Fe_2O_3 by

A. Reduction with Al

B. Smelting with carbon

C. Calcination

D. Water gas

Answer: b

Watch Video Solution

6. The deep red colour of $Fe(SCN)_3$ and $Fe(SCN)_4^{\theta}$ is destroyed by addition of :

A. F^{Θ}

В. СЛ ө

C. SCN[€]

D. Fe^{Θ}

Answer: a

Exercises Single Correct Compounds Of Transition Elements Copper

1. On heating copper nitrate strongly the compound obtained is

A. Copper

B. Copper oxide

C. Copper nirtite

D. Copper nitride

Answer: b

Watch Video Solution

2. Which of the following is fool's gold?

A. $CuFeS_2$

 $\mathsf{B.}\, Cu(OH)_2$

 $C. Cu(OH)_2. CuCO_3$

 $\mathsf{D.}\, Cu_2S$

Answer: a

Watch Video Solution

3. Bordeaux mixture consists of lime and

A. $FeSO_4$

- $\mathsf{B.}\, CuSO_4$
- $\mathsf{C}.\,Cu(NO_3)_2$
- $\mathsf{D.}\,AgNO_3$

Answer: b

4. In vapour state $Cu(NO_3)_2$ and $Cu_2(CH_3COO)_4.2H_2O$ exist as

A. Dimer, monomer

B. monomer, dimer

C. monomer, monomer

D. dimer, dimer

Answer: b

Watch Video Solution

Exercises Single Correct Compounds Of Transition Elements Silver

1. Which of the following is known as WORM SILVER?

A. AgCl

B. A specific horn, type article made of silver

 $\mathsf{C}.Ag_2S$

D. PbS containing traces of Ag_2S

Answer: a

Watch Video Solution


2. It is always advisable not to cover egg yolk or mustard with silver cutlery because

- A. Silver reacts with water of egg yolk to form AgOH
- B. Silver reacts with sulphur of egg yolk forming black Ag_2S
- C. silver reacts with egg yolk forming Ag_2SO_4 which is a poisonous

substance.

D. Silver attracts UV light of the atmosphere, thereby spoiling the food.

Answer: b



B. Soluble in hyposolution

C. Soluble in NH_4OH

D. Insoluble in acids

Answer: a

Watch Video Solution

4. Which of the following false?


A. molten lead and zinc are miscible

B. silver is more soluble in molten zinc than lead

C. zinc-silver alloy is volatile

D. zinc-silver alloy is heavier and gets solidified later than lead

Answer: d



Answer: d

Exercises Single Correct Compounds Of Transition Elements Gold

1. percentage of gold in 21.6 carat gold is

A. 21.8~%

 $\mathbf{B.\,90~\%}$

 $\mathsf{C}.\,10\,\%$

D. 70 %

Answer: b

Watch Video Solution

2. gold dissolves in aqua regia to give

A. $Au(NO_3)_2$

 $\mathsf{B.}\, H[AuCl_4]$

 $\mathsf{C}.\,AuCl$

D. $AuNO_3$

Answer: b



3. The process of extraction of Au and Ag ores is besed on their solubility

in

A. NH_3

 $\mathsf{B}.\,HCl$

 $C. HNO_3$

D. KCN

Answer: d

Watch Video Solution

Exercises Single Correct Compounds Of Transition Elements Mercury

1. Which of the following is used as purgative in medicine?

A. $ZnCl_2$

B. $HgCl_2$

 $\mathsf{C.}\,Hg_2Cl_2$

D. $ZnSO_{4.7}H_2O$

Answer: c

Watch Video Solution

2. The roasting of HgS in air produces

A. HgO

 $\mathsf{B.}\,HgCl_2$

 $\mathsf{C}.\,HgSO_4$

 $\mathsf{D}.\,Hg$

Answer: d Watch Video Solution 3. Which of the following is used as a white pigment? A. ZnOB. Na_2ZnO_2 C.ZnSD. $ZnCO_3$

Answer: a



4. Mercury (I) chloride sublimes when its compound Is heated and the vapours it gives off are cooled the substance collected consists of

A. Mercury (II) chloride

B. Mercury (I) and mercury (II) chlorides

C. Mercury (II) chloride andmercury

D. mercury

Answer: d

Watch Video Solution

Exercises Single Correct Compounds Of Transition Elements Zinc

1. White vitriol is

A. $CuSO_{4.5}H_2O$

B. $FeSO_{4.7}H_2O$

C. $ZnSO_{4.7}H_2O$

D. $NiSO_{4.5}H_2O$

Answer: c
O Watch Video Solution
2. When zinc reacts with very dilute nitric acid it produces
A. NO
B. NH_4 . NO_3
C. NO_2
D. H_2
Answer: b
Vatch Video Solution

3. Lithapone is a mixture of

A. $ZnSO_4, BaSO_4$

B. ZnS, $BaSO_4$

C. $ZnCO_3$, $BaSO_3$

D. ZnS, HNO_3

Answer: b

Watch Video Solution

Exercises Single Correct Compounds Of Transition Elements Miscellaneous Compounds

1. Iron is rendered passive by treatment with

A. H_2SO_4 is a strong oxidising agent and it reacts with $KMnO_4$

during titration

B. H_3PO_4

 $\mathsf{C}.\,HCl$

D. conc. HNO_3

Answer: d



2. Nitriding is a process of hardening steel by treating it in an atmosphere of

A. NH_3

 $B.O_3$

 $\mathsf{C}.\,N_2$

D. H_2O

Answer: a



3. Which of the following is used for joining the broken pieces of glass

and stones?

A. Heamatite

B. Sindoor

C. Massicot

D. German silver

Answer: c

Watch Video Solution

4. Which method is based on distribution principle?


A. Mond's process

B. Plattner's principle

C. Glower's process

D. Parke's process

Answer: d



Answer: b

Watch Video Solution

6. When H_2S has is passed through an orange red solution (X), the solution turns milky. When an alkali is added to this orange red solution it turns yellow and on acidifying this yellow solution again turns orange red. X is

A. $K_2 CrO_4$

B. $KMnO_4$

 $\mathsf{C.}\, Na_2 Cr_2 O_7$

D. $K_2 Cr_2 O_7$

Answer: d

Watch Video Solution

7. Which of the following statement is incorrect?

A. Basic copper carbonate is $CuCO_3$. $Cu(OH)_2$

B. On strong heating potassium dichromate decomposes with

evolution of oxygen

C. CuS is white in colour

D. $KMnO_4$ exists as dark purple black prismatic crystals

Answer: c

8. A man made white silvery metal, radioactive innature has a strong tendency to form oxocations and complexes. It is used as nuclear fuel in atomic reactors. The metal is a

A. Lanthanide

B. Actinide

C. Transition metal representative element

D. S block element

Answer: b

Watch Video Solution

9. From a solution of $CuSO_4$ the metal used to recover copper is

B.Ag

 $\mathsf{C}.\,Hg$

 $\mathsf{D}.\,Fe$

Answer: d

Watch Video Solution

10. Iron, once dipped in concentrated H_2SO_4 , does not displace copper

from copper sulphate solution, because

A. It is lessreactive than copper

B. A layer of sulphate is deposited on it

C. An inert layer of iron oxide s deposited on it

D. All valence electrons of iron are consumed

Answer: c

11. Which of the followig ions will finally give a black precipitate with Ag^{\oplus} ion?

A. $SO_3^{2\,-}$

 $\mathsf{B.}\,Br^{\, \Theta}$

 $\mathsf{C.}\, CrO_4^{2\,-}$

D. $S_2 O_3^{2\,-}$

Answer: d

Watch Video Solution

12. The compound which gives off oxygen on moderate heating is

A. Zinc oxide

B. Mercuric oxide

C. Aluminium oxide

D. Ferric oxide

Answer: b



13. Identify the statement which is not correct regarding $CuSO_4$?

A. It reacts with KI to give iodine

B. It reacts with KCl to give Cu_2Cl_2

C. It reacts with NaOH and glucose to give Cu_2O

D. It gives CuO on stron heating in air

Answer: b

Watch Video Solution

14. KI and $CuSO_4$ solutions when mixed give

A.
$$CuI_2 + K_2SO_4$$

- $\mathsf{B.}\,Cu_2I_2+K_2SO_4$
- $\mathsf{C}.\,K_2SO_4+Cu_2I_2+I_2$

D.
$$K_2SO_4+CuI_2+I_2$$

Answer: c



15. Bordeaux used a fungicide is a mixture of

A. $CuSO_4 + Ca(OH)_2$

- $\mathsf{B.} \operatorname{CaSO}_4 + \operatorname{Cu}(OH)_2$
- $C. CuCO_3 + Cu(OH)_2$
- $\mathsf{D.}\, CuO+CaO$

Answer: a

16. Mercury is the only metal which is liquid at $0^\circ C$ this is due to its

A. Very high ionisation energy and weakly metallic bond

B. Low ionisation potential

C. High atomic weight

D. High vapour pressure

Answer: a

Watch Video Solution

17. Which of the following is false for Hg?

A. It can evolve hydrogen from H_2S

B. It is metal

C. It has high specific heat

D. It is less reactive than hydrogen

Answer: a



18. Which compound is volatile on heating?

A. $MgCl_2$

B. $HgCl_2$

 $C. ZnCl_2$

D. None of these

Answer: b

19. Which one of the following pairs of substances on reaction will not evolve H_2 gas ?

A. Iron and H_2SO_4

B. Iron and steam

C. Copper and HCl

D. Sodium and ethyl alcohol

Answer: c

Watch Video Solution

20. Which of the following is not the characteristic of zinc?

A. It is volatile metal

B. It dissolves in alkali forming sodium zincate

C. It is brittle at very high temperature

D. Zinc dust is used as a reducing agent

Answer: c



21. The correct formula of permaganic acid is

A. $HMnO_4$

B. $HMnO_5$

 $\mathsf{C}.\,H_2MnO_4$

D. H_2MnO_3

Answer: a

Watch Video Solution

Exercises Assertion Reasoning

1. Assertion: The second ionisation energies of V, Cr and Mn are in the order V < Cr < Mn.

Reason: Ionisation energies show a regular increase along a period with increase of atomic number.

A. If both (A) and (R) are correct and (R) is the correct explanation of

(A).

B. If both (A) and (R) are correct, but (R) is not the correct explanation

of (A).

C. If (A) is correct, but (R) is incorrect.

D. If (A) is incorrect but (R) is correct.

Answer: d



2. Metal-metal bonding is more frequent in 4d or 5d series than in 3d

series due to Tyndall effect. True/False

A.		
В.		
C.		
D.		

Answer: a



3. Assertion: Mercuryis not considered as a transition element.

Reason: Mercury is liquid.

A. If both (A) and (R) are correct and (R) is the correct explanation of

(A).

B. If both (A) and (R) are correct, but (R) is not the correct explanation

of (A).

C. If (A) is correct, but (R) is incorrect.

D. If (A) is incorrect but (R) is correct.

Answer: b



- **4.** Assertion: Ti^{3+} salts are coloured whereas Ti^{4+} salts are white. Reason: Ti^{3+} is less stable than Ti^{4+}
 - A. If both (A) and (R) are correct and (R) is the correct explanation of

(A).

B. If both (A) and (R) are correct, but (R) is not the correct explanation

of (A).

- C. If (A) is correct, but (R) is incorrect.
- D. If (A) is incorrect but (R) is correct.

Answer: b

5. Assertion: Cu, Ag and Au lie in Group 11 of the periodic table. Hence their atomic radii are in the order Cu < Ag < Au

Reason: In any group iof the transition elements, the atomic radii increase as we move down the group.

A. If both (A) and (R) are correct and (R) is the correct explanation of

(A).

B. If both (A) and (R) are correct, but (R) is not the correct explanation of (A).

C. If (A) is correct, but (R) is incorrect.

D. If (A) is incorrect but (R) is correct.

Answer: d

6. Assertion : $K_2Cr_2O_7$ is used as primary standard in volumetric analysis.

Reason : It has a good solubility in water.

- A. If both (A) and (R) are correct and (R) is the correct explanation of
 - (A).
- B. If both (A) and (R) are correct, but (R) is not the correct explanation

of (A).

- C. If (A) is correct, but (R) is incorrect.
- D. If (A) is incorrect but (R) is correct.

Answer: c

Watch Video Solution

7. Statement-1: Tungsten has very high melting point

Statament-2: Tungsten is a convalent compounds

A. If both (A) and (R) are correct and (R) is the correct explanation of

(A).

B. If both (A) and (R) are correct, but (R) is not the correct explanation

of (A).

- C. If (A) is correct, but (R) is incorrect.
- D. If (A) is incorrect but (R) is correct.

Answer: c

Watch Video Solution

8. Assertion: Manganese shows a number of oxidation states.

Reason: The difference of energy between 3d and 4s subshells is large.

A. If both (A) and (R) are correct and (R) is the correct explanation of

(A).

B. If both (A) and (R) are correct, but (R) is not the correct explanation

of (A).

C. If (A) is correct, but (R) is incorrect.

D. If (A) is incorrect but (R) is correct.

Answer: c

Watch Video Solution

9. Assertion: Equivalent mass of $KMnO_4$ is same in the neutral as well as alkaline medium.

Reason: The product formed in both cases in MnO_2

A. If both (A) and (R) are correct and (R) is the correct explanation of

(A).

B. If both (A) and (R) are correct, but (R) is not the correct explanation

of (A).

C. If (A) is correct, but (R) is incorrect.

D. If (A) is incorrect but (R) is correct.

Answer: a

Watch Video Solution

10. Assertion: In $Cr_2O_7^{2-}$ ion, all the Cr-O bond lengths are equal.

Reason: In $Cr_2O_7^{2-}$ ion all the O - Cr - O bond angles are equal.

A. If both (A) and (R) are correct and (R) is the correct explanation of

B. If both (A) and (R) are correct, but (R) is not the correct explanation

of (A).

- C. If (A) is correct, but (R) is incorrect.
- D. If (A) is incorrect but (R) is incorrect.

Answer: d

⁽A).



11. Assertion: In any transition series the magnetic moment of $M^{2\,+}$

ions first increases then decreases

Reason: In a transition series, the number of unpaired electrons first increases and then decreases.

A. If both (A) and (R) are correct and (R) is the correct explanation of

(A).

B. If both (A) and (R) are correct, but (R) is not the correct explanation of (A).

C. If (A) is correct, but (R) is incorrect.

D. If (A) is incorrect but (R) is correct.

Answer: a

12. Assertion: MnO is basic whereas Mn_2O_7 is acidic.

Reason: Higher the oxidation state of a transition metal in its oxide, greater is the acidic character.

A. If both (A) and (R) are correct and (R) is the correct explanation of

(A).

B. If both (A) and (R) are correct, but (R) is not the correct explanation

of (A).

C. If (A) is correct, but (R) is incorrect.

D. If (A) is incorrect but (R) is correct.

Answer: a

Watch Video Solution

13. Assertion : Ce^{4+} is used as an oxidising agent in volumetric analysis.

Reason : Ce^{4+} has the tendency of attaining +3 oxidation state.

A. If both (A) and (R) are correct and (R) is the correct explanation of

(A).

B. If both (A) and (R) are correct, but (R) is not the correct explanation

of (A).

- C. If (A) is correct, but (R) is incorrect.
- D. If (A) is incorrect but (R) is correct.

Answer: a

Watch Video Solution

14. Statament-1: Mn atom loses ns electrons first during ionisation as compared to (n-1) d-electrons.

Statement-1: The effective nuclear charge experienced by (n-1) d elctrons is greater than that by ns electrons.

A. If both (A) and (R) are correct and (R) is the correct explanation of

(A).

B. If both (A) and (R) are correct, but (R) is not the correct explanation

of (A).

C. If (A) is correct, but (R) is incorrect.

D. If (A) is incorrect but (R) is correct.

Answer: a

Watch Video Solution

15. Assertion: Actinoids show greater number of oxidation states than lanthanoids. Reason: Actinoids are radioactive.

A. If both (A) and (R) are correct and (R) is the correct explanation of

(A).

B. If both (A) and (R) are correct, but (R) is not the correct explanation

of (A).

C. If (A) is correct, but (R) is incorrect.

D. If (A) is incorrect but (R) is correct.

Answer: b

Watch Video Solution

16. Assertion : E° for Mn^{3+}/Mn^{2+} is more positve than Cr^{3+}/Cr^{2+} . Reason : The third ionization energy of Mn is larger than that of Cr.

A. If both (A) and (R) are correct and (R) is the correct explanation of

(A).

B. If both (A) and (R) are correct, but (R) is not the correct explanation

of (A).

- C. If (A) is correct, but (R) is incorrect.
- D. If (A) is incorrect but (R) is correct.

Answer: a

17. Assertion: The spin only magnetic moment of Sc^{3+} is 1.73 BM. Reason: The spin only magnetic momentum in (BM) is equal to $\sqrt{n(n+2)}$.

A. If both (A) and (R) are correct and (R) is the correct explanation of

(A).

B. If both (A) and (R) are correct, but (R) is not the correct explanation of (A).

C. If (A) is correct, but (R) is incorrect.

D. If (A) is incorrect but (R) is correct.

Answer: d



18. Assertion: An aqueous solution of $FeCl_3$ is acidic.

Reason: It is due to cationic hydrolysis of Fe^{3+} ion to give $Fe(OH)_3$ and H^{\oplus} ions.

- A. If both (A) and (R) are correct and (R) is the correct explanation of
 - (A).
- B. If both (A) and (R) are correct, but (R) is not the correct explanation

of (A).

- C. If (A) is correct, but (R) is incorrect.
- D. If (A) is incorrect but (R) is correct.

Answer: a



19. Assertion: La should be the f-block element according to Aufbau principle.

Reason: 57th electron exceptionally enters in 5d orbital.

If both (A) and (R) are correct and (R) is the correct explanation of (A).

If both (A) and (R) are correct, but (R) is not the correct explanation of (A).

If (A) is correct, but (R) is incorrect.

If (A) is incorrect but (R) is correct.

A. If both (A) and (R) are correct and (R) is the correct explanation of

(A).

B. If both (A) and (R) are correct, but (R) is not the correct explanation

of (A).

C. If (A) is correct, but (R) is incorrect.

D. If (A) is incorrect but (R) is correct.

Answer: b

20. Assertion: Mercurous compounds are diamagnetic.

Reason: Two Hg^{\oplus} species which have $6s^1$ configuration are bonded together using s-electrons.

- A. If both (A) and (R) are correct and (R) is the correct explanation of
 - (A).
- B. If both (A) and (R) are correct, but (R) is not the correct explanation

of (A).

- C. If (A) is correct, but (R) is incorrect.
- D. If (A) is incorrect but (R) is correct.

Answer: a



1. What is the value of x in the following equation.

$$Cr_2O_7^{2\,-} + 8H^{\,\oplus} + xS_2O_3^{2\,-}
ightarrow 2Cr^{3\,+} + 3SO_4^{2\,-} + 3S + 4H_2O$$



2. What is the oxidation states of Cr in butterfly structure.



3. What is the value of x in the following equation:

$$2MnO_4^{ heta}+3Mn^{2+}+2H_2O
ightarrow xMnO_2+4H^{\oplus}$$

Watch Video Solution

4. Out of the following how many oxides are acidic.

 $MnO, Mn_2O_3, MnO_2, MnO_3, Mn_2O_7$

5. Out of the following how many of them have magnetic moment value

 $\sqrt{24}BM.$

 $Ti^{2+}, Ti^{3+}, V^{2+}, Cr^{2+}, Cr^{3+}, Mn^{2+}, Fe^{2+}.$

Watch Video Solution


6. Out of the following how many of them are coloured compounds.

 $MnO_{4}^{\, \Theta}, Cr_{2}O_{7}^{2\, -}, CrO_{4}^{2\, -}, Sc^{3\, +}, Ti^{4\, +}, Zn^{2\, +}, Mn^{3\, +}, Cu^{2\, +}, Fe^{2\, +}, Fe^{3\, +}$

Watch Video Solution

7. Out of the following how many oxides are basic:

 $TiO, Sc_2O_3, Ti_2O_3, VO, V_2O_5, CrO_3, Cr_2O_3, CuO, TiO_2$



Watch Video Solution		
2 ion show maximum paramagnetic character among the		
bivalent ions of first transition series.		
Watch Video Solution		
3. Transition elements exhibit oxidation states.		
Watch Video Solution		
4. The chromate ion in acidic medium changes to		
Watch Video Solution		
5. The transition metal present in the alooy gun metal is		
Watch Video Solution		

6. $AgNO_3$ on heating forms, and
Watch Video Solution
7. Iron is copper in the electrochemical series and hence displaces from a solution of copper sulphate.
Watch Video Solution
8. When a solution of iron (II) salt is addedto potassium ferricyanide solution, a coloured product known as is formed.
Watch Video Solution
9. Red form of HgS is used as a pigment under the name of
Watch Video Solution

Exercises True False

1. Mn_2O_7 is a basic oxide.

Watch Video Solution

2. Fe_2O_4 is a mixed oxide of FeO and F_2O_3 .



3. The orange colour of dichromate solution changes of blue on heating

with alkalies due to the formation of chromate ions.

4. Transition metal ions containing $(n-1)d^0$ or $(n-1)d^{10}$ configuration are only coloured.



5. The transition elements are those elements which have partially filled penultimate d-subshells in their elementary form or in their commonly occurring oxidation states.

Watch Video Solution

6. Steel contains more carbon than wrought iron



7. When iron is heated in oxygen, it forms ferroso ferric oxide (Fe_3O_4) .

8. Ferric chloride gives dark blue colour with ammonium thiocyanate solution.

Watch Video Solution		
9. Blister copper is $98~\%$ copper. (T/F)		
Watch Video Solution		
10. silver nitrate is used for making inks and hair dyes.		
Watch Video Solution		

Archives Linked Comprehension

1. p-Amino-N,N-dimethylaniline is added to a strongly acidic solution of X. The resulting solution is treated with a few drops of aqueous solution of Y to yield blue colouration due to the formation of methylene blue. Treatment of aqueous solution of Y with reagent potassium hexacyanoferrate (II) leads to the formation of an intense blue precipitate. The precipitate dissolves on excess addition of the reagent. Similarly, the treatment of the solution of Y with the solution of potassium hexacyanoferrate (III) leads to a brown colouration due to the formation of Z.

Q. Compound X is

A. $NaNO_3$

 $\mathsf{B.}\, NaCl$

 $C. Na_2SO_4$

D. Na_2S

Answer: D

2. p-Amino-N,N-dimethylaniline is added to a strongly acidic solution of X. The resulting solution is treated with a few drops of aqueous solution of Y to yield blue colouration due to the formation of methylene blue. Treatment of aqueous solution of Y with reagent potassium hexacyanoferrate (II) leads to the formation of an intense blue precipitate. The precipitate dissolves on excess addition of the reagent. Similarly, the treatment of the solution of Y with the solution of potassium hexacyanoferrate (III) leads to a brown colouration due to the formation of Z.

Q. Compound Y is

A. $MgCl_2$

B. $FeCl_2$

 $C. FeCl_3$

D. $ZnCl_2$

Answer: C

3. p-Amino-N,N-dimethylaniline is added to a strongly acidic solution of X. The resulting solution is treated with a few drops of aqueous solution of Y to yield blue colouration due to the formation of methylene blue. Treatment of aqueous solution of Y with reagent potassium hexacyanoferrate (II) leads to the formation of an intense blue precipitate. The precipitate dissolves on excess addition of the reagent. Similarly, the treatment of the solution of Y with the solution of potassium hexacyanoferrate (III) leads to a brown colouration due to the formation of Z.

Q. Compound Z is

- A. $Mg_2 \big[Fe(CN)_6\big]$
- $\mathsf{B.}\,Fe\big[Fe(CN)_6\big]$
- $\mathsf{C}.\,Fe_4\big[Fe(CN)_6\big]_3$
- $\mathsf{D}.\,K_2Zn_3\big[Fe(CN)_6\big]_2$

Answer: B

Watch Video Solution

4. Copper is the most noble of first row transition metals and occurs in small deposits in serveral countries. Ores of copper include chalcanthite $(CuSO_4.5H_2O)$, atacanite $[Cu_2Cl(OH)_3]$, cuprite (Cu_2O) , copper glance (Cu_2S) , and malachite $[Cu_2(OH)_2CO_3]$. However, 80 % of the world copper production comes from the ore chalcopyrite $(CuFeS_2)$. The extraction of copper from chalcopyrite involves partial roasting, removal of iron and self-reduction.

Partial roasting of chalcopyrite produces

A. Cu_2S and FeO

B. Cu_2O and FeO

C. CuS and Fe_2O_3

D. Cu_2O and Fe_2O_3

Answer: A

Watch Video Solution

5. Copper is the most noble of first row transition metals and occurs in small deposits in serveral countries. Ores of copper include chalcanthite $(CuSO_4.5H_2O)$, atacanite $[Cu_2Cl(OH)_3]$, cuprite (Cu_2O) , copper glance (Cu_2S) , and malachite $[Cu_2(OH)_2CO_3]$. However, 80 % of the world copper production comes from the ore chalcopyrite $(CuFeS_2)$. The extraction of copper from chalcopyrite involves partial roasting, removal of iron and self-reduction.

Iron is removed from chalcopyrite as.

A. FeO

 $\mathsf{B}.\,FeS$

 $\mathsf{C}. Fe_2O_3$

D. $FeSiO_3$

Answer: D

Watch Video Solution

6. Copper is the most noble of first row transition metals and occurs in small deposits in serveral countries. Ores of copper include chalcanthite $(CuSO_4.5H_2O)$, atacanite $[Cu_2Cl(OH)_3]$, cuprite (Cu_2O) , copper glance (Cu_2S) , and malachite $[Cu_2(OH)_2CO_3]$. However, 80 % of the world copper production comes from the ore chalcopyrite $(CuFeS_2)$. The extraction of copper from chalcopyrite involves partial roasting, removal of iron and self-reduction.

In self-reduction, the reducing species is.

A. S

- $\mathsf{B}.O^{2-}$
- C. S^{2-}

D. SO_2

Answer: C

Watch Video Solution

7. When a metal rod M is dipped into an aqueous colourless concetrated solution of compound N, the solution turns light blue. Addition of aqueous NaCl to the blue solution gives a white precipitate O. Addition of aqueous NH_3 dissolves O and gives an intense blue solution. Q. The metal rod M is

Fe

Cu

Ni

Co

A. Fe

 $\mathsf{B.}\,Cu$

 $\mathsf{C}.\,Ni$

 $\mathsf{D}.\,Co$

Answer: B

> Watch Video Solution

8. When a metal rod M is dipped into an aqueous colourless concetrated solution of compound N, the solution turns light blue. Addition of aqueous NaCl to the blue solution gives a white precipitate O. Addition of aqueous NH_3 dissolves O and gives an intense blue solution. Q. The compound N is

- A. $AgNO_3$
- B. $Zn(NO_3)_2$
- $\mathsf{C}. Al(NO_3)_3$
- D. $Ph(NO_3)_2$

Answer: A



9. When a metal rod M is dipped into an aqueous colourless concetrated solution of compound N, the solution turns light blue. Addition of aqueous NaCl to the blue solution gives a white precipitate O. Addition of aqueous NH_3 dissolves O and gives an intense blue solution. Q. The final solution contains.

A.
$$[Ph(NH_3)_4]^{2+}$$
 and $[ClCl_4]^{2-}$
B. $[Al(NH_3)_4]^{3+}$ and $[Cu(NH_3)_4]^{2+}$
C. $[Ag(NH_3)_2]^{\oplus}$ and $[Cu(NH_3)_4]^{4+}$
D. $[Ag(NH_3)_2]^{\oplus}$ and $[Ni(NH_3)_6]^{2+}$

Answer: C



Archives Multiple Correct

1. Potassium manganate $(K_2 M n O_4)$ is formed when

A. Chlorine is passed through aqueous $KMnO_4$ solution

B. Manganese dioxide is fused with potassium hydroxide in air.

C. Formaldehyde reacts with potassium permanganate in the

presence of a strong alkali.

D. Potassium permanganate reacts with concentrated sulphuric acid.

Answer: B::C

Watch Video Solution

2. The aqueous solution of the following salts will be colored in the case

of

A. $Zn(NO_3)_2$

B. $LiNO_3$

 $C.Co(NO_3)_2$

D. $CrCl_3$

Answer: C::D
Vatch Video Solution
3. Which of the following alloys contain (s) Cu and Zn?
A. Bronze
B. Brass
C. Gun metal
D. Type metal
Answer: A::B::C Watch Video Solution

4. Which of the following statements is/are correct, when a mixture of

NaCl and $K_2Cr_2O_7$ is gently warmed with concentrated H_2SO_4 ?

A. Deep red vapours are evolved.

B. The vapours when passed into NaOH solution give a yellow

solution of Na_2CrO_4 .

C. Chlorine gas is evloved.

D. Chromyl chloride is formed.

Answer: A::B::D

Watch Video Solution

5. The addition of high proportions of maganese makes steel useful in

making rails or railroads because manganese

A. Gives hardness to steel

B. Helph in the formation of oxides of iron

C. Can remove oxygen and sulphur

D. Can show the highest oxidation state of +7

Answer: A::C



6. Reduction of the metal centre in aqueous permanganate ion involves

A. 3 electrons in neutral medium

B. 5 electrons in neutral medium

C. 3 electrons in alkaline medium

D. 5 electrons in acidic medium

Answer: A::D



7. The equilibrium $2Cu^I \Leftrightarrow Cu + Cu^{II}$

In aqueous medium at $25^{\,\circ}\,C$ shifts towards the left in the presence of

А. *NO*^ө

 $\mathsf{B}.\,Cl^{\, \Theta}$

C. SCN^e

 $\mathsf{D.}\, CN^{\, \theta}$

Answer: B::C::D

Watch Video Solution

8. Which of the following arrangements does not represent the correct order of the property stated against it ?

A. $Ni^{2+} < Co^{2+} < Fe^{2+} < Mn^{2+}$: ionic size

B. $Co^{3+} < Fe^{3+} < Cr^{3+} < Sc^{3+}$: Stability in aqueous solution.

C. Sc < Ti < Cr < Mn: number of oxidation states.

D. $V^{2+} < Cr^{2+} < Mn^{2+} < Fe^{2+}$: paramagnetic behaviour

Answer: B::D



Archives Single Correct

1. One of the consituents of German silver is

A. Ag

 $\mathsf{B}.\,Cu$

 $\mathsf{C}.\,Mg$

D. Al

Answer: B

Watch Video Solution

2. Which one of the following dissolves in hot concentrated NaOH?

B. Zn

C. Cu

D. Ag

Answer: B

O Watch Video Solution

3. How many unpaired electrons are present in Ni²⁺ cation? (At. No. = 28)
A. 0
B. 2
C. 4
D. 4

Answer: B

4. In the metallurgy of iron, when limestone is added to the blast furnace, the calcium ions end up in

A. Slag

B. Gangue

C. Metallic calcium

D. Calcium carbonate

Answer: A

Watch Video Solution

5. Iron is rendered passive by treatment with

A. H_2SO_4

 $\mathsf{B.}\,H_3PO_4$

 $\mathsf{C}.\,HCl$

 $D.HNO_3$

Answer: D



6. The types of bonds present in $CuSO_4.5H_2O$ are only

A. Electrovalent and covalent

B. Electrovalent and coordinate covalent

C. Electrovalent, covalent, and coordinate covalent

D. Covalent and coordinate covalent.

Answer: C

7. Zinc-copper couple that can be used as a reducing agent is obtained

by

A. Mixing zinc dust and copper gauze

B. Zinc coated with copper

C. Copper coated with zinc

D. Zinc and copper wires welded together.

Answer: B

Watch Video Solution

8. The reaction which proceeds in the forward direction is

A.
$$Fe_2O_3+ 6\mathrm{HCl}
ightarrow 2FeCl_3+ 3H_2O$$

 $\mathsf{B.} \, NH_3 + H_2O + \mathrm{NaCl} \rightarrow NH_4Cl + NaOH$

C. $SnCl_4 + Hg_2Cl_2
ightarrow SnCl_2 + 2HgCl_2$

D.
$$2CuI + I_2 + 4H^{\oplus}
ightarrow 2Cu^{2+} + 4KI$$

Answer: A



9. Which of the following is formed when execess of KCN is added to an aqueous solution of copper sulphate ? .

A. $Cu(CN)_2$

 $\mathsf{B.}\, K_2\big[Cu(CN)_4\big]$

 $\mathsf{C}.\,K\big[Cu(CN)_2\big]$

D. $K_3ig[Cu(CN)_4ig]$

Answer: D

10. An aqueous solution of $FeSO_4$. $Al_2(SO_4)_3$ and chrome alum is heated with excess of Na_2O_2 and filtered. The materials obtained are

A. A colourless filtrate and a green residue

B. A yellow filtrate and a green residue

C. A yellow filtrate and a brown residue

D. A green filtrate and a brown residue

Answer: C

Watch Video Solution

11. Among the following sulphides , the one that does not dissolve in dil

 HNO_3 is

A. HgS

 $\mathsf{B}.\, PbS$

 $\mathsf{C}.\,CuS$

 $\mathsf{D.}\, CdS$

Answer: A



12. Ammonium dichromate is used in some fire works. The green coloured power blown in the air is

A. CrO_3

 $\mathsf{B.}\, Cr_2O_3$

 $\mathsf{C}.\,Cr$

D. $CrO(O_2)$

Answer: B

13. Which of the following compounds is expected to be coloured?

A. Ag_2SO_4

 $\mathsf{B.}\, CuF_2$

 $\mathsf{C}.\,MgF_2$

 $\mathsf{D.}\, CuCl$

Answer: B

Watch Video Solution

14. Which of the following is an organometallic compound ?

A. Lithium methoxide

B. Lithium acetate

C. Lithium dimethylamide

D. methyl lithium

Answer: D



15. In the dichromate anion $\left(Cr_2O_7^{2-}\right)$

A. Four Cr-O bonds are equivalent

B. six Cr - O bonds are equivalent

C. All Cr-O bonds are equivalent

D. All Cr - O bonds are non-equivalent.

Answer: B

Watch Video Solution

16. On heating ammonium dichromate, the gas evolved is

A. Oxygen

B. Ammonia

C. Nitrous oxide

D. Nitrogen

Answer: D

Watch Video Solution

17. Which of the following species has an atom with +6 oxidation state?

- A. MnO_4^{Θ}
- $\mathsf{B.}\left[Cr(CN)_6\right]^{3-}$
- $\mathsf{C.}\left[NiF_{6}\right]^{2-}$
- $\mathsf{D.} \mathit{CrO}_2 \mathit{Cl}_2$

Answer: D

18. Anhydrous feric chloride is prepared by:

A. Heating hydrated ferric chloride at a high temperature in a stream

of air.

B. Heating metallic iron in astream of dry chlorine gas

C. Reaction of ferric oxide with HCl

D. Reaction of metallic iron with HCl

Answer: B

Watch Video Solution

19. When MnO_2 fused with KOH, a coloured compound is formed . The

product and its colour is:

A. $K_2 MnO_4$, purple

B. $KMnO_4$, purple

C. Mn_2O_3 , brown

D. Mn_3O_4 , black

Answer: A

Watch Video Solution

20. $(NH_4)_2 Cr_2 O_7$ on heating gives a gas which is also given by

A. heating NH_4NO_2

B. Heating NH_4NO_3

 $\mathsf{C.}\, Mg_3N_2+H_2O$

 $\mathsf{D}.\,Na+H_2O_2$

Answer: A

21. Which of the following pair of compounds is expected to exhibit same

colour in aqueous solution?

A. $FeCl_3$ and $CuCl_2$

B. $VOCl_2$ and $CuCl_2$

C. $VOCl_2$ and $FeCl_2$

D. $FeCl_2$ and $MnCl_2$

Answer: B



22. A solution of a metal ion when treated with KI gives a red precipitate which dissolves in excess KI to give a colourless solution. Moreover, the solution of metal ion on treatment with a solution of cobalt (II) thiocyanate gives rise to a deep-blue crystalline precipitate. The metal ions is

A. Pb^{2+}

 $\mathsf{B}.\,Hg^{2\,+}$

 $\mathsf{C.}\, Cu^{2\,+}$

D. CO^{2+}

Answer: B

Watch Video Solution

23. The colour of light absorbed by an aqueous solution of $CuSO_4$ is

A. Orange red

B. Blute green

C. Yellow

D. Violet

Answer: A

24. In the cyanide extraction process of silver form argentite ore , the oxidizing and reducing agents used are

A. O_2 and CO_2

B. O_2 and Zn dust

C. HNO_3 and CO

D. HNO_3 and Zn dust

Answer: B

Watch Video Solution

25. Four successive members of the first series of the transition metals are listed below. For which one of them the standard potential $\left(E^{\,\circ}_{M^{2+}\,/\,M}
ight)$ value has a positive sign

A. Mn(Z=25)

B. Fe(Z=26)C. Co(Z=27)D. Cr(Z=24)

Answer: C

Watch Video Solution

26. Which of the following statements about the interstitial compounds is incorrect?

A. They are much harder than the pure metal.

B. They have higher melting points than the pure metal.

C. The retain metallic conductivity

D. They are chemically ractive.

Answer: D

27. $KMnO_4$ can be prepared from K_2MnO_4 as per the reaction:

The reaction can go the completion by removing OH^{θ} ions by adding.

A. CO_2

 $\mathsf{B.}\,SO_2$

 $\mathsf{C}.\,HCl$

 $\mathsf{D}.KOH$

Answer: A

Watch Video Solution

28. A magnetic moment of 1. 73 BM will be shown by one among the

following

A. $TiCl_4$

 $\mathsf{B.}\left[CoCl_{6}\right]^{4\,-}$

C.
$$\left[Cu(NH_3)_4\right]^{2+}$$

D. $\left[Ni(CN)_4\right]^{2-}$

Answer: C

Watch Video Solution

29. Which of the following lanthanoid ions is diamagnetic? (At. No. Ce=58,

Sm=62, Eu=63, Yb=70)

A. $Eu^{2\,+}$

 $\mathsf{B}.\,Yb^{2\,+}$

C. Ce^{2+}

D. Sm^{2+}

Answer: B

1. (a). Assertion (A) is true, Reason (R) is also true, Reason (R) is the correct explanation for assertion (A).

(b). Assertion (A) is true, Reason (R) is true, Reason (R) is not the correct explanation for Assertion (A).

(c). Assertion (A) is true, Reason (R) is false.

(d). Assertion (A) is false, Reason (R) is true.

Q. Assertion (A): to a solution of potassium chromate, if a strong acid is

added, it changes its colour from yellow to orange.

Reason (R): The colour change is due to the change in oxidation state of potassium chromate.



2. (a). Assertion (A) is true, Reason (R) is also true, Reason (R) is the correct explanation for assertion (A).

(b). Assertion (A) is true, Reason (R) is true, Reason (R) is not the correct

explanation for Assertion (A).

(c). Assertion (A) is true, Reason (R) is false.

(d). Assertion (A) is false, Reason (R) is true.

Q. Assertion (A): Zn^{2+} is diamagnetic.

Reason (R): The electrons are lost from 4s-orbital to form Zn^{2+} .

Watch Video Solution

3. (a). Assertion (A) is true, Reason (R) is also true, Reason (R) is the correct explanation for assertion (A).

(b). Assertion (A) is true, Reason (R) is true, Reason (R) is not the correct explanation for Assertion (A).


(c). Assertion (A) is true, Reason (R) is false.

(d). Assertion (A) is false, Reason (R) is true.

Q. Assertion (A): $\left[Fe(H_2O)_5NO
ight]SO_4$ paramagnetic.


Reason (R): The Fe in $[Fe(H_2O)_5NO]SO_4$ has three unpaired electrons.

1. Oxidation state of Mn in the compound formed by alkaline oxidative fusion of MnO_2 by KOH is :




2. Galvanisation of iron denotes coating with
Watch Video Solution
3. Silver chloride is sparingly soluble in water because its lattice energy is
greater thanenegry.
Watch Video Solution
4. The salts and
$(FeSO_{4.7}H_2O, CuSO_{4.5}H_2O, MnSO_{4.4}H_2O, ZnSO_{4.7}H_2O)$ are
isostructural.
Watch Video Solution
5. Fehling solution A consist of an aqueous solution of copper sulphate
while Fehling solution B consists of an alkaline solution of

l



Archives True False



1. What happens when SO_2 gas is bubbled through an aqueous solution of copper sulphate in the presence of potassium thiocyanate?

Watch Video Solution

2. Given reasons for the following in one or two sentences: silver bromide is used in photography?

Watch Video Solution

3. Complete and balance the following reactions:

(i).
$$Zn + NO_3^{ullet} o Zn^{2+} + NH_4^{\oplus}$$

(ii).
$$Cr_2O_7^{2-} + C_2H_4O o C_2H_4O_2 + Cr^{3+}$$

4. State the conditions under which the following preparations are carried out. Give the necessary equations which need not be balanced. Potassium permanganate from manganese dioxide.



5. Show with balanced equations what happens when the following are mixed.

(i). Aqueous solution of potassium manganate and acid.

(ii). Aqueous solution of potassium chromate and acid.

Watch Video Solution

6. Write the balanced equations for the reactions when

(i). Potassium permanganate interacts with manganese dioxide in the

presence of potassium hydroxide.

(ii). Potassium ferrocyanide is heated with concentrated sulphuric acid.

7. Mention the products formed in the following

(i) Zinc oxide is treated with excess of sodium hydroxide solution

(ii). Iodine is added to a solution of stannous chloride.

Watch Video Solution

8. Write balanced chemical equations for the following :

"Gold is dissolved in aqua regia".

Watch Video Solution

9. Write the balanced equation for the following "potassium permanganate is reacted with warm solution of oxalic acid in the presence of sulphuric acid"

10. Complete and balance the following reactions:

(i).
$$Mn^{2\,+} + PbO_2
ightarrow MnO_4^{\,m \Theta} + H_2O$$

(ii). $Ag^{\,\oplus} + AsH_3
ightarrow H_3AsO_3 + H^{\,\oplus}$



11. Answer the following questions briefly.

What is the actual reducing agent of haematite in blast furnace ?

Watch Video Solution

12. Write the balanced equations for extraction of silver from glance by

cyanide process.



13. Write the balancexd chemical equation for the following

i. Silver chloride is reacted with sodium cyanide and the product thus

formed is allowed to react with zine in an alkline medium .

ii Cobalt (II) solution reacts with KNO_2 in acetic acid medium

Watch Video Solution

14. Write balanced equations for "the extraction of copper from pyrites by self-reduction".

Watch Video Solution

15. Write the balanced chemical equations for the following reactions:

(i). A mixture of potassium dichromate and sodium chloride is heated

with concentrated H_2SO_4 .

(ii). Potassium peramanganate is added to a hot solution of manganous

sulphate.

16. A light bluish-green crystalline compound responds to the following tests:

(i). Its aqueous solution gives a brown precipitate on reaction with alkaline $K_2[HgI_4]$ solution.

(ii). Its aqueous solution gives a blue colour with $K_3[Fe(CN)_6]$ solution.

(iii). Its solution in hydrochloric acid gives a white precipitate with $BaCl_2$ solution. Identify the ions present and suggest the formula of the compound.

Watch Video Solution

- 17. Complete and balace the following chemical reaction:
- (i). Na_2CO_3 is added to a solution of copper sulphate.

 $CuSO_4 + Na_2O_3 + H_2O
ightarrow \ldots + Na_2SO_4$

(ii). Potassium dichromate and concentrated HCl are heated together.

 $K_2Cr_2O_7 + \mathrm{HCl} \rightarrow KCl + \dots + \dots + \dots H_2O$

(iii). Copper reacts with HNO_3 to give NO and NO_2 in molar ratio of

2:1

 $Cu + HNO_3 \rightarrow \ldots + NO + NO_2 + \ldots$

Watch Video Solution

18. Compelete and balance the following:

 $(NH_4)_2S_2O_8+H_2O+MnSO_4
ightarrow$

Watch Video Solution

19. A sample of ferrous oxide has actual formula $Fe_{0.93}O_{1.00}$.In this sample what fraction of metal ions are Fe^{2+} ions ? What type of non - stoichiometric defect is present in this sample ?



20. Complete and balance the following reaction:

(i). ${[MnO_4]}^{2-} + H^{\,\oplus}
ightarrow (ii).$ SO_2 (aq)+Cr_2O_7^(2-)+2H^(o+)to

21. Write a balanced equation for the reaction of argentite with KCN

and name the products in the solution .

Watch Video Solution

22. Write a balancfed equation for the following:

"Reaction of zinc with dilute nitric acid"



23. Thionyl chloride can be synthesised by chlorinating SO_2 using PCl_5 . Thionyl chloride is used to prepare anhydrous ferric chloride starting from its hexahydrated salt. Alternatively, the anhydrous ferric chloride can also be prepared from its hexahydrated salt by treating with 2,2-dimethoxypropane. Discuss all this using balanced chemical equations.



24. When the ore haematite is burnt in air with coke around 2000K along with lime, the process not only produces steel but also produces a silicate slag that is useful in making building materials such as cement. Discuss the same and show through balanced chemical equation.



25. Work out the following using chemical equation.

In moist air, copper corrodes to produce a green layer on the surface?

Watch Video Solution

26. Give reasons for following in one or two sentences:

 CrO_3 is an acid anhydride?

27. An aqueous blue-coloured solution of a transition metal sulphate reacts with H_2S I acidic medium to give a black precipitate A which is insoluble in warm auqous solution of KOH. The blue solution on treatment with KI in weakly acidic medium turns yellow and produces a white precipitate B. Identify the transition metal ion. write the chemical reactions involved in the formation of A and B.

Watch Video Solution

28. When a white crystalline compound X is heated with $K_2Cr_2O_7$ and concentrated H_2SO_4 a reddish-brown gas A is evolved. On passing A into caustic soda solution, a yellow-coloured solution B is obtained neutralising the of lead acetate, a yellow precipitate C is obtained. When X is heated with NaOH solution a colourless gas is evolved and on passing this gas into K_2HgI_4 solution a reddish-brown precipitate D is formed. Identify A,B,C,D, and X. Write the equations of the reactions involved.

29. A_1 and A_2 are two ores of metal M. A_1 on calcination gives a black precipitate, CO_2 and water. Identify A_1 and A_2 .

