



MATHS

BOOKS - SHRI BALAJI MATHS (ENGLISH)

COMPLEX NUMBERS

Exercise 1 Single Choice Problems

1. Let t_1, t_2, t_3 be the three distinct points on circle $|t|=1$. if θ_1, θ_2 and θ_3 be the arguments of t_1, t_2, t_3 respectively then $\cos(\theta_1 - \theta_2) + \cos(\theta_2 - \theta_3) + \cos(\theta_3 - \theta_1)$

A. $\geq -\frac{3}{2}$

B. $\leq -\frac{3}{2}$

C. $\geq \frac{3}{2}$

D. ≤ 2

Answer: A



Watch Video Solution

2. The number of points of intersection of the curves represented by

$$\arg(z - 2 - 7i) = \cot^{-1}(2) \text{ and } \arg\left(\frac{z - 5i}{z + 2 - i}\right) = \pm \frac{\pi}{2}$$

A. 0

B. 1

C. 2

D. None of these

Answer: A



Watch Video Solution

3. Let z_1 and z_2 be the roots of the equation $z^2 + az + b = 0$ z being complex. Further, assume that the origin z_1 and z_2 form an equilateral

triangle then

- A. All a, b, c, d have the same sign
- B. a, b, c have same sign
- C. a, b, d have same sign
- D. b, c, d have same sign

Answer: C



[Watch Video Solution](#)

4. Let z_1 and z_2 be the roots of the equation $z^2 + az + b = 0$ z being complex. Further, assume that the origin z_1 and z_2 form an equilateral triangle then

- A. $a^2 = b$
- B. $a^2 = 2b$
- C. $a^2 = 3b$

D. $a^2 = 4b$

Answer: C



Watch Video Solution

5. Let Z and w be two complex number such that $|zw| = 1$ and $arg(z) - arg(w) = \pi/2$ then

A. 1

B. -1

C. i

D. $-i$

Answer: D



Watch Video Solution

6. If ω is a complex n th root of unity, then $\sum_{r=1}^n (a + b)\omega^{r-1}$ is equal to $\frac{n(n+1)a}{2}$ b. $\frac{nb}{1+n}$ c. $\frac{na}{\omega-1}$ d. none of these

A. $\frac{n(n+1)a}{2\omega}$

B. $\frac{nb}{1-n}$

C. $\frac{na}{\omega-1}$

D. None of these

Answer: C



Watch Video Solution

7. If α and β are complex numbers then the maximum value of

$$\frac{\alpha\bar{\beta} + \bar{\alpha}\beta}{|\alpha\beta|} = \text{(A) 1 (B) 2 (C) } > 2 \text{ (D) } < 1$$

A. 1

B. 2

C. greater than 2

D. less than 1

Answer: B



[Watch Video Solution](#)

8. let z_1, z_2, z_3 and z_4 be the roots of the equation $z^4 + z^3 + 2 = 0$,

then the value of $\prod_{r=1}^4 (2z_r + 1)$ is equal to :

A. 28

B. 29

C. 30

D. 31

Answer: D



[Watch Video Solution](#)

9. If $\arg\left(\frac{z - 6 - 3i}{z - 3 - 6i}\right) = \frac{\pi}{4}$, then maximum value of $|z|$:

A. A) $6\sqrt{2} + 3$

B. B) $6\sqrt{3} + 3$

C. C) $\sqrt{2} + 3$

D. D) 6

Answer: A



Watch Video Solution

10. If $z_1 \neq -z_2$ and $|z_1 + z_2| = \left| \frac{1}{z_1} + \frac{1}{z_2} \right|$ then :

A. at least one of z_1, z_2 is unimodular

B. both z_1, z_2 are unimodular

C. $z_1 \cdot z_2$ is unimodular

D. $z_1 - z_2$ is unimodular

Answer: C



Watch Video Solution

11. If $|z-i| \leq 2$ and $z_0 = 5 + 3i$, the maximum value of $|iz + z_0|$ is

A. $5 + \sqrt{13}$

B. $5 + \sqrt{2}$

C. 7

D. 8

Answer: C



Watch Video Solution

12. about to only mathematics

A. $\pm \frac{\pi}{3}$

B. 0

C. $\pm \frac{\pi}{2}$

D. $\pm \frac{\pi}{6}$

Answer: C



[Watch Video Solution](#)

13. If all the roots of $z^3 + az^2 + bz + c = 0$ are of unit modulus, then (A)

$|a| \leq 3$ (B) $|b| \leq 3$ (C) $|c| = 1$ (D) none of these

A. $|a| \leq 3$

B. $|b| \leq 3$

C. $|c| = 1$

D. All of the above

Answer: D



[Watch Video Solution](#)

14. Let z be a complex number satisfying $\frac{1}{2} \leq |z| \leq 4$, then sum of greatest and least values of $\left|z + \frac{1}{z}\right|$ is :

A. $\frac{65}{4}$

B. $\frac{65}{16}$

C. $\frac{17}{4}$

D. 17

Answer: C

 [Watch Video Solution](#)

15. Let $x - \frac{1}{x} = (\sqrt{2})i$ where $i = \sqrt{-1}$. Then the value of $x^{2187} - \frac{1}{x^{2187}}$ is :

A. $i\sqrt{2}$

B. $-i\sqrt{2}$

C. -2

D. $\frac{i}{\sqrt{2}}$

Answer: A



[Watch Video Solution](#)

16. If $z = re^{i\theta}$ ($r > 0$ & $0 \leq \theta < 2\pi$) is a root of the equation $z^8 - z^7 + z^6 - z^5 + z^4 - z^3 + z^2 - z + 1 = 0$ then number of value of ' θ ' is :

A. 6

B. 7

C. 8

D. 9

Answer: C



[Watch Video Solution](#)

17. Let P and Q be two points on the circle $|w|=r$ represented by w_1 and w_2 respectively, then the complex number representing the point of intersection of the tangents of P and Q is :

A. $\frac{w_1 w_2}{2(w_1 + w_2)}$

B. $\frac{2w_1 \bar{w}_2}{w_1 + w_2}$

C. $\frac{2w_1 w_2}{w_1 + w_2}$

D. $\frac{2\bar{w}_1 w_2}{w_1 + w_2}$

Answer: C



Watch Video Solution

18. If z_1, z_2, z_3 are complex numbers such that $|z_1| = |z_2| = |z_3| = 1$, then $|z_1 - z_2|^2 + |z_2 - z_3|^2 + |z_3 - z_1|^2$ cannot exceed.

A. 58

B. 29

C. 87

D. None of these

Answer: C



Watch Video Solution

19. If $Z = \frac{7 + i}{3 + 4i}$, then find Z^{14} .

A. 2^7

B. $(-2)^7$

C. $(2^7)i$

D. $(-2^7)i$

Answer: C



Watch Video Solution

20. If $|Z-4| + |Z+4|=10$, then the difference between the maximum and the minimum values of $|Z|$ is :

A. 2

B. 3

C. $\sqrt{41} - 5$

D. 0

Answer: A



[Watch Video Solution](#)

Exercise 2 One Or More Than One Answer Is Are Correct

1. Let Z_1 and Z_2 are two non-zero complex number such that $|Z_1 + Z_2| = |Z_1| = |Z_2|$, then $\frac{Z_1}{Z_2}$ may be :

A. $1 + \omega$

B. $1 + \omega^2$

C. ω

D. ω^2

Answer: C::D



Watch Video Solution

2. Let z_1, z_2 and z_3 be three distinct complex numbers , satisfying $|z_1| = |z_2| = |z_3| = 1$ find maximum value of $|z_1 - z_2|^2 + |z_2 - z_3|^2 + |z_3 - z_1|^2$



Watch Video Solution

3.

The triangle formed by complex numbers z, iz, i^2z is

(a) Equilateral

(b) Isosceles

(c) Right angle

(d) Scalene

A. equilateral

B. isosceles

C. right angled

D. isosceles but not right angled

Answer: B::C



Watch Video Solution

4. if $A(z_1), B(z_2), C(z_3), D(z_4)$ lies on $|z|=4$ (taken in order) , where

$z_1 + z_2 + z_3 + z_4 = 0$ then :

A. A) Max. area of quadrilateral ABCD=32

B. B) Max. area of quadrilateral ABCD=16

C. C) The triangle ΔABC is right angled

D. D) The quadrilateral ABCD is rectangle

Answer: A::C::D



[Watch Video Solution](#)

5. If $z_1 = a+ib$ and $z_2 = c+id$ are complex numbers such that $|z_1| = |z_2| = 1$ and $\operatorname{Re}(z_1 z_2) = 0$, then the pair of complex numbers $w_1 = a+ic$ and $w_2 = b+id$ satisfy

A. $|w_1| = 1$

B. $|w_2| = 1$

C. $\operatorname{Re}(w_1 w_2) = 0$

D. All the above

Answer: A::B::C



[Watch Video Solution](#)

6. The solutions of the equation $z^4 + 4iz^3 - 6z^2 - 4iz - i = 0$ represent vertices of a convex polygon in the complex plane. The area of the polygon is :

A. $2^{1/2}$

B. $2^{3/2}$

C. $2^{5/2}$

D. $2^{5/4}$

Answer: D



[Watch Video Solution](#)

7. Least positive argument of the 4^{th} root of the complex number $2 - i\sqrt{12}$ is :

A. $\frac{\pi}{6}$

B. $\frac{\pi}{12}$

C. $\frac{5\pi}{12}$

D. $\frac{7\pi}{12}$

Answer: C

 [Watch Video Solution](#)

8. Let ω be the imaginary cube root of unity and $(a + b\omega + c\omega^2)^{2015} = (a + b\omega^2 + c\omega)$ where a, b, c are unequal real numbers. Then the value of $a^2 + b^2 + c^2 - ab - bc - ca$ equals.

A. 0

B. 1

C. 2

D. 3

Answer: B

 [Watch Video Solution](#)

9. Let n be a positive integer and a complex number with unit modulus is a solution of the equation $Z^n + Z + 1 = 0$, then the value of n can be

A. 62

B. 155

C. 221

D. 196

Answer: A::B::C



[Watch Video Solution](#)

Exercise 3 Comprehension Type Problems

1. Let $f(z)$ is of the form $\alpha z + \beta$, where α, β, z are complex numbers such that $|\alpha| \neq |\beta|$. $f(z)$ satisfies following properties :

(i) If imaginary part of z is non zero, then $f(z) + \overline{f(z)} = f(\bar{z}) + \overline{f(\bar{z})}$

(ii) If real part of z is zero, then $f(z) + \overline{f(z)} = 0$

(iii) If z is real, then $\overline{f(z)}f(z) > (z + 1)^2 \forall z \in \mathbb{R}$

$\frac{4x^2}{(f(1) - f(-1))^2} + \frac{y^2}{(f(0))^2} = 1, x, y \in \mathbb{R}$, in (x, y) plane will represent

:

A. hyperbola

B. circle

C. ellipse

D. pair of line

Answer: A



[Watch Video Solution](#)

2. Let z_1 and z_2 be complex numbers such that $z_1^2 - 4z_2 = 16 + 20i$ and the roots α and β of $x^2 + z_1x + z_2 + m = 0$ for some complex number m satisfies $|\alpha - \beta| = 2\sqrt{7}$.

The locus of the complex number m is a curve

A. a square with side 7 and centre (4,5)

B. a circle with radius 7 and centre (4,5)

C. a circle with radius 7 and centre (-4,5)

D. a square with side 7 and centre (-4,5)

Answer: B



Watch Video Solution

3. Let z_1 and z_2 be complex numbers such that $z_1^2 - 4z_2 = 16 + 20i$ and the roots α and β of $x^2 + z_1x + z_2 + m = 0$ for some complex number m satisfies $|\alpha - \beta| = 2\sqrt{7}$. The value of $|m|$,

A. $5\sqrt{21}$

B. $5 + \sqrt{23}$

C. $7 + \sqrt{43}$

D. $7 + \sqrt{41}$

Answer: D



Watch Video Solution

4. Let z_1 and z_2 be complex numbers such that $z_1^2 - 4z_2 = 16 + 20i$ and the roots α and β of $x^2 + z_1x + z_2 + m = 0$ for some complex number m satisfies $|\alpha - \beta| = 2\sqrt{7}$.

The locus of the complex number m is a curve

A. $7 - \sqrt{41}$

B. $7 - \sqrt{43}$

C. $5 - \sqrt{23}$

D. $5 + \sqrt{21}$

Answer: A



Watch Video Solution

5. Let $z_1 = 3$ and $z_2 = 7$ represent two points A and B respectively on complex plane . Let the curve C_1 be the locus of pint P(z) satisfying $|z - z_1|^2 + |z - z_2|^2 = 10$ and the curve C_2 be the locus of point P(z) satisfying $|z - z_1|^2 + |z - z_2|^2 = 16$

Least distance between curves C_1 and C_2 is :

A. A) 4

B. B) 3

C. C) 2

D. D) 1

Answer: D



[Watch Video Solution](#)

6. Let $z_1 = 3$ and $z_2 = 7$ represent two points A and B respectively on complex plane . Let the curve C_1 be the locus of pint P(z) satisfying $|z - z_1|^2 + |z - z_2|^2 = 10$ and the curve C_2 be the locus of point P(z)

satisfying $|z - z_1|^2 + |z - z_2|^2 = 16$

The locus of point from which tangents drawn to C_1 and C_2 are perpendicular, is :

A. $|z-5|=4$

B. $|z-3|=2$

C. $|z-5|=3$

D. $|z-5|=\sqrt{5}$

Answer: D



Watch Video Solution

7. In an Agrad plane z_1, z_2 and z_3 are, respectively, the vertices of an isosceles triangle ABC with $AC=BC$ and $\angle CAB = \theta$. If z_4 is incentre of triangle, then

The value of $AB \times AC / (IA)^2$ is

A. $\left| \frac{(Z_2 - Z_1)(Z_1 - Z_3)}{(Z_4 - Z_1)} \right|$

- B. $\left| \frac{(Z_2 - Z_1)(Z_3 - Z_1)}{(Z_4 - Z_1)} \right|$
- C. $\left| \frac{(Z_2 - Z_1)(Z_3 - Z_1)}{(Z_4 - Z_1)^2} \right|$
- D. $\left| \frac{(Z_2 - Z_1)(Z_3 - Z_1)}{(Z_4 + Z_1)} \right|$

Answer: C

 [Watch Video Solution](#)

8. In an Argand plane z_1, z_2 and z_3 are, respectively, the vertices of an isosceles triangle ABC with $AC = BC$ and $\angle CAB = \theta$. If z_4 is incentre of triangle, then

The value of $(z_4 - z_1)^2(\cos \theta + 1)\sec \theta$ is

- A. $(Z_2 - Z_1)(Z_3 - Z_1)$
- B. $\frac{(Z_2 - Z_1)(Z_3 - Z_1)}{Z_4 - Z_1}$
- C. $\frac{(Z_2 - Z_1)(Z_3 - Z_1)}{(Z_4 - Z_1)^2}$
- D. $(Z_2 - Z_1)(Z_3 - Z_1)^2$

Answer: A



Watch Video Solution

Exercise 4 Matching Type Problems

1. Let ABCDEF is a regular hexagon $A(z_1), B(z_2), C(z_3), D(z_4), E(z_5), F(z_6)$ in argand plane where A,B,C,D,E and F are taken in anticlockwise manner. If $z_1 = -2, z_3 = 1 - \sqrt{3}i$.

	Column-I		Column-II
(A)	If $z_2 = a + ib$, then $2a^2 + b^2$ is equal to	(P)	8
(B)	The square of the inradius of hexagon is	(Q)	7
(C)	The area of region formed by point $P(z)$ lying inside the incircle of hexagon and satisfying $\frac{\pi}{3} \leq \arg(z) \leq \frac{5\pi}{6}$ is $\frac{m}{n} \pi$, where m, n are relatively prime natural numbers, then $m + n$ is equal to	(R)	5
(D)	The value of $z_4^2 - z_1^2 - z_2^2 - z_3^2 - z_5^2 - z_6^2$ is equal to	(S)	3
		(T)	2



Watch Video Solution

2. Match the following Column I to Column II

Column-I		Column-II	
(A)	Let ω be a non real cube root of unity then the number of distinct elements in the set $\{(1 + \omega + \omega^2 + \dots + \omega^n)^m; n, m \in \mathbb{N}\}$ is :	(P)	3
(B)	Let ω and ω^2 be non real cube root of unity. The least possible degree of a polynomial with real co-efficients having roots $2\omega(2 + 3\omega), (2 + 3\omega)^2, (2 - \omega - \omega^2)$ is	(Q)	4
(C)	Let $\alpha = 6 + 4i$ and $\beta = 2 + 4i$ are two complex numbers on Argand plane. A complex number z satisfying $\text{amp} \left(\frac{z - \alpha}{z - \beta} \right) = \frac{\pi}{6}$ moves on a major segment of a circle whose radius is	(R)	5
(D)	Let z_1, z_2, z_3 are complex numbers denoting the vertices of an equilateral triangle ABC having circumradius equals to unity. If P denotes any arbitrary point on its circumcircle then the value of $\frac{1}{2}((PA)^2 + (PB)^2 + (PC)^2)$ equals to	(S)	7



Watch Video Solution

Exercise 5 Subjective Type Problems

1. Let complex number 'z' satisfy the inequality $2 \leq |x| \leq 4$. A point P is selected in this region at random. The probability that argument of P lies in the interval $\left[-\frac{\pi}{4}, \frac{\pi}{4} \right]$ is $\frac{1}{K}$, then $K =$

 [Watch Video Solution](#)

2. Let Z be a complex number satisfying

$$|Z - 1| \leq |Z - 3|, |Z - 3| \leq |Z - 5|, |Z + i| \leq |Z - i|, |Z - i| \leq |Z - 5i|$$

. Then area of region in which Z lies is A square units, Where A is equal to

:

 [Watch Video Solution](#)

3. If z_1 and z_2 both satisfy $z + \bar{z}r = 2|z - 1|$ and $\arg(z_1 - z_2) = \frac{\pi}{4}$,

then find $\operatorname{Im}(z_1 + z_2)$.

 [Watch Video Solution](#)

4. If

$$|z_1| = 1, |z_2| = 2, |z_3| = 3 \text{ and } |z_1 + z_2 + z_3| = 1, \text{ then } |9z_1z_2 + 4z_3z_1 + z_2z_3|$$

is equal to

A. 3

B. 6

C. 36

D. 216

Answer: 6

 [Watch Video Solution](#)

5. If $|z_1|$ and $|z_2|$ are the distance of points on the curve $5z\bar{z} - 2i(z^2 - \bar{z}^2) - 9 = 0$ which are at maximum and minimum distance from the origin, then the value of $|z_1| + |z_2|$ is equal to :

 [Watch Video Solution](#)

6. Let $\frac{1}{a_1 + \omega} + \frac{1}{a_2 + \omega} + \frac{1}{a_3 + \omega} + \dots + \frac{1}{a_n + \omega} = i$

where $a_1, a_2, a_3 \dots a_n \in \mathbb{R}$ and ω is imaginary cube root of unity , then

evaluate $\sum_{r=1}^n \frac{2a_r - 1}{a_r^2 - a_r + 1}$.



Watch Video Solution

7. If $|z_1| = 2$, $|z_2| = 3$, $|z_3| = 4$ and $|2z_1 + 3z_2 + 4z_3| = 9$, then value of $|8z_2z_3 + 27z_3z_1 + 64z_1z_2|^{1/3}$ is :



Watch Video Solution

8. The sum of maximum and minimum modulus of a complex number z satisfying $|z - 25i| \leq 15$, $i = \sqrt{-1}$ is S , then $\frac{S}{10}$ is :



Watch Video Solution