



India's Number 1 Education App

MATHS

BOOKS - SHRI BALAJI MATHS (ENGLISH)

INVERSE TRIGONOMETRIC FUNTIONS

Exercise 1 Single Choice Problems

1. If $\sin^{-1} x \in \left(0, \frac{\pi}{2}\right)$, then the value of $\tan\left(\frac{\cos^{-1}(\sin(\cos^{-1} x)) + \sin^{-1}(\cos(\sin^{-1} x))}{2}\right)$ is :

A. 1

B. 2

C. 3

D. 4

Answer: A



Watch Video Solution

2. The solution set of inequality $(\cot^{-1} x)(\tan^{-1} x) + \left(2 - \frac{\pi}{2}\right)\cot^{-1} x - 3\tan^{-1} x - 3\left(2 - \frac{\pi}{2}\right) > 0$, is

- A. $x \in (\tan 2, \tan 3)$
- B. $x \in (\cot 3, \cot 2)$
- C. $x \in (-\infty, \tan 2) \cup (\tan 3, \infty)$
- D. $x \in (-\infty, \cot 3) \cup (\cot 2, \infty)$

Answer: B



Watch Video Solution

3. $\sec^2(\tan^{-1} 2) + \cos ec^2(\cot^{-1} 3)$ is equal to

A. 14

B. 15

C. 16

D. 17

Answer: B



Watch Video Solution

4. Sum the series :

$$\tan^{-1}\left(\frac{4}{1+3.4}\right) + \tan^{-1}\left(\frac{6}{1+8.9}\right) + \tan^{-1}\left(\frac{8}{1+15.16}\right) + \dots \infty$$

is :

A. $\cot^{-1}(2)$

B. $\tan^{-1}(2)$

C. $\frac{\pi}{2}$

D. $\frac{\pi}{4}$

Answer: A



Watch Video Solution

5. If $\cot^{-1}(\sqrt{\cos \alpha}) - \tan^{-1}(\sqrt{\cos \alpha}) = x$, then $\sin x$ is $\frac{\tan^2 \alpha}{2}$ (b)
 $\frac{\cot^2 \alpha}{2}$ (c) $\tan^2 \alpha$ (d) $\frac{\cot \alpha}{2}$

A. $\tan^2\left(\frac{\alpha}{2}\right)$

B. $\cot^2\left(\frac{\alpha}{2}\right)$

C. $\tan \alpha$

D. $\cot\left(\frac{\alpha}{2}\right)$

Answer: A



Watch Video Solution

6. The sum of the infinite series
 $\cot^{-1}\left(\frac{7}{4}\right) + \cot^{-1}\left(\frac{19}{4}\right) + \cot^{-1}\left(\frac{39}{4}\right) \dots \infty$

A. $\frac{\pi}{4} - \cot^{-1}(3)$

B. $\frac{\pi}{4} - \tan^{-1}(3)$

C. $\frac{\pi}{4} + \cot^{-1}(3)$

D. $\frac{\pi}{4} + \tan^{-1}(3)$

Answer: C



Watch Video Solution

7. the number of solutions of $\cos^{-1}(1-x) + m \cos^{-1} x = \frac{n\pi}{2}$ where

$m > 0, n \leq 0$

A. 0

B. 1

C. 2

D. none of these

Answer: A



Watch Video Solution

8. Number of solution(s) of the equation $2 \tan^{-1}(2x - 1) = \cos^{-1}(x)$ is

:

A. 1

B. 2

C. 3

D. infinitely many

Answer: A



Watch Video Solution

9. $\sin^{-1}\left(\frac{x^2}{4} + \frac{y^2}{9}\right) + \cos^{-1}\left(\frac{x}{2\sqrt{2}} + \frac{y}{3\sqrt{2}} - 2\right)$

A. $\frac{\pi}{2}$

B. π

C. $\frac{\pi}{\sqrt{2}}$

D. $\frac{3\pi}{2}$

Answer: D



Watch Video Solution

10. The complete solution set of the inequality

$$(\cos^{-1} x)^2 - (\sin^{-1} x)^2 > 0 \text{ is :}$$

A. $\left[0, \frac{1}{\sqrt{2}}\right)$

B. $\left[-1, \frac{1}{\sqrt{2}}\right)$

C. $(-1, 1)$

D. $\left[-1, \frac{1}{2}\right)$

Answer: B



Watch Video Solution

11. Let α, β are the roots of the equation $x^2 + 7x + k(k - 3) = 0$, where $k \in (0, 3)$ and k is a constant. Then the value of $\tan^{-1} \alpha + \tan^{-1} \beta + \tan^{-1} \frac{1}{\alpha} + \tan^{-1} \frac{1}{\beta}$ is :

A. π

B. $\frac{\pi}{2}$

C. 0

D. $-\frac{\pi}{2}$

Answer: C



Watch Video Solution

12. Let $f(x) = a + 2b \cos^{-1} x, b > 0$. If domain and range of $f(x)$ are the same set, then $(b - a)$ is equal to :

A. $1 - \frac{1}{\pi}$

B. $\frac{2}{\pi}$

C. $\frac{2}{\pi} + 1$

D. $1 + \frac{1}{\pi}$

Answer: D



Watch Video Solution

13. about to only mathematics

A. -1

B. 4

C. 5

D. 6

Answer: A



Watch Video Solution

14. The total number of ordered pairs (x, y) satisfying

$|y| = \cos x$ and $y = \sin^{-1}(\sin x)$, where $x \in [-2\pi, 3\pi]$ is equal to :

A. 2

B. 4

C. 5

D. 6

Answer: C



Watch Video Solution

15. If $[\sin^{-1}(\cos^{-1}(\sin^{-1}(\tan^{-1} x)))] = 1$ where $[.]$ denotes integer

function, then complete set of values of x is :

A. $[\tan(\sin(\cos 1)), \tan(\cos(\sin 1))]$

B. $[\tan(\sin(\cos 1)), \tan(\sin(\cos(\sin 1)))]$

C. $[\tan(\cos(\sin 1)), \tan(\sin(\cos(\sin 1)))]$

D. $[\tan(\sin(\cos 1)), 1]$

Answer: B



Watch Video Solution

16. The number of ordered pair(s) (x, y) of real numbers satisfying the equation $1 + x^2 + 2x \sin(\cos^{-1} y) = 0$, is :

A. 0

B. 1

C. 2

D. 3

Answer: B



Watch Video Solution

17. The value of $\tan^{-1} 1 + \tan^{-1} 2 + \tan^{-1} 3$ is

A. (a) $\frac{\pi}{2}$

B. (b) π

C. (c) $\frac{3\pi}{4}$

D. (d) $\frac{5\pi}{8}$

Answer: B



Watch Video Solution

18. The complete set of values of x for which

$2\tan^{-1}x + \cos^{-1}\left(\frac{1-x^2}{1+x^2}\right)$ is independent of x is :

A. $(-\infty, 0]$

B. $[0, \infty)$

C. $(-\infty, -1]$

D. $[1, \infty)$

Answer: A



Watch Video Solution

19. The number of ordered pair(s) (x, y) which satisfy $y = \tan^{-1} \tan x$ and $16(x^2 + y^2) - 48\pi x + 16\pi y + 31\pi^2 = 0$, is :

A. 0

B. 1

C. 2

D. 3

Answer: D



Watch Video Solution

20. Domain (D) and range (R) of $f(x) = \sin^{-1}(\cos^{-1}[x])$, where $[.]$ denotes the greatest integer function, is $D \equiv x \in [1, 2], R \in \{0\}$ $D \equiv x \in [0, 1], R \equiv \{-1, 0, 1\}$

$$\equiv x \in [-1, 1], R \equiv \left\{0, \sin^{-1}\left(\frac{\pi}{2}\right), \sin^{-1}(\pi)\right\}$$

$$\equiv x \in [-1, 1], R \equiv \left\{-\frac{\pi}{2}, 0, \frac{\pi}{2}\right\}$$

A. $D \equiv [1, 2], R \equiv \{0\}$

B. $D \equiv [0, 1], R \equiv \{-1, 0, 1\}$

C. $D \equiv [-1, 1), R \equiv \left\{0, \frac{\pi}{2}, \pi\right\}$

D. $D \equiv [-1, 1], R \equiv \left\{-\frac{\pi}{2}, 0, \frac{\pi}{2}\right\}$

Answer: A



Watch Video Solution

21. If $2\sin^{-1}x + \{\cos^{-1}x\} > \frac{\pi}{2} + \{\sin^{-1}x\}$, then $x \in :$ (where $\{\cdot\}$ denotes fractional part function)

A. $(\cos 1, 1]$

B. $[\sin 1, 1]$

C. $(\sin 1, 1]$

D. none of these

Answer: B



Watch Video Solution

22. If $f(x) = x^{11} + x^9 - x^7 + x^3 + 1$ and $f(\sin^{-1}(\sin 8)) = \alpha$, α is constant, then $f(\tan^{-1}(\tan 8))$ is equal to (a) α (b) $\alpha - 2$ (c) $\alpha + 2$ (d) $2 - \alpha$

A. 2

B. 3

C. 4

D. 1

Answer: A



Watch Video Solution

23. The number of real values of x satisfying the equation

$3\sin^{-1}x + \pi x - \pi = 0$ is/are :

A. 0

B. 1

C. 2

D. -3

Answer: B



Watch Video Solution

24. Range of $f(x) = \sin^{-1}x + x^2 + 4x + 1$ is :

A. $\left[-\frac{\pi}{2} - 2, \frac{\pi}{2} + 6 \right]$

B. $\left[0, \frac{\pi}{2} + 6 \right]$

$$\subset \left[-\frac{\pi}{2} - 2, \infty \right)$$

D. $(-3, \infty)$

Answer: A



Watch Video Solution

25. The solution set of the inequality $(\cosec^{-1}x)^2 - 2\cosec^{-1}x \geq \frac{\pi}{6}(\cosec^{-1}x - 2)$ is $(-\infty, a] \cup [b, \infty)$, then $(a + b)$ equals

A. 0

B. 1

C. 2

D. -3

Answer: B



Watch Video Solution

26. Number of solution of the equation $2\sin^{-1}(x+2) = \cos^{-1}(x+3)$ is :

- A. 0
- B. 1
- C. 2
- D. None of these

Answer: B



Watch Video Solution

27. Find the sum series:

$$\tan^{-1}\left(\frac{1}{3}\right) + \tan^{-1}\left(\frac{1}{7}\right) + \tan^{-1}\left(\frac{1}{13}\right) + \dots \rightarrow \infty$$

A. $\frac{\pi}{4}$

B. $\frac{\pi}{2}$

C. $\frac{\pi}{3}$

D. $\frac{\pi}{6}$

Answer: A



Watch Video Solution

28. If $\tan^{-1} \frac{1}{4} + \tan^{-1} \frac{2}{9} = \frac{1}{2} \cos^{-1} x$ then x is equal to :

A. $\frac{1}{2}$

B. $\frac{2}{5}$

C. $\frac{3}{5}$

D. none of these

Answer: C



Watch Video Solution

29. The set of value of x , satisfying the equation $\tan^2(\sin^{-1} x) > 1$ is :

A. $(-1, 1)$

B. $\left(-\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}\right)$

C. $[-1, 1] - \left(-\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}\right)$

D. $(-1, 1) - \left[-\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}\right]$

Answer: D



Watch Video Solution

30. The sum of the series

$\cot^{-1}\left(\frac{9}{2}\right) + \cot^{-1}\left(\frac{33}{4}\right) + \cot^{-1}\left(\frac{129}{8}\right) + \dots \infty$ is equal to :

A. $\cot^{-1}(2)$

B. $\cot^{-1}(3)$

C. $\cot^{-1}(-1)$

D. $\cot^{-1}(1)$

Answer: A



Watch Video Solution

31. If $\int \frac{\ln(\cot x)}{\sin x \cos x} dx = \frac{-1}{k} \ln^2(\cot x) + C$

(where C is a constant), then the value of k is :

A. 1

B. 2

C. 3

D. $\frac{1}{2}$

Answer: B



Watch Video Solution

32. The number of solutions of $\sin^{-1} x + \sin^{-1}(1+x) = \cos^{-1} x$ is/are

:

- A. 0
- B. 1
- C. 2
- D. infinite

Answer: B



Watch Video Solution

33. The value of x satisfying the equation

$$(\sin^{-1} x)^3 - (\cos^{-1} x)^3 + (\sin^{-1} x)(\cos^{-1} x)(\sin^{-1} x - \cos^{-1} x) = \frac{\pi^3}{16}$$

is : (a) $\cos \frac{\pi}{5}$ (b) $\cos \frac{\pi}{4}$ (c) $\cos \frac{\pi}{8}$ (d) $\cos \frac{\pi}{12}$

A. $\cos \frac{\pi}{5}$

B. $\cos \frac{\pi}{4}$

C. $\cos \frac{\pi}{8}$

D. $\cos \frac{\pi}{12}$

Answer: C



Watch Video Solution

34. Let $f(x) = \tan^{-1} \left(\frac{\sqrt{1+x^2}-1}{x} \right)$ then which of the following is

correct :

A. (a) $f(x)$ has only one integer in its range

B. (b) Range of $f(x)$ is $\left(-\frac{\pi}{4}, \frac{\pi}{4} \right) - \{0\}$

C. (c) Range of $f(x)$ is $\left(-\frac{\pi}{2}, \frac{\pi}{2} \right) - \{0\}$

D. (d) Range of $f(x)$ is $\left[-\frac{\pi}{4}, \frac{\pi}{4} \right] - \{0\}$

Answer: B



Watch Video Solution

35. If $\tan^{-1} \frac{1}{4} + \tan^{-1} \frac{2}{9} = \frac{1}{2} \cos^{-1} x$ then x is equal to

- A. (a) $\frac{1}{2}$
- B. (b) $\frac{2}{5}$
- C. (c) $\frac{3}{5}$
- D. (d) None of these

Answer: C



Watch Video Solution

36. The set of value of x , satisfying the equation $\tan^2(\sin^{-1} x) > 1$ is :

- A. $(-1, 1)$
- B. $\left(-\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}} \right)$
- C. $[-1, 1] - \left(-\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}} \right)$
- D. $(-1, 1) - \left[-\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}} \right]$

Answer: D



Watch Video Solution

37. The sum of the series

$$\cot^{-1}\left(\frac{9}{2}\right) + \cot^{-1}\left(\frac{33}{4}\right) + \cot^{-1}\left(\frac{129}{8}\right) + \dots \dots \infty \text{ is equal to :}$$

A. $\cot^{-1}(2)$

B. $\cot^{-1}(3)$

C. $\cot^{-1}(-1)$

D. $\cot^{-1}(1)$

Answer: A



Watch Video Solution

38. The number of real values of x satisfying

$$\tan^{-1}\left(\frac{x}{1-x^2}\right) + \tan^{-1}\left(\frac{1}{x^3}\right) = \frac{(3\pi)}{4}$$

A. 0

B. 1

C. 2

D. infinitely many

Answer: B



Watch Video Solution

39. Number of integral values of λ such that the equation $\cos^{-1} x + \cot^{-1} x = \lambda$ possesses solution is :

A. 2

B. 8

C. 5

D. 10

Answer: C



Watch Video Solution

40. If the equation $x^3 + bx^2 + cx + 1 = 0$, (b

A. $-\frac{\pi}{2}$

B. $-\pi$

C. $\frac{\pi}{2}$

D. π

Answer: B



Watch Video Solution

41. Range of the function
 $f(x) = \cot^{-1}\{-x\} + \sin^{-1}\{x\} + \cos^{-1}\{x\}$, where $\{.\}$ denotes fractional part function:

A. $\left(\frac{3\pi}{4}, \pi\right)$

- B. $\left[\frac{3\pi}{4}, \pi \right)$
- C. $\left[\frac{3\pi}{4}, \pi \right]$
- D. $\left(\frac{3\pi}{4}, \pi \right]$

Answer: D



Watch Video Solution

42. If $3 \leq a < 4$ then the value of $\sin^{-1}(\sin[a]) + \tan^{-1}(\tan[a]) + \sec^{-1}(\sec[a])$, where $[x]$ denotes greatest integer function less than or equal to x, is equal to :

- A. 3
- B. $2\pi - 9$
- C. $2\pi - 3$
- D. $9 - 2\pi$

Answer: A



43. The number of real solutions of

$y + y^2 = \sin x$ and $y + y^3 = \cos^{-1}(\cos x)$ is/are (a) 0 (b) 1 (c) 3 (d) infinite

A. 0

B. 1

C. 3

D. Infinite

Answer: D



44. Range of $f(x) = \sin^{-1}[x - 1] + 2\cos^{-1}[x - 2]$ ([.] denotes greatest integer function)

A. $\left\{ -\frac{\pi}{2}, 0 \right\}$

B. $\left\{ \frac{\pi}{2}, 2\pi \right\}$

C. $\left\{ \frac{\pi}{4}, \frac{\pi}{2} \right\}$

D. $\left\{ \frac{3\pi}{2}, 2\pi \right\}$

Answer: D



Watch Video Solution

Exercise 2 One Or More Than One Answer Is Are Correct

1. If $f(x) = \sin^{-1}(\sin x)$, $g(x) = \cos^{-1}(\cos x)$ and

$h(x) = \cot^{-1}(\cot x)$, then which of the following is/are correct ?

A. $f(x) = g(x)$ if $x \in \left(0, \frac{\pi}{4}\right)$

B. $f(x) < g(x)$ if $x \in \left(\frac{\pi}{2}, \frac{3\pi}{4}\right)$

C. $f(x) < g(x)$ if $\left(\pi, \frac{5\pi}{4}\right)$

D. $f(x) > g(x)$ if $x \in \left(\pi, \frac{5\pi}{4}\right)$

Answer: A::B::C



Watch Video Solution

2. The solution(s) of the equation $\cos^{-1} x = \tan^{-1} x$ satisfy

A. $x^2 = \frac{\sqrt{5} - 1}{2}$

B. $x^2 = \frac{\sqrt{5} + 1}{2}$

C. $\sin(\cos^{-1} x) = \frac{\sqrt{5} - 1}{2}$

D. $\tan(\cos^{-1} x) = \frac{\sqrt{5} - 1}{2}$

Answer: A::C



Watch Video Solution

3. If the numerical value of $\tan\left(\cos^{-1}\left(\frac{4}{5}\right) + \tan^{-1}\left(\frac{2}{3}\right)\right)$ is $\left(\frac{a}{b}\right)$,

where a, b are two positive integers and their H.C.F. is 1

A. $a + b = 23$

B. $a - b = 11$

C. $3b = a + 1$

D. $2a = 3b$

Answer: A::B::C



Watch Video Solution

4. A solution of the equation

$\cot^{-1} 2 = \cot^{-1} x + \cot^{-1}(10 - x)$ where $1 < x < 9$ is : (a) 7 (b) 3

(c) 2 (d) 5

A. 7

B. 3

C. 2

D. 5

Answer: A::B



Watch Video Solution

5. Consider the equation $\sin^{-1} \left(x^2 - 6x + \frac{17}{2} \right) + \cos^{-1} k = \frac{\pi}{2}$, then :

- A. the largest value of k for which equation has 2 distinct solution is 1
- B. the equation must have real root if $k \in \left(-\frac{1}{2}, 1 \right)$
- C. the equation must have real root if $k \in \left(-1, \frac{1}{2} \right)$
- D. the equation has unique solution if $k = -\frac{1}{2}$

Answer: A::B::D



Watch Video Solution

6. The value of x satisfying the equation

$$(\sin^{-1} x)^3 - (\cos^{-1} x)^3 + (\sin^{-1} x)(\cos^{-1} x)(\sin^{-1} x - \cos^{-1} x) = \frac{\pi^3}{16}$$

is : (a) $\cos \frac{\pi}{5}$ (b) $\cos \frac{\pi}{4}$ (c) $\cos \frac{\pi}{8}$ (d) $\cos \frac{\pi}{12}$

A. $\cos \frac{\pi}{5}$

B. $\cos \frac{\pi}{4}$

C. $\cos \frac{\pi}{8}$

D. $\cos \frac{\pi}{12}$

Answer: A::B::D



Watch Video Solution

Exercise 3 Comprehension Type Problems

1. Let $\cos^{-1}(4x^3 - 3x) = a + b \cos^{-1} x$

Q. If $x \in \left[-\frac{1}{2}, \frac{1}{2} \right]$, then $\sin^{-1}\left(\sin \frac{a}{b}\right)$ is :

A. $-\frac{\pi}{3}$

B. $\frac{\pi}{3}$

C. $-\frac{\pi}{6}$

D. $\frac{\pi}{6}$

Answer: A



Watch Video Solution

2. Let $\cos^{-1}(4x^3 - 3x) = a + b \cos^{-1} x$

If $x \in \left(\frac{1}{2}, 1\right]$, then the value of $\lim_{y \rightarrow a} b \cos(y)$ is

A. $-\frac{1}{3}$

B. -3

C. $\frac{1}{3}$

D. 3

Answer: D



Watch Video Solution

1.

	Column-I		Column-II
(A)	$\sin^{-1} \frac{4}{5} + 2 \tan^{-1} \frac{1}{3} =$	(P)	$\frac{\pi}{6}$
(B)	$\sin^{-1} \frac{12}{13} + \cos^{-1} \frac{4}{5} + \tan^{-1} \frac{63}{16} =$	(Q)	$\frac{\pi}{2}$
(C)	If $A = \tan^{-1} \frac{x\sqrt{3}}{2\lambda - x}$, $B = \tan^{-1} \left(\frac{2x - \lambda}{\lambda\sqrt{3}} \right)$ then $A - B$ can be equal to	(R)	$\frac{\pi}{4}$
(D)	$\tan^{-1} \frac{1}{7} + 2 \tan^{-1} \frac{1}{3} =$	(S)	π
		(T)	$\frac{\pi}{3}$



Watch Video Solution

Exercise 5 Subjective Type Problems

1. The complete set of values of x satisfying the inequality $\sin^{-1}(\sin 5) > x^2 - 4x$ is $(2 - \sqrt{\lambda - 2\pi}, 2 + \sqrt{\lambda - 2\pi})$, then $\lambda =$



Watch Video Solution

2. If $2\tan^{-1} \frac{1}{5} - \sin^{-1} \frac{3}{5} = -\cos^{-1} \frac{63}{\lambda}$, then $\lambda =$



Watch Video Solution

3. If $2\tan^{-1}\frac{1}{5} - \sin^{-1}\frac{3}{5} = -\cos^{-1}\frac{9\lambda}{65}$, then $\lambda =$

 Watch Video Solution

4. If $\sum_{n=0}^{\infty} 2 \cot^{-1} \left(\frac{n^2 + n + 4}{2} \right) = k\pi$, then find the value of k .

 Watch Video Solution

5. Find number of solutions of the equation $\sin^{-1}(|\log_6(\cos x) - 1|) + \cos^{-1}(|3\log_6(\cos x) - 7|) = \frac{\pi}{2}$, if $x \in [0, 4\pi]$.

 Watch Video Solution