




## CHEMISTRY

### BOOKS - VK JAISWAL ENGLISH


### CHEMICAL BONDING (BASIC)

#### Level 1


1. The correct order of boiling point is :



(I)



(II)



(III)

A.  $I > II > III$

B.  $III > II > I$

C.  $II > I > III$

D.  $III > I > II$

Answer: B

 [Watch Video Solution](#)

2. Which of the following is not true about  $H_2O$  molecule ?

A. The molecule has  $\mu = 0$

B. The molecule can act as a base



C. Shows abnormally high boiling point in comparison to the hydrides  
of other elements of oxygen group

D. The molecule has a bent shape

Answer: A

 [Watch Video Solution](#)

3. The boiling points at atmospheric pressure of  $HF$ ,  $H_2S$ ,  $NH_3$  can be arranged in the following order :







**Answer: A**



[Watch Video Solution](#)

4. The correct order of strength of  $H -$  bond in the following compound :









**Answer: D**

 [Watch Video Solution](#)

5. Which compound has electrovalent, covalent, co-ordinate as well as hydrogen bond ?



**Answer: A**

 [Watch Video Solution](#)

6. Which statement is correct ?

- A. m.p. of  $H_2O$ ,  $NH_3$  are maximum in their respective group due to intermolecular H-Bonding
- B. b.p. of  $CH_4$  out of  $CH_4$ ,  $SiH_4$ ,  $GeH_4$  and  $SnH_4$  is least due to weak intermolecular force of attraction
- C. formic acid forms dimer by H-bonding
- D. all are correct

**Answer: D**

 [Watch Video Solution](#)

7. Which of the following molecules are expected to exhibit intermolecular H-bonding ?

(I) Acetic acid (II) o-nitrophenol (III) m-nitrophenol (IV) o-boric acid

Select correct alternate :

A. I, II, III

B. I,II,IV

C. I,III,IV

D. II,III,IV

**Answer: C**

 [Watch Video Solution](#)

8. Which of the following compounds can form H-bonding with each other ?

A.  $CH_3COOH$  and  $H_2O$

B. Phenol and  $CH_4$

C.  $CHF_3$  and acetone

D.  $PH_3$  and HF

**Answer: A**

 [Watch Video Solution](#)

9.  $BF_3$  and  $NF_3$  both are covalent compounds but  $NF_3$  is polar whereas  $BF_3$  is non-polar. This is because :

- A. Nitrogen atom is smaller than boron atom
- B. N-F bond is more polar than B-F bond
- C.  $NF_3$  is pyramidal whereas  $BF_3$  is planar triangular
- D.  $BF_3$  is electron deficient whereas  $NF_3$  is not

**Answer: C**

 [Watch Video Solution](#)

10. Which of the following molecules will have polar bonds but zero dipole moment?

- A.  $O_2$

B.  $CHCl_3$



C.  $CF_4$

D. none of these


**Answer: C**

 **Watch Video Solution**


11. Which has maximum dipole moment?







C.



D.


Answer: B

 [Watch Video Solution](#)

12. Which of the following compound is planar and non-polar ?

A.  $XeO_4$

B.  $SF_4$



**Answer: C**

 [Watch Video Solution](#)

13.  $H_2O$  has net dipole moment while  $BeF_2$  has zero dipole moment because

A. F is more electronegativity than oxygen

B. Be is more electronegativity than oxygen

C.  $H_2O$  molecule is linear and  $BeF_2$  is bent

D.  $BeF_2$  molecule is linear and  $H_2O$  is bent

**Answer: D**

 [Watch Video Solution](#)

14. Correct set of species with zero dipole moment is :

(i)  $CO_2$     (ii)  $COCl_2$     (iii)  $CH_2Cl_2$     (iv)  $BCl_3$

A. I and iv

B. ii and iv

C. iii and iv

D. I, iii and iv

Answer: A



Watch Video Solution

15. Which pair of molecules are polar species ?

A.  $CO_2$  and  $H_2O$

B.  $BF_3$  and  $PCl_3$

C.  $SO_2$  and  $SCl_2$

D.  $CS_2$  and  $SO_3$

**Answer: C**

 [Watch Video Solution](#)





**16.** In which molecule does the chlorine have the most positive partial charge?

- A. HCl
- B. BrCl
- C.  $OCl_2$
- D.  $SCl_2$

**Answer: C**

 [Watch Video Solution](#)

**17.** Which of the following attraction is strongest ?



**Answer: D**

 [Watch Video Solution](#)

**18. Which is distilled first ?**

A. Liquid  $H_2$

B. Liquid  $CO_2$

C. Liquid  $O_2$

D. Liquid  $N_2$

**Answer: A**

 [Watch Video Solution](#)

19. The molecular size of  $\text{Icl}$  and  $\text{Br}_2$  is approximately same, but b.p. of  $\text{Icl}$  is about  $40^\circ\text{C}$  higher than that of  $\text{Br}_2$ . It is because :

- A.  $\text{Icl}$  bond is stronger than  $\text{Br-Br}$  bond
- B. IE of iodine  $<$  IE of bromine
- C.  $\text{Icl}$  is polar while  $\text{Br}_2$  is nonpolar
- D. I has larger size than Br

Answer: C

 Watch Video Solution

20. Which of the following order of molecular force of attraction among given species is incorrect ?

- A.  $\text{HI} > \text{HBr} > \text{Cl}_2$
- B.  $\text{CH}_3\text{Cl} > \text{CCl}_4 > \text{CH}_4$

C. n-pentane > iso-pentane > neo-pentane

D.  $\text{OH}_2 > \text{O}(\text{CH}_3)_2 > \text{OBr}_2$

**Answer: D**

 [Watch Video Solution](#)

21. Which gas should not be collected over water because of its high solubility in water ?

A.  $\text{H}_2$

B.  $\text{N}_2$

C.  $\text{CH}_4$

D.  $\text{HCl}$

**Answer: D**

 [Watch Video Solution](#)

22. Low melting point is expected for a solid :

- A. Ionic solid
- B. Metallic solid
- C. Molecular solid
- D. Covalent solid

Answer: C



Watch Video Solution

23. Which substance has the strongest London dispersion forces ?

- A.  $SiH_4$
- B.  $CH_4$
- C.  $SnH_4$
- D.  $GeH_4$



**Answer: C**

 [Watch Video Solution](#)

**24.** Which of the following compounds has the lowest boiling point ?

A. HF

B. HCl

C. HBr

D. HI

**Answer: B**

 [Watch Video Solution](#)

**25.** When the substances *Si*, *KCl*, *CH<sub>3</sub>OH* and *C<sub>2</sub>H<sub>6</sub>* are arranged in order of increasing melting point, what is the correct order ?

A.  $Si$ ,  $KCl$ ,  $CH_3OH$ ,  $C_2H_6$

B.  $CH_3OH$ ,  $C_2H_6$ ,  $Si$ ,  $KCl$

C.  $KCl$ ,  $Si$ ,  $C_2H_6$ ,  $CH_3OH$

D.  $C_2H_6$ ,  $CH_3OH$ ,  $KCl$ ,  $Si$

**Answer: D**

 [Watch Video Solution](#)

**26.** Which substance has the highest melting point ?

A.  $CO$

B.  $CO_2$


C.  $SiO_2$

D.  $P_2O_5$

**Answer: C**

 [View Text Solution](#)

27. How many  $sp^2$  and  $sp$ -hybridised carbon atoms are present respectively in the following compound ?



A. 4,2

B. 6,0

C. 3,3

D. 5,1

**Answer: B**



**Watch Video Solution**

28. Which of the following is a correct set with respect to molecule, hybridization, and shape?

- A.  $BeCl_2$ ,  $sp^2$ , linear
- B.  $BeCl_2$ ,  $sp^2$ , triangular planar
- C.  $BCl_3$ ,  $sp^2$ , triangular planar
- D.  $BCl_3$ ,  $sp^3$ , tetrahedral

Answer: C

 [Watch Video Solution](#)

29. Hybridisation of central atom in  $ICl_2^+$  is

- A.  $dsp^2$
- B.  $sp$
- C.  $sp^2$

D.  $sp^3$

**Answer: D**



**Watch Video Solution**

**30.** The state of hybridization of the central atom is not the same as in the others :

A. B in  $BF_3$

B. O in  $H_3O^+$

C. N in  $NH_3$

D. P in  $PCl_3$

**Answer: A**



**Watch Video Solution**

31. The number of  $sp^2 - s$  sigma bonds in benzene are

- A. 3
- B. 6
- C. 12
- D. none of these

**Answer: B**



[Watch Video Solution](#)

32. The hybridization of the central atom will change when :

- A.  $NH_3$  combines with  $H^+$
- B.  $H_3BO_3$  combines with  $OH^-$
- C.  $NH_3$  forms  $NH_2^-$
- D.  $H_2O$  combines with  $H^+$

Answer: B

 [Watch Video Solution](#)

33.  $CH_3 - CH_2 - CH = CH_2$  has hybridisation :

A.  $sp, sp, sp^2, sp^2$

B.  $sp^3, sp^3, sp^2, sp$

C.  $sp^3, sp^3, sp^2, sp^2$



D.  $sp^3, sp^2, sp^2, sp$

Answer: C

 [Watch Video Solution](#)

34. The state of hybridization of xenon of  $XeF_6$  is





A.  $sp^3d^3$



**Answer: B**

 [Watch Video Solution](#)

35. During the complete combustion of methane  $CH_4$ , what change in hybridisation does the carbon atom undergo ?







**Answer: A**

 [Watch Video Solution](#)





36. The hybridisation of central iodine atom in  $IF_5$ ,  $I_3^-$  and  $I_3^+$  are respectively :




**Answer: A**

 [Watch Video Solution](#)

37. In which of the following combination hybridisation of central atom ( \* ) does not change ?





D. none of these

**Answer: C**

 [Watch Video Solution](#)

**38.** Which of the following species used both axial set of d-orbitals in hybridisation of central atom ?



D. none of these

**Answer: C**

 [Watch Video Solution](#)


39. Which bonds are formed by a carbon atom with  $sp^2$ -hybridisation ?

- A.  $4\pi$ -bonds
- B.  $2\pi$ -bonds and  $2\sigma$ -bonds
- C.  $1\pi$ -bonds and  $3\sigma$ -bonds
- D.  $4\sigma$ -bonds

Answer: C

 [Watch Video Solution](#)

40. What are the hybridisation of the carbon atoms labeled  $C_1$  and  $C_2$ , respectively in glycine ?



- A.  $C_1$   $C_2$   
 $sp^2$   $sp^2$
- B.  $C_1$   $C_2$   
 $sp^2$   $sp^3$
- C.  $C_1$   $C_2$   
 $sp^3$   $sp^2$
- D.  $C_1$   $C_2$   
 $sp^3$   $sp^3$

**Answer: C**



**Watch Video Solution**

**41.** The H-O-H bond angles in  $H_3O^+$  are approximately  $107^\circ$ . The orbitals used by oxygen in these bonds are best described as :

- A. p-orbitals
- B. sp-hybrid orbitals
- C.  $sp^2$ -hybrid orbital
- D.  $sp^3$ -hybrid orbital

**Answer: D**

 [Watch Video Solution](#)

42. Which pair of elements can form multiple bond with itself and oxygen ?

A. F,N

B. N, Cl

C. N,P

D. N,C



**Answer: D**

 [Watch Video Solution](#)

43. Which of the following is a covalent compound ?

A.  $Al_2O_3$


B.  $AlF_3$



**Answer: C**

 [Watch Video Solution](#)

**44.** Which of the following is an example of super octet molecule ?







D. All the three

**Answer: D**

 [Watch Video Solution](#)

45. Which of the following molecule is theoretically not possible ?




Answer: C



Watch Video Solution


46. The phosphate of a metal has the formula  $MHPO_4$ . The formula of its chloride would be



**Answer: B**

 [Watch Video Solution](#)

**47.** The compound that has the highest ionic character associated with the X-Cl bond is :







**Answer: D**

 [Watch Video Solution](#)

**48.** The bond having the highest bond energy is :






**Answer: C**

 [Watch Video Solution](#)


**49.** Which of the following species is neither hypervalent nor hypovalent ?



**Answer: D**

 [Watch Video Solution](#)

50. In which of the following species central atom is NOT surrounded by exactly 8 valence electrons ?



**Answer: D**



[Watch Video Solution](#)

51. Which atom can have more than eight valence electrons when it is forming covalent bonds ?



C. F

D. Cl

**Answer: D**



**Watch Video Solution**

**52. Which bond is expected to be the least polar?**

A. O-F

B. P-F

C. Si-N





D. B-F

**Answer: A**



**Watch Video Solution**

53. Which set contains only covalently bonded molecules ?







Answer: A



Watch Video Solution

54. Which molecule does not exist ?



**Answer: B**

 [Watch Video Solution](#)

**55.** Solid NaCl is a bad conductor of electricity because

- A. in solid NaCl there are no ions
- B. solid NaCl is covalent
- C. in solid NaCl there is no mobility of ions
- D. in solid NaCl there are no electrons

**Answer: C**

 [Watch Video Solution](#)

**56.** An ionic compound  $A^+ B^-$  is most likely to be formed when :




- A. the ionization energy of A is high and electron affinity of B is low

- B. the ionization energy of A is low and electron affinity of B is high
- C. both, the ionization energy of A and electron affinity of B are high
- D. both, the ionization energy of A and electron affinity of B are low

**Answer: B**

 [Watch Video Solution](#)

57. A compound contains atom A, B and C . The oxidation number of A is +2, of B is +5 and C is -2. The possible formula of the compound is



**Answer: B**

 [Watch Video Solution](#)

58. Which pair of atoms form strongest ionic bond ?

A. Al and As

B. Al and N

C. Al and Se

D. Al and O

**Answer: D**


 [Watch Video Solution](#)

59. The correct order of increasing  $C - O$  bond length of  $CO$ ,  $CO_3^{2-}$ ,  $CO_2$  is

A.  $CO_3^{2-} < CO_2 < CO$

B.  $CO_2 < CO_3^{2-} < CO$

C.  $CO < CO_3^{2-} < CO_2$




**Answer: A**



**Watch Video Solution**

**60.** Resonance structures can be written for .







**Answer: A**



**Watch Video Solution**

**61.** The correct order of Cl-O bond order is :






Answer: C



Watch Video Solution

62. How many resonance structures can be drawn for the nitrate ion,



A. 1

B. 2

C. 3

D. 4

Answer: C

 [Watch Video Solution](#)

63. Among given species identify the isostructural pairs :

- A. [ $NF_3$  and  $BF_3$ ]
- B. [ $BF_4^-$  and  $NH_4^+$ ]
- C. [ $BCl_3$  and  $BrCl_3$ ]
- D. [ $NH_3$  and  $NO_3^-$ ]

**Answer: B**

 [Watch Video Solution](#)

64. If 0.1 mole  $H_3PO_x$  is completely neutralised by 5.6g KOH then select the true statement.

- A.  $x=3$  and given acid is dibasic
- B.  $x=2$  and given acid is monobasic

C.  $x=3$  and given acid is monobasic

D.  $x=4$  and given acid forms three series of salt

**Answer: B**

 [Watch Video Solution](#)

65. The solid  $PCl_5$  exists as

A.  $PCl_5$

B.  $PCl_4^+ Cl^-$

C.  $PCl_4^+ PCl_6^-$

D.  $PCl_5, Cl_2$

**Answer: C**

 [Watch Video Solution](#)

66. The ratio of  $\sigma$  – bond and  $\pi$  – bond in tetracyano ethylene is :

A. 2 : 1

B. 1 : 1

C. 1 : 2

D. none of these

**Answer: B**



[Watch Video Solution](#)

67. The bonds present in  $N_2O_5$  are .

A. only ionic

B. only covalent





C. covalent and co-ordinate

D. covalent and ionic

**Answer: C**

 [Watch Video Solution](#)

**68.** The pair of species with similar shape is :



**Answer: A**

 [Watch Video Solution](#)

**69.** Which of the following statements is correct in the context of the allene molecule,  $C_3H_4$  ?

- A. The central carbon is  $sp$  hybridized
- B. The terminal carbon atoms are  $sp^2$  hybridized
- C. The planes containing the  $CH_2$  groups are mutually perpendicular to permit the formations of two separate  $\pi$ -bonds
- D. All are correct

**Answer: D**

 [Watch Video Solution](#)

70. Number of S-S bond in  $H_2S_nO_6$  is :

- A.  $n$
- B.  $(n-1)$
- C.  $(n-2)$
- D.  $(n+1)$

**Answer: B**



Watch Video Solution

71. How many S-S bonds, S-O-S bonds,  $\sigma$ -bonds,  $\pi$ -bonds are present in trimer of sulphur trioxide ?

A. 0,3,16,2

B. 0,3,12,6

C. 0,6,12,16

D. 0,4,12,6

Answer: B



Watch Video Solution

72. Number of identical Cr-O bonds in dichromate ion  $Cr_2O_7^{2-}$  is :

A. 4

B. 6

C. 7

D. 8

**Answer: B**

 [Watch Video Solution](#)

**73.** The nodal plane in the  $\pi$ -bond of ethene is located in:

A. the molecular plane

B. a plane parallel to the molecular plane

C. a plane perpendicular to the molecular plane which bisects the carbon-carbon  $\sigma$  bond at right angle





D. a plane perpendicular to the molecular plane which contains the carbon-carbon bond

**Answer: A**

 [Watch Video Solution](#)



74. Which of the following pairs of ions are isoelectronic and isostructural?



**Answer: A**



**Watch Video Solution**

75. In the electronic structure of  $\text{H}_2\text{SO}_4$ , the total number of unshared electrons is :

A. 20

B. 16

C. 12

D. 8

**Answer: B**

 [Watch Video Solution](#)

76. Which of the following xenon compound has the same number of lone pairs as in  $I_3^-$  ?

A.  $XeO_4$

B.  $XeF_4$

C.  $XeF_2$

D.  $XeO_3$

**Answer: C**

 [Watch Video Solution](#)

77. The shape of  $XeF_3^+$  is :

- A. Trigonal planar
- B. Pyramidal
- C. Bent T-shape
- D. See-saw

**Answer: C**



[Watch Video Solution](#)

78. Which of the following shape are not possible for possible value of  $n$  in  $XeF_n$  molecule ?


- A. Linear
- B. Square planar
- C. Trigonal planar
- D. Capped octahedral

Answer: C



Watch Video Solution

79.  $BeCl_2$  is not isostructural with



Answer: D



Watch Video Solution

80. Which statement is true about the most stable Lewis structure for  $CS_2$ ?

- A. There are no lone pairs in molecule
- B. All bonds are double bonds
- C. The central atom does not have an octet of electrons
- D. A sulfur atom must be the central atom for the structure to be stable

**Answer: B**

 [Watch Video Solution](#)

81. Shape of the compounds  $XeF_3^+$  and  $XeF_5^+$  are respectively


- A. Square pyramidal, T-shpaed
- B. Bent-T-shape, square pyramidal
- C. See-saw, square pyramidal
- D. Square pyramidal, see -saw

**Answer: B**



Watch Video Solution

82. In which of the following species maximum atom can lie in same plane ?



Answer: D



Watch Video Solution

83. Correct statement regarding molecules  $SF_4$ ,  $CF_4$  and  $XeF_4$  are :

A. 2,0 and 1 lone pairs of central atom respectively

B. 1,0 and 1 lone pairs of central atom respectively

C. 0,0 and 2 lone pairs of central atom respectively

D. 1,0 and 2 lone pairs of central atom respectively

**Answer: D**

 [Watch Video Solution](#)

**84.** The geometrical arrangement and shape of  $I_3^-$  are respectively

A. trigonal bipyramidal geometry, linear shape

B. hexagonal geometry, T-shape

C. triangular planar geometry, triangular shape

D. tetrahedral geometry, pyramidal shape

**Answer: A**

 [Watch Video Solution](#)

85. Which of the following statements is incorrect for  $PCl_5$  ?

- A. Its three P-Cl bond lengths are equal
- B. It involves  $sp^3d$  hybridization
- C. It has a regular geometry
- D. Its shape is trigonal bipyramidal

Answer: C



Watch Video Solution

86. Molecular shapes of  $SF_4$ ,  $CF_4$ ,  $XeF_4$  are

- A. the same with 2,0 and 1 lone pair of electrons respectively
- B. the same with 1,1 and 1 lone pair of electrons respectively
- C. the same with 0,1 and 2 lone pair of electrons respectively
- D. the same with 1,0 and 2 lone pair of electrons respectively



**Answer: D**



[Watch Video Solution](#)

**87.** The structure of the noble gas compound  $XeF_4$  is :


- A. square planar
- B. distorted tetrahedral
- C. tetrahedral
- D. octahedral

**Answer: A**



[Watch Video Solution](#)





**88.** The molecule exhibiting maximum number of non-bonding electron pairs (l.p.) around the central atom is :



**Answer: C**

 [Watch Video Solution](#)

**89.** Which is the following pairs of species have identical shapes ?



**Answer: C**

 [Watch Video Solution](#)

90. The shapes of  $XeF_4$ ,  $XeF_5^-$  and  $SnCl_2$  are :

- A. octahedral, trigonal bipyramidal and bent
- B. square pyramidal, pentagonal planar and linear
- C. square planar, pentagonal planar and angular
- D. see-saw, T-shaped and linear

Answer: C



Watch Video Solution

91. Which is not correctly matched ?


- A.  $XeO_3$  -Trigonal bipyramidal
- B.  $ClF_3$ - bent T-shape
- C.  $XeOF_4^-$  Square pyramidal

D.  $XeF_2$ - Linear shape

Answer: A

 [Watch Video Solution](#)

92. Amongst  $NO_3^-$ ,  $AsO_3^{3-}$ ,  $CO_3^{2-}$ ,  $ClO_3^-$ ,  $SO_3^{2-}$  and  $BO_3^{3-}$ , the non-planar species are :



Answer: B

 [Watch Video Solution](#)

93. The geometry of ammonia molecule can be best described as :

- A. Nitrogen at one vertex of a regular tetrahedron, the other three vertices being occupied by three hydrogens
- B. Nitrogen at the centre of the tetrahedron, three of the vertices being occupied by three hydrogens
- C. Nitrogen at the centre of an equilateral triangle, three corners being occupied by three hydrogens
- D. Nitrogen at the junction of a T, three open ends being occupied by three hydrogens

**Answer: B**

 [Watch Video Solution](#)

94. Which molecular geometry is least likely to result from a trigonal bipyramidal electron geometry ?

A. Trigonal planar

B. See-saw

C. Linear

D. T-shpaed

**Answer: A**

 [Watch Video Solution](#)

**95.** Give the correct order of initials T or F for following statements. Use T if statement is true and F if it is false :

(I) The order of repulsion between different pair of electron is

$$I_p - I_p > I_p - b_p > b_p - b_p$$

(II) In general, as the number of lone pair of electron on central atom increases, value of bond angle from normal bond angle also increases

(III) The number of lone pair on O in  $H_2O$  is 2 while on N in  $NH_3$  is 1

(IV) The structures of xenon fluorides and xenon oxyfluorides could not be explained on the basis of VSEPR theory

A. T T T F

B. T F T F

C. T F T T

D. T F F F

**Answer: B**

 [Watch Video Solution](#)

**96. Which species is planar ?**



**Answer: A**

 [Watch Video Solution](#)

97. What is the geometry of the  $IBr_2^-$  ion ?

- A. Linear
- B. Bent shape with bond angle of about  $90^\circ$
- C. Bent shape with bond angle of about  $109^\circ$
- D. Bent shape with bond angle of about  $120^\circ$

**Answer: A**



[Watch Video Solution](#)

98. Molecular shape of  $ClF_3$ ,  $I_3^-$  and  $XeO_3$  respectively are

- A. Trigonal planar
- B. Trigonal pyramidal
- C. T-shaped




D. Tetrahedral

Answer: C



Watch Video Solution

99. Which species has the same shape as  $NH_3$ ?



Answer: A



Watch Video Solution

100. According to VSEPR theory, in which species do all the atoms lie in the same plane ?



A. 1 only

B. 2 only

C. both 1 and 2


D. neither 1 nor 2


**Answer: A**



[Watch Video Solution](#)

101. Which of the following species / molecules does not have same number of bond pairs and lone pairs ?





**Answer: D**

 [Watch Video Solution](#)

**102. Least stable hydride is**

A. stannane

B. silane



C. plumbane

D. germane

**Answer: C**

 [Watch Video Solution](#)

103. The lowest O-O bond length in the following molecule is :



**Answer: B**



[Watch Video Solution](#)

104. The fluorine molecules is formed by :

A. p-p orbitals (sideways overlap)

B. p-p orbitals (end-to-end overlap)





C. sp-sp orbitals

D. s-s orbitals

Answer: B

 Watch Video Solution

105. Which of the following leads to bonding?



Answer: B

 Watch Video Solution

106. Which of the following overlaps is incorrect (assuming Z-axis is internuclear axis) ?

(A)  $2p_y + 2p_y \rightarrow \pi$ - Bond formation      (B)  $2p_x + 2p_x \rightarrow \sigma$ - Bond formation

(C)  $3d_{xy} + 3d_{xy} \rightarrow \pi$ - Bond formation      (D)  $2s + 2p_y \rightarrow \pi$ - Bond formation

(E)  $3d_{xy} + 3d_{xy} \rightarrow \delta$ - Bond formation      (F)  $2p_x + 2p_x \rightarrow \sigma$ -Bond formation

A. A,B,C

B. C,F

C. B,E





D. B,C,D

**Answer: D**



**Watch Video Solution**

107. Which of the following overlapping is not present in  $XeO_3$  molecule ?



Answer: D



Watch Video Solution

108. How many sigma bonds are in a molecule of diethyl ether,  $C_2H_5OC_2H_5$  ?

A. 14

B. 12





C. 8

D. 16

Answer: A

 [Watch Video Solution](#)

109. The lattice energies of KF, KCl, KBr and KI follow the order :



Answer: A

 [Watch Video Solution](#)



110. Which set of compounds in the following pair of ionic compounds has the higher lattice energy ?

(i)  $KCl$  or  $MgO$     (ii)  $LiF$  or  $LiBr$     (iii)  $Mg_3N_2$  or  $NaCl$

A.  $KCl$ ,  $LiBr$ ,  $Mg_3N_2$

B.  $MgO$ ,  $LiBr$ ,  $Mg_3N_2$

C.  $MgO$ ,  $LiF$ ,  $NaCl$

D.  $MgO$ ,  $LiF$ ,  $Mg_3N_2$

Answer: D



Watch Video Solution

111. The incorrect order of lattice energy is :

A.  $AlF_3 > MgF_2$

B.  $Li_3N > Li_2O$

C.  $NaCl > LiF$

D.  $TiC > ScN$

**Answer: C**

 [Watch Video Solution](#)

**112.** Which ionic compound has the largest amount of lattice energy ?

A. NaF

B.  $AlF_3$



C.  $AlN$

D.  $MgF_2$

**Answer: C**

 [Watch Video Solution](#)

113. Which of the following compounds has the smallest bond angle in its molecule ?




**Answer: B**



[Watch Video Solution](#)

114. Maximum bond angle is present in :







D. none of these

Answer: D

 [Watch Video Solution](#)


115. The correct order of H-M-H bonds angle is :



Answer: D

 [Watch Video Solution](#)

116. The correct increasing order of adjacent bond angle among  $BF_3$ ,  $PF_3$  and  $ClF_3$  :



**Answer: C**



**Watch Video Solution**

**117.** Among the following species, the least angle around the central atom is in :



**Answer: D**

 [Watch Video Solution](#)

118. Compare the H-N-H bond angle in the following species.



**Answer: B**

 [Watch Video Solution](#)

119. The H-C-H bond angle in  $CH_4$  is  $109.5^\circ$ , due to lone pair repulsion, the H-O-H angle in  $H_2O$  will :

A. remain the same

B. increase

C. decrease

D. become  $180^\circ$

**Answer: C**

 [Watch Video Solution](#)

**120.** The molecule having the largest bond angle is :

A.  $H_2O$

B.  $H_2S$

C.  $H_2Se$

D.  $H_2Te$

**Answer: A**

 [Watch Video Solution](#)

121. The compound  $MX_4$  is tetrahedral. The number of  $\angle XMX$  angles in the compound is :

- A. three
- B. four
- C. five
- D. six

Answer: D




Watch Video Solution

122. Which of the following is the correct order for increasing bond angle ?

- A.  $Nh_3 < PH_3 < AsH_3 < SbH_3$
- B.  $H_2O < OF_2 < Cl_2O$
- C.  $H_2Te^+ < H_3Se^+ < H_3S^+ < H_3O^+$








Answer: C

 Watch Video Solution

123. The correct order of boiling point is :



A.  $I > II > III$

B.  $III > II > I$

C.  $II > I > III$

D.  $III > I > II$

Answer: B

 Watch Video Solution

124. Which of the following is not true about  $H_2O$  molecule ?

- A. The molecule has  $\mu = 0$
- B. The molecule can act as a base
- C. Shows abnormally high boiling point in comparison to the hydrides of other elements of oxygen group
- D. The molecule has a bent shape

Answer: A



Watch Video Solution

125. The boiling points at atmospheric pressure of  $HF$ ,  $H_2S$ ,  $NH_3$  can be arranged in the following order :





- A.  $HF > NH_3 > H_2S$
- B.  $HF > H_2S > NH_3$
- C.  $HF < H_2S < NH_3$



Answer: A

 [Watch Video Solution](#)





126. The correct order of strength of  $H -$  bond in the following compound :



Answer: D

 [Watch Video Solution](#)

127. Which compound has electrovalent, covalent, co-ordinate as well as hydrogen bond ?



Answer: A



Watch Video Solution

128. Which statement is correct ?

A. m.p. of  $H_2O$ ,  $NH_3$  are maximum in their respective group due to intermolecular H-Bonding

B. b.p. of  $CH_4$  out of  $CH_4$ ,  $SiH_4$ ,  $GeH_4$  and  $SnH_4$  is least due to weak intermolecular force of attraction

C. formic acid forms dimer by H-bonding

D. all are correct

**Answer: D**

 [Watch Video Solution](#)

**129.** Which of the following molecules are expected to exhibit intermolecular H-bonding ?

(I) Acetic acid (II) o-nitrophenol (III) m-nitrophenol (IV) o-boric acid

Select correct alternate :

A. I, II, III

B. I,II,IV

C. I,III,IV

D. II,III,IV

**Answer: C**

 [Watch Video Solution](#)



Watch Video Solution

130. Which of the following compounds can form H-bonding with each other ?

A.  $CH_3COOH$  and  $H_2O$

B. Phenol and  $CH_4$

C.  $CHF_3$  and acetone

D.  $PH_3$  and HF

Answer: A



Watch Video Solution

131.  $BF_3$  and  $NF_3$  both are covalent compounds but  $NF_3$  is polar whereas  $BF_3$  is non-polar. This is because :

A. Nitrogen atom is smaller than boron atom

B. N-F bond is more polar than B-F bond

C.  $NF_3$  is pyramidal whereas  $BF_3$  is planar triangular

D.  $BF_3$  is electron deficient whereas  $NF_3$  is not

**Answer: C**

 [Watch Video Solution](#)

**132.** Dipole moment of  $NF_3$  is smaller than :

A.  $NH_3$

B.  $CO_2$

C.  $BF_3$

D.  $CCl_4$

**Answer: A**

 [Watch Video Solution](#)

133. Which of the following molecules will have polar bonds but zero dipole moment ?

A.  $O_2$

B.  $CHCl_3$


C.  $CF_4$

D. none of these

Answer: C


 Watch Video Solution

134. Which has maximum dipole moment ?




A.






B.



C.






D.

**Answer: B**

 [Watch Video Solution](#)

**135.** Which of the following compound is planar and non-polar ?



**Answer: C**

 [Watch Video Solution](#)

**136.**  $H_2O$  has net dipole moment while  $BeF_2$  has zero dipole moment because

A. F is more electronegativity than oxygen

B. Be is more electronegativity than oxygen

C.  $H_2O$  molecule is linear and  $BeF_2$  is bent

D.  $BeF_2$  molecule is linear and  $H_2O$  is bent

**Answer: D**

 [Watch Video Solution](#)

137. Correct set of species with zero dipole moment is :

(i)  $CO_2$     (ii)  $COCl_2$     (iii)  $CH_2Cl_2$     (iv)  $BCl_3$

A. I and iv

B. ii and iv

C. iii and iv

D. I, iii and iv

Answer: A

 [Watch Video Solution](#)

138. Which pair of molecules are polar species ?

A.  $CO_2$  and  $H_2O$

B.  $BF_3$  and  $PCl_3$

C.  $SO_2$  and  $SCl_2$

D.  $CS_2$  and  $SO_3$

**Answer: C**

 [Watch Video Solution](#)

**139.** In which molecule does the chlorine atom has positive partial charge

?

A. HCl

B. BrCl





C.  $OCl_2$

D.  $SCl_2$

**Answer: C**

 [Watch Video Solution](#)

140. Which of the following attraction is strongest ?



Answer: D

 [Watch Video Solution](#)

141. Which is distilled first ?

A. Liquid  $\text{H}_2$

B. Liquid  $\text{CO}_2$

C. Liquid  $\text{O}_2$

D. Liquid  $\text{N}_2$

**Answer: A**



[Watch Video Solution](#)

**142.** The molecular size of  $\text{Icl}$  and  $\text{Br}_2$  is approximately same, but b.p. of  $\text{Icl}$  is about  $40^\circ\text{C}$  higher than that of  $\text{Br}_2$ . It is because :


- A.  $\text{Icl}$  bond is stronger than  $\text{Br-Br}$  bond
- B. IE of iodine  $<$  IE of bromine
- C.  $\text{Icl}$  is polar while  $\text{Br}_2$  is nonpolar
- D. I has larger size than Br

**Answer: C**




[Watch Video Solution](#)

**143.** Which of the following order of molecular force of attraction among given species is incorrect ?



C. n-pentane > iso-pentane > neo-pentane



**Answer: D**

 [Watch Video Solution](#)

**144.** Which gas should not be collected over water because of its high solubility in water ?



**Answer: D**

 [Watch Video Solution](#)

145. Low melting point is expected for a solid :

- A. Ionic solid
- B. Metallic solid
- C. Molecular solid
- D. Covalent solid

**Answer: C**

 [Watch Video Solution](#)

146. Which substance has the strongest London dispersion forces ?

- A.  $SiH_4$
- B.  $CH_4$
- C.  $SnH_4$



D.  $GeH_4$

**Answer: C**



**Watch Video Solution**

**147.** Which of the following compounds has the lowest boiling point ?

A. HF

B. HCl

C. HBr

D. HI

**Answer: B**



**View Text Solution**

148. When the substances  $Si$ ,  $KCl$ ,  $CH_3OH$  and  $C_2H_6$  are arranged in order of increasing melting point, what is the correct order ?

A.  $Si$ ,  $KCl$ ,  $CH_3OH$ ,  $C_2H_6$

B.  $CH_3OH$ ,  $C_2H_6$ ,  $Si$ ,  $KCl$

C.  $KCl$ ,  $Si$ ,  $C_2H_6$ ,  $CH_3OH$

D.  $C_2H_6$ ,  $CH_3OH$ ,  $KCl$ ,  $Si$

Answer: D



Watch Video Solution

149. Which substance has the highest melting point ?

A.  $CO$

B.  $CO_2$


C.  $SiO_2$

D.  $P_2O_5$

Answer: C

 Watch Video Solution

150. How many  $sp^2$  and  $sp$ -hybridised carbon atoms are present respectively in the following compound ?



A. 4,2

B. 6,0

C. 3,3

D. 5,1

**Answer: B**



**Watch Video Solution**

**151.** Which one of the following is the correct set with respect to molecule, hybridization and shape ?

- A.  $BeCl_2$ ,  $sp^2$ , linear
- B.  $BeCl_2$ ,  $sp^2$ , triangular planar
- C.  $BCl_3$ ,  $sp^2$ , triangular planar
- D.  $BCl_3$ ,  $sp^3$ , tetrahedral

**Answer: C**



**Watch Video Solution**

**152.** The hybridization of the central atom in  $ICl_2^+$  is :

A.  $dsp^2$

B.  $sp$

C.  $sp^2$

D.  $sp^3$

**Answer: D**



**Watch Video Solution**

**153.** The state of hybridization of the central atom is not the same as in the others :

A. B in  $BF_3$

B. O in  $H_3O^+$

C. N in  $NH_3$

D. P in  $PCl_3$

**Answer: A**

 [Watch Video Solution](#)

154. The number of  $sp^2$ -s sigma bonds in benzene are :

- A. 3
- B. 6
- C. 12
- D. none of these

**Answer: B**

 [Watch Video Solution](#)

155. The hybridization of the central atom will change when :

- A.  $NH_3$  combines with  $H^+$
- B.  $H_3BO_3$  combines with  $OH^-$
- C.  $NH_3$  forms  $NH_2^-$

D.  $H_2O$  combines with  $H^+$

**Answer: B**

 [Watch Video Solution](#)

156.  $CH_3 - CH_2 - CH = CH_2$  has hybridisation :

A.  $sp, sp, sp^2, sp^2$

B.  $sp^3, sp^3, sp^2, sp$




C.  $sp^3, sp^3, sp^2, sp^2$

D.  $sp^3, sp^2, sp^2, sp$

**Answer: C**

 [Watch Video Solution](#)

157. What is the state of hybridisation of Xe in cationic part of solid  $XeF_6$  ?






**Answer: B**



[Watch Video Solution](#)

158. During the complete combustion of methane  $CH_4$ , what change in hybridisation does the carbon atom undergo ?





D.  $sp^2$  to  $sp^3$

**Answer: A**

 [Watch Video Solution](#)

**159.** The hybridisation of central iodine atom in  $IF_5$ ,  $I_3^-$  and  $I_3^+$  are respectively :

A.  $sp^3d^2$ ,  $sp^3d$ ,  $sp^3$

B.  $sp^3d$ ,  $sp^3d$ ,  $sp^3$




C.  $sp^3d^2$ ,  $sp^3d^2$ ,  $sp^3$

D.  $sp^3d$ ,  $sp^3d^2$ ,  $sp^3$

**Answer: A**

 [Watch Video Solution](#)

160. In which of the following combination hybridisation of central atom ( \* ) does not change ?



D. none of these

Answer: C



Watch Video Solution

161. Which of the following species used both axial set of d-orbitals in hybridisation of central atom ?



D. none of these

**Answer: C**

 [Watch Video Solution](#)

**162.** Which bonds are formed by a carbon atom with  $sp^2$ -hybridisation ?

A.  $4\pi$ -bonds

B.  $2\pi$ -bonds and  $2\sigma$ -bonds


C.  $1\pi$ -bonds and  $3\sigma$ -bonds

D.  $4\sigma$ -bonds

**Answer: C**

 [Watch Video Solution](#)

163. What are the hybridisation of the carbon atoms labeled  $C_1$  and  $C_2$ , respectively in glycine ?



- A.  $C_1$   $C_2$   
 $sp^2$   $sp^2$
- B.  $C_1$   $C_2$   
 $sp^2$   $sp^3$
- C.  $C_1$   $C_2$   
 $sp^3$   $sp^2$
- D.  $C_1$   $C_2$   
 $sp^3$   $sp^3$

Answer: C



Watch Video Solution

164. The H-O-H bond angles in  $H_3O^+$  are approximately  $107^\circ$ . The orbitals used by oxygen in these bonds are best described as :

- A. p-orbitals
- B. sp-hybrid orbitals
- C.  $sp^2$ -hybrid orbital
- D.  $sp^3$ -hybrid orbital

**Answer: D**



[Watch Video Solution](#)

165. Which pair of elements can form multiple bond with itself and oxygen ?




- A. F,N
- B. N, Cl
- C. N,P

D. N,C

**Answer: D**

 [Watch Video Solution](#)


**166.** Which of the following is a covalent compound ?



**Answer: C**

 [Watch Video Solution](#)

**167.** Which of the following is an example of super octet molecule ?







D. All the three

**Answer: D**

 [Watch Video Solution](#)

**168.** Which of the following molecule is theoretically not possible ?



**Answer: C**

 [Watch Video Solution](#)

169. The phosphate of a metal has the formula  $MHPO_4$ . The formula of its chloride would be :

A.  $MCl$

B.  $MCl_2$

C.  $MCl_3$

D.  $M_2Cl_3$

**Answer: B**




[Watch Video Solution](#)

170. The compound that has the highest ionic character associated with the X-Cl bond is :

A.  $PCl_5$

B.  $BCl_3$









**Answer: D**

 [Watch Video Solution](#)


171. The bond having the highest bond energy is :



**Answer: C**

 [Watch Video Solution](#)

172. Which of the following species is neither hypervalent nor hypovalent ?



Answer: D



Watch Video Solution

173. In which of the following species central atom is NOT surrounded by exactly 8 valence electrons ?



D.  $SF_4$

**Answer: D**



**Watch Video Solution**

**174.** Which atom can have more than eight valence electrons when it is forming covalent bonds ?

A. H

B. N

C. F

D. Cl

**Answer: D**



**Watch Video Solution**

175. Which bond is expected to be the least polar ?

A. O-F

B. P-F

C. Si-N

D. B-F

Answer: A



Watch Video Solution

176. Which set contains only covalently bonded molecules ?

A.  $BCl_3$ ,  $SiCl_4$ ,  $PCl_3$

B.  $NH_4Br$ ,  $N_2H_4$ ,  $HBr$





C.  $I_2$ ,  $H_2S$ ,  $NaI$

D.  $Al$ ,  $O_3$ ,  $As_4$

**Answer: A**

 [Watch Video Solution](#)

**177.** Which molecule does not exist ?



**Answer: B**

 [Watch Video Solution](#)

**178.** Solid NaCl is a bad conductor of electricity because :

A. in solid NaCl there are no ions

B. solid NaCl is covalent

C. in solid NaCl there is no mobility of ions

D. in solid NaCl there are no electrons

**Answer: C**

 [Watch Video Solution](#)

**179.** An ionic compound  $A^+ B^-$  is most likely to be formed when :

A. the ionization energy of A high and electron affinity of B is low

B. the ionization energy of A is low and electron affinity of B is high




C. both, the ionization energy of A and electron affinity of B are high

D. both, the ionization energy of A and electron affinity of B are low

**Answer: B**

 [Watch Video Solution](#)

180. A compound contains three elements A, B and C, if the oxidation of  $A = +2$ ,  $B = +5$  and  $C = -2$ , the possible formula of the compound is :



**Answer: B**



[Watch Video Solution](#)

181. Which pair of atoms form strongest ionic bond ?

A. Al and As

B. Al and N




C. Al and Se

D. Al and O

Answer: D

 Watch Video Solution

182. The correct order of C-O bond length among  $CO$ ,  $CO_3^{2-}$ ,  $CO_2$  is





Answer: A

 Watch Video Solution

183. Resonance structures can be written for .







**Answer: A**

 [Watch Video Solution](#)

**184.** How many resonance structures can be drawn for the nitrate ion,  $NO_3^-$ ?

A. 1

B. 2





C. 3

D. 4

**Answer: C**

 [Watch Video Solution](#)

185. Among given species identify the isostructural pairs :



**Answer: B**



[Watch Video Solution](#)

186. 0.01 mole  $H_3PO_x$  is completely neutralised by 0.56 gram of KOH

hence :

A.  $x=3$  and given acid is dibasic

B.  $x=2$  and given acid is monobasic




C.  $x=3$  and given acid is monobasic

D.  $x=4$  and given acid forms three series of salt

Answer: B

 [Watch Video Solution](#)

187. Phosphorus pentachloride in the solid exists as :



Answer: C

 [Watch Video Solution](#)

188. The ratio of  $\sigma$  - bond and  $\pi$  - bond in tetracyano ethylene is :

A. 2:1

B. 1:1

C. 1:2

D. none of these

**Answer: B**

 [Watch Video Solution](#)

**189.** The bonds present in  $N_2O_5$  are .

A. only ionic

B. only covalent





C. covalent and co-ordinate

D. covalent and ionic

**Answer: C**

 [Watch Video Solution](#)

190. The pair of species with similar shape is



Answer: A



Watch Video Solution

191. Which of the following statements is correct in the context of the allene molecule,  $C_3H_4$  ?

A. The central carbon is  $sp$  hybridized

B. The terminal carbon atoms are  $sp^2$  hybridized

- C. The planes containing the  $CH_2$  groups are mutually perpendicular to permit the formations two separate  $\pi$ -bonds
- D. all are correct

**Answer: D**

 [Watch Video Solution](#)

**192.** Number of S-S bond is  $H_2S_nO_6$  :

- A. n
- B. (n-1)
- C. (n-2)
- D. (n+1)

**Answer: B**

 [Watch Video Solution](#)

193. How many S-S bonds, S-O-S bonds,  $\sigma$ -bonds,  $\pi$ -bonds are present in trimer of sulphur trioxide ?

A. 0,3,16,2

B. 0,3,12,6

C. 0,6,12,16

D. 0,4,12,6

**Answer: B**



[Watch Video Solution](#)

194. Number of identical Cr-O bonds in dichromate ion  $Cr_2O_7^{2-}$  is :

A. 4

B. 6

C. 7

D. 8

**Answer: B**

 [Watch Video Solution](#)

**195.** The nodal plane in the  $\pi$ -bond of ethene is located in:





- A. the molecular plane
- B. a plane parallel to the molecular plane
- C. a plane perpendicular to the molecular plane which bisects the carbon-carbon  $\sigma$  bond at right angle
- D. a plane perpendicular to the molecular plane which contains the carbon-carbon bond

**Answer: A**

 [Watch Video Solution](#)

**196.** Which of the following are isoelectronics and isostructural ?





**Answer: A**



**Watch Video Solution**

**197.** In the electronic structure of  $\text{H}_2\text{SO}_4$ , the total number of unshared electrons is

A. 20

B. 16


C. 12

D. 8

**Answer: B**

 [Watch Video Solution](#)

198. Which of the following xenon compound has the same number of lone pairs as in  $I_3^-$  ?



**Answer: C**

 [Watch Video Solution](#)

199. The shape of  $XeF_3^+$  is :

A. Trigonal planar

B. Pyramidal

C. Bent T-shpae

D. See-saw

**Answer: C**

 [Watch Video Solution](#)

**200.** Which of the following shape are not possible for possible value of  $n$  in  $XeF_n$  molecule ?

A. Linear

B. Square planar


C. Trigonal planar

D. Capped octahedral

**Answer: C**

 [Watch Video Solution](#)

201.  $BeCl_2$  is not isostructural with



Answer: D



Watch Video Solution

202. Which statement is true about the most stable Lewis structure for  $CS_2$  ?

A. There are no lone pairs in molecule

B. All bonds are double bonds

C. The central atom does not have an octet of electrons

D. A sulfur atom must be the central atom for the structure to be stable

**Answer: B**

 [Watch Video Solution](#)

203.  $SbF_5$  reacts with  $XeF_4$  and  $XeF_6$  to form ionic compounds

$[XeF_3^+][SbF_6^-]$  and  $[XeF_5^+][SbF_6^-]$  then the shape of


$[XeF_3^+]$  ion and  $[XeF_5^+]$  ion respectively :

- A. Square pyramidal, T-shaped
- B. Bent-T-shape, square pyramidal
- C. See-saw, square pyramidal
- D. Square pyramidal, see-saw

**Answer: B**

 [Watch Video Solution](#)

204. In which of the following species maximum atom can lie in same plane ?



Answer: D



Watch Video Solution

205. Correct statement regarding molecules  $SF_4$ ,  $CF_4$  and  $XeF_4$  are :

A. 2,0 and 1 lone pairs of central atom respectively

B. 1,0 and 1 lone pairs of central atom respectively

C. 0,0 and 2 lone pairs of central atom respectively

D. 1,0 and 2 lone pairs of central atom respectively

**Answer: D**

 [Watch Video Solution](#)

**206.** The geometrical arrangement of orbitals and shape of  $I_3^-$  are respectively :

- A. trigonal bipyramidal geometry, linear shape
- B. hexagonal geometry, T-shape
- C. triangular planar geometry, triangular shape
- D. tetrahedral geometry, pyramidal shape

**Answer: A**

 [Watch Video Solution](#)

207. Which of the following statements is incorrect for  $PCl_5$  ?

- A. Its three P-Cl bond lengths are equal
- B. It involves  $sp^3d$  hybridization
- C. It has an regular geometry
- D. Its shape is trigonal bipyramidal

Answer: C



Watch Video Solution

208. Molecular shapes of  $SF_4$ ,  $CF_4$ ,  $XeF_4$  are

- A. the same with 2,0 and 1 lone pair of electrons respectively
- B. the same with 1,1 and 1 lone pair of electrons respectively
- C. the same with 0,1 and 2 lone pair of electrons respectively
- D. the same with 1,0 and 2 lone pair of electrons respectively



**Answer: D**

 [Watch Video Solution](#)


**209.** The structure of the noble gas compound  $XeF_4$  is :

- A. square planar
- B. distorted tetrahedral
- C. tetrahedral
- D. octahedral

**Answer: A**

 [Watch Video Solution](#)





**210.** The molecule exhibiting maximum number of non-bonding electron pairs (l.p.) around the central atom is :



**Answer: C**

 [Watch Video Solution](#)

**211.** Which is the following pairs of species have identical shapes ?



**Answer: C**

 [Watch Video Solution](#)

212. The shapes of  $XeF_4$ ,  $XeF_5^-$  and  $SnCl_2$  are :

- A. octahedral, trigonal bipyramidal and bent
- B. square pyramidal, pentagonal planar and linear
- C. square planar, pentagonal planar and angular
- D. see-saw, T-shaped and linear

Answer: C



Watch Video Solution

213. Which is not correctly matched ?


- A.  $XeO_3$  -Trigonal bipyramidal
- B.  $ClF_3$  - bent T-shape
- C.  $XeOF_4^-$  - Square pyramidal

D.  $XeF_2$ - Linear shape

**Answer: A**

 [Watch Video Solution](#)

214. Amongst  $NO_3^-$ ,  $AsO_3^{3-}$ ,  $CO_3^{2-}$ ,  $ClO_3^-$ ,  $SO_3^{2-}$  and  $BO_3^{3-}$ , the non-planar species are :



**Answer: B**

 [Watch Video Solution](#)

**215.** The geometry of ammonia molecule can be best described as :

- A. Nitrogen at one vertex of a regular tetrahedron, the other three vertices being occupied by three hydrogens
- B. Nitrogen at the centre of the tetrahedron, three of the vertices being occupied by three hydrogens
- C. Nitrogen at the centre of an equilateral triangle, three corners being occupied by three hydrogens
- D. Nitrogen at the junction of a T, three open ends being occupied by three hydrogens

**Answer: B**



**Watch Video Solution**

**216.** Which molecular geometry is least likely to result from a trigonal bipyramidal electron geometry ?

A. Trigonal planar

B. See-saw

C. Linear

D. T-shaped

**Answer: A**



**Watch Video Solution**

**217.** Give the correct order of initials T or F for following statements. Use T if statement is true and F if it is false :

(I) The order of repulsion between different pair of electron is

$$I_p - I_p > I_p - b_p > b_p - b_p$$

(II) In general, as the number of lone pair of electron on central atom increases, value of bond angle from normal bond angle also increases

(III) The number of lone pair on O in  $H_2O$  is 2 while on N in  $NH_3$  is 1

(IV) The structures of xenon fluorides and xenon oxyfluorides could not be explained on the basis of VSEPR theory

A. T T T F

B. T F T F

C. T F T T

D. T F F F

**Answer: B**

 [Watch Video Solution](#)

**218.** Which species is planar ?



**Answer: A**

 [Watch Video Solution](#)

219. What is the geometry of the  $IBr_2^-$  ion ?

- A. Linear
- B. Bent shape with bond angle of about  $90^\circ$
- C. Bent shape with bond angle of about  $109^\circ$
- D. Bent shape with bond angle of about  $120^\circ$

**Answer: A**



[Watch Video Solution](#)

220. What is the shape of the  $ClF_3$  molecule?

- A. Trigonal planar
- B. Trigonal pyramidal
- C. T-shaped




D. Tetrahedral

Answer: C



Watch Video Solution

221. Which species below has the same general shape as  $NH_3$ ?



Answer: A



Watch Video Solution

222. According to VSEPR theory, in which species do all the atoms lie in the same plane ?



A. 1 only

B. 2 only

C. both 1 and 2


D. neither 1 nor 2


**Answer: A**



[Watch Video Solution](#)

223. Which of the following species / molecules does not have same number of bond pairs and lone pairs ?





**Answer: D**



**Watch Video Solution**

**224.** Least stable hydride is

A. stannane

B. silane

C. plumbane



D. germane

**Answer: C**



**Watch Video Solution**

225. The lowest O-O bond length in the following molecule is :



**Answer: B**



[Watch Video Solution](#)

226. The fluorine molecules is formed by :

A. p-p orbitals (sideways overlap)

B. p-p orbitals (end-to-end overlap)





C. sp-sp orbitals

D. s-s orbitals

Answer: B

 Watch Video Solution

227. Which of the following leads to bonding?



Answer: B

 Watch Video Solution

228. Which of the following overlaps is incorrect (assuming Z-axis is internuclear axis) ?

(A)  $2P_y + 2p_y \rightarrow \pi$ - Bond formation      (B)  $2p_x + 2p_x \rightarrow \sigma$ - Bond formation

(C)  $3d_{xy} + 3d_{xy} \rightarrow \pi$ - Bond formation      (D)  $2s + 2p_y \rightarrow \pi$ - Bond formation

(E)  $3d_{xy} + 3d_{xy} \rightarrow \delta$ - Bond formation

A. A,B,C

B. C,F

C. B,E

D. B,C,D

Answer: D



Watch Video Solution

229. Which of the following overlapping is not present in  $XeO_3$  molecule ?

A.  $sp^3 + p_x$

B.  $sp^3 + p_y$

C.  $d_{xz} + p_x$


D.  $sp^3 + s$

**Answer: D**



**Watch Video Solution**

**230.** How many sigma bonds are in a molecule of diethyl ether,



A. 14

B. 12





C. 8

D. 16

**Answer: A**

 [Watch Video Solution](#)

231. The lattice energies of KF, KCl, KBr and KI follow the order :




Answer: A

 [Watch Video Solution](#)

232. Which set of compounds in the following pair of ionic compounds has the higher lattice energy ?

(i)  $KCl$  or  $MgO$     (ii)  $LiF$  or  $LiBr$     (iii)  $Mg_3N_2$  or  $NaCl$





B.  $MgO$ ,  $LiBr$ ,  $Mg_3N_2$

C.  $MgO$ ,  $LiF$ ,  $NaCl$

D.  $MgO$ ,  $LiF$ ,  $Mg_3N_2$

**Answer: D**

 [Watch Video Solution](#)

**233.** The incorrect order of lattice energy is :

A.  $AlF_3 > MgF_2$

B.  $Li_3N > Li_2O$

C.  $NaCl > LiF$

D.  $TiC > ScN$

**Answer: C**

 [Watch Video Solution](#)

234. Which ionic compound has the largest amount of lattice energy ?

A. NaF

B.  $AlF_3$

C.  $AlN$

D.  $MgF_2$

Answer: C



Watch Video Solution

235. Which of the following compounds has the smallest bond angle?

A.  $OH_2$

B.  $SH_2$

C.  $NH_3$


D.  $SO_2$

**Answer: B**



[Watch Video Solution](#)

**236.** Maximum bond angle is present in case of




D. none of these




**Answer: D**



[Watch Video Solution](#)

**237.** The correct order of H-M-H bonds angle is :






**Answer: D**



**Watch Video Solution**

**238.** The correct increasing bond angle among  $BF_3$ ,  $PF_3$  and  $ClF_3$  follow the order



**Answer: C**



**Watch Video Solution**

239. Among the following species, the least angle around the central atom is in :



Answer: D

 [Watch Video Solution](#)

240. The bond angles of  $NH_3$ ,  $NH_4^+$  and  $NH_2^-$  are in the order





**Answer: B**

 [Watch Video Solution](#)

**241.** The H-C-H bond angle in  $CH_4$  is  $109.5^\circ$ , due to lone pair repulsion, the H-O-H angle in  $H_2O$  will :

A. remain the same

B. increase





C. decrease

D. become  $180^\circ$

**Answer: C**

 [Watch Video Solution](#)

242. The molecule having the largest bond angle is :



Answer: A



Watch Video Solution

243. The compound  $MX_4$  is tetrahedral. The number of  $\angle XMX$  angles formed in the compound is

A. three

B. four


C. five

D. six

**Answer: D**

 [Watch Video Solution](#)

**244.** The  $O - N - O$  bond angle in the nitrite ion,  $NO_2^-$ , is closest to :







**Answer: B**

 [Watch Video Solution](#)

**245.** Which of the following is the correct order for increasing bond angle

?






Answer: C

 [View Text Solution](#)

1.  $O - N - O$  bond angle is maximum in :







Answer: B

 [Watch Video Solution](#)

## Level 2

1. The incorrect order of boiling point is :



Answer: B,D

 [Watch Video Solution](#)

2. Iodine molecules are held in the crystals lattice by \_\_\_\_

- A. London forces
- B. dipole-dipole interactions
- C. covalent bonds
- D. coulombic force

**Answer: A**

 [Watch Video Solution](#)

3. At room temperature,  $CO_2$  is a gas while  $SiO_2$  is a solid because

- A.  $CO_2$  is a linear molecule, while  $SiO_2$  is angular
- B. van der Waals's forces are very strong in  $SiO_2$
- C.  $CO_2$  is covalent, while  $SiO_2$  is ionic
- D. Si cannot form stable bonds with O, hence Si has to form a 3D lattice

**Answer: D**



Watch Video Solution

4. Choose the correct code of characteristics for the given order of hybrid orbitals of same atom,

$$sp < sp^2 < sp^3$$

(i) Electronegativity (ii) Bond angle between same hybrid orbitals

(iii) Size (iv) Energy level

A. ii, iii and iv

B. iii, iv

C. ii and iv

D. I, ii, iii and iv

**Answer: B**



Watch Video Solution

5. Which is correct statement ?

As the s-character of a hybrid orbital decreases

(I) The bond angle decreases (II) The bond strength increases

(III) The bond length increases (IV) Size of orbitals increases

A. I, III and IV

B. II, III and IV

C. I and II

D. all are correct

**Answer: A**

 [Watch Video Solution](#)

6. Which of the following is incorrectly match ?

A.	Hybridisation $sp^3d$	Geometry Trigonal bipyramidal	Orbitals use $s + p_x + p_y + p_z + d_{s^2}$
----	--------------------------	----------------------------------	---

B.

Hybridisation	Geometry	Orbitals use
$sp^3d^3$	Pentagonal bipyramidal	$s + p_x + p_y + p_z + d_{x^2-y^2} + d_{xy} + d_{xz} + d_{yz}$

C.

Hybridisation	Geometry	Orbitals use
$sp^3d^2$	Capped octahedral	$s + p_x + p_y + p_z + d_{x^2-y^2} + d_{xy} + d_{xz} + d_{yz}$

D.

Hybridisation	Geometry	Orbitals use
$sp^3$	Tetrahedral	$s + p_x + p_y + p_z$

**Answer: C**

 [Watch Video Solution](#)

7. The ionic bond  $X^+Y^-$  are formed when :

(I) electron affinity of Y is high (II) ionization energy of X is low

(III) lattice energy of XY is high (IV) lattice energy of XY is low

Choose the correct code :

A. I and II

B. I and III

C. I, II and III

D. All

**Answer: C**

 [Watch Video Solution](#)

8. In the Born-Haber cycle for the formation of solid common salt (NaCl), the largest contribution comes from :

A. the low ionization potential of Na

B. the high electron affinity of Cl



C. the low  $\Delta H_{\text{vap}}$  of Na(s)

D. the lattice energy

**Answer: D**

 [Watch Video Solution](#)

9. Species having maximum 'Cl-O' bond order is :




**Answer: B**



[Watch Video Solution](#)

10. Which of the following species contains minimum number of atoms in XY plane ?





**Answer: B**

 [Watch Video Solution](#)


11. The molecule  $ML_x$  is planar with 7 pairs of electrons around M in the valence shell. The value of x is :

- A. 6
- B. 5
- C. 4
- D. 3

**Answer: B**

 [Watch Video Solution](#)

12. Choose the correct option for the following molecule in view of chemical bonding :



- A. non-planar
- B.  $\mu \neq 0$
- C. both a and b
- D.  $\mu = 0$

**Answer: D**



[Watch Video Solution](#)

13. Which of the following statement is correct about  $I_3^+$  and  $I_3^-$  molecular ions ?

- A. Number of lone pairs at central atoms are same in both molecular ions
- B. Hybridization of central atoms in both ions are same

C. Both are polar species

D. Both are planar species

**Answer: D**

 [Watch Video Solution](#)

14. In which of the following molecular shape  $d_{z^2}$  orbital must not be involved in bonding ?

A. Pentagonal planar

B. Trigonal planar

C. Linear

D. Square planar

**Answer: B**

 [Watch Video Solution](#)

15. The correct statement regarding  $SO_2$  molecule is :

- A. two  $p\pi - d\pi$  bonds
- B. molecule has 2 lone pair,  $2\sigma$  bonds and  $2\pi$  bonds
- C. two  $p\pi - p\pi$  bonds
- D. one  $p\pi - p\pi$  and one  $p\pi - d\pi$  bond

**Answer: D**



[Watch Video Solution](#)

16. A molecule  $XY_2$  contains two  $\sigma$  bonds two  $\pi$  bond and one lone pair of electrons in the valence shell of  $X$ . The arrangement of lone pair as well as bond pairs is

- A. square pyramidal
- B. linear
- C. Trigonal planar




D. unpredictable

Answer: C

 Watch Video Solution

17. In which of the following pairs, both the species have the same hybridisation ?

(I)



A. I,II

B. II, III

C. II, IV

D. I,II,III

Answer: C

 Watch Video Solution

18. Which of the following possess two lone pair of electrons on the central atom and square planar in shape ?

(I)  $SF_4$       (II)  $XeO_4$       (III)  $XeF_4$       (IV)  $ICl_4^-$

A. I,III

B. II,IV

C. III, IV

D. All

**Answer: C**





[Watch Video Solution](#)

19. Select pair of compounds in which both have different hybridization but have same molecular geometry :

A.  $BF_3$ ,  $BrF_3$

B.  $ICl_2^\ominus$ ,  $BeCl_2$



**Answer: B**



[Watch Video Solution](#)

20. The species having no  $p\pi - p\pi$  bond but its bond order equal to that of  $O_2^-$



**Answer: D**



[Watch Video Solution](#)

21. Which of the following fact is directly explained by the statement oxygen is a smaller atom than sulphur ?

- A.  $H_2O$  boils at a much higher temperature than  $H_2S$
- B.  $H_2O$  undergoes intermolecular hydrogen bonding
- C.  $H_2O$  is liquid and  $H_2S$  is gas at room temperature
- D. S-H bond is longer than O-H bond

Answer: D



Watch Video Solution

22. Which of the following compound has maximum "C-C" single bond length ?

- A.  $CH_2CHCCH$
- B.  $HC \ C \ C \ CH$
- C.  $CH_3CHCH_2$





Answer: C

 [Watch Video Solution](#)

23. If two different non-axial d-orbitals having 'xz' nodal plane form  $\pi$ -bond by overlapping each other, then internuclear axis will be :

A. x

B. y

C. z

D. They don't form  $\pi$ -bond

Answer: D

 [Watch Video Solution](#)

24. Assuming pure 2s and 2p orbitals of carbon are used in forming  $CH_4$  molecule, which of the following statement is false ?

- A. Three C-H bonds will be at right angle
- B. One C-H bond will be weaker than other three C-H bonds
- C. The shape of molecule will be tetrahedral
- D. The angle of C-H bond formed by s-s overlapping will be uncertain with respect to other three bonds.

Answer: C

 [Watch Video Solution](#)

25. The strength of bonds by 2s-2s, 2p-2p and 2p-2s overlap has the order

- A.  $s - s > p - p > p - s$
- B.  $s - s > p - s > p - p$
- C.  $p - p > p - s > s - s$

$$D. p - p > s - s > p - s$$

**Answer: C**

 [Watch Video Solution](#)

26. Which of the following statement is not correct for sigma and pi bonds formed between two carbon atoms ?

A. Sigma-bond is stronger than a  $\pi$ -bond

B. Bond energies of sigma and  $\pi$  -bonds are of the order of 264 kJ / mol and 347 kJ / mol

C. Free rotation of surrounding atoms about a sigma -bond is allowed but not in case of a  $\pi$ -bond

D. Sigma-bond determines the direction between carbon atoms but a  $\pi$ -bond has no primary effect in this regard

**Answer: B**

 [Watch Video Solution](#)

27. Assuming the bond direction to the z-axis, which of the overlapping of atomic orbitals of two atom (A) and (B) will result in bonding ?

- (I) s-orbital of A and  $p_x$  orbital of B (II) s-orbital of A and  $p_z$  orbital of B  
(III)  $p_y$ -orbital of A and  $p_z$  orbital of B (IV) s-orbital of both (A) and (B)

A. I and IV

B. I and II

C. III and IV

D. II and IV

**Answer: D**

 [Watch Video Solution](#)

28. Which of the following orbital (s) cannot form  $\delta$ -bond ?

A.  $d_{xy}$

B.  $d_{z^2}$

C.  $d_{x^2-y^2}$

D.  $d_{yz}$

**Answer: B**



[Watch Video Solution](#)

**29. Incorrect statement is :**

A.  $AlF_3 > MgO > MgF_2$  : Lattice energy

B.  $Li > Na > Al > Mg$  : Electron affinity

C.  $SF_6 > PF_5 > SiF_4$  : Lewis acidic character

D.  $SiCl_4 > SiBr_4 > SiI_4$  : Decreasing order of electronegativity of Si

**Answer: C**



[Watch Video Solution](#)

30. Which of the following set contains species having same angle around the central atom ?






Answer: D



Watch Video Solution

31. Which of the following compound has the smallest ( $X - A - X$ ) bond angle in each series repectively.



A.  $OSF_2$ ,  $SbCl_3$  and  $PI_3$

B.  $OSBr_2$ ,  $SbI_3$  and  $PI_3$

C.  $OSF_2$ ,  $SbI_3$  and  $PI_3$

D.  $OSF_2$ ,  $SbCl_3$  and  $SbI_3$

**Answer: D**

 [Watch Video Solution](#)

**32.** The incorrect order of boiling point is :

A.  $H_2O > CH_3OH$

B.  $N(CH_3)_3 > NH(CH_3)_2$

C.  $H_3PO_4 > Mg_3PO_4$

D.  $CH_3N_3 > HN_3$

**Answer: D**

 [Watch Video Solution](#)

33. Iodine molecules are held in the solid lattice by \_\_\_\_\_

- A. London forces
- B. dipole-dipole interactions
- C. covalent bonds
- D. coulombic force

Answer: A



Watch Video Solution

34. Carbon dioxide is a gas but silica is a solid because:

- A.  $CO_2$  is a linear molecule, while  $SiO_2$  is angular
- B. van der Waals's forces are very strong in  $SiO_2$
- C.  $CO_2$  is covalent, while  $SiO_2$  is ionic



D. Si cannot form stable bonds with O, hence Si has to form a 3D lattice

**Answer: D**

 [Watch Video Solution](#)

**35.** Choose the correct code of characteristics for the given order of hybrid orbitals of same atom,

$$sp < sp^2 < sp^3$$

(i) Electronegativity (ii) Bond angle between same hybrid orbitals  
(iii) Size (iv) Energy level

A. ii, iii and iv

B. iii, iv

C. ii and iv

D. I, ii, iii and iv

**Answer: B**



[Watch Video Solution](#)

**36.** Which is correct statement ?

As the s-character of a hybrid orbital decreases

(I) The bond angle decreases (II) The bond strength increases

(III) The bond length increases (IV) Size of orbitals increases

A. I, III and IV

B. II, III and IV

C. I and II

D. all are correct

**Answer: A**



[Watch Video Solution](#)

**37.** Which of the following is incorrectly matched ?

A. Hybridisation Geometry Orbitals use  
 $sp^3d$  Trigonal bipyramidal  $s + p_x + p_y + p_z + d_{s^2}$

B.

Hybridisation Geometry Orbitals use  
 $sp^3d^3$  Pentagonal bipyramidal  $s + p_x + p_y + p_z + d_{x^2-y^2} + d_{z^2}$

C.

Hybridisation Geometry Orbitals use  
 $sp^3d^2$  Capped octahedral  $s + p_x + p_y + p_z + d_{x^2-y^2} + d_{z^2}$

D. Hybridisation Geometry Orbitals use  
 $sp^3$  Tetrahedral  $s + p_x + p_y + p_z$

**Answer: C**



[Watch Video Solution](#)

38. The ionic bond  $X^+Y^-$  are formed when :

(I) electron affinity of Y is high (II) ionization energy of X is low

(III) lattice energy of XY is high (IV) lattice energy of XY is low

Choose the correct code :

A. I and II

B. I and III

C. I, II and III

D. All

**Answer: C**

 [Watch Video Solution](#)

**39.** In the Born-Haber cycle for the formation of solid common salt (NaCl), the largest contribution comes from :

A. the low ionization potential of Na

B. the high electron affinity of Cl



C. the low  $\Delta H_{\text{vap}}$  of Na(s)

D. the lattice energy

**Answer: D**

 [Watch Video Solution](#)

40. Species having maximum 'Cl-O' bond order is :





Answer: B



Watch Video Solution

41. Which of the following species contains minimum number of atoms in XY plane ?



D. All

**Answer: B**

 [Watch Video Solution](#)

42. The molecule  $ML_x$  is planar with 7 pairs of electrons around M in the valence shell. The value of x is :

A. 6

B. 5


C. 4

D. 3

**Answer: B**

 [Watch Video Solution](#)

43. Choose the correct option for the following molecule in view of chemical bonding :



- A. non-planar
- B.  $\mu \neq 0$
- C. both a and b
- D.  $\mu = 0$

**Answer: D**

 [Watch Video Solution](#)

44. Which of the following statement is correct about  $I_3^+$  and  $I_3^-$  molecular ions ?

- A. Number of lone pairs at central atoms are same in both molecular ions
- B. Hybridization of central atoms in both ions are same

C. Both are polar species

D. Both are planar species

**Answer: D**

 [Watch Video Solution](#)

45. In which of the following molecular shape  $d_{z^2}$  orbital must not be involved in bonding ?

A. Pentagonal planar

B. Trigonal planar

C. Linear

D. Square planar

**Answer: B**

 [Watch Video Solution](#)



46. The correct statement regarding  $SO_2$  molecule is :

A. two  $p\pi - d\pi$  bonds

B. molecule has 2 lone pair,  $2\sigma$  bonds and  $2\pi$  bonds

C. two  $p\pi - p\pi$  bonds

D. one  $p\pi - p\pi$  and one  $p\pi - d\pi$  bond

Answer: D



Watch Video Solution

47. A molecule  $XY_2$  contains two  $\sigma$  bonds two  $\pi$  bond and one lone pair of electrons in the valence shell of  $X$ . The arrangement of lone pair as well as bond pairs is

A. square pyramidal

B. linear

C. Trigonal planar




D. unpredictable

Answer: C

 Watch Video Solution

48. In which of the following pairs, both the species have the same hybridisation ?

(I)



A. I,II

B. II, III

C. II, IV

D. I,II,III

Answer: C

 Watch Video Solution

49. Which of the following possess two lone pair of electrons on the central atom and square planar in shape ?

(I)  $SF_4$       (II)  $XeO_4$       (III)  $XeF_4$       (IV)  $ICl_4^-$

A. I,III

B. II,IV

C. III, IV

D. All

Answer: C





Watch Video Solution

50. Select pair of compounds in which both have different hybridization but have same molecular geometry :

A.  $BF_3$ ,  $BrF_3$


B.  $ICl_2^\ominus$ ,  $BeCl_2$



**Answer: B**

 [Watch Video Solution](#)

51. The species having no  $p\pi - p\pi$  bond but its bond order equal to that of  $O_2^-$



**Answer: D**

 [Watch Video Solution](#)

52. Which of the following fact is directly explained by the statement oxygen is a smaller atom than sulphur ?

- A.  $H_2O$  boils at a much higher temperature than  $H_2S$
- B.  $H_2O$  undergoes intermolecular hydrogen bonding
- C.  $H_2O$  is liquid and  $H_2S$  is gas at room temperature
- D. S-H bond is longer than O-H bond

Answer: D



Watch Video Solution

53. Which of the following compound has maximum "C-C" single bond length ?

- A.  $CH_2CHCCH$
- B. HC C C CH
- C.  $CH_3CHCH_2$



Answer: C

 [Watch Video Solution](#)

54. If two different non-axial d-orbitals having 'xz' nodal plane form  $\pi$ -bond by overlapping each other, then internuclear axis will be :

A. x

B. y

C. z

D. They don't form  $\pi$ -bond

Answer: D

 [Watch Video Solution](#)

55. Assuming pure 2s and 2p orbitals of carbon are used in forming  $CH_4$  molecule, which of the following statement is false ?

- A. Three C-H bonds will be at right angle
- B. One C-H bond will be weaker than other three C-H bonds
- C. The shape of molecule will be tetrahedral
- D. The angle of C-H bond formed by s-s overlapping will be uncertain with respect to other three bonds.

**Answer: C**



[Watch Video Solution](#)

56. Which of the following is correct order of  $\sigma$  - bond strength ?

- I. 2s-2s
- II. 2s-2p
- III. 2p-2p
- IV. 3s-3s

A.  $s - s > p - p > p - s$

B.  $s - s > p - s > p - p$

C.  $p - p > p - s > s - s$

D.  $p - p > s - s > p - s$

**Answer: C**

 [Watch Video Solution](#)

57. Which of the following statements is incorrect for sigma and  $\pi$  -bonds formed between two carbon atoms ?

A. Sigma-bond is stronger than a  $\pi$ -bond

B. Bond energies of sigma and  $\pi$  -bonds are of the order of 264 kJ / mol and 347 kJ / mol

C. Free rotation of surrounding atoms about a sigma -bond is allowed but not in case of a  $\pi$ -bond



D. Sigma-bond determines the direction between carbon atoms but a

$\pi$ -bond has no primary effect in this regard

**Answer: B**

 [Watch Video Solution](#)

58. Assuming the bond direction to the z-axis, which of the overlapping of atomic orbitals of two atom (A) and (B) will result in bonding ?

- (I) s-orbital of A and  $p_x$  orbital of B (II) s-orbital of A and  $p_z$  orbital of B  
(III)  $p_y$ -orbital of A and  $p_z$  orbital of B (IV) s-orbital of both (A) and (B)

A. I and IV

B. I and II


C. III and IV

D. II and IV

**Answer: D**

 [Watch Video Solution](#)

59. Which of the following orbital can not form  $\pi$  as well as  $\delta$ -Bond ?





Answer: B



Watch Video Solution

60. Incorrect statement is :



D.  $SiCl_4 > SiBr_4 > SiI_4$  : Decreasing order of electronegativity of Si

**Answer: C**

 [Watch Video Solution](#)

61. Which of the following set contains species having same angle around the central atom ?

A.  $SF_4, CH_4, NH_3$

B.  $NF_3, BCl_3, NH_3$



C.  $BF_3, NF_3, AlCl_3$

D.  $BF_3, BCl_3, BBr_3$

**Answer: D**

 [Watch Video Solution](#)

62. Which of the following compound has the smallest bond angle (X-A-X) in each series respectively ?



**Answer: D**



**Watch Video Solution**

Level 3 (Passive 1)

1. According to VSEPR model, molecules adopt geometries in which their valence electron pairs position themselves as far from each other as possible. The VSEPR model considers double and triple bonds to have slightly greater repulsive effects than single bonds because of the repulsive effect of  $\pi$  -electrons. However the lone pair creates the maximum repulsive effect.

Which of the following statement is false ?

A.  $SbF_4^-$  and  $SF_4$  are isostructural

B. In  $IOF_5$  the hybridization of central atom is  $sp^3d^2$

C. Double bond(s) in  $SOF_4$  and  $XeO_3F_2$ , is / are occupying equatorial position(s) of their respective geometry

D. none of these




**Answer: D**



**Watch Video Solution**

2. According to VSEPR model, molecules adopt geometries in which their valence electron pairs position themselves as far from each other as possible. The VSEPR model considers double and triple bonds to have slightly greater repulsive effects than single bonds because of the repulsive effect of  $\pi$ -electrons. However the lone pair creates the maximum repulsive effect.

Which of the following does not represent the isostructural pair ?



D. None

**Answer: D**



**Watch Video Solution**

3. According to VSEPR model, molecules adopt geometries in which their valence electron pairs position themselves as far from each other as possible. The VSEPR model considers double and triple bonds to have slightly greater repulsive effects than single bonds because of the repulsive effect of  $\pi$ -electrons. However the lone pair creates the maximum repulsive effect.

Select the incorrect statement with respect to  $SO_2Cl_2$  molecule :

- A. It gives  $H_2SO_4$  and HCl on hydrolysis at room temperature
- B. It has two  $d\pi - p\pi$  bonds between S and O bonded atoms
- C. It is a polar molecule
- D. None

**Answer: D**



[Watch Video Solution](#)

4. According to VSEPR model, molecules adopt geometries in which their valence electron pairs position themselves as far from each other as possible. The VSEPR model considers double and triple bonds to have slightly greater repulsive effects than single bonds because of the repulsive effect of  $\pi$  -electrons. However the lone pair creates the maximum repulsive effect.

Which of the following statement is false ?

A.  $SbF_4^-$  and  $SF_4$  are isostructural

B. In  $IOF_5$  the hybridization of central atom is  $sp^3d^2$

C. Double bond(s) in  $SOF_4$  and  $XeO_3F_2$ , is / are occupying equatorial position(s) of their respective geometry

D. none of these

**Answer: D**






**Watch Video Solution**



5. According to VSEPR model, molecules adopt geometries in which their valence electron pairs position themselves as far from each other as possible. The VSEPR model considers double and triple bonds to have slightly greater repulsive effects than single bonds because of the repulsive effect of  $\pi$ -electrons. However the lone pair creates the maximum repulsive effect.

Which of the following does not represent the isostructural pair ?



D. None

**Answer: D**



**Watch Video Solution**

6. According to VSEPR model, molecules adopt geometries in which their valence electron pairs position themselves as far from each other as possible. The VSEPR model considers double and triple bonds to have slightly greater repulsive effects than single bonds because of the repulsive effect of  $\pi$ -electrons. However the lone pair creates the maximum repulsive effect.

Select the incorrect statement with respect to  $SO_2Cl_2$  molecule :

- A. It gives  $H_2SO_4$  and HCl on hydrolysis at room temperature
- B. It has two  $d\pi - p\pi$  bonds between S and O bonded atoms
- C. It is a polar molecule
- D. None

**Answer: D**



[Watch Video Solution](#)

1. According to VBT any covalent bond will be formed by overlapping of atomic orbitals of bonded atoms provided atomic orbitals must be half-filled and electrons be in opposite spin. According to type of overlapping covalent bonds can be classified as (a)  $\sigma$ -bond (b)  $\pi$ -bond (c)  $\delta$ -bond :

Which of the following set of orbitals does not produce nodal plane in xz-plane ?

A.  $d_{yz} + d_{yz}$

B.  $d_{xy} + d_{xy}$

C.  $p_y + d_{xy}$

D. none of these

**Answer: D**

 [View Text Solution](#)

2. According to VBT any covalent bond will be formed by overlapping of atomic orbitals of bonded atoms provided atomic orbitals must be half-

filled and electrons be in opposite spin. According to type of overlapping covalent bonds can be classified as (a) $\sigma$ -bond (b)  $\pi$ -bond (c )  $\delta$ -bond :

The combination of orbital that can not produce non-bonding molecular orbital is (internuclear axis is z-axis ) :

A.  $p_y + d_{x^2 - y^2}$

B.  $p_z + d_{yz}$

C.  $s + d_{xz}$

D.  $d_{xy} + d_{xy}$

**Answer: D**



[View Text Solution](#)

3. According to VBT any covalent bond will be formed by overlapping of atomic orbitals of bonded atoms provided atomic orbitals must be half-filled and electrons be in opposite spin. According to type of overlapping covalent bonds can be classified as (a) $\sigma$ -bond (b)  $\pi$ -bond (c )  $\delta$ -bond :

If  $F_2C_1 = C_2$  part of  $F_2C_1 = C_2 = C_3 = C_4F_2$  lies in yz-plane, then incorrect statement is :

- A. Nodal plane of  $\pi$ -bond between  $C_1$  and  $C_2$  lies in yz-plane, formed by sideways overlapping of  $p_x$ -orbitals
- B. Nodal plane of  $\pi$ -bond between  $C_2$  and  $C_3$  lies in xz-plane, formed by sideways overlapping of  $p_y$ -orbitals
- C. Nodal plane of  $\pi$ -bond between  $C_3$  and  $C_4$  lies in yz-plane, formed by sideways overlapping of  $p_y$ -orbitals
- D. Nodal plane of  $\pi$ -bond between  $C_2$  and  $C_3$  lies in xy-plane, formed by sideways overlapping of  $p_x$ -orbitals

**Answer: C**



[View Text Solution](#)

4. According to VBT any covalent bond will be formed by overlapping of atomic orbitals of bonded atoms provided atomic orbitals must be half-

filled and electrons be in opposite spin. According to type of overlapping covalent bonds can be classified as (a)  $\sigma$ -bond (b)  $\pi$ -bond (c)  $\delta$ -bond :

Which of the following set of orbitals does not produce nodal plane in xz-plane ?

A.  $d_{yz} + d_{yz}$

B.  $d_{xy} + d_{xy}$

C.  $p_y + d_{xy}$

D. none of these

**Answer: D**



[View Text Solution](#)

5. According to VBT any covalent bond will be formed by overlapping of atomic orbitals of bonded atoms provided atomic orbitals must be half-filled and electrons be in opposite spin. According to type of overlapping covalent bonds can be classified as (a)  $\sigma$ -bond (b)  $\pi$ -bond (c)  $\delta$ -bond :

The combination of orbital that can not produce non-bonding molecular orbital is (internuclear axis is z-axis) :

A.  $p_y + d_{x^2 - y^2}$

B.  $p_z + d_{yz}$

C.  $s + d_{xz}$

D.  $d_{xy} + d_{xy}$

**Answer: D**



[View Text Solution](#)

6. According to VBT any covalent bond will be formed by overlapping of atomic orbitals of bonded atoms provided atomic orbitals must be half-filled and electrons be in opposite spin. According to type of overlapping covalent bonds can be classified as (a)  $\sigma$ -bond (b)  $\pi$ -bond (c)  $\delta$ -bond :

If  $F_2C_1 = C_2$  part of  $F_2C_1 = C_2 = C_3 = C_4F_2$  lies in yz-plane, then incorrect statement is :

- A. Nodal plane of  $\pi$ -bond between  $C_1$  and  $C_2$  lies in yz-plane, formed by sideways overlapping of  $p_x$ -orbitals
- B. Nodal plane of  $\pi$ -bond between  $C_2$  and  $C_3$  lies in xz-plane, formed by sideways overlapping of  $p_y$ -orbitals
- C. Nodal plane of  $\pi$ - bond between  $C_3$  and  $C_4$  lies in yz-plane, formed by sideways overlapping of  $p_y$  -orbitals
- D. Nodal plane of  $\pi$ -bond between  $C_2$  and  $C_3$  lies in xy-plane, formed by sideways overlapping of  $p_x$  - orbitals

**Answer: C**



[View Text Solution](#)

### Level 3 (Passive 3)

1. If the central atom is of third row or below this in the periodic table, then lone pair will occupy a stereochemically inactive s-orbital and



bonding will be through almost pure p-orbitals and bond angles are nearly  $90^\circ$ , if the substituent's electronegativity value is  $\leq 2.5$ .

In which of the following option, covalent bond is having maximum s% character ?

- A. S-H bond in  $H_2S$
- B. P-H bond in  $PH_3$
- C. N-H bond in  $NH_3$
- D. All have equal s% character

**Answer: C**

 [Watch Video Solution](#)

2. If the central atom is of third row or below this in the periodic table, then lone pair will occupy a stereochemically inactive s-orbital and bonding will be through almost pure p-orbitals and bond angles are nearly  $90^\circ$ , if the substituent's electronegativity value is  $\leq 2.5$ .

Select incorrect statement regarding  $P_4$  molecule.

- A. Each P atom is joined with three P-atoms
- B.  $P_4$  molecule contains total 12 bond angles
- C. Lone pair of each P atom is present in almost pure s-orbital
- D. Lone pair of each P atom present in hybrid orbital

**Answer: D**

 [Watch Video Solution](#)

3. If the central atom is of third row or below this in the periodic table, then lone pair will occupy a stereochemically inactive s-orbital and bonding will be through almost pure p-orbitals and bond angles are nearly  $90^\circ$ , if the substituent's electronegativity value is  $\leq 2.5$ .

The hybridisation of atomic orbitals of central atom "Xe" in  $XeO_4$ ,  $XeO_2F_2$  and  $XeOF_4$  respectively.

A.  $sp^3$ ,  $sp^3d^2$ ,  $sp^3d^2$

B.  $sp^3d$ ,  $sp^3d$ ,  $sp^3d^2$

C.  $sp^3$ ,  $sp^3d^2$ ,  $sp^3d$

D.  $sp^3$ ,  $sp^3d$ ,  $sp^3d^2$

**Answer: D**



**Watch Video Solution**

4. If the central atom is of third row or below this in the periodic table, then lone pair will occupy a stereochemically inactive s-orbital and bonding will be through almost pure p-orbitals and bond angles are nearly  $90^\circ$ , if the substituent's electronegativity value is  $\leq 2.5$ .

In which of the following option, covalent bond is having maximum s% character ?

A. S-H bond in  $H_2S$

B. P-H bond in  $PH_3$

C. N-H bond in  $NH_3$

D. All have equal s% character

**Answer: C**

 [Watch Video Solution](#)

5. If the central atom is of third row or below this in the periodic table, then lone pair will occupy a stereochemically inactive s-orbital and bonding will be through almost pure p-orbitals and bond angles are nearly  $90^\circ$ , if the substituent's electronegativity value is  $\leq 2.5$ .

Select incorrect statement regarding  $P_4$  molecule.





- A. Each P atom is joined with three P-atoms
- B.  $P_4$  molecule contains total 12 bond angles
- C. Lone pair of each P atom is present in almost pure s-orbital
- D. Lone pair of each P atom present in hybrid orbital

**Answer: D**

 [Watch Video Solution](#)

6. If the central atom is of third row or below this in the periodic table, then lone pair will occupy a stereochemically inactive s-orbital and bonding will be through almost pure p-orbitals and bond angles are nearly  $90^\circ$ , if the substituent's electronegativity value is  $\leq 2.5$ .

The hybridisation of atomic orbitals of central atom "Xe" in  $XeO_4$ ,  $XeO_2F_2$  and  $XeOF_4$  respectively.



**Answer: D**



[Watch Video Solution](#)

Level 3 (Passive 4)

1. According to V.B.T., atoms of element form bond only to pair up their unpaired electrons present in ground state or excited state. This pairing of unpaired electron will take place by overlapping of orbitals each one having one unpaired electron with opposite spin.

Which of the following orbital combination does not form  $\pi$ -bond ?

- A.  $p_x + p_x$  sideways overlapping
- B.  $d_{x^2-y^2} + p_y$  sideways overlapping
- C.  $d_{xy} + d_{xy}$  sideways overlapping
- D.  $d_{yz} + p_y$  sideways overlapping

**Answer: B**



[Watch Video Solution](#)

2. According to V.B.T., atoms of element form bond only to pair up their unpaired electrons present in ground state or excited state. This pairing of unpaired electron will take place by overlapping of orbitals each one

having one unpaired electron with opposite spin.

Which of the following orbital cannot form  $\delta$ -bond ?

A.  $d_{x^2-y^2}$  orbital

B.  $d_{xy}$  orbital

C.  $d_{z^2}$  orbital

D.  $d_{zx}$  orbital

**Answer: C**



[Watch Video Solution](#)

3. According to V.B.T., atoms of element form bond only to pair up their unpaired electrons present in ground state or excited state. This pairing of unpaired electron will take place by overlapping of orbitals each one having one unpaired electron with opposite spin.

Which of the following combination of orbitals does not form any type of covalent bond (if z-axis is molecular axis)?

A.  $p_s + p_z$

B.  $p_y + p_y$

C.  $s + p_y$

D.  $s + s$

**Answer: C**

 [Watch Video Solution](#)

4. According to V.B.T., atoms of element form bond only to pair up their unpaired electrons present in ground state or excited state. This pairing of unpaired electron will take place by overlapping of orbitals each one having one unpaired electron with opposite spin.

Which of the following orbital cannot form  $\delta$ -bond ?

A.  $d_{x^2-y^2}$  orbital

B.  $d_{xy}$  orbital

C.  $d_{x^2}$  orbital



D.  $d_{zx}$  orbital

**Answer: C**

 [Watch Video Solution](#)

5. According to V.B.T., atoms of element form bond only to pair up their unpaired electrons present in ground state or excited state. This pairing of unpaired electron will take place by overlapping of orbitals each one having one unpaired electron with opposite spin.

Which of the following combination of orbitals does not form any type of covalent bond (if z-axis is molecular axis)?

A.  $p_s + p_z$

B.  $p_y + p_y$

C.  $s + p_y$

D.  $s + s$

**Answer: C**



Watch Video Solution

### Level 3 (Passive 5)

1. The space model which is obtained by joining the points representing various bonded atoms gives the shape of the molecule. The geometry of the molecule is definite relative arrangement of the bonded atoms in a molecule. The shape and geometry of a molecule is explained by valence shell electron pair repulsion theory given by Gillespie and Nyholm.

Select the correct code for the following repulsion orders, according to VSEPR theory :

(I) lone pair -lone pair > lone pair-bond pair

(II) lone pair-bond pair > bond pair -bond pair

(III) lone pair -lone pair > bond pair-bond pair

(IV) lone pair - bond pair > lone pair-lone pair

A. I,II and III

B. II and IV

C. I,II and IV

D. All

**Answer: A**

 [Watch Video Solution](#)

2. The space model which is obtained by joining the points representing various bonded atoms gives the shape of the molecule. The geometry of the molecule is definite relative arrangement of the bonded atoms in a molecule. The shape and geometry of a molecule is explained by valence shell electron pair repulsion theory given by Gillespie and Nyholm.

Which molecule has both shape and geometry identical ?

(I)  $SnCl_2$     (II)  $NH_3$     (III)  $PCl_5$     (IV)  $SF_6$

A. I, III and IV

B. II,III and IV

C. III and IV

D. All

**Answer: C**



**Watch Video Solution**

3. The space model which is obtained by joining the points representing various bonded atoms gives the shape of the molecule. The geometry of the molecule is definite relative arrangement of the bonded atoms in a molecule. The shape and geometry of a molecule is explained by valence shell electron pair repulsion theory given by Gillespie and Nyholm.

Which is not the electron geometry of covalent molecules ?

A. Pentagonal bipyramidal

B. Octahedral

C. Hexagonal

D. Tetrahedral

**Answer: C**



4. The space model which is obtained by joining the points representing various bonded atoms gives the shape of the molecule. The geometry of the molecule is definite relative arrangement of the bonded atoms in a molecule. The shape and geometry of a molecule is explained by valence shell electron pair repulsion theory given by Gillespie and Nyholm.

Select the correct code for the following repulsion orders, according to VSEPR theory :

- (I) lone pair -lone pair > lone pair-bond pair
- (II) lone pair-bond pair > bond pair -bond pair
- (III) lone pair -lone pair > bond pair-bond pair
- (IV) lone pair - bond pair > lone pair-lone pair

A. I,II and III

B. II and IV

C. I,II and IV

D. All

**Answer: A**



**Watch Video Solution**

5. The space model which is obtained by joining the points representing various bonded atoms gives the shape of the molecule. The geometry of the molecule is definite relative arrangement of the bonded atoms in a molecule. The shape and geometry of a molecule is explained by valence shell electron pair repulsion theory given by Gillespie and Nyholm.

Which molecule has both shape and geometry identical ?

(I)  $SnCl_2$     (II)  $NH_3$     (III)  $PCl_5$     (IV)  $SF_6$

A. I, III and IV

B. II, III and IV

C. III and IV

D. All

**Answer: C**





Watch Video Solution

6. The space model which is obtained by joining the points representing various bonded atoms gives the shape of the molecule. The geometry of the molecule is definite relative arrangement of the bonded atoms in a molecule. The shape and geometry of a molecule is explained by valence shell electron pair repulsion theory given by Gillespie and Nyholm.

Which is not the electron geometry of covalent molecules ?

- A. Pentagonal bipyramidal
- B. Octahedral
- C. Hexagonal
- D. Tetrahedral

**Answer: C**



Watch Video Solution

1. When hybridisation involving d-orbitals are considered then all the five d-orbitals are not degenerate, rather  $d_{x^2-y^2}$ ,  $d_{z^2}$  and  $d_{xy}$ ,  $d_{zx}$  form two different sets of orbitals and orbitals of appropriate set is involved in the hybridisation.

In  $sp^3d^2$  hybridisation, which sets of d-orbitals is involved ?

A.  $d_{x^2-y^2}$ ,  $d_{z^2}$

B.  $d_{z^2}$ ,  $d_{xy}$

C.  $d_{xy}$ ,  $d_{yz}$

D.  $d_{x^2-y^2}$ ,  $d_{xy}$

**Answer: A**



[Watch Video Solution](#)

2. The d orbitals involved in  $sp^3d^3$  hybridization are ?

A.  $d_{x^2-y^2}$ ,  $d_{z^2}$ ,  $d_{xy}$



B.  $d_{xy}$ ,  $d_{yz}$ ,  $d_{zx}$

C.  $d_{x^2-y^2}$ ,  $d_{xy}$ ,  $d_{xz}$

D.  $d_{x^2}$ ,  $d_{yz}$ ,  $d_{zx}$

**Answer: A**



**Watch Video Solution**

3. When hybridisation involving d-orbitals are considered then all the five d-orbitals are not degenerate, rather  $d_{x^2-y^2}$ ,  $d_{s^2}$  and  $d_{xy}$ ,  $d_{zx}$  form two different sets of orbitals and orbitals of appropriate set is involved in the hybridisation.

Molecule having trigonal bipyramidal geometry and  $sp^3d$  hybridisation, d-orbitals involved is :

A.  $d_{xy}$

B.  $d_{yz}$

C.  $d_{x^2-y^2}$

D.  $d_{z^2}$

**Answer: D**



**Watch Video Solution**

4. When hybridisation involving d-orbitals are considered then all the five d-orbitals are not degenerate, rather  $d_{x^2-y^2}$ ,  $d_{s^2}$  and  $d_{xy}$ ,  $d_{zx}$  form two different sets of orbitals and orbitals of appropriate set is involved in the hybridisation.

Which of the following orbitals can not undergo hybridisation amongst themselves.

(I)  $3d, 4s$       (II)  $3d, 4d$

(III)  $3d, 4s \& 4p$       (IV)  $3s, 3p \& 4s$

A. only II

B. II and III

C. I, II and IV





D. II and IV

**Answer: D**

 [Watch Video Solution](#)

5. When hybridisation involving d-orbitals are considered then all the five d-orbitals are not degenerate, rather  $d_{x^2-y^2}$ ,  $d_{s^2}$  and  $d_{xy}$ ,  $d_{zx}$  form two different sets of orbitals and orbitals of appropriate set is involved in the hybridisation.

In  $sp^3d^2$  hybridisation, which sets of d-orbitals is involved ?



**Answer: A**

 [Watch Video Solution](#)

6. When hybridisation involving d-orbitals are considered then all the five d-orbitals are not degenerate, rather  $d_{x^2-y^2}$ ,  $d_{s^2}$  and  $d_{xy}$ ,  $d_{zx}$  form two different sets of orbitals and orbitals of appropriate set is involved in the hybridisation.

In  $sp^3d^3$  hybridisation, which orbitals are involved ?

A.  $d_{x^2-y^2}$ ,  $d_{z^2}$ ,  $d_{xy}$

B.  $d_{xy}$ ,  $d_{yz}$ ,  $d_{zx}$

C.  $d_{x^2-y^2}$ ,  $d_{xy}$ ,  $d_{xz}$

D.  $d_{x^2}$ ,  $d_{yz}$ ,  $d_{zx}$

**Answer: A**



[Watch Video Solution](#)

7. When hybridisation involving d-orbitals are considered then all the five d-orbitals are not degenerate, rather  $d_{x^2-y^2}$ ,  $d_{s^2}$  and  $d_{xy}$ ,  $d_{zx}$  form two different sets of orbitals and orbitals of appropriate set is involved in the

hybridisation.

Molecule having trigonal bipyramidal geometry and  $sp^3d$  hybridisation, d-orbitals involved is :

A.  $d_{xy}$

B.  $d_{yz}$

C.  $d_{x^2-y^2}$

D.  $d_{z^2}$

**Answer: D**



[Watch Video Solution](#)

8. When hybridisation involving d-orbitals are considered then all the five d-orbitals are not degenerate, rather  $d_{x^2-y^2}$ ,  $d_{z^2}$  and  $d_{xy}$ ,  $d_{zx}$  form two different sets of orbitals and orbitals of appropriate set is involved in the hybridisation.

Which of the following orbitals can not undergo hybridisation amongst themselves.

(I)  $3d, 4s$       (II)  $3d, 4d$

(III)  $3d, 4s \& 4p$       (IV)  $3s, 3p \& 4s$

A. only II

B. II and III

C. I, II and IV

D. II and IV

**Answer: D**



[Watch Video Solution](#)

### Level 3 (Passive 7)

1. Ionic bond is defined as the electrostatic force of attraction holding the oppositely charged ions. Ionic compounds are mostly crystalline solid having high melting and boiling points, electrical conductivity in molten state, solubility in water etc. Covalent bond is defined as the force which binds atoms of same or different elements by mutual sharing of electrons

in a covalent bond. Covalent compounds are solids, liquids or gases. They have low melting and boiling points compounds. They are more soluble in non-polar solvents.

The valence electrons are involved in formation of covalent bonds is /are called :

A. non-bonding electrons

B. lone pairs

C. unshared pairs

D. none of these

**Answer: D**



[Watch Video Solution](#)

2. Ionic bond is defined as the electrostatic force of attraction holding the oppositely charged ions. Ionic compounds are mostly crystalline solid having high melting and boiling points, electrical conductivity in molten state, solubility in water etc. Covalent bond is defined as the force which

binds atoms of same or different elements by mutual sharing of electrons in a covalent bond. Covalent compounds are solids, liquids or gases. They have low melting and boiling points compounds. They are more soluble in non-polar solvents.

The amount of energy released when one mole of ionic solid is formed by packing of gaseous ion is called :

- A. Ionisation energy
- B. Solvation energy
- C. Lattice energy
- D. Hydration energy

**Answer: C**







[Watch Video Solution](#)

**3.** Ionic bond is defined as the electrostatic force of attraction holding the oppositely charged ions. Ionic compounds are mostly crystalline solid having high melting and boiling points, electrical conductivity in molten



state, solubility in water etc. Covalent bond is defined as the force which binds atoms of same or different elements by mutual sharing of electrons in a covalent bond. Covalent compounds are solids, liquids or gases. They have low melting and boiling points compounds. They are more soluble in non-polar solvents.

Which of the following is arranged order of increasing boiling point ?



**Answer: B**



**Watch Video Solution**

4. Ionic bond is defined as the electrostatic force of attraction holding the oppositely charged ions. Ionic compounds are mostly crystalline solid having high melting and boiling points, electrical conductivity in molten

state, solubility in water etc. Covalent bond is defined as the force which binds atoms of same or different elements by mutual sharing of electrons in a covalent bond. Covalent compounds are solids, liquids or gases. They have low melting and boiling points compounds. They are more soluble in non-polar solvents.

The valence electrons are involved in formation of covalent bonds is /are called :

- A. non-bonding electrons
- B. lone pairs
- C. unshared pairs
- D. none of these

**Answer: D**



[Watch Video Solution](#)

5. Ionic bond is defined as the electrostatic force of attraction holding the oppositely charged ions. Ionic compounds are mostly crystalline solid

having high melting and boiling points, electrical conductivity in molten state, solubility in water etc. Covalent bond is defined as the force which binds atoms of same or different elements by mutual sharing of electrons in a covalent bond. Covalent compounds are solids, liquids or gases. They have low melting and boiling points compounds. They are more soluble in non-polar solvents.

The amount of energy released when one mole of ionic solid is formed by packing of gaseous ion is called :

- A. Ionisation energy
- B. Solvation energy
- C. Lattice energy
- D. Hydration energy





**Answer: C**



**Watch Video Solution**

6. Ionic bond is defined as the electrostatic force of attraction holding the oppositely charged ions. Ionic compounds are mostly crystalline solid having high melting and boiling points, electrical conductivity in molten state, solubility in water etc. Covalent bond is defined as the force which binds atoms of same or different elements by mutual sharing of electrons in a covalent bond. Covalent compounds are solids, liquids or gases. They have low melting and boiling points compounds. They are more soluble in non-polar solvents.

Which of the following is arranged order of increasing boiling point ?



**Answer: B**



**Watch Video Solution**

1. When an ionic compound is dissolved in water (polar solvent), it breaks up into its constituent ions. The given ionic compound will be dissolved in water if its hydration energy is more than lattice energy . IF hydration energy is less than lattice energy then ionic compound is usually either sparingly soluble or insoluble in water.

Which of the following ionic compound is having maximum lattice energy

:

A.  $\text{NaF}$

B.  $\text{MgF}_2$

C.  $\text{AlF}_3$

D.  $\text{KF}$





**Answer: C**



[Watch Video Solution](#)

2. When an ionic compound is dissolved in water (polar solvent), it breaks up into its constituent ions. The given ionic compound will be dissolved in water if its hydration energy is more than lattice energy . IF hydration energy is less than lattice energy then ionic compound is usually either sparingly soluble or insoluble in water.

Most hydrated cation is :



**Answer: A**



**Watch Video Solution**

3. When an ionic compound is dissolved in water (polar solvent), it breaks up into its constituent ions. The given ionic compound will be dissolved

in water if its hydration energy is more than lattice energy . IF hydration energy is less than lattice energy then ionic compound is usually either sparingly soluble or insoluble in water.

Which of the following ionic compound is having maximum lattice energy :

A. NaF

B.  $MgF_2$

C.  $AlF_3$

D. KF

**Answer: C**







[Watch Video Solution](#)

4. When an ionic compound is dissolved in water (polar solvent), it breaks up into its constituent ions. The given ionic compound will be dissolved in water if its hydration energy is more than lattice energy . IF hydration energy is less than lattice energy then ionic compound is usually either

sparingly soluble or insoluble in water.

Most hydrated cation is :



**Answer: A**



[Watch Video Solution](#)

### Level 3 (Passive 9)

1. A covalent bond will be formed by the overlapping of atomic orbitals having single electron of opposite spin, according to the overlapping of atomic orbitals the covalent bond may be of two types :



(i) Sigma bond ( $\sigma$ )      (ii) Pi bond ( $\pi$ )

Sigma bond is stronger bond than the Pi-bond. If atomic orbitals overlap



about the nuclear axis then sigma bond is formed but when atomic orbitals overlap sideway then Pi-bond is formed.

The correct order of increasing C-O bond length of  $CO$ ,  $CO_3^{2-}$ ,  $CO_2$  is :



**Answer: D**



**Watch Video Solution**


2. A covalent bond will be formed by the overlapping of atomic orbitals having single electron of opposite spin, according to the overlapping of atomic orbitals the covalent bond may be of two types :

(i) Sigma bond ( $\sigma$ )      (ii) Pi bond ( $\pi$ )

Sigma bond is stronger bond than the Pi-bond. If atomic orbitals overlap about the nuclear axis then sigma bond is formed but when atomic

orbitals overlap sideway then Pi-bond is formed.

Compound having maximum bond angle is :



D. none of these

**Answer: D**



[Watch Video Solution](#)

3. A covalent bond will be formed by the overlapping of atomic orbitals having single electron of opposite spin, according to the overlapping of atomic orbitals the covalent bond may be of two types :

(i) Sigma bond ( $\sigma$ )      (ii) Pi bond ( $\pi$ )

Sigma bond is stronger bond than the Pi-bond. If atomic orbitals overlap about the nuclear axis then sigma bond is formed but when atomic orbitals overlap sideway then Pi-bond is formed.

The strength of bonds formed by  $2s-2s$ ,  $2p-2p$  and  $2p-2s$  overlap has the order :

A.  $s - s > p - p > p - s$

B.  $s - s > p - s > p - p$

C.  $p - p > p - s > s - s$

D.  $p - p > s - s > p - s$

**Answer: C**



**Watch Video Solution**



4. A covalent bond will be formed by the overlapping of atomic orbitals having single electron of opposite spin, according to the overlapping of atomic orbitals the covalent bond may be of two types :

(i) Sigma bond ( $\sigma$ )      (ii) Pi bond ( $\pi$ )

Sigma bond is stronger bond than the Pi-bond. If atomic orbitals overlap about the nuclear axis then sigma bond is formed but when atomic

orbitals overlap sideway then Pi-bond is formed.

The correct order of increasing C-O bond length of  $CO$ ,  $CO_3^{2-}$ ,  $CO_2$  is :



**Answer: D**



**Watch Video Solution**


5. A covalent bond will be formed by the overlapping of atomic orbitals having single electron of opposite spin, according to the overlapping of atomic orbitals the covalent bond may be of two types :

(i) Sigma bond ( $\sigma$ )      (ii) Pi bond ( $\pi$ )

Sigma bond is stronger bond than the Pi-bond. If atomic orbitals overlap about the nuclear axis then sigma bond is formed but when atomic

orbitals overlap sideway then Pi-bond is formed.

Compound having maximum bond angle is :



D. none of these

**Answer: D**



[Watch Video Solution](#)

6. A covalent bond will be formed by the overlapping of atomic orbitals having single electron of opposite spin, according to the overlapping of atomic orbitals the covalent bond may be of two types :

(i) Sigma bond ( $\sigma$ )      (ii) Pi bond ( $\pi$ )

Sigma bond is stronger bond than the Pi-bond. If atomic orbitals overlap about the nuclear axis then sigma bond is formed but when atomic orbitals overlap sideway then Pi-bond is formed.

The strength of bonds formed by  $2s-2s$ ,  $2p-2p$  and  $2p-2s$  overlap has the order :

A.  $s - s > p - p > p - s$

B.  $s - s > p - s > p - p$

C.  $p - p > p - s > s - s$

D.  $p - p > s - s > p - s$

**Answer: C**





**Watch Video Solution**

### Level 3 (Passive 10)

1. According to VBT the extent of overlapping depends upon types of orbitals involved in overlapping and nature of overlapping. More will be the overlapping and the bond energy will also be high.

The incorrect order of bond dissociation energy will be :







**Answer: B**

 [Watch Video Solution](#)

2. According to VBT the extent of overlapping depends upon types of orbitals involved in overlapping and nature of overlapping. More will be the overlapping and the bond energy will also be high.

Which of the following combination of orbitals does not form covalent bond (x-axis is inter nuclear axis) :






**Answer: A**

 [Watch Video Solution](#)

3. According to VBT the extent of overlapping depends upon types of orbitals involved in overlapping and nature of overlapping. More will be the overlapping and the bond energy will also be high.

Which of the following compound does not form  $p\pi - p\pi$  bond ?







**Answer: C**

 [Watch Video Solution](#)



4. According to VBT the extent of overlapping depends upon types of orbitals involved in overlapping and nature of overlapping. More will be the overlapping and the bond energy will also be high.

The incorrect order of bond dissociation energy will be :







**Answer: B**



[Watch Video Solution](#)

5. According to VBT the extent of overlapping depends upon types of orbitals involved in overlapping and nature of overlapping. More will be the overlapping and the bond energy will also be high.

Which of the following combination of orbitals does not form covalent bond (x-axis is inter nuclear axis) :




**Answer: A**



[Watch Video Solution](#)

6. According to VBT the extent of overlapping depends upon types of orbitals involved in overlapping and nature of overlapping. More will be the overlapping and the bond energy will also be high.

Which of the following compound does not form  $p\pi - p\pi$  bond ?





**Answer: C**



**Watch Video Solution**

### Level 3 (Passive 11)

1. Consider the following elements with their period number and valence electrons.

Elements	Period number	Total valence <sup>e</sup> <sup>-</sup>
<i>P</i>	2	4
<i>Q</i>	2	6
<i>R</i>	3	7
<i>S</i>	3	3
<i>T</i>	3	6
<i>U</i>	3	4

According to the given informations, answer the following questions :

Choose incorrect statement :

A. R exhibits maximum covalency among all elements given

B. Q does not exhibit variable covalency

C. R exhibits minimum covalency among all elements given

D. R and S combine each other and form  $SR_5$  type of compound

**Answer: D**

 [Watch Video Solution](#)

2. Consider the following elements with their period number and valence electrons.

Elements	Period number	Total valencee <sup>-</sup>
<i>P</i>	2	4
<i>Q</i>	2	6
<i>R</i>	3	7
<i>S</i>	3	3
<i>T</i>	3	6
<i>U</i>	3	4

According to the given informations, answer the following questions :

Choose the correct statement :

A. Q has maximum value of electron affinity

B. R has maximum value of electronegativity

C. S has maximum atomic size

D. T and U are same group elements

**Answer: C**



**Watch Video Solution**

3. Consider the following elements with their period number and valence electrons.

Elements	Period number	Total valencee <sup>-</sup>
<i>P</i>	2	4
<i>Q</i>	2	6
<i>R</i>	3	7
<i>S</i>	3	3
<i>T</i>	3	6
<i>U</i>	3	4

According to the given informations, answer the following questions :

Choose incorrect statement :

A.  $SR_3$  is a hypovalent compound

B.  $UR_4$  can act as a Lewis acid

C.  $PQ_2$  can not acts as Lewis acid

D.  $UR_4 > SR_3$  : Lewis acidic character

**Answer: C**

 [Watch Video Solution](#)

4. Consider the following elements with their period number and valence electrons.

Elements	Period number	Total valencee <sup>-</sup>
<i>P</i>	2	4
<i>Q</i>	2	6
<i>R</i>	3	7
<i>S</i>	3	3
<i>T</i>	3	6
<i>U</i>	3	4

According to the given informations, answer the following questions :

Choose the incorrect statement :

A. R exhibits maximum covalency among all elements given

B. Q does not exhibit variable covalency

C. R exhibits minimum covalency among all elements given

D. R and S combine each other and form  $SR_5$  type of compound

**Answer: D**

 [Watch Video Solution](#)

5. Consider the following elements with their period number and valence electrons.

Elements	Period number	Total valencee <sup>-</sup>
<i>P</i>	2	4
<i>Q</i>	2	6
<i>R</i>	3	7
<i>S</i>	3	3
<i>T</i>	3	6
<i>U</i>	3	4

According to the given informations, answer the following questions :

Choose the correct statement :

A. Q has maximum value of electron affinity

B. R has maximum value of electronegativity

C. S has maximum atomic size

D. T and U are same group elements

**Answer: C**

 [Watch Video Solution](#)

6. Consider the following elements with their period number and valence electrons.

Elements	Period number	Total valencee <sup>-</sup>
<i>P</i>	2	4
<i>Q</i>	2	6
<i>R</i>	3	7
<i>S</i>	3	3
<i>T</i>	3	6
<i>U</i>	3	4

According to the given informations, answer the following questions :

Choose the incorrect statement :

A.  $SR_3$  is a hypovalent compound



B.  $UR_4$  can act as a Lewis acid

C.  $PQ_2$  can not acts as Lewis acid

D.  $UR_4 > SR_3$  : Lewis acidic character

**Answer: C**

 [Watch Video Solution](#)

### Level 3 (Passive 12)

1. Hybridisation involves the mixing of orbitals having comparable energies of same atom. Hybridised orbitals perform efficient overlapping than overlapping by pure s, p or orbitals.

Which of the following is not correctly match between given species and type of overlapping ?

A.  $XeO_3$  : Three ( $d\pi - p\pi$ ) bonds

B.  $H_2SO_4$  : Two ( $d\pi - p\pi$ ) bonds

C.  $SO_3$  : Three ( $d\pi - p\pi$ ) bonds

D.  $HClO_4$  : Three ( $d\pi - p\pi$ ) bonds

**Answer: C**

 [Watch Video Solution](#)

2. Hybridisation involves the mixing of orbitals having comparable energies of same atom. Hybridised orbitals perform efficient overlapping than overlapping by pure s, p or orbitals.

Consider the following compounds and select the incorrect statement from the following :

$NH_3, PH_3, H_2S, SO_2, BF_3, PCl_3, IF_7, P_4, H_2$

A. Six molecules out of given compounds involves hybridisation

B. Three molecules are hypervalent compounds

C. Six molecules out of above compounds are non-planar in structure

D. Two molecules out of given compounds involves ( $d\pi - p\pi$ )

bonding as well as also involves ( $p\pi - p\pi$ ) bonding

**Answer: C**



[View Text Solution](#)

3. Hybridisation involves the mixing of orbitals having comparable energies of same atom. Hybridised orbitals perform efficient overlapping than overlapping by pure s, p or orbitals.

Which of the following is not correctly match between given species and type of overlapping ?

A.  $XeO_3$  : Three ( $d\pi - p\pi$ ) bonds

B.  $H_2SO_4$  : Two ( $d\pi - p\pi$ ) bonds

C.  $SO_3$  : Three ( $d\pi - p\pi$ ) bonds

D.  $HClO_4$  : Three ( $d\pi - p\pi$ ) bonds

**Answer: C**



[Watch Video Solution](#)

4. Hybridisation involves the mixing of orbitals having comparable energies of same atom. Hybridised orbitals perform efficient overlapping than overlapping by pure s, p or orbitals.

Consider the following compounds and select the incorrect statement from the following :

$NH_3, PH_3, H_2S, SO_2, BF_3, PCl_3, IF_7, P_4, H_2$

- A. Six molecules out of given compounds involves hybridisation
- B. Three molecules are hypervalent compounds
- C. Six molecules out of above compounds are non-planar in structure
- D. Two molecules out of given compounds involves  $(d\pi - p\pi)$  bonding as well as also involves  $(p\pi - p\pi)$  bonding

**Answer: C**



[View Text Solution](#)

1. In which of the following there is intermolecular hydrogen bonding ?

- A. Water
- B. Ethanol
- C. Acetic acid
- D. H-F

Answer: A::B::C::D

 [Watch Video Solution](#)

2. Correct order of decreasing boiling points is :



- A.  $HF > HI > HBr > HCl$
- B.  $H_2O > H_2Te > H_2Se > H_2S$
- C.  $Br_2 > Cl_2 > F_2$



Answer: A::B::C

 Watch Video Solution

3. In which species the hybrid state of central atom is / are  $sp^3 d$  ?



Answer: A::B::C

 Watch Video Solution

4. Select correct statement(s) is /are :

- A. In  $AsH_3$  molecule lone pair at central atom is present in almost pure s-orbital
- B. Number of  $p\pi - d\pi$  bond in  $SO_3$  and  $SO_2$  are same
- C.  $NF_3$  is better Lewis base than  $NCl_3$
- D. Stable oxidation state of Lead is +2

**Answer: A::D**

 [Watch Video Solution](#)

5. Which of the following species does / do not exist ?




- A.  $OF_4$
- B.  $NH_2^-$
- C.  $NCl_5$
- D.  $ICl_3^{2-}$

**Answer: A::C::D**



Watch Video Solution

6. Which of the following species is /are superoctet molecule ?



Answer: C::D



Watch Video Solution

7. Which of the following statements is incorrect ?

A.  $\sigma$  - bond is weaker than  $\pi$ -bond

B. There are four co-ordinate bonds in the  $NH_4^+$  ions

C. The covalent bond is directional in nature



D. HF is less polar than HCl

**Answer: A::B::D**

 [Watch Video Solution](#)

**8.** Which of the following species is /are capable of forming a coordinate bond with  $BF_3$  ?

A.  $PH_3$

B.  $NH_4^+$

C.  $OH^-$

D.  $Mg^{2+}$

**Answer: A::C**

 [Watch Video Solution](#)

9. Ionic compounds in general do not possess :





- A. high melting points and non-directional bonds
- B. high melting points and low-boiling points
- C. directional bonds and low-boiling points
- D. high solubilities in polar and non-polar solvents

Answer: B::C::D



Watch Video Solution

10. Correct stability order of metal cation is /are :



**Answer: B::D**

 [Watch Video Solution](#)

11. Consider the following two molecules and according to the given information select correct statement(s) about  $AX_2$  and  $AY_2$  :

where A: 16th group of 3rd period element

X: more electronegative than (A) and same group number of (A) ltbgt Y:

Less atomic size than (A) and same period number of (A)

- A. The hybridization of central atoms are different in both compounds
- B. The shape of both molecules are same
- C. Both compounds are planar
- D. The X-A-X bond angle is less than Y-A-Y bond angle

**Answer: A::B::C**

 [Watch Video Solution](#)

12. Which of the following statements are correct about sulphur hexafluoride ?

A. all S-F bonds are equivalent

B.  $SF_6$  is a planar molecule

C. oxidation number of sulphur is the same as number of electrons of sulphur involved in bonding

D. sulphur has acquired the electronic structure of the gas argon

Answer: A:C

 [Watch Video Solution](#)

13. If  $AB_4^n$  types species are tetrahedral, then which of the following is /are correctly match ?

A.  $A$   $B$   $n$   
 $Xe$   $O$   $0$

B.  $A$   $B$   $n$   
 $Se$   $F$   $0$

- C.  $\begin{matrix} A & B & n \\ P & O & -3 \end{matrix}$
- D.  $\begin{matrix} A & B & n \\ N & H & +1 \end{matrix}$

**Answer: A::C::D**

 [Watch Video Solution](#)

14. Which of the following statements is correct ?

- A.  $ClF_3$  molecule is bent T-shape
- B. In  $SF_4$  molecule, F-S-F equatorial bond angle is  $103^\circ$  due to lp-lp repulsion
- C. In  $[ICl_4]^-$  molecular ion, Cl-I-Cl bond angle is  $90^\circ$
- D. In  $OBr_2$ , the bond angle is less than  $OCl_2$

**Answer: A::C**

 [Watch Video Solution](#)

15. Which of the following combination of bond pair (b.p.) and lone pair (l.p.) give same shape ?

(i) 3 b.p.+1 l.p. (ii) 2 b.p.+2 l.p. (iii) 2 b.p.1 l.p. (iv) 2 b.p. + 0 l. p.

(v) 3 b.p. +2 l.p. (vi) 2 b.p.+3 l.p.

A. ii and iii

B. iv and v

C. iv and vi

D. iii and vi

**Answer: A:C**



**Watch Video Solution**

16. Select the true statement(s) among the following :

A. Pure overlapping of two  $d_{xy}$  orbitals along x-axis results in the formation of  $\pi$ -bond

B.  $NO_2^+ > NO_3^- > NO_2^-$  is the correct order of bond angle as well as N-O bond order

C.  $NF_3 < NCl_3 < NBr_3 < NI_3$  is the correct order of Lewis basic character as well as bond angle

D.  $HF > HCl > HBr > HI$  is the correct order of dipole moment as well as boiling point

**Answer: A::C**

 [Watch Video Solution](#)

17.  $p_y$ -orbital can not form  $\pi$ -bond by lateral overlap with :

A.  $d_{xz}$ -orbital

B.  $d_{x^2-y^2}$ -orbitals

C.  $d_{xy}$ -orbital

D.  $p_z$ -orbital

Answer: A::B::D



Watch Video Solution

18. Which of the following orbital (s) cannot form  $\delta$ -bond ?

A.  $d_{x^2-y^2}$ -orbital

B.  $d_{xy}$ -orbital

C.  $d_{z^2}$ -orbital

D.  $p_x$ -orbital

Answer: C::D



Watch Video Solution

19. Select correct statement(s) regarding  $\sigma$  and  $\pi$  bonds :

A.  $\sigma$  - bond lies on the line joining the nuclei of bonded atoms



B.  $\pi$  - electron cloud lies on either side to the line joining the nuclei of bonded atoms

C.  $(2p_{\pi} - 3d_{\pi})\pi$  - bond is stronger than  $(2p_{\pi} - 3p_{\pi})\pi$  - bond.

D.  $\sigma$ -bond has primary effect to decide direction of covalent bond, while  $\pi$ - bond has no primary effect in direction of bond

**Answer: A::B::C::D**

 [Watch Video Solution](#)

20. Which of the following statements is / are correct ?

A. All carbon to carbon bonds contain a sigma bond and one or more  $\pi$ - bonds

B. All carbon to carbon bonds are sigma bonds




C. All oxygen to hydrogen bonds are hydrogen bonds

D. All carbon to hydrogen bonds are sigma bonds

Answer: D

 Watch Video Solution

21. Consider the following three orbitals :







Correct statement(s) regarding given information is /are :

- A. Orbitals (i) and (ii) can never form any type of covalent bond
- B. If internuclear axis is x, then combination of (ii) and (iii) orbitals can form  $\pi$ -bond
- C. Orbital (iii) can form  $\delta$ -bond with other orbital having same orientation of lobes
- D. If internuclear axis is 'x', then combination of (i) and (iii) orbitals can form  $\pi$ -bond

**Answer: A::C::D**

 [Watch Video Solution](#)


**22.** Which of the following combination of orbitals can not form bond. (If x axis in internuclear axis)



**Answer: A::C**

 [Watch Video Solution](#)

23. Consider the following atomic orbitals :



Which of the following statement(s) is /are correct regarding given orbital ?

- A. It is a gerade atomic orbital
- B. It has zero nodal plane
- C. Circular electron density is present in XY plane
- D. Opposite lobes of orbital have same sign of wave function ( $\psi$ )

**Answer: A::B::C::D**



[View Text Solution](#)

24. In which of the following there is intermolecular hydrogen bonding ?

- A. Water
- B. Ethanol
- C. Acetic acid
- D. H-F

Answer: A::B::C::D



[Watch Video Solution](#)

25. Correct order of decreasing boiling points is :



- A.  $HF > HI > HBr > HCl$
- B.  $H_2O > H_2Te > H_2Se > H_2S$
- C.  $Br_2 > Cl_2 > F_2$



Answer: A::B::C

 Watch Video Solution

26. In which species the hybrid state of central atom is / are  $sp^3 d$  ?



Answer: A::B::C

 Watch Video Solution

27. Select correct statement(s) is /are :

- A. In  $AsH_3$  molecule lone pair at central atom is present in almost pure s-orbital
- B. Number of  $p\pi - d\pi$  bond in  $SO_3$  and  $SO_2$  are same
- C.  $NF_3$  is better Lewis base than  $NCl_3$
- D. Stable oxidation state of Lead is +2

**Answer: A::D**

 [Watch Video Solution](#)

**28. Which of the following species does / do not exist ?**



- A.  $OF_4$
- B.  $NH_2^-$
- C.  $NCl_5$
- D.  $ICl_3^{2-}$

**Answer: A::C::D**



Watch Video Solution

29. Which of the following species is /are superoctet molecule ?



Answer: C::D



Watch Video Solution

30. Which of the following statements is incorrect ?

A.  $\sigma$  - bond is weaker than  $\pi$ -bond

B. There are four co-ordinate bonds in the  $NH_4^+$  ions

C. The covalent bond is directional in nature



D. HF is less polar than HCl

Answer: A::B::D

 [Watch Video Solution](#)

31. Which of the following species is /are capable of forming a coordinate bond with  $BF_3$  ?

A.  $PH_3$

B.  $NH_4^+$

C.  $OH^-$

D.  $Mg^{2+}$

Answer: A::C

 [Watch Video Solution](#)

32. Ionic compounds in general do not possess :

- A. high melting points and non-directional bonds
- B. high melting points and low-boiling points
- C. directional bonds and low-boiling points
- D. high solubilities in polar and non-polar solvents

Answer: B::C::D



Watch Video Solution

33. Correct stability order of metal cation is /are :

- A.  $Pb^{2+} < Sn^{2+}$
- B.  $Pb^{4+} < Pb^{2+}$
- C.  $Sn^{4+} < Sn^{2+}$
- D.  $Pb^{4+} < Sn^{4+}$

**Answer: B::D**

 [Watch Video Solution](#)

**34.** Consider the following two molecules and according to the given information select correct statement(s) about  $AX_2$  and  $AY_2$  :

where A: 16th group of 3rd period element

X: more electronegative than (A) and same group number of (A) ltbgt Y:

Less atomic size than (A) and same period number of (A)

- A. The hybridization of central atoms are different in both compounds
- B. The shape of both molecules are same
- C. Both compounds are planar
- D. The X-A-X bond angle is less than Y-A-Y bond angle

**Answer: A::B::C**

 [Watch Video Solution](#)

35. Which of the following statements are correct about sulphur hexafluoride ?

A. all S-F bonds are equivalent

B.  $SF_6$  is a planar molecule

C. oxidation number of sulphur is the same as number of electrons of sulphur involved in bonding

D. sulphur has acquired the electronic structure of the gas argon

Answer: A::C

 [Watch Video Solution](#)

36. If  $AB_n$  types species are tetrahedral, then which of the following is /are correctly match ?

A.  $A$   $B$   $n$   
 $Xe$   $O$   $0$

B.  $A$   $B$   $n$   
 $Se$   $F$   $0$

- C.  $\begin{matrix} A & B & n \\ P & O & -3 \end{matrix}$
- D.  $\begin{matrix} A & B & n \\ N & H & +1 \end{matrix}$

**Answer: A::C::D**

 [Watch Video Solution](#)

**37. Which of the following statements is correct ?**

- A.  $ClF_3$  molecule is bent T-shape
- B. In  $SF_4$  molecule, F-S-F equatorial bond angle is  $103^\circ$  due to lp-lp repulsion
- C. In  $[ICl_4]^-$  molecular ion, Cl-I-Cl bond angle is  $90^\circ$
- D. In  $OBr_2$ , the bond angle is less than  $OCl_2$

**Answer: A::C**

 [Watch Video Solution](#)

38. Which of the following combination of bond pair (b.p.) and lone pair (l.p.) give same shape ?

(i) 3 b.p.+1 l.p. (ii) 2 b.p.+2 l.p. (iii) 2 b.p.1 l.p. (iv) 2 b.p. + 0 l. p.

(v) 3 b.p. +2 l.p. (vi) 2 b.p.+3 l.p.

A. ii and iii

B. iv and v

C. iv and vi

D. iii and vi

Answer: A:C



Watch Video Solution

39. Select the true statement(s) among the following :

A. Pure overlapping of two  $d_{xy}$  orbitals along x-axis results in the formation of  $\pi$ -bond

B.  $NO_2^+ > NO_3^- > NO_2^-$  is the correct order of bond angle as well

as N-O bond order

C.  $NF_3 < NCl_3 < NBr_3 < NI_3$  is the correct order of Lewis basic

character as well as bond angle

D.  $HF > HCl > HBr > HI$  is the correct order of dipole moment

as well as boiling point

**Answer: A::C**



**Watch Video Solution**

40.  $p_y$ -orbital can not form  $\pi$ -bond by lateral overlap with :

A.  $d_{xz}$ -orbital

B.  $d_{x^2-y^2}$ -orbitals

C.  $d_{xy}$ -orbital

D.  $p_z$ -orbital

**Answer: A::B::D**

 [Watch Video Solution](#)

**41.** Which of the following orbital (s) cannot form  $\delta$ -bond ?

A.  $d_{x^2-y^2}$ -orbitals

B.  $d_{xy}$ -orbital

C.  $d_{z^2}$ -orbital

D.  $p_x$ -orbital

**Answer: C::D**

 [Watch Video Solution](#)

**42.** Select correct statements regarding  $\sigma$  and  $\pi$ -bonds

A.  $\sigma$  - bond lies on the line joining the nuclei of bonded atoms



B.  $\pi$  - electron cloud lies on either side to the line joining the nuclei of bonded atoms

C.  $(2p_{\pi} - 3d_{\pi})\pi$  - bond is stronger than  $(2p_{\pi} - 3p_{\pi})\pi$  - bond.

D.  $\sigma$ -bond has primary effect to decide direction of covalent bond, while  $\pi$ - bond has no primary effect in direction of bond

**Answer: A::B::C::D**

 [Watch Video Solution](#)

**43.** Which of the following statements is / are correct ?

A. All carbon to carbon bonds contain a sigma bond and one or more  $\pi$ - bonds

B. All carbon to carbon bonds are sigma bonds




C. All oxygen to hydrogen bonds are hydrogen bonds

D. All carbon to hydrogen bonds are sigma bonds

Answer: D

 Watch Video Solution

44. Consider the following three orbitals :







Correct statement(s) regarding given information is /are :

- A. Orbitals (i) and (ii) can never form any type of covalent bond
- B. If internuclear axis is x, then combination of (ii) and (iii) orbitals can form  $\pi$ -bond
- C. Orbital (iii) can form  $\delta$ -bond with other orbital having same orientation of lobes
- D. If internuclear axis is 'x', then combination of (i) and (iii) orbitals can form  $\pi$ -bond

**Answer: A::C::D**

 [Watch Video Solution](#)


**45.** Which of the following combination of orbitals do / does not form bond (if x-axis is internuclear axis) ?



**Answer: A::C**

 [Watch Video Solution](#)

46. Consider the following atomic orbitals :



Which of the following statement(s) is /are correct regarding given orbital ?

- A. It is a gerade atomic orbital
- B. It has zero nodal plane
- C. Circular electron density is present in XY plane
- D. Opposite lobes of orbital have same sign of wave function ( $\psi$ )

**Answer: A::B::C::D**



Watch Video Solution

## MATCH THE COLUMN

1. Column-I and Column -II contains four entries each. Entries of Column-I are to be matched with some entries of Column-II. One or more than one entries of Column-I may have the matching with the same entries of Column-II.

Column-I	Column-II
(A) $B_3N_3H_6$	(P) Planar geometry
(B) $I_3^-$	(Q) Non-planar geometry
(C) $B_2Cl_4$ (Solid)	(R) Compound having coordinate bond
(D) $SiF_4$	(S) Compound having back bond
	(T) Non-polar compound



View Text Solution

2. Column-I and Column -II contains four entries each. Entries of Column-I are to be matched with some entries of Column-II. One or more than one entries of Column-I may have the matching with the same entries of

## Column-II.

Column-I (Shape)	Column-II (Hybridisation)
(A) Linear	(P) $sp^3$
(B) Angular	(Q) $sp^3d^2$
(C) Square planar	(R) $sp^2$
(D) Trigonal planar	(S) $sp^3d$

 [Watch Video Solution](#)

3. Column-I and Column -II contains four entries each. Entries of Column-I are to be matched with some entries of Column-II. One or more than one entries of Column-I may have the matching with the same entries of Column-II.

Column-I	Column-II
(A) $SO_3$	(P) Largest bond angle
(B) $BeCl_2$	(Q) Lowest bond angle
(C) $NH_3$	(R) $sp^2$ -hybridisation
(D) $NO_2^-$	(S) $sp^3$ -hybridisation

 [Watch Video Solution](#)

4. Column-I and Column -II contains four entries each. Entries of Column-I are to be matched with some entries of Column-II. One or more than one entries of Column-I may have the matching with the same entries of

## Column-II.

Column-I	Column-II
(A) Hypo phosphoric acid	(P) All hydrogen are ionizable in water
(B) Pyro phosphorous acid	(Q) Lewis acid in water
(C) Boric acid	(R) Monobasic
(D) Hypo phosphorous acid	(S) $sp^3$ -hybridised central atom

 [View Text Solution](#)

5. Column-I and Column -II contains four entries each. Entries of Column-I are to be matched with some entries of Column-II. One or more than one entries of Column-I may have the matching with the same entries of Column-II.

Column-I	Column-II
(A) $\text{NH}_2^-$	(P) Square pyramidal
(B) $\text{XeOF}_2$	(Q) V-shaped
(C) $\text{ICl}_4^-$	(R) T-shaped
(D) $[\text{SbF}_5]^{2-}$	(S) Square planar

 [Watch Video Solution](#)

6. Column-I and Column -II contains four entries each. Entries of Column-I are to be matched with some entries of Column-II. One or more than one

entries of Column-I may have the matching with the same entries of Column-II.

Column-I	Column-II
(A) $\text{ICl}_2$	(P) Linear
(B) $\text{BrF}_2^+$	(Q) Pyramidal
(C) $\text{ClF}_4^-$	(R) Tetrahedral
(D) $\text{AlCl}_4^-$	(S) Square planar
	(T) Angular

 [Watch Video Solution](#)

7. Column-I and Column -II contains four entries each. Entries of Column-I are to be matched with some entries of Column-II. One or more than one entries of Column-I may have the matching with the same entries of Column-II.

Column-I	Column-II
(A) $\text{Re}_2\text{Cl}_8^{2-}$	(P) $p\pi - p\pi$ bonding
(B) $\text{NO}_3^-$	(Q) $p\pi - d\pi$ bonding
(C) $\text{SO}_4^{2-}$	(R) $d\pi - d\pi$ bonding
(D) $\text{SO}_3$	(S) $\delta$ -bonding

 [Watch Video Solution](#)



8. Match the refining methods (Column I) with (Column II).

Column I (Refining methods)	Column II (Metals)
I. Liquation	(a) Zr
II. Zone Refining	(b) Ni
III. Mond Process	(c) Sn
IV. Van Arkel Method	(d) Ga

 [Watch Video Solution](#)

9. Column-I and Column -II contains four entries each. Entries of Column-I are to be matched with some entries of Column-II. One or more than one entries of Column-I may have the matching with the same entries of Column-II.

Column-I	Column-II
(A) $\text{AsO}_4^{3-}$	(P) All three $p$ -orbitals used in hybridisation
(B) $\text{ICl}_2^+$	(Q) Tetrahedral shape
(C) $\text{SOF}_4$	(R) Axial $d$ -orbital with two nodal cones used in hybridisation
(D) $\text{XeOF}_4$	(S) All bond lengths are identical
	(T) $p\pi - d\pi$ bond(s) present

 [Watch Video Solution](#)

10. Column-I and Column -II contains four entries each. Entries of Column-I are to be matched with some entries of Column-II. One or more than one entries of Column-I may have the matching with the same entries of Column-II.

Column-I
$\sigma$ -bond pairs + lone pairs around central atom of $AB_2$ type compound
(A) 2 + 1
(B) 2 + 3
(C) 4 + 2
(D) 2 + 2

Column-II
Characteristics/shape of compound
(P) Linear
(Q) Angular
(R) Polar
(S) Non-polar
(T) Planar



[Watch Video Solution](#)

11. Column-I and Column -II contains four entries each. Entries of Column-I are to be matched with some entries of Column-II. One or more than one entries of Column-I may have the matching with the same entries of

## Column-II.

Column-I (Type of bond formed)	Column-II (Combining orbitals (Internuclear axis])
(A) $\pi$ -bond	(P) $d_{yz} + p_y, (z)$
(B) $\sigma$ -bond	(Q) $s + p_x, (y)$
(C) $\delta$ -bond	(R) $d_{yz} + d_{yz}, (x)$
(D) Non-bonding	(S) $s + s, (z)$
	(T) $s + d_{xy}, (y)$

 [Watch Video Solution](#)

12. Column-I and Column-II contains four entries each. Entries of Column-I are to be matched with some entries of Column-II. One or more than one entries of Column-I may have the matching with the same entries of Column-II.

Column-I	Column-II
(A) $\text{XeF}_5^-$	(P) $d$ -orbital with zero nodal plane is used in hybridisation
(B) $\text{PBr}_4^+$	(Q) Non-axial $d$ -orbital is used in hybridisation
(C) $\text{IOF}_3$	(R) Planar species
(D) $\text{NH}_2^-$	(S) Non-planar species
	(T) Bond angle $109^\circ 28'$ or less than $109^\circ 28'$

 [View Text Solution](#)

13. Column-I and Column -II contains four entries each. Entries of Column-I are to be matched with some entries of Column-II. One or more than one entries of Column-I may have the matching with the same entries of Column-II.

Column-I	Column-II
(A) $\text{IO}_2\text{F}_2^-$	(P) Trigonal pyramidal shape
(B) $\text{IOF}_4^-$	(Q) Square pyramidal shape
(C) $\text{SeOF}_2$	(R) See-saw shape
(D) $\text{XeOF}_2$	(S) Non-planar
	(T) One of the bond angle $< 90^\circ$

 [View Text Solution](#)

14. Column-I and Column -II contains four entries each. Entries of Column-I are to be matched with some entries of Column-II. One or more than one entries of Column-I may have the matching with the same entries of Column-II.

Column-I	Column-II
(A) $\text{B}_3\text{N}_3\text{H}_6$	(P) Planar geometry
(B) $\text{I}_3^-$	(Q) Non-planar geometry
(C) $\text{B}_2\text{Cl}_4$ (Solid)	(R) Compound having coordinate bond
(D) $\text{SiF}_4$	(S) Compound having back bond
	(T) Non-polar compound

 [Watch Video Solution](#)

15. Column-I and Column -II contains four entries each. Entries of Column-I are to be matched with some entries of Column-II. One or more than one entries of Column-I may have the matching with the same entries of Column-II.

Column-I (Shape)	Column-II (Hybridisation)
(A) Linear	(P) $sp^3$
(B) Angular	(Q) $sp^3d^2$
(C) Square planar	(R) $sp^2$
(D) Trigonal planar	(S) $sp^3d$



Watch Video Solution

16. Column-I and Column -II contains four entries each. Entries of Column-I are to be matched with some entries of Column-II. One or more than one entries of Column-I may have the matching with the same entries of Column-II.

Column-I	Column-II
(A) $SO_3$	(P) Largest bond angle
(B) $BeCl_2$	(Q) Lowest bond angle
(C) $NH_3$	(R) $sp^2$ -hybridisation
(D) $NO_2^-$	(S) $sp^3$ -hybridisation



Watch Video Solution

17. Column-I and Column -II contains four entries each. Entries of Column-I are to be matched with some entries of Column-II. One or more than one entries of Column-I may have the matching with the same entries of Column-II.

Column-I	Column-II
(A) Hypo phosphoric acid	(P) All hydrogen are ionizable in water
(B) Pyro phosphorous acid	(Q) Lewis acid in water
(C) Boric acid	(R) Monobasic
(D) Hypo phosphorous acid	(S) $sp^3$ -hybridised central atom

 [Watch Video Solution](#)

18. Column-I and Column -II contains four entries each. Entries of Column-I are to be matched with some entries of Column-II. One or more than one entries of Column-I may have the matching with the same entries of Column-II.

Column-I	Column-II
(A) $\text{NH}_2^-$	(P) Square pyramidal
(B) $\text{XeOF}_2$	(Q) V-shaped
(C) $\text{ICl}_4^-$	(R) T-shaped
(D) $[\text{SbF}_5]^{2-}$	(S) Square planar

 [Watch Video Solution](#)

19. Column-I and Column -II contains four entries each. Entries of Column-I are to be matched with some entries of Column-II. One or more than one entries of Column-I may have the matching with the same entries of Column-II.

Column-I	Column-II
(A) $\text{ICl}_2$	(P) Linear
(B) $\text{BrF}_2^+$	(Q) Pyramidal
(C) $\text{ClF}_4^-$	(R) Tetrahedral
(D) $\text{AlCl}_4^-$	(S) Square planar
	(T) Angular

 [Watch Video Solution](#)

20. Column-I and Column -II contains four entries each. Entries of Column-I are to be matched with some entries of Column-II. One or more than one entries of Column-I may have the matching with the same entries of Column-II.

Column-I	Column-II
(A) $\text{Re}_2\text{Cl}_8^{2-}$	(P) $p\pi - p\pi$ bonding
(B) $\text{NO}_3^-$	(Q) $p\pi - d\pi$ bonding
(C) $\text{SO}_4^{2-}$	(R) $d\pi - d\pi$ bonding
(D) $\text{SO}_3$	(S) $\delta$ -bonding

21. Column-I and Column -II contains four entries each. Entries of Column-I are to be matched with some entries of Column-II. One or more than one entries of Column-I may have the matching with the same entries of Column-II.

Column-I (Bond order range)	Column-II (Oxyanions)
(A) 1.0 to 1.30	(P) $\text{NO}_3^-$
(B) 1.31 to 1.55	(Q) $\text{ClO}_4^-$
(C) 1.56 to 1.70	(R) $\text{PO}_4^{3-}$
(D) 1.71 to 2.0	(S) $\text{ClO}_3^-$
	(T) $\text{SO}_4^{2-}$

22. Column-I and Column -II contains four entries each. Entries of Column-I are to be matched with some entries of Column-II. One or more than one entries of Column-I may have the matching with the same entries of



## Column-II.

Column-I	Column-II
(A) $\text{AsO}_4^{3-}$	(P) All three $p$ -orbitals used in hybridisation
(B) $\text{ICl}_2^+$	(Q) Tetrahedral shape
(C) $\text{SOF}_4$	(R) Axial $d$ -orbital with two nodal cones used in hybridisation
(D) $\text{XeOF}_4$	(S) All bond lengths are identical
	(T) $p\pi - d\pi$ bond(s) present

 [Watch Video Solution](#)

23. Column-I and Column -II contains four entries each. Entries of Column-I are to be matched with some entries of Column-II. One or more than one entries of Column-I may have the matching with the same entries of Column-II.

Column-I	Column-II
$\sigma$ -bond pairs + lone pairs around central atom of $\text{AB}_n$ type compound	Characteristics/shape of compound
(A) 2 + 1	(P) Linear
(B) 2 + 3	(Q) Angular
(C) 4 + 2	(R) Polar
(D) 2 + 2	(S) Non-polar
	(T) Planar

 [Watch Video Solution](#)

24. Column-I and Column -II contains four entries each. Entries of Column-I are to be matched with some entries of Column-II. One or more than one entries of Column-I may have the matching with the same entries of Column-II.

Column-I (Type of bond formed)	Column-II (Combining orbitals (Internuclear axis])
(A) $\pi$ -bond	(P) $d_{yz} + p_y, (z)$
(B) $\sigma$ -bond	(Q) $s + p_x, (y)$
(C) $\delta$ -bond	(R) $d_{yz} + d_{yz}, (x)$
(D) Non-bonding	(S) $s + s, (z)$
	(T) $s + d_{xy}, (y)$

 [Watch Video Solution](#)

25. Column-I and Column -II contains four entries each. Entries of Column-I are to be matched with some entries of Column-II. One or more than one entries of Column-I may have the matching with the same entries of Column-II.

Column-I	Column-II
(A) $\text{XeF}_5^-$	(P) $d$ -orbital with zero nodal plane is used in hybridisation
(B) $\text{PBr}_4^+$	(Q) Non-axial $d$ -orbital is used in hybridisation
(C) $\text{IOF}_3$	(R) Planar species
(D) $\text{NH}_2^-$	(S) Non-planar species
	(T) Bond angle $109^\circ 28'$ or less than $109^\circ 28'$

26. Column-I and Column -II contains four entries each. Entries of Column-I are to be matched with some entries of Column-II. One or more than one entries of Column-I may have the matching with the same entries of Column-II.

Column-I	Column-II
(A) $\text{IO}_2\text{F}_2^-$	(P) Trigonal pyramidal shape
(B) $\text{IOF}_4^-$	(Q) Square pyramidal shape
(C) $\text{SeOF}_2$	(R) See-saw shape
(D) $\text{XeOF}_2$	(S) Non-planar
	(T) One of the bond angle $< 90^\circ$

## ASSERTION-REASON TYPE QUESTIONS

1. Assertion : Multiple bond between two bonded atoms can have more than three bonds.

Reason : Multiple bond between two bonded atoms can not have more than two  $\pi$ -bonds.

- A. If assertion is true but the reason is false
- B. If assertion is false but reason is true
- C. IF both assertion and reason are true and the reason is the correct explanation of assertion
- D. If both assertion and reason are true but reason is not the correct explanation of assertion

**Answer: D**



**Watch Video Solution**

2. Assertion :  $2^{nd}$  period elements do not involve in excitation of electron.

Reason :  $2^{nd}$  period elements do not have vacant 2d-orbitals.

- A. If assertion is true but the reason is false
- B. If assertion is false but reason is true

C. IF both assertion and reason are true and the reason is the correct explanation of assertion

D. If both assertion and reason are true but reason is not the correct explanation of assertion

**Answer: B**

 [Watch Video Solution](#)

3. Assertion : In  $SO_3$  molecule bond dissociation energy of all S=O bonds are not equivalent.

Reason :  $SO_3$  molecule is having two types of  $2p\pi - 3p\pi$  and  $2p\pi - 3d\pi$  pi-bonds.

A. If assertion is true but the reason is false

B. If assertion is false but reason is true

C. IF both assertion and reason are true and the reason is the correct explanation of assertion

D. If both assertion and reason are true but reason is not the correct explanation of assertion

**Answer: B**

 [Watch Video Solution](#)

4. Assertion : All diatomic molecules with polar bond have dipole moment.

Reason : Dipole moment is a vector quantity.

A. If assertion is true but the reason is false

B. If assertion is false but reason is true

C. IF both assertion and reason are true and the reason is the correct explanation of assertion

D. If both assertion and reason are true but reason is not the correct explanation of assertion

**Answer: D**



**Watch Video Solution**

**5. Assertion :** Water is a good solvent for ionic compounds but poor one for covalent compounds.

**Reason :**Hydrogen energy of ions releases sufficient energy to overcome lattice energy and break hydrogen bonds in water, while covalent bonded compound interact so weakly that even van der Waals force between molecule of covalent compounds cannot be broken .

A. If assertion is true but the reason is false

B. If assertion is false but reason is true

C. IF both assertion and reason are true and the reason is the correct explanation of assertion

D. If both assertion and reason are true but reason is not the correct explanation of assertion

**Answer: C**



**Watch Video Solution**

**6. Assertion :** Xe-atom in  $XeF_2$  assumes sp-hybrid state.

**Reason :**  $XeF_2$  molecule does not follow octet rule.

- A. If assertion is true but the reason is false
- B. If assertion is false but reason is true
- C. IF both assertion and reason are true and the reason is the correct explanation of assertion
- D. If both assertion and reason are true but reason is not the correct explanation of assertion

**Answer: B**



**Watch Video Solution**



7. Assertion : The atoms in a covalent molecule are said to share electrons, yet some covalent molecule are polar.

Reason :In a polar covalent molecule , the shared electron spend more time on the average near one of the atoms .

- A. If assertion is true but the reason is false
- B. If assertion is false but reason is true
- C. IF both assertion and reason are true and the reason is the correct explanation of assertion
- D. If both assertion and reason are true but reason is not the correct explanation of assertion

**Answer: C**



[Watch Video Solution](#)

8. Assertion :  $CCl_4$  is a non-polar molecule.

Reason :  $CCl_4$  has polar bonds.

- A. If assertion is true but the reason is false
- B. If assertion is false but reason is true
- C. IF both assertion and reason are true and the reason is the correct explanation of assertion
- D. If both assertion and reason are true but reason is not the correct explanation of assertion

**Answer: D**



**Watch Video Solution**

**9. Assertion :** Geometry of  $ICl_3$  is tetrahedral.

**Reason :** Its shape is T-shape, due to the presence of two lone pairs.

- A. assertion is true but the reason is false
- B. assertion is false but reason is true

C. both assertion and reason are true and the reason is the correct explanation of assertion

D. both assertion and reason are true but reason is not the correct explanation of assertion

**Answer: B**

 [Watch Video Solution](#)

**10.** Assertion : The covalency of carbon is four in excited state.

Reason : The four half-filled pure orbitals of carbon form same kind of bonds with an atom as those are with hybridised orbitals.

A. If assertion is true but the reason is false

B. If assertion is false but reason is true

C. IF both assertion and reason are true and the reason is the correct explanation of assertion

D. If both assertion and reason are true but reason is not the correct explanation of assertion

**Answer: A**

 [Watch Video Solution](#)

**11.** Assertion : The shape of  $XeF_4$  is square-planar.

Reason : In an octahedral geometry, a single lone pair can occupy any position but a second lone pair will occupy the opposite position to the first lone pair.

A. If assertion is true but the reason is false

B. If assertion is false but reason is true

C. If both assertion and reason are true and the reason is the correct explanation of assertion

D. If both assertion and reason are true but reason is not the correct explanation of assertion

**Answer: C**

 [Watch Video Solution](#)

**12.** Assertion : Multiple bond between two bonded atoms can have more than three bonds.

Reason : Multiple bond between two bonded atoms can not have more than two  $\pi$ -bonds.

A. If assertion is true but the reason is false

B. If assertion is false but reason is true

C. IF both assertion and reason are true and the reason is the correct explanation of assertion

D. If both assertion and reason are true but reason is not the correct explanation of assertion

**Answer: D**

 [Watch Video Solution](#)

13. Assertion :  $2^{nd}$  period elements do not involve in excitation of electron.

Reason :  $2^{nd}$  period elements do not have vacant 2d-orbitals.

A. If assertion is true but the reason is false

B. If assertion is false but reason is true

C. IF both assertion and reason are true and the reason is the correct explanation of assertion

D. If both assertion and reason are true but reason is not the correct explanation of assertion

**Answer: B**



**Watch Video Solution**

14. Assertion : In  $SO_3$  molecule bond dissociation energy of all S=O bonds are not equivalent.

Reason :  $SO_3$  molecule is having two types of  $2p\pi - 3p\pi$  and  $2p\pi - 3d\pi$  pi-bonds.

- A. If assertion is true but the reason is false
- B. If assertion is false but reason is true
- C. IF both assertion and reason are true and the reason is the correct explanation of assertion
- D. If both assertion and reason are true but reason is not the correct explanation of assertion

**Answer: B**



[Watch Video Solution](#)

15. Assertion :  $PH_4^+$  ion is having tetrahedron geometry.

Reason : P-atom is unhybridised in  $PH_4^+$  ion.

A. If assertion is true but the reason is false

B. If assertion is false but reason is true

C. IF both assertion and reason are true and the reason is the correct explanation of assertion

D. If both assertion and reason are true but reason is not the correct explanation of assertion

**Answer: A**



[Watch Video Solution](#)

16. Assertion : All diatomic molecules with polar bond have dipole moment.

Reason : Dipole moment is a vector quantity.



- A. If assertion is true but the reason is false
- B. If assertion is false but reason is true
- C. IF both assertion and reason are true and the reason is the correct explanation of assertion
- D. If both assertion and reason are true but reason is not the correct explanation of assertion

**Answer: D**

 [Watch Video Solution](#)

**17. Assertion :** Water is a good solvent for ionic compounds but poor for covalent compounds.

**Reason :** Hydration energy of ions releases sufficient energy to overcome lattice energy and break hydrogen bonds in water while covalent compounds interact so weakly that even van der Waal's forces between molecules of covalent compounds cannot be broken.

- A. If assertion is true but the reason is false
- B. If assertion is false but reason is true
- C. IF both assertion and reason are true and the reason is the correct explanation of assertion
- D. If both assertion and reason are true but reason is not the correct explanation of assertion

**Answer: C**

 [Watch Video Solution](#)

**18.** Assertion : Xe-atom in  $XeF_2$  assumes sp-hybrid state.

Reason :  $XeF_2$  molecule does not follow octet rule.

- A. If assertion is true but the reason is false
- B. If assertion is false but reason is true

C. IF both assertion and reason are true and the reason is the correct explanation of assertion

D. If both assertion and reason are true but reason is not the correct explanation of assertion

**Answer: B**

 [Watch Video Solution](#)

**19.** Assertion : The atoms in a covalent molecule are said to share electrons, yet some covalent molecules are polar.

Reason : In polar covalent molecule, the shared electrons spend more time on the average near one of the atoms.

A. If assertion is true but the reason is false

B. If assertion is false but reason is true

C. IF both assertion and reason are true and the reason is the correct explanation of assertion

D. If both assertion and reason are true but reason is not the correct explanation of assertion

**Answer: C**



**Watch Video Solution**

**20.** Assertion :  $CCl_4$  is a non-polar molecule.

Reason :  $CCl_4$  has polar bonds.

A. If assertion is true but the reason is false

B. If assertion is false but reason is true

C. IF both assertion and reason are true and the reason is the correct explanation of assertion

D. If both assertion and reason are true but reason is not the correct explanation of assertion

**Answer: D**

 [Watch Video Solution](#)

21. Assertion : Geometry of  $ICl_3$  is tetrahedral.

Reason : Its shape is T-shape, due to the presence of two lone pairs.

- A. If assertion is true but the reason is false
- B. If assertion is false but reason is true
- C. IF both assertion and reason are true and the reason is the correct explanation of assertion
- D. If both assertion and reason are true but reason is not the correct explanation of assertion

**Answer: B**

 [Watch Video Solution](#)

**22. Assertion :** The covalency of carbon is four in excited state.

**Reason :** The four half-filled pure orbitals of carbon form same kind of bonds with an atom as those are with hybridised orbitals.

A. If assertion is true but the reason is false

B. If assertion is false but reason is true

C. IF both assertion and reason are true and the reason is the correct explanation of assertion

D. If both assertion and reason are true but reason is not the correct explanation of assertion

**Answer: A**



[Watch Video Solution](#)

**23. Assertion :** The shape of  $XeF_4$  is square-planar.

**Reason :** In an octahedral geometry, a single lone pair can occupy any

position but a second lone pair will occupy the opposite position to the first lone pair.


- A. If assertion is true but the reason is false
- B. If assertion is false but reason is true
- C. IF both assertion and reason are true and the reason is the correct explanation of assertion
- D. If both assertion and reason are true but reason is not the correct explanation of assertion


**Answer: C**

 [Watch Video Solution](#)

## SUBJECTIVE PROBLEMS

1. Consider following compounds A to E :





If value of  $n$  is 4, then calculate value of  $p \div q$  here, 'p' is total number of bond pair and 'q' is total number of lone pair on central atoms of compounds (A) to (E).

 [Watch Video Solution](#)

2. Consider the following five group (According to modern periodic table) of elements with their increasing order to atomic numbers :

Group 1  $\rightarrow A, B, C, D, E$       Group 2  $\rightarrow F, G, H, I, J$

Group

13  $\rightarrow K, L, M, N, O$       Group 15  $\rightarrow P, Q, R, S, T$       Group 17  $\rightarrow U$

If first and last element of each group belongs to 2nd and 6th period respectively and Z represents to carbonate ion ( $CO_3^{2-}$ ) then consider the following orders.

(i)  $O^+ > H^{2+}$ , Polarising power

(ii)  $T^{3+} > S^{3+} > R^{3+}$ , Stability of cation

(iii)  $U^-(aq) > V^-(aq) > W^-(aq) > X^-(aq)$ , Size

(iv)  $JV_2 < IV_2 < GV_2 < LV_3$ , Covalent character



(v)  $GZ > IZ > JZ$ , Thermal stability

(vi)  $AV > BV > CV > DV > EV$ , Thermal stability

(vii)  $C_3P > B_3P > A_3P$ , Lattice energy

(viii)  $KU_3 < KV_3 < KW_3 < KX_3$ , Melting point

Then calculate value of  $|p - q|^2$ , here p and q are correct and incorrect orders in the given eight orders respectively.



[View Text Solution](#)

3. Consider the following species and find out total number of species which are polar and can act as Lewis acid

$CCl_4, CO_2, SO_2, AlCl_3, HCHO, SO_3, SiCl_4, BCl_3, CF_4$



[Watch Video Solution](#)

4. Consider the following table regarding interhalogen compounds,  $XY_n$  (where Y is more electronegative than X)

Value of n for respective interhalogen compound	Total number of d-orbitals used in hybridization of central atom	Polarity	Planarity
$P_1$	1	Polar	Planar
$P_2$	$Q_1$	Polar	Non-Planar
$P_3$	$Q_2$	Non-Polar	Non-Planar

Then according to given information calculate value of expression

$$P_2 \times \left( \frac{P_3 - P_1}{(Q_1 + Q_2)} \right)$$

 [View Text Solution](#)

5. What is covalency of chlorine atom in second excited state ?

 [Watch Video Solution](#)

6. Sum of  $\sigma$  and  $\pi$  bonds in  $NH_4^+$  cation is ..

 [Watch Video Solution](#)

7. Calculate the value of X-Y, for  $XeOF_4$ . (X=Number of  $\sigma$  bond pair and Y=Number of lone pair on central atom)

 [Watch Video Solution](#)

[Watch Video Solution](#)

8. The molecule  $AB_n$  is planar with six pairs of electrons around A in the valence shell. The value of n is

[Watch Video Solution](#)

9. Calculate value of  $\frac{X + Y + Z}{10}$ , here X is O-N-O bond angle in  $NO_3^-$  Y is O-N-O bond angle in  $NO_2^+$  and Z is F-Xe-F adjacent bond angle in  $XeF_4$ .

[Watch Video Solution](#)

10. Calculate  $x+y+z$  for  $H_3PO_3$  acid, where x is no. of lone pairs, y is no. of  $\sigma$  bonds and z is no. of  $\pi$  bonds.

[Watch Video Solution](#)

11. How many right angle, bond angles are present in  $TeF_5^-$  molecular ion ?

 [Watch Video Solution](#)

12. How many possible  $\angle FSeF$  bond angles are present in  $SeF_4$  molecule ?


 [Watch Video Solution](#)

13. In  $IF_6^-$  and  $TeF_5^-$ , sum of axial d-orbitals which are used in hybridisation in both species.

 [Watch Video Solution](#)

14. Among the following, total no. of planar species is :

(i)  $SF_4$     (ii)  $BrF_3$     (iii)  $XeF_2$     (iv)  $IF_5$



 [Watch Video Solution](#)

15. Calculate the value of "x+y-z" here x,y and z are total number of non-bonded electron pair (s),pie ( $\pi$ ) bond(s) and sigma ( $\sigma$ ) bonds in hydrogen phosphite ion respectively.

 [Watch Video Solution](#)

16. Consider the following table

Total number of electron pairs (l.p. + $\sigma$ -bond)	Total number of lone pairs	Shape
5	...p...	linear
...q...	1	see-saw
4	...r...	Bent shape
...s...	2	Square planar
5	...t...	Bent 'T' shape

Then calculate value of "p+q+r-s-t".

 [Watch Video Solution](#)

17. In phosphorus acid, if X is number of non bonding electron pairs. Y is number of  $\sigma$ -bonds and Z is number of  $\pi$ - bonds. Then, calculate value of  $Y \times Z - X$ .

 [Watch Video Solution](#)

18. Calculate the number of  $p_{\pi} - d_{\pi}$  bond(s) present in  $SO_4^{2-}$  :

 [Watch Video Solution](#)

19. Sum of  $\sigma$  and  $\pi$  bonds in  $NH_4^+$  cation is ..

 [Watch Video Solution](#)

20. Consider the following orbitals (i)  $3p_x$  (ii)  $4d_{z^2}$  (iii)  $3d_{x^2 - y^2}$  (iv)  $3d_{yz}$

Then, calculate value of " $x + y - z$ " here x is total number of gerade

orbital and y is total number of ungerade orbitals and z is total number of axial orbitals in given above orbitals.

 [Watch Video Solution](#)

21. Calculate value of  $|x-y|$ , here x and y are the total number of bonds in benzene and benzyne respectively which are formed by overlapping of hybridized orbitals.

 [Watch Video Solution](#)


22. Consider the following compounds:

(i)  $IF_5$  (ii)  $ClI_4^-$  (iii)  $XeO_2F_2$  (iv)  $NH_2^-$  (v)  $BCl_3$  (vi)  $BeCl_2$  (vii)  $AsCl_4^+$  (viii)  $B(OH)_3$  (ix)  $NO_2^-$  (x)  $ClO_2^+$

Then, calculate value of " $x + y - z$ ", here x, y and z are total number of compounds in given compounds in which central atom used their all three p-orbitals, only two p-orbitals and only one p-orbital in hybridisation respectively:


 [Watch Video Solution](#)

23. Total number of species which used all three p-orbitals in hybridisation of central atom and should be non-polar also.



 Watch Video Solution

24. Consider the following species



Then calculate value of  $|x-y|$ , where

x : Total number of species which have bond order 1.5 or greater than 1.5

y : Total number of species which have bond order less than 1.5

 Watch Video Solution

25. Consider the following orbitals


$3s, 2p_x, 4d_{xy}, 4d_{z^2}, 3d_{x^2-y^2}, 3p_y, 4s, 4p_z$  and find total number of orbital



(s) having even number of nodal plane.

 [Watch Video Solution](#)

26. For the following molecules :



Calculate the value of  $\frac{a+b}{c}$

a = Number of species having  $sp^3$  d-hybridisation


b = Number of species which are planar


c = Number of species which are non-planar

 [View Text Solution](#)

27. Find out number of transformation among following which involves the change of hybridisation of underlined atom.



(a)





 [Watch Video Solution](#)

**28.** Consider following compounds A to E :




If value of n is 4, then calculate value of  $p \div q$  here, 'p' is total number of bond pair and 'q' is total number of lone pair on central atoms of compounds (A) to (E).

 [Watch Video Solution](#)

**29.** Consider the following five group (According to modern periodic table) of elements with their increasing order to atomic numbers :



Group



If first and last element of each group belongs to 2nd and 6th period respectively and Z represents to carbonate ion ( $\text{CO}_3^{2-}$ ) then consider the following orders.

(i)  $\text{O}^+ > \text{H}^{2+}$ , Polarising power

(ii)  $\text{T}^{3+} > \text{S}^{3+} > \text{R}^{3+}$ , Stability of cation

(iii)  $\text{U}^-(\text{aq}) > \text{V}^-(\text{aq}) > \text{W}^-(\text{aq}) > \text{X}^-(\text{aq})$ , Size

(iv)  $\text{JV}_2 < \text{IV}_2 < \text{GV}_2 < \text{LV}_3$ , Covalent character

(v)  $\text{GZ} > \text{IZ} > \text{JZ}$ , Thermal stability

(vi)  $\text{AV} > \text{BV} > \text{CV} > \text{DV} > \text{EV}$ , Thermal stability

(vii)  $\text{C}_3\text{P} > \text{B}_3\text{P} > \text{A}_3\text{P}$ , Lattice energy

(viii)  $\text{KU}_3 < \text{KV}_3 < \text{KW}_3 < \text{KX}_3$ , Melting point

Then calculate value of  $|p - q|^2$ , here p and q are correct and incorrect orders in the given eight orders respectively.



[View Text Solution](#)

**30.** Consider the following species and find out total number of species which are polar and can act as Lewis acid

$\text{CCl}_4, \text{CO}_2, \text{SO}_2, \text{AlCl}_3, \text{HCHO}, \text{SO}_3, \text{SiCl}_4, \text{BCl}_3, \text{CF}_4$

 [Watch Video Solution](#)

31. Consider the following table regarding interhalogen compounds,  $XY_n$  (where Y is more electronegative than X)

Value of n for respective interhalogen compound	Total number of d-orbitals used in hybridization of central atom	Polarity	Planarity
$P_1$	1	Polar	Planar
$P_2$	$Q_1$	Polar	Non-Planar
$P_3$	$Q_2$	Non-Polar	Non-Planar

Then according to given information calculate value of expression

$$P_2 \times \left( \frac{P_3 - P_1}{(Q_1 + Q_2)} \right)$$

 [View Text Solution](#)

32. What is covalency of chlorine atom in second excited state ?

 [Watch Video Solution](#)

33. Sum of  $\sigma$  and  $\pi$  bonds in  $NH_4^+$  cation is ..

 [Watch Video Solution](#)

34. Calculate the value of X-Y, for  $XeOF_4$ . (X=Number of  $\sigma$  bond pair and Y=Number of lone pair on central atom )

 [Watch Video Solution](#)

35. The molecule  $ML_x$  is planar with 6 electron pairs around M in the valence shell. The value of x is :

 [Watch Video Solution](#)

36. Calculate value of  $\frac{X + Y + Z}{10}$ , here X is O-N-O bond angle in  $NO_3^-$  Y is O-N-O bond angle in  $NO_2^+$  and Z is F-Xe-F adjacent bond angle in  $XeF_4$ .

 [Watch Video Solution](#)

37. Calculate  $x+y+z$  for  $H_3PO_3$  acid, where  $x$  is no. of lone pairs,  $y$  is no. of  $\sigma$  bonds and  $z$  is no. of  $\pi$  bonds.

 [Watch Video Solution](#)

38. How many right angle, bond angles are present in  $TeF_5^-$  molecular ion ?

 [Watch Video Solution](#)

39. How many possible  $\angle FSeF$  bond angles are present in  $SeF_4$  molecule ?

 [Watch Video Solution](#)



40. In  $IF_6^-$  and  $TeF_5^-$ , sum of axial d-orbitals which are used in hybridisation in both species.





Watch Video Solution

41. Among the following, total no. of planar species is :



Watch Video Solution

42. Calculate the value of "  $x+y-z$ " here  $x,y$  and  $z$  are total number of non-bonded electron pair (s), pie ( $\pi$ ) bond(s) and sigma ( $\sigma$ ) bonds in hydrogen phosphite ion respectively.



Watch Video Solution

43. Consider the following table

Total number of electron pairs (l.p. + $\sigma$ -bond)	Total number of lone pairs	Shape
5	...p...	linear
...q...	1	see-saw
4	...r...	Bent shape
...s...	2	Square planar
5	...t...	Bent 'T' shape

Then calculate value of "p+q+r-s-t".

 [Watch Video Solution](#)

44. In phosphorus acid, if X is number of non bonding electron pairs. Y is number of  $\sigma$ -bonds and Z is number of  $\pi$ - bonds. Then, calculate value of  $Y \times Z - X$ .

 [Watch Video Solution](#)

45. Calculate the number of  $p_{\pi} - d_{\pi}$  bond(s) present in  $SO_4^{2-}$  :

 [Watch Video Solution](#)



46. Sum of  $\sigma$  and  $\pi$  bonds in  $NH_4^+$  cation is ..

 [Watch Video Solution](#)

47. Consider the following orbitals (i)  $3p_x$  (ii)  $4d_{z^2}$  (iii)  $3d_{x^2-y^2}$  (iv)  $3d_{yz}$



Then, calculate value of " $x + y - z$ " here x is total number of gerade orbital and y is total number of ungerade orbitals and z is total number of axial orbitals in given above orbitals.

 [Watch Video Solution](#)

48. Calculate value of  $|x-y|$ , here x and y are the total number of bonds in benzene and benzyne respectively which are formed by overlapping of hybridized orbitals.

 [Watch Video Solution](#)


49. Consider the following compounds :



Then calculate value of "x+y-z", here, x,y and z are total number of compounds in given compounds in which central atom used their all three p-orbitals, only two p-orbitals and only one p-orbital in hybridisation respectively .


 [Watch Video Solution](#)

50. Total number of species which used all three p-orbitals in hybridisation of central atom and should be non-polar also.



 [Watch Video Solution](#)

51. Consider the following species



Then calculate value of  $|x-y|$ , where

x : Total number of species which have bond order 1.5 or greater than 1.5

y : Total number of species which have bond order less than 1.5

 [Watch Video Solution](#)


52. Consider the following orbitals

$3s, 2p_x, 4d_{xy}, 4d_{z^2}, 3d_{x^2-y^2}, 3p_y, 4s, 4p_z$  and find total number of orbital

(s) having even number of nodal plane.

 [Watch Video Solution](#)

53. For the following molecules :



Calculate the value of  $\frac{a+b}{c}$

a = Number of species having  $sp^3$  d-hybridisation




b= Number of species which are planar

c= Number of species which are non-planar

 [Watch Video Solution](#)

54. Find out number of transformation among following which involves the change of hybridisation of underlined atom.

(a)



 [Watch Video Solution](#)