



MATHS

BOOKS - VK JAISWAL ENGLISH

APPLICATION OF DERIVATIVES

Exercise Single Choice Problems

1. The difference between the maximum and minimum value of the function $f(x) = 3 \sin^4 x - \cos^6 x$ is :

A. $\frac{3}{2}$

B. $\frac{5}{2}$

C. 3

D. 4

Answer: D



Watch Video Solution

2. A function $y = f(x)$ has a second-order derivative $f''(x) = 6(x - 1)$. It its graph passes through the point (2,1) and at that point tangent to the graph is $y = 3x - 5$, then the value of $f(0)$ is 1 (b) -1 (c) 2 (d) 0

A. $(x - 1)^2$

B. $(x - 1)^3$

C. $(x + 1)^3$

D. $(x + 1)^2$

Answer: B



Watch Video Solution

3. If the subnormal at any point on the curve $y = 3^{1-k} \cdot x^k$ is of constant length the k equals to :

A. $\frac{1}{2}$

B. 1

C. 2

D. 0

Answer: A



Watch Video Solution

4. If $x^5 - 5qx + 4r$ is divisible by $(x - c)^2$ then which of the following must hold true

a) $q=r$ b) $q+r=0$ c) $q^5 + r=0$ d) $q^4 = r^5$

A. $q = r$

B. $q + r = 0$

C. $q^5 + r = 0$

D. $q^5 = r^4$

Answer: C



Watch Video Solution

5. A spherical iron ball 10cm in radius is coated with a layer of ice of uniform thickness that melts at a rate of $50\text{cm}^3/\text{min}$. When the thickness of ice is 5cm, then find the rate at which the thickness of ice decreases.

A. $\frac{1}{36\pi}\text{cm}/\text{min}$

B. $\frac{1}{18\pi}\text{cm}/\text{min}$

C. $\frac{1}{54\pi}\text{cm}/\text{min}$

D. $\frac{5}{6\pi}\text{cm}/\text{min}$

Answer: B



Watch Video Solution

6. If $f(x) = \frac{(x-1)(x-2)}{(x-3)(x-4)}$, then number of local extremas for $g(x)$,

where $g(x) = f(|x|)$: (a) 3 (b) 4 (c) 5 (d) none of these

A. 3

B. 4

C. 5

D. None of these

Answer: C



[Watch Video Solution](#)

7. Two straight roads OA and OB intersect at an angle 60° . A car approaches O from A, where $OA = 700m$ at a uniform of 20 m/s, Simultaneously, a runner starts running from O towards B at a uniform speed of 5 m/s. The time after start when the car and the runner are closest is:

A. 10 sec

B. 15 sec

C. 20 sec

D. 30 sec

Answer: D



Watch Video Solution

8. Let $f(x) = \begin{cases} a - 3x & -2 \leq x < 0 \\ 4x \pm 3 & 0 \leq x < 1 \end{cases}$, if $f(x)$ has smallest value at $x = 0$, then range of a , is

A. $(-\infty, 3)$

B. $(-\infty, 3]$

C. $(3, \infty)$

D. $[3, \infty)$

Answer: D



Watch Video Solution

9. If $f(x) = \begin{cases} 3 + |x - k| & x \leq k \\ a^2 - 2 + \frac{\tan(x-k)}{x-k} & x > k \end{cases}$ has minimum at $x=k$, then

(a) $|a| \leq 2$ (b) $|a| < 2$ (c) $|a| > 2$ (d) $|a| \geq 2$

A. $a \in R$

B. $|a| < 2$

C. $|a| > 2$

D. $1 < |a| < 2$

Answer: C



Watch Video Solution

10. For certain curve $y = f(x)$ satisfying $\frac{d^2y}{dx^2} = 6x - 4$, $f(x)$ has local minimum value 5 when $x = 1$

Global maximum value of $y = f(x)$ for $x \in [0, 2]$ is

A. -2

B. 2

C. 12

D. -12

Answer: B



Watch Video Solution

11. The tangent to $y = ax^2 + bx + \frac{7}{2}at(1, 2)$ is parallel to the normal at the point $(-2, 2)$ on the curve $y = x^2 + 6x + 10$. Then the value of $\frac{a}{2} - b$ is:

A. 2

B. 0

C. 3

D. 1

Answer: C



Watch Video Solution

12. If (a, b) be the point on the curve $9y^2 = x^3$ where normal to the curve make equal intercepts with the axis, then the value of $(a + b)$ is:

A. 0

B. $\frac{10}{3}$

C. $\frac{20}{3}$

D. None of these

Answer: C



Watch Video Solution

13. The curve $y = f(x)$ satisfies $\frac{d^2y}{dx^2} = 6x - 4$ and $f(x)$ has a local minimum value 5 when $x = 1$. Then $f'(0)$ is equal to :

A. 1

B. 0

C. 5

D. None of these

Answer: C



Watch Video Solution

14. Let A be the point where the curve $5\alpha^2x^3 + 10\alpha x^2 + x + 2y - 4 = 0$ ($\alpha \in R, \alpha \neq 0$) meets the y-axis, then the equation of tangent to the curve at the point where normal at A meets the curve again, is:

A. $x - \alpha y + 2\alpha = 0$

B. $\alpha x + y - 2 = 0$

C. $2x - y + 2 = 0$

D. $x + 2y - 4 = 0$

Answer: C



Watch Video Solution

15. The difference between the greatest and least value of the functions,

$$f(x) = \cos x + \frac{1}{2}\cos 2x - \frac{1}{3}\cos 3x \text{ is}$$

A. $\frac{11}{5}$

B. $\frac{13}{6}$

C. $\frac{9}{4}$

D. $\frac{7}{3}$

Answer: C



Watch Video Solution

16. The x co-ordinate of the point on the curve $y = \sqrt{x}$ which is closest to the point (2, 1) is :

A. $\frac{2 + \sqrt{3}}{2}$

B. $\frac{1 + \sqrt{2}}{2}$

C. $\frac{-1 + \sqrt{3}}{2}$

D. 1

Answer: A



Watch Video Solution

17. The tangent at a point P on the curve

$$y = \ln\left(\frac{2 + \sqrt{4 - x^2}}{2 - \sqrt{4 - x^2}}\right) - \sqrt{4 - x^2}$$

meets the y-axis at T, then PT^2

equals to :

A. 2

B. 4

C. 8

D. 16

Answer: B



Watch Video Solution

18. Let $f(x) = \int_{x^2}^{x^3} \frac{dt}{\ln t}$ for $x > 1$ and $g(x) = \int_1^x (2t^2 - \ln t) f(t) dt (x > 1)$, then: (a) g is increasing on $(1, \infty)$ (b) g is decreasing on $(1, \infty)$ (c) g is increasing on $(1, 2)$ and decreasing on $(2, \infty)$ (d) g is decreasing on $(1, 2)$ and increasing on $(2, \infty)$

A. g is increasing on $(1, \infty)$

B. g is decreasing on $(1, \infty)$

C. g is increasing on $(1, 2)$ and decreasing on $(2, \infty)$

D. g is decreasing on $(1, 2)$ and increasing on $(2, \infty)$

Answer: A



Watch Video Solution

19. Let $f(x) = x^3 + 6x^2 + ax + 2$, if $(-3, -1)$ is the largest possible interval for which $f(x)$ is decreasing function, then $a =$

A. 3

B. 9

C. -2

D. 1

Answer: B



[Watch Video Solution](#)

20. Let $f(x) = \tan^{-1}\left(\frac{1-x}{1+x}\right)$. Then difference of the greatest and least value of $f(x)$ on $[0, 1]$ is:

A. $\pi/2$

B. $\pi/4$

C. π

D. $\pi/3$

Answer: B



[Watch Video Solution](#)

21. The number of integral values of a for which

$f(x) = x^3 + (a + 2)x^2 + 3ax + 5$ is monotonic in $\forall x \in R$.

A. 2

B. 4

C. 6

D. 7

Answer: B



[Watch Video Solution](#)

22. The number of critical points of

$$f(x) = \left(\int_0^x (\cos^2 t - \sqrt[3]{t}) dt \right) + \frac{3}{4}x^{4/3} - \frac{x+1}{2} \text{ in } (0, 6\pi] \text{ is:}$$

A. 10

B. 8

C. 6

D. 12

Answer: D



[Watch Video Solution](#)

23. Let $f(x) = \min \left(\frac{1}{2} - \frac{3x^2}{4}, \frac{5x^2}{4} \right)$ for $0 \leq x \leq 1$, then maximum

value of $f(x)$ is:

A. 0

B. $\frac{5}{64}$

C. $\frac{5}{4}$

D. $\frac{5}{16}$

Answer: D



Watch Video Solution

24. Let $f(x) = \begin{cases} 2 - |x^2 + 5x + 6| & x \neq -2 \\ b^2 + 1 & x = -2 \end{cases}$

Has relative maximum at $x = -2$, then complete set of values b can take is:

A. $|b| \geq 1$

B. $|b| < 1$

C. $b > 1$

D. $b < 1$

Answer: A



Watch Video Solution

25. Let for function $f(x) = \begin{cases} \cos^{-1} x & -1 \leq x \leq 0 \\ mx + c & 0 < x \leq 1 \end{cases}$, Lagrange's mean value theorem is applicable in $[-1, 1]$ then ordered pair (m, c) is:

A. $\left(1, -\frac{\pi}{2}\right)$

B. $\left(1, \frac{\pi}{2}\right)$

C. $\left(-1, -\frac{\pi}{2}\right)$

D. $\left(-1, \frac{\pi}{2}\right)$

Answer: D



Watch Video Solution

26. Tangents are drawn to $y = \cos x$ from origin then points of contact for these tangents will always lie on :

A. $\frac{1}{x^2} = \frac{1}{y^2} + 1$

B. $\frac{1}{x^2} = \frac{1}{y^2} - 2$

C. $\frac{1}{y^2} = \frac{1}{x^2} + 1$

$$D. \frac{1}{y^2} = \frac{1}{x^2} - 2$$

Answer: C



Watch Video Solution

27. Least natural number a for which

$$x + ax^{-2} > 2, \forall x \in (0, \infty) \text{ is}$$

A. 1

B. 2

C. 5

D. None of these

Answer: B



Watch Video Solution

28. Angle between the tangents to the curve $y = x^2 - 5x + 6$ at points $(2, 0)$, and $(3, 0)$ is:

A. $\frac{\pi}{6}$

B. $\frac{\pi}{4}$

C. $\frac{\pi}{3}$

D. $\frac{\pi}{2}$

Answer: D



[Watch Video Solution](#)

29. Difference between the greatest and least values of the function

$f(x) = \int_0^x (\cos^2 t + \cos t + 2) dt$ in the interval $[0, 2\pi]$ is $K\pi$, then K is

equal to:

A. 1

B. 3

C. 5

D. None of these

Answer: C



[Watch Video Solution](#)

30. The range of the function $f(\theta) = \frac{\sin \theta}{\theta} + \frac{\theta}{\tan \theta}$, $\theta \in \left(0, \frac{\pi}{2}\right)$ is equal to :

A. $(0, \infty)$

B. $\left(\frac{1}{\pi}, 2\right)$

C. $(2, \infty)$

D. $\left(\frac{2}{\pi}, 2\right)$

Answer: D



[Watch Video Solution](#)

31. Number of integers in the range of c so that the equation $x^3 - 3x + c = 0$ has all its roots real and distinct is:

A. 2

B. 3

C. 4

D. 5

Answer: B



[Watch Video Solution](#)

32. Let $f(x) = \int e^x (x - 1)(x - 2) dx$. Then f decreases in the interval $(-\infty, -2)$ (b) $(-2, -1)$ (c) $(1, 2)$ (d) $(2, +\infty)$

A. $(2, \infty)$

B. $(-2, -1)$

C. $(1, 2)$

D. $(-\infty, 1) \cup (2, \infty)$

Answer: C



[Watch Video Solution](#)

33. If the cubic polynomial $y = ax^3 + bx^2 + cx + d$ ($a, b, c, d \in R$) has only one critical point in its entire domain and $ac = 2$, then the value of $|b|$ is:

A. $\sqrt{2}$

B. $\sqrt{3}$

C. $\sqrt{5}$

D. $\sqrt{6}$

Answer: D



[Watch Video Solution](#)

34. On the curve $y = \frac{1}{1+x^2}$, the point at which $\left| \frac{dy}{dx} \right|$ is greatest in the first quadrant is :

- A. $\left(\frac{1}{2}, \frac{4}{5} \right)$
- B. $\left(1, \frac{1}{4} \right)$
- C. $\left(\frac{1}{\sqrt{2}}, \frac{2}{3} \right)$
- D. $\left(\frac{1}{\sqrt{3}}, \frac{3}{4} \right)$

Answer: D

 [Watch Video Solution](#)

35. If $f(x) = 2x$, $g(x) = 3 \sin x - x \cos x$, then for $x \in \left(0, \frac{\pi}{2} \right)$:

- A. $f(x) > g(x)$
- B. $f(x) < g(x)$
- C. $f(x) = g(x)$ has exactly one real root.
- D. $f(x) = g(x)$ has exactly two real roots

Answer: A



Watch Video Solution

36. let $f(x) = \sin^{-1}\left(\frac{2g(x)}{1+g(x)^2}\right)$, then which are correct ?

(i) $f(x)$ is decreasing if $g(x)$ is increasing and $|g(x)| > 1$

(ii) $f(x)$ is an increasing function if $g(x)$ is increasing and $|g(x)| \leq 1$

(iii) $f(x)$ is decreasing function if $f(x)$ is decreasing and $|g(x)| > 1$

A. (i) and (iii)

B. (i) and (ii)

C. (i) (ii) and (iii)

D. (iii)

Answer: B



Watch Video Solution

37. The graph of function $y = f(x)$ has a unique tangent at $(e^a, 0)$ through which the graph passes, then

$$\lim_{x \rightarrow e^a} \frac{\log(1 + 7f(x)) - \sin(f(x))}{3f(x)} \text{ equals}$$

- A. 1
- B. 3
- C. 2
- D. 7

Answer: C



[Watch Video Solution](#)

38. Let $f(x)$ be a function such that $f'(x) = \log_{1/3}(\log_3(\sin x + a))$.

The complete set of values of 'a' for which $f(x)$ is strictly decreasing for all real values of x is:

- A. $[4, \infty)$

B. $[3, 4]$

C. $(-\infty, 4)$

D. $[2, \infty)$

Answer: A



Watch Video Solution

39. If $f(x) = a \ln|x| + bx^2 + x$ has extremas at $x = 1$ and $x = 3$ then:

A. $a = \frac{3}{4}, b = -\frac{1}{8}$

B. $a = \frac{3}{4}, b = \frac{1}{8}$

C. $a = -\frac{3}{4}, b = -\frac{1}{8}$

D. $a = -\frac{3}{4}, b = \frac{1}{8}$

Answer: C



Watch Video Solution

40. Let $f(x) = \begin{cases} 1 + \sin x, & x < 0 \\ x^2 - x + 1, & x \geq 0 \end{cases}$, then:

- A. f has a local maximum at $x = 0$
- B. f has a local minimum at $x = 0$
- C. f is increasing everywhere
- D. f is decreasing everywhere

Answer: A

 [Watch Video Solution](#)

41. If m and n are positive integers and

$$f(x) = \int_1^x (t - a)^{2n} (t - a)^{2m+1} dt, \quad a \neq b, \text{ then}$$

- A. $x = b$ is a point of local minimum
- B. $x = b$ is a point of local maximum
- C. $x = a$ is a point of local minimum
- D. $x = a$ is a point of local maximum

Answer: A



[Watch Video Solution](#)

42. For any *real* θ , the maximum value of $\cos^2(\cos \theta) + \sin^2(\sin \theta)$ is

A. 1

B. $1 + \sin^2 1$

C. $1 + \cos^2 1$

D. Does not exist

Answer: B



[Watch Video Solution](#)

43. If the tangent at P of the curve $y^2 = x^3$ intersects the curve again at Q and the straight line OP, OQ have inclinations α, β where O is origin, then $\left(\frac{\tan \alpha}{\tan \beta} \right)$ has the value, equals to:

A. -1

B. -2

C. 2

D. $\sqrt{2}$

Answer: B



Watch Video Solution

44. If $x + 4y = 14$ is a normal to the curve $y^2 = \alpha x^3 - \beta$ at $(2,3)$, then the value of $\alpha + \beta$ is 9 (b) -5 (c) 7 (d) -7

A. 9

B. -5

C. 7

D. -7

Answer: A


 Watch Video Solution

45. The tangent to the curve $y = e^{kx}$ at a point $(0,1)$ meets the x-axis at $(a,0)$, where $a \in [-2, -1]$. Then $k \in$ (a) $\left[-\frac{1}{2}, 0\right]$ (b) $\left[-1, -\frac{1}{2}\right]$ (c) $[0, 1]$ (d) $\left[\frac{1}{2}, 1\right]$
- A. $\left[-\frac{1}{2}, 0\right]$
- B. $\left[-1 - \frac{1}{2}\right]$
- C. $[0, 1]$
- D. $\left[\frac{1}{2}, 1\right]$


Answer: D

 Watch Video Solution


46. Which of the following graph represent the function $f(x) = \int_0^{\sqrt{x}} e^{\frac{u^2}{x}}$ du, for $x > 0$ and $f(0) = 0$




A.



B.



C.



D.

Answer: B



Watch Video Solution

47. Let $f(x) = (x - a)(x - b)(x - c)$ be a rational function where $a < bc$ ($a, b, c \in \mathbb{R}$) such that $f''(\alpha) = 0$. Then if $\alpha \in (c_1, c_2)$, which one of the following is correct ?

A. $\alpha < c_1 < b$ and $b < c_2 < c$

B. $\alpha < c_1, c_2 < b$

C. $b < c_1, c_2 < c$

D. None of these

Answer: A



[Watch Video Solution](#)

48. $f(x) = x^6 - x - 1, x \in [1, 2]$. Consider the following statements :

A. f is increasing on $[1, 2]$

B. f has a root in $[1, 2]$

C. f is decreasing on $[1, 2]$

D. f has no root in $[1, 2]$

Answer: A



[Watch Video Solution](#)

49. Which one of the following curves is the orthogonal trajectory of straight lines passing through a fixed point (a,b) ?

A. $x - a = k(y - b)$

B. $(x - a)(y - b) = k$

C. $(x - a)^2 = k(y - b)$

D. $(x - a)^2 + (y - b)^2 = k$

Answer: D



[Watch Video Solution](#)

50. The function $f(x) = \sin^3 x - m \sin x$ is defined on open interval $\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$ and if assumes only 1 maximum value and only 1 minimum value on this interval. Then, which one of the must be correct? (a) $0 < m < 3$ (b) $-3 < m < 0$ (c) $m > 3$ (d) $m < -3$

A. $0 < m < 3$

B. $-3 < m < 0$

C. $m > 3$

D. $m < -3$

Answer: A



Watch Video Solution

51. The greatest of the numbers $1, 2^{1/2}, 4^{1/4}, 5^{1/5}, 6^{1/6},$ and $7^{1/7}$ is:

A. $2^{1/2}$

B. $3^{1/3}$

C. $7^{1/7}$

D. $6^{1/6}$

Answer: B



[Watch Video Solution](#)

52. Let l be the line through $(0, 0)$ and tangent to the curve $y = x^3 + x + 16$. Then the slope of l equal to :

A. 10

B. 11

C. 17

D. 13

Answer: D



[Watch Video Solution](#)

53. The slope of the tangent at the point of inflection of $y = x^3 - 3x^2 + 6x + 2009$ is equal to :

A. 2

B. 3

C. 1

D. 4

Answer: B



[Watch Video Solution](#)

54. If $f(x)$ is a differentiable real valued function satisfying $f''(x) - 3f'(x) > 3 \forall x \geq 0$ and $f'(0) = -1$, then $f(x) + x \forall x > 0$ is

A. strictly increasing

B. strictly decreasing

C. non monotonic

D. data insufficient

Answer: A



[Watch Video Solution](#)

55. If the line joining the points $(0, 3)$ and $(5, -2)$ is a tangent to the curve $y = \frac{C}{x+1}$, then the value of C is (a) 1 (b) -2 (c) 4 (d) none of these

A. 2

B. 3

C. 4

D. 5

Answer: C



[Watch Video Solution](#)

56. Find the number of solutions to $\log_e |\sin x| = -x^2 + 2x$ in $\left[-\frac{\pi}{2}, \frac{3\pi}{2}\right]$.

A. 2

B. 4

C. 6

D. 8

Answer: B



[Watch Video Solution](#)

57. Find the values of a for which $\sin^{-1}(-1)x = |x - a|$ will have at least one solution.

A. $[-1, 1]$

B. $\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$

C. $\left[1 - \frac{\pi}{2}, 1 + \frac{\pi}{2}\right]$

D. $\left[\frac{\pi}{2} - 1, \frac{\pi}{2} + 1\right]$

Answer: C



Watch Video Solution

58. For any real number b , let $f(b)$ denotes the maximum of $\left|\sin x + \frac{2}{3 + \sin x} + b\right| \forall x \in R$. Then the minimum value of $f(b) \forall b \in R$ is:

A. $\frac{1}{2}$

B. $\frac{3}{4}$

C. $\frac{1}{4}$

D. 1

Answer: B



Watch Video Solution

59. Which of the following are correct

A. $x^4 + 2x^2 - 6x + 2 = 0$ has exactly four real solution

B. $x^3 + 5x + 1 = 0$ has exactly three real solutions

C. $x^n + ax + b = 0$ where n is an even natural number has atmost two real solution a, b , in \mathbb{R} .

D. $x^3 - 3x + c = 0, x > 0$ has two real solutin for $x \in (0, 1)$

Answer: C



[Watch Video Solution](#)

60. For any real number b , let $f(b)$ denotes the maximum of

$\left| \sin x + \frac{2}{3 + \sin x} + b \right| \forall x \in \mathbb{R}$. Then the minimum value of

$f(b) \forall b \in \mathbb{R}$ is:

A. $\frac{1}{2}$

B. $\frac{3}{4}$

C. $\frac{1}{4}$

D. 1

Answer: B



Watch Video Solution

61. Find the coordinates of the point on the curve $y = \frac{x}{1+x^2}$ where the tangent to the curve has the greatest slope.

A. (0, 0)

B. $\left(\sqrt{3}, \frac{\sqrt{3}}{4}\right)$

C. $\left(-\sqrt{3}, -\frac{\sqrt{3}}{4}\right)$

D. $\left(1, \frac{1}{2}\right)$

Answer: A



Watch Video Solution

62. Let $f: [0, 2\pi] \rightarrow [-3, 3]$ be a given function defined at $f(x) = 3 \cos \frac{x}{2}$. The slope of the tangent to the curve $y = f^{-1}(x)$ at the point where the curve crosses the y-axis is:

A. -1

B. $-\frac{2}{3}$

C. $-\frac{1}{6}$

D. $-\frac{1}{3}$

Answer: B



Watch Video Solution

63. Number of stationary points in $[0, \pi]$ for the function $f(x) = \sin x + \tan x - 2x$ is:

A. 0

B. 1

C. 2

D. 3

Answer: C



[Watch Video Solution](#)

64. If $a, b, c, d \in R$ such that $\frac{a + 2c}{b + 3d} + \frac{4}{3} = 0$, then the equation $ax^3 + bx^3 + cx + d = 0$ has

A. atleast one root in $(-1, 0)$

B. atleast one root in $(0, 1)$

C. no root in $(-1, 1)$

D. no root in $(0, 2)$

Answer: B



[Watch Video Solution](#)

65. If $f'(x)\phi(x)(x-2)^2$. Were $\phi(2) \neq 0$ and $\phi(x)$ is continuous at $x = 2$ then in the neighbourhood of $x = 2$

- A. f is increasing if $\phi(2) < 0$
- B. f is decreasing if $\phi(2) > 0$
- C. f is neither increasing nor decreasing
- D. f is increasing if $\phi(2) > 0$

Answer: D



Watch Video Solution

66. If the function $f(x) = x^3 - 6x^2 + ax + b$ defined on $[1,3]$ satisfies

Rolle's theorem for $c = \frac{2\sqrt{3} + 1}{\sqrt{3}}$ then find the value of a and b

- A. $a = -11, b = 5$
- B. $a = -11, b = -6$

$$C. a = 11, b \in R$$

$$D. 1 = 22, b = -6$$

Answer: C

 [Watch Video Solution](#)

67. For which of the following function 9s) Lagrange's mean value theorem is not applicable in $[1, 2]$?

$$A. f(x) = \begin{cases} \frac{3}{2} - x, & x < \frac{3}{2} \\ \left(\frac{3}{2} - x\right)^2, & x \geq \frac{3}{2} \end{cases}$$

$$B. f(x) = \begin{cases} \frac{\sin(x-1)}{x-1}, & x \neq 1 \\ 1, & x = 1 \end{cases}$$

$$C. f(x) = (x - 1)|x + 1|$$

$$D. f(x) = |x - 1|$$

Answer: A

 [Watch Video Solution](#)

68. If the curves $\frac{x^2}{a^2} + \frac{y^2}{4} = 1$ and $y^3 = 16x$ intersect at right angles, then:

A. $a = \pm 1$

B. $a = \pm \sqrt{3}$

C. $a = \pm \sqrt{3}$

D. $a = \pm \sqrt{2}$

Answer: D



Watch Video Solution

69. If the line $x \cos \alpha + y \sin \alpha = P$ touches the curve $4x^3 = 27ay^2$, then

$$\frac{P}{a} =$$

A. $\cot^2 \alpha \cos \alpha$

B. $\cot^2 \alpha \sin \alpha$

C. $\tan^2 \alpha \cos \alpha$

D. $\tan^2 \alpha \sin \alpha$

Answer: A



Watch Video Solution

Exercise One Or More Than Answer Is Are Correct

1. Common tangent (s) to $y = x^3$ and $x = y^3$ is/are

A. $x - y = \frac{1}{\sqrt{3}}$

B. $x - y = -\frac{1}{\sqrt{3}}$

C. $x - y = \frac{2}{3\sqrt{3}}$

D. $x - y = \frac{-2}{3\sqrt{3}}$

Answer: C::D



Watch Video Solution

2. Let $f: [0, 8] \rightarrow \mathbb{R}$ be differentiable function such that $f(0) = 0, f(4) = 1, f(8) = 1$, then which of the following hold(s) good ?

A. There exist some $c_1 \in (0, 8)$ where $f(c_1) = \frac{1}{4}$

B. There exist some $x \in (0, 8)$ where $f'(c) = \frac{1}{12}$

C. There exist $c_1, c_2 \in [0, 8]$ where $8f'(c_1)f(c_2) = 1$

D. There exist some $\alpha, \beta \in (0, 2)$ such that

$$\int_0^8 f(t)dt = 3(\alpha^2 f(\alpha^3) + \beta^2 f(\beta^3))$$

Answer: A::C::D



Watch Video Solution

3. If $f(x) = \begin{cases} \sin^{-1}(\sin x) & x > 0 \\ \frac{\pi}{2} & x = 0 \\ \cos^{-1}(\cos x) & x < 0 \end{cases}$, then

A. $x = 0$ is a point of maxima

B. $f(x)$ is continuous $\forall x \in \mathbb{R}$

C. global maximum value of $f(x) \forall x \in \mathbb{R}$ is π

D. global minimum value of $f(x) \forall x \in \mathbb{R}$ is 0

Answer: A::C::D



[Watch Video Solution](#)

4. A function $f: \mathbb{R} \rightarrow \mathbb{R}$ is given by $f(x) = \begin{cases} x^4 \left(2 + \sin \frac{1}{x} \right) & x \neq 0 \\ 0 & x = 0 \end{cases}$,

then

A. f has a continuous derivative $\forall x \in \mathbb{R}$

B. f is a bounded function

C. f has a global minimum at $x = 0$

D. f'' is continuous $\forall x \in \mathbb{R}$

Answer: A::C::D



[Watch Video Solution](#)

5. If $f''(x) \leq 1 \forall x \in R$, and $f(0) = 0 = f'(0)$, then which of the following can not be true ?

A. $f\left(-\frac{1}{2}\right) = \frac{1}{6}$

B. $f(2) = -4$

C. $f(-2) = 3$

D. $f\left(\frac{1}{2}\right) = \frac{1}{5}$

Answer: A::B::C::D



Watch Video Solution

6. Let $f: [-3, 4] \rightarrow R$ such that $f''(x) > 0$ for all $x \in [-3, 4]$, then which of the following are always true ?

A. $f(x)$ has a relative minimum on $(-3, 4)$

B. $f(x)$ has a minimum on $[3, 4]$

C. $f(x)$ has a maximum on $[-3, 4]$

D. if $f(3) = f(4)$, then $f(x)$ has a critical point on $[-3, 4]$

Answer: B::C::D



Watch Video Solution

7. Let $f(x)$ be twice differentiable function such that $f''(x) > 0$ in $[0, 2]$.

Then :

A. $f(0) + f(2) = 2f(x)$, for atleast one $c, c \in (0, 2)$

B. $f(0) + f(2) < 2f(1)$

C. $f(0) + f(2) > 2f(1)$

D. $2f(0) + f(2) > 3f\left(\frac{2}{3}\right)$

Answer: C::D



Watch Video Solution

8. Let $g(x)$ be a cubic polynomial having local maximum at $x = -1$ and $g'(x)$ has a local minimum at $x = 1$, If $g(-1) = 10$, $g(3) = -22$, then

- A. perpendicular distance between its two horizontal tangents is 12
- B. perpendicular distance between its two horizontal tangents is 32
- C. $g(x) = 0$ has at least one real root lying in interval $(-2, 0)$
- D. $g(x) = 0$, has 3 distinct real roots

Answer: B::D



[Watch Video Solution](#)

9. Let S be the set of real values of parameter λ for which the equation $f(x) = 2x^3 - 3(2 + \lambda)x^2 + 12\lambda x$ has exactly one local maximum and exactly one local minimum.

Then, S is a subset of

- A. $\lambda \in (-4, \infty)$

B. $\lambda \in (-\infty, 0)$

C. $\lambda \in (-3, 3)$

D. $\lambda \in (1, \infty)$

Answer: A::B::C::D



Watch Video Solution

10. The function $f(x) = 1 + x \ln(x + \sqrt{1 + x^2}) - \sqrt{1 - x^2}$ is:

A. strictly increasing $\forall x \in (0, 1)$

B. strictly decreasing $\forall x \in (-1, 0)$

C. strictly decreasing for $x \in (-1, 0)$

D. strictly decreasing for $x \in (0, 1)$

Answer: A::C::D



Watch Video Solution

11. Let m and n be positive integers and $x, y > 0$ and $x + y = k$, where k is constant. Let $f(x, y) = x^m y^n$, then: (a) $f(x, y)$ is maximum when $x = \frac{mk}{m+n}$ (b) $f(x, y)$ is maximum where $x = y$ (c) maximum value of $f(x, y)$ is $\frac{m^n n^m k^{m+n}}{(m+n)^{m+n}}$ (d) maximum value of $f(x, y)$ is $\frac{k^{m+n} m^m n^n}{(m+n)^{m+n}}$

A. $f(x, y)$ is maximum when $x = \frac{mk}{m+n}$

B. $f(x, y)$ is maximum where $x = y$

C. maximum value of $f(x, y)$ is $\frac{m^n n^m k^{m+n}}{(m+n)^{m+n}}$

D. maximum value of $f(x, y)$ is $\frac{k^{m+n} m^m n^n}{(m+n)^{m+n}}$

Answer: A:D



Watch Video Solution

12. The straight line which is both tangent and normal to the curve

$$x = 3t^2, y = 2t^3 \text{ is:}$$

A. $y + \sqrt{3}(x - 1) = 0$

$$B. y - \sqrt{3}(x - 1) = 0$$

$$C. y + \sqrt{2}(x - 2) = 0$$

$$D. y - \sqrt{2}(x - 2) = 0$$

Answer: C::D



Watch Video Solution

13. A curve is such that the ratio of the subnormal at any point to the sum of its co-ordinates is equal to the ratio of the ordinate of this point to its abscissa. If the curve passes through $(1, 0)$, then possible equation of the curve (s) is:

$$A. y = x \ln x$$

$$B. y = \frac{\ln x}{x}$$

$$C. y = \frac{2(x - 1)}{x^2}$$

$$D. y = \frac{1 - x^2}{2x}$$

Answer: A::D



Watch Video Solution

14. A parabola of the form $y = ax^2 + bx + c$ ($a > 0$) intersects the graph of $f(x) = \frac{1}{x^2 - 4}$. The number of possible distinct intersection (s) of these graphs can be:

A. 0

B. 2

C. 3

D. 4

Answer: B::C::D



Watch Video Solution

15. Gradient of the line passing through the point $(2, 8)$ and touching the curve $y = x^3$, can be:

A. 3

B. 6

C. 9

D. 12

Answer: A::D



[Watch Video Solution](#)

16. The equation $x + \cos x = a$ has exactly one positive root, then:

A. $a \in (0, 1)$

B. $a \in (2, 3)$

C. $a \in (1, \infty)$

D. $a \in (-\infty, 1)$

Answer: B::C



Watch Video Solution

17. Given that $f(x)$ is a non-constant linear function. Then the curves :

A. $y = f(x)$ and $y = f^{-1}(x)$ are orthogonal

B. $y = f(x)$ and $y = f^{-1}(-x)$ are orthogonal

C. $y = f(-x)$ and $y = f^{-1}(x)$ are orthogonal

D. $y = f(-x)$ and $y = f^{-1}(-x)$ are orthogonal

Answer: B::C



Watch Video Solution

18. $d(x) = \int_0^x e^{t^3} (t^2 - 1)(t + 1)^{2011} dt$ ($x > 0$) then :

A. The number of point of inflections is atleast 1

B. The number of point of inflectins is 0

C. The number of point of local maxima is 1

D. The number of point of local minima is 1

Answer: A::D



Watch Video Solution

19. Let $f(x) = \sin x + ax + b$. Then which of the following is/are true?

(a) $f(x) = 0$ has only one real root which is positive if $a > 1, b < 0$. (b)

$f(x) = 0$ has only one real root which is negative if $a > 1, b < 0$. (c)

$f(x) = 0$ has only one real root which is negative if $a > 1, b > 0$. (d)

none of these

A. only one real root which is positive if $a > 1, b < 0$

B. only one real root which is negative if $a > 1, b > 0$


C. only one real root which is negative if $a < -1, b < 0$

D. only one real root which is positive if $a < -1, b < 0$


Answer: A::B::C

 Watch Video Solution


20. Which of the following graphs represent function whose derivatives have a maximum in the interval $(0, 1)$?




A.



B.



C.



Answer: A::B::D

[▶ Watch Video Solution](#)

21. Consider $f(x) = \sin^5 x - 1$, $x \in \left[0, \frac{\pi}{2}\right]$, which of the following is/are correct ?

- A. f is strictly decreasing in $\left[0, \frac{\pi}{4}\right]$
- B. f is strictly increasing in $\left[\frac{\pi}{4}, \frac{\pi}{2}\right]$
- C. There exist a number 'c' in $\left(0, \frac{\pi}{2}\right)$ such that $f(c) = 0$
- D. The equation $f(x) = 0$ has only two roots in $\left[0, \frac{\pi}{2}\right]$

Answer: A::B::C::D

[▶ Watch Video Solution](#)

22. Let $f(x) = \begin{cases} x^{2\alpha+1} \ln x & x > 0 \\ 0 & x = 0 \end{cases}$ If $f(x)$ satisfies Rolle's theorem in interval $[0, 1]$, then α can be:

A. $-\frac{1}{2}$

B. $-\frac{1}{3}$

C. $-\frac{1}{4}$

D. -1

Answer: B::C



Watch Video Solution

23. Which of the following is/are true for the function

$$f(x) = \int_0^x \frac{\cos t}{t} dt (x > 0) ?$$

A. $f(x)$ is monotonically increasing in

$$\left((4n - 1), \frac{\pi}{2}, (4n + 1) \frac{\pi}{2} \right) \forall n \in \mathbb{N}$$

B. $f(x)$ has a local minima at $x = (4n - 1) \frac{\pi}{2} \forall n \in \mathbb{N}$

C. The point of inflection of the curve $y = f(x)$ lie on the curve

$$x \tan x + 1 = 0$$

D. Number of critical points of $y = f(x)$ in $(0, 10\pi)$ are 19

Answer: A::B::C



Watch Video Solution

24. Let $F(x) = (f(x))^2 + (f'(x))^2$, $F(0) = 6$, where $f(x)$ is a thrice differentiable function such that $|f(x)| \leq 1 \forall x \in [-1, 1]$, then choose the correct statement (s)

A. there is atleast one point in each of the intervals

$$(-1, 0) \text{ and } (0, 1) \text{ where } |f'(x)| \leq 2$$

B. there is atleast one point in each of the intervals

$$(-1, 0) \text{ and } (0, 1) \text{ where } F(x) \leq 5$$

C. there is no point of local maxima of $F(x)$ in $(-1, 1)$

D. for some $c \in (-1, 1)$, $F(c) \geq 6$, $F'(c) = 0$ and $f''(c) \leq 0$

Answer: A::B::D

 Watch Video Solution

25. Let $f(x) = \begin{cases} x^3 + x^2 - 10x & -1 \leq x < 0 \\ \sin x & 0 \leq x < \pi/2 \\ 1 + \cos x & \pi/2 \leq x \leq \pi \end{cases}$ then $f(x)$ has

A. local maximum at $x = \frac{\pi}{2}$

B. local minimum at $x = \frac{\pi}{2}$

C. absolute maximum at $x = 0$

D. absolute maximum at $x = -1$

Answer: A::D

 Watch Video Solution

26. Minimum distance between the curves

$y^2 = x - 1$ and $x^2 = x - 1$ and $x^2 = y - 1$ is equal to :

A. $\frac{\sqrt{2}}{4}$

B. $\frac{3\sqrt{2}}{4}$

C. $\frac{5\sqrt{2}}{4}$

D. $\frac{7\sqrt{2}}{4}$

Answer: B



Watch Video Solution

27. For the equation $\frac{e^{-x}}{1+x} = \lambda$ which of the following statement (s) is/are correct ?

A. When $\lambda \in (0, \infty)$ equation has 2 real and distinct roots

B. When $\lambda, \in (-\infty, -e^2)$ equation has 2 real and distinct roots

C. When $\lambda \in (0, \infty)$ equation has 1 real root

D. When $\lambda \in (-e, 0)$ equation has no real root

Answer: B::C::D



Watch Video Solution

28. If $y = mx + 5$ is a tangent to the curve $x^3y^3 = ax^3 + by^3$ at $P(1, 2)$, then

A. $a + b = \frac{18}{5}$

B. $a > b$

C. $a < b$

D. $a + b = \frac{19}{5}$

Answer: A:D



Watch Video Solution

29. If $(f(x) - 1)(x^2 + x + 1)^2 - (f(x) + 1)(x^4 + x^2 + 1) = 0$

$\forall x \in R - \{0\}$ and $f(x) \neq \pm 1$, then which of the following statement (s) is/are correct ?

A. $|f(x)| \geq 2 \forall x \in \mathbb{R} - \{0\}$

B. $f(x)$ has a local maximum at $x = -1$

C. $f(x)$ has a local minimum at $x = 1$

D. $\int_{-\pi}^{\pi} (\cos x) f(x) dx = 0$

Answer: A::B::C::D

 [Watch Video Solution](#)

Exercise Comprehension Type Problems

1. Let $y = f(x)$ such that

$$xy = x + y + 1, x \in \mathbb{R} - \{1\} \text{ and } g(x) = xf(x)$$

The minimum value of $g(x)$ is:

A. $3 - \sqrt{2}$

B. $3 + \sqrt{2}$

C. $3 - 2\sqrt{2}$

D. $3 + 2\sqrt{2}$

Answer: D



Watch Video Solution

2. Let $y = f(x)$ such that

$$xy = x + y + 1, x \in \mathbb{R} - \{1\} \text{ and } g(x) = xf(x)$$

There exist two values of x, x_1 and x_2 where $g'(x) = \frac{1}{2}$, then

$$|x_1| + |x_2| =$$

A. 1

B. 2

C. 4

D. 5

Answer: C



Watch Video Solution

3. Let $f(x) = \begin{cases} 1 - x & 0 \leq x \leq 1 \\ 0 & 1 < x \leq 2 \\ (2 - x)^2 & 2 < x \leq 3 \end{cases}$ and $g(x) = \int_0^x f(t) dt$.

Let the tangent to the curve $y = g(x)$ at point P whose abscissa is $\frac{5}{2}$ cuts x-axis in point Q.

Let the perpendicular from point Q on x-axis meets the curve $y = g(x)$ in point R. Find equation of tangent at to $y=g(x)$ at P. Also the value of $g(1) =$

A. 0

B. $\frac{1}{2}$

C. 1

D. 2

Answer: B



Watch Video Solution

4. Let $f(x) = \begin{cases} 1 - x & 0 \leq x \leq 1 \\ 0 & 1 < x \leq 2 \\ (2 - x)^2 & 2 < x \leq 3 \end{cases}$ and $g(x) = \int_0^x f(t)dt$.

Let the tangent to the curve $y = g(x)$ at point P whose abscissa is $\frac{5}{2}$ cuts x-axis in point Q.

Let the perpendicular from point Q on x-axis meets the curve $y = g(x)$ in point R .Find equation of tangent at to $y=g(x)$ at P .Also the value of $g(1) =$

A. $3y = 12x - 1$

B. $3y = 12x - 1$

C. $12y = 3x - 1$

D. $12y = 3x + 1$

Answer: C



Watch Video Solution

5. Let $f(x) = \begin{cases} 1 - x & 0 \leq x \leq 1 \\ 0 & 1 < x \leq 2 \\ (2 - x)^2 & 2 < x \leq 3 \end{cases}$ and $g(x) = \int_0^x f(t) dt$.

Let the tangent to the curve $y = g(x)$ at point P whose abscissa is $\frac{5}{2}$ cuts x-axis in point Q.

Let the perpendicular from point Q on x-axis meets the curve $y = g(x)$ in point R. Find equation of tangent at to $y=g(x)$ at P. Also the value of $g(1) =$

A. $\frac{5}{6}$

B. $\frac{5}{14}$

C. $\frac{5}{7}$

D. $\frac{5}{12}$

Answer: B



Watch Video Solution

6. Let $f(x) < 0 \forall x \in (-\infty, 0)$ and $f(x) > 0, \forall x \in (0, \infty)$ also

$f(0) = 0,$ Again

$f'(x) < 0, \forall x \in (-\infty, -1)$ and $f'(x) > 0, \forall x \in (-1, \infty)$ also

$f'(-1) = 0$ given $\lim_{x \rightarrow -\infty} f(x) = 0$ and $\lim_{x \rightarrow \infty} f(x) = \infty$ and

function is twice differentiable.

If $f'(x) < 0 \forall x \in (0, \infty)$ and $f'(0) = 1$ then number of solutions of equation $f(x) = x^2$ is : (a) 1 (b) 2 (c) 3 (d) 4

A. 2

B. 3

C. 4

D. None of these

Answer: D



Watch Video Solution

7. Let $f(x) < 0 \forall x \in (-\infty, 0)$ and $f(x) > 0, \forall x \in (0, \infty)$ also
 $f(0) = 0,$ Again

$f'(x) < 0, \forall x \in (-\infty, -1)$ and $f'(x) > 0, \forall x \in (-1, \infty)$ also

$f'(-1) = 0$ given $\lim_{x \rightarrow -\infty} f(x) = 0$ and $\lim_{x \rightarrow \infty} f(x) = \infty$ and

function is twice differentiable.

If $f'(x) < 0 \forall x \in (0, \infty)$ and $f'(0) = 1$ then number of solutions of
equation $f(x) = x^2$ is : (a) 1 (b) 2 (c) 3 (d) 4

A. 1

B. 2

C. 3

D. 4

Answer: B



Watch Video Solution

8. Let $f(x) < 0 \forall x \in (-\infty, 0)$ and $f(x) > 0 \forall x \in (0, \infty)$ also

$f(0) = 0$, Again

$f'(x) < 0 \forall x \in (-\infty, -1)$ and $f'(x) > 0 \forall x \in (-1, \infty)$ also

$f'(-1) = 0$ given $\lim_{x \rightarrow -\infty} f(x) = 0$ and $\lim_{x \rightarrow \infty} f(x) = \infty$ and

function is twice differentiable.

The minimum number of points where $f'(x)$ is zero is: (a) 1 (b) 2 (c) 3 (d)

4

A. 1

B. 2

C. 3


D. 4

Answer: A



Watch Video Solution

9. In the given figure graph of $y = p(x) = x^4 + ax^3 + bx^2 + cx + d$ is given



The product of all imaginary roots of $p(x) = 0$ is (a) 1 (b) 2 (c) $\frac{1}{3}$ (d) $\frac{1}{4}$

- A. -2
- B. -1
- C. $-1/2$
- D. none of these


Answer: D



Watch Video Solution

10. In the given figure graph of :

$$y = p(x) = x^n + a_1x^{n-1} + a_2x^{n-2} + \dots + a_n \text{ is given.}$$



If $p(x) + k = 0$ has 4 distinct real roots $\alpha, \beta, \gamma, \delta$ then

$[\alpha] + [\beta] + [\gamma] + [\delta]$, (where $[.]$ denotes greatest integer function) is

equal to:

A. -1

B. -2

C. 0


D. 1

Answer: A



Watch Video Solution

11. In the given figure graph of $y = p(x) = x^4 + ax^3 + bx^2 + cx + d$ is given



The product of all imaginary roots of $p(x) = 0$ is (a) 1 (b) 2 (c) $\frac{1}{3}$ (d) $\frac{1}{4}$

A. 1

B. 4

C. 5

D. 6

Answer: B



Watch Video Solution

12. The differentiable function $y = f(x)$ has a property that the chord joining any two points $A(x_1, f(x_1))$ and $B(x_2, f(x_2))$ always intersects y-axis at $(0, 2x_1x_2)$. Given that $f(1) = -1$. then:

$\int_0^{1/2} f(x)dx$ is equal to : (a) $\frac{1}{6}$ (b) $\frac{1}{8}$ (c) $\frac{1}{12}$ (d) $\frac{1}{24}$

A. $\frac{1}{6}$

B. $\frac{1}{8}$

C. $\frac{1}{12}$

D. $\frac{1}{24}$

Answer: D



Watch Video Solution

13. The differentiable function $y = f(x)$ has a property that the chord joining any two points $A(x_1, f(x_1))$ and $B(x_2, f(x_2))$ always intersects

y-axis at $(0, 2x_1x_2)$. Given that $f(1) = -1$. then:

The largest interval in which $y = f(x)$ is monotonically increasing, is : (a)

$\left(-\infty, \frac{1}{2}\right]$ (b) $\left[\frac{-1}{2}, \infty\right)$ (c) $\left(-\infty, \frac{1}{4}\right]$ (d) $\left[\frac{-1}{4}, \infty\right)$

A. $\left(-\infty, \frac{1}{2}\right]$

B. $\left[\frac{-1}{2}, \infty\right)$

C. $\left(-\infty, \frac{1}{4}\right]$

D. $\left[\frac{-1}{4}, \infty\right)$

Answer: C



Watch Video Solution

14. The differentiable function $y = f(x)$ has a property that the chord joining any two points $A(x_1, f(x_1))$ and $B(x_2, f(x_2))$ always intersects y-axis at $(0, 2x_1x_2)$. Given that $f(1) = -1$. then:

In which of the following intervals, the Rolle's theorem is applicable to the function $F(x) = f(x) + x$? (a) $[-1, 0]$ (b) $[0, 1]$ (c) $[-1, 1]$ (d)

$[0, 2]$

A. $0 - 1, 0]$

B. $[0, 1]$

C. $[-1, 1]$

D. $[0, 2]$

Answer: B

 [Watch Video Solution](#)

15. If $f(x) = x + \int_0^1 (xy^2 + x^2y)(f(y))dy$, find $f(x)$ if x and y are independent.

A. $\frac{8}{25}$

B. $\frac{16}{25}$

C. $\frac{14}{25}$

D. $\frac{4}{5}$

Answer: A



Watch Video Solution

Exercise Matching Type Problems

1. Column-1 gives pair of curves and column-II gives the angle θ between the curves at their intersection point.

Column-I		Column-II	
(A)	$y = \sin x, y = \cos x$	(P)	$\frac{\pi}{4}$
(B)	$x^2 = 4y, y = \frac{8}{x^2 + 4}$	(Q)	$\frac{\pi}{2}$
(C)	$\frac{x^2}{18} + \frac{y^2}{8} = 1, x^2 - y^2 = 5$	(R)	$\tan^{-1} 3$
(D)	$xy = 1, x^2 - y^2 = 5$	(S)	$\tan^{-1} 5$
		(T)	$\tan^{-1}(2\sqrt{2})$



Watch Video Solution

2.

Let

$$f(x) = \frac{x^3 - 4}{(x - 1)^3} \forall x \neq 1, g(x) = \frac{x^4 - 2x^2}{4} \forall x \in \mathbb{R}, h(x) = \frac{x^3 + 4}{(x + 1)^3} \forall x$$

Column-I		Column-II	
(A)	The number of possible distinct real roots of equation $f(x) = c$ where $c \geq 4$ can be	(P)	0
(B)	The number of possible distinct real roots of equation $g(x) = c$, where $c \geq 0$ can be	(Q)	1
(C)	The number of possible distinct real roots of equation $h(x) = c$, where $c \geq 1$ can be	(R)	2

(D)	The number of possible distinct real roots of equation $(x) = c$ where $-1 < c < 0$ can be	(S)	3
		(T)	4

 [Watch Video Solution](#)

3. Consider the function $f(x) = \frac{\ln x}{8} - ax + x^2$ and $a \geq 0$ is a real constant :

Column-I		Column-II	
(A)	$f(x)$ gives a local maxima at	(P)	$a = 1; x = \frac{1}{4}$
(B)	$f(x)$ gives a local minima at	(Q)	$a > 1; x = \frac{a - \sqrt{a^2 - 1}}{4}$
(C)	$f(x)$ gives a point of inflection for	(R)	$0 \leq a < 1$
(D)	$f(x)$ is strictly increasing for all $x \in \mathbb{R}^+$	(S)	$a > 1; x = \frac{a + \sqrt{a^2 - 1}}{4}$

 [Watch Video Solution](#)

4. The function $f(x) = \sqrt{ax^3 + bx^2 + cx + d}$ has its non-zero local minimum and local maximum values at $x = -2$ and $x = 2$, respectively. If a is a root of $x^2 - x - 6 = 0$, then find a, b, c and d .

 [Watch Video Solution](#)

Column-I		Column-II	
(A)	The ratio of altitude to the radius of the cylinder of maximum volume that can be inscribed in a given sphere is	(P)	$\frac{1}{\sqrt{2}}$
(B)	The ratio of radius to the altitude of the cone of the greatest volume which can be inscribed in a given sphere is	(Q)	$\sqrt{2}$
(C)	The cone circumscribing the sphere of radius 'r' has the maximum volume if its semi vertical angle is θ , then $33 \sin \theta =$	(R)	$\frac{32}{3}$
(D)	The greatest value of $x^3 y^4$ if $2x + 3y = 7$, $x \geq 0, y \geq 0$ is	(S)	11

5.

 [Watch Video Solution](#)

Exercise Subjective Type Problems

1. A conical vessel is to be prepared out of a circular sheet of metal of unit radius in order that the vessel has maximum value, the sectorial area that must be removed from the sheet is A_1 and the area of the given sheet is A_2 , then $\frac{A_2}{A_1}$ is equal to

 [Watch Video Solution](#)

2. On $[1, e]$, then least and greatest values of $f(x) = x^2 \ln x$ are m and M respectively, then $[\sqrt{M + m}]$ is : (where $[\]$ denotes greatest integer function)

 [Watch Video Solution](#)

3. If $f(x) = \frac{px}{e^x} - \frac{x^2}{2} + x$ is a decreasing function for every $x \leq 0$. Find the least value of p^2 .

 [Watch Video Solution](#)

4. Let $f(x) = \begin{cases} xe^{ax}, & x \leq 0 \\ x + ax^2 - x^3, & x > 0 \end{cases}$ where a is a positive constant .

The interval in which $f'(x)$ is increasing is $\left[\frac{k}{a}, \frac{a}{l}\right]$, Then $k + l$ is equal to



[Watch Video Solution](#)

5. Find sum of all possible values of the real parameter 'b' if the difference between the largest and smallest values of the function

$f(x) = x^2 - 2bx + 1$ in the interval $[0, 1]$ is 4.



[Watch Video Solution](#)

6. Let ' θ ' be the angle in radians between the curves

$\frac{x^2}{36} + \frac{y^2}{4} = 1$ and $x^2 + y^2 = 12$. If $\theta = \tan^{-1}\left(\frac{a}{\sqrt{3}}\right)$, Find the value

of a.



[Watch Video Solution](#)

7. Let set of all possible values of λ such that $f(x) = e^{2x} - (\lambda + 1)e^x + 2x$ is monotonically increasing for $\forall x \in \mathbb{R}$ is $(-\infty, k]$. Find the value of k .

 [Watch Video Solution](#)

8. Let a, b, c and d be non-negative real number such that $a^5 + b^5 \leq 1$ and $c^5 + d^5 \leq 1$. Find the maximum value of $a^2c^3 + b^2d^3$.

 [Watch Video Solution](#)

9. There is a point (p, q) on the graph of $f(x) = x^2$ and a point (r, s) on the graph of $g(x) = \frac{-8}{x}$, where $q > 0$ and $r > 0$. If the line through (p, q) and (r, s) is also tangent to both the curves at these points, respectively, then the value of $p + r$ is _____

 [Watch Video Solution](#)

10. If $f(x) = \max |2 \sin y - x|$, (where $y \in R$), then find the minimum value of $f(x)$.

 [Watch Video Solution](#)

11. Let $f(x) = \int_0^x \left((a-1)(t^2 + t + 1)^2 - (a+1)(t^4 + t^2 + 1) \right) dt$. Then the total number of integral values of 'a' for which $f'(x) = 0$ has no real roots is

 [Watch Video Solution](#)

12. The number of real roots of the equation $x^{2013} + e^{20144x} = 0$ is

 [Watch Video Solution](#)

13. Let the maximum value of expression $y = \frac{x^4 - x^2}{x^6 + 2x^3 - 1}$ for $x > 1$ is $\frac{p}{1}$, where p and $1q$ are relatively prime natural numbers, then

$$p + q =$$



Watch Video Solution

14. The least positive value of the parameter 'a' for which there exist at least one line that is tangent to the graph of the curve $y = x^3 - ax$, at one point and normal to the graph at another point is $\frac{p}{q}$, where p and q are relatively prime positive integers. Find product pq.



Watch Video Solution

15. Let $f(x) = x^2 + 2x - t^2$ and $f(x) = 0$ has two roots $\alpha(t)$ and $\beta(t)$ ($\alpha < \beta$) where t is a real parameter. Let $I(t) = \int_{\alpha}^{\beta} f(x) dx$. If the maximum value of $I(t)$ be λ and $|\lambda| = \frac{p}{q}$ where p and q are relatively prime positive integers. Find the product (pq).



Watch Video Solution

16. A tank contains 100 litres of fresh water. S solution containing 1 gm/litre of salt runs into the tank at the rate of 1 lit/min. The homogenised mixture is pumped out of the tank at the rate of 3 lit/min. If T be the time when the amount of salt in the tank is maximum.

Find $[T]$ (where $[.]$ denotes greatest integer function)



[Watch Video Solution](#)

17. If $f(x)$ is continuous and differentiable in $[-3, 9]$ and $f'(x) \in [-2, 8] \forall x \in (-3, 9)$. Let N be the number of divisors of the greatest possible value of $f(9) - f(-3)$, then find the sum of digits of N.



[Watch Video Solution](#)

18. It is given that $f(x)$ is defined on \mathbb{R} satisfying $f(1) = 1$ and for $\forall x \in \mathbb{R}$,

$$f(x + 5) \geq f(x) + 5 \text{ and } f(x + 1) \leq f(x) + 1.$$

If

$$g(x) = f(x) + 1 - x, \text{ then } g(2002) = \underline{\hspace{2cm}}$$

 [Watch Video Solution](#)

19. The number of normals to the curve $3y^3 = 4x$ which passes through the point $(0, 1)$ is

 [Watch Video Solution](#)

20. Find the number of real root (s) of the equation $ae^x = 1 + x + \frac{x^2}{2}$, where a is positive constant.

 [Watch Video Solution](#)

21. Let $f(x) = ax + \cos 2x + \sin x + \cos x$ is defined for $\forall x \in R$ and $a \in R$ and is strictly increasing function. If the range of a is $\left[\frac{m}{n}, \infty\right)$, then find the minimum value of $(m - n)$.



[Watch Video Solution](#)

22. If p_1 and p_2 are the lengths of the perpendiculars from origin on the tangent and normal drawn to the curve $x^{2/3} + y^{2/3} = 6^{2/3}$ respectively.

Find the value of $\sqrt{4p_1^2 + p_2^2}$.



[Watch Video Solution](#)