



MATHS

BOOKS - VK JAISWAL ENGLISH

AREA UNDER CURVES

Exercise Single Choice Problems

1. The area enclosed by the curve

$$[x + 3y] = [x - 2] \text{ where } x \in [3, 4] \text{ is :}$$

(where $[.]$ denotes greatest integer function)

A. $\frac{2}{3}$

B. $\frac{1}{3}$

C. $\frac{1}{4}$

D. 1

Answer: B

 [Watch Video Solution](#)

2. The area of region(s) enclosed by the curve $y = x^2$ and $y = \sqrt{|x|}$ is

A. $\frac{1}{3}$

B. $\frac{2}{3}$

C. $\frac{4}{3}$

D. $\frac{16}{3}$

Answer: B

 [Watch Video Solution](#)

3. Find the area enclosed by the figure described by the equation

$$x^4 + 1 = 2x^2 + y^2.$$

A. 2

B. $\frac{16}{3}$

C. $\frac{8}{3}$

D. $\frac{4}{3}$

Answer: C



Watch Video Solution

4. The area defined by $|y| \leq e^{|x|} - \frac{1}{2}$ in cartesian co-ordinate system,

is :

A. $(2 - 2 \ln 2)$

B. $(4 - \ln 2)$

C. $(2 - \ln 2)$

D. $(2 - 2 \ln 2)$

Answer: D



Watch Video Solution

5. For each positive integer $n > a$, A_n represents the area of the region restricted to the following two inequalities :

$$\frac{x^2}{n^2} + y^2 \text{ and } x^2 + \frac{y^2}{n^2} < 1. \text{ Find } \lim_{n \rightarrow \infty} A_n.$$

A. 4

B. 1

C. 2

D. 3

Answer: A



Watch Video Solution

6. Find the ratio in which the area bounded by the curves $y^2 = 12x$ and $x^2 = 12y$ is divided by the line $x = 3$.

A. 7: 15

B. 15: 49

C. 1: 3

D. 17: 49

Answer: B



[Watch Video Solution](#)

7. The value of positive real parameter 'a' such that area of region bounded by parabolas $y = x - ax^2$, $ay = x^2$ attains its maximum value is equal to :

A. $\frac{1}{2}$

B. 2

C. $\frac{1}{3}$

D. 1

Answer: D

 [Watch Video Solution](#)

8. For $0 < r < 1$, let n_r denotes the line that is normal to the curve $y = x^r$ at the point $(1, 1)$ Let S_r denotes the region in the first quadrant bounded by the curve $y = x^r$, the x-axis and the line n_r . Then the value of r that minimizes the area of S_r is :

A. $\frac{1}{\sqrt{2}}$

B. $\sqrt{2} - 1$

C. $\frac{\sqrt{2} - 1}{2}$

D. $\sqrt{2} - \frac{1}{2}$

Answer: B

 [Watch Video Solution](#)

9. The area bounded by $|x| = 1 - y^2$ and $|x| + |y| = 1$ is:

A. $\frac{1}{3}$

B. $\frac{1}{2}$

C. $\frac{2}{3}$

D. 1

Answer: C



Watch Video Solution

10. Point A lies on curve $y = e^{-x^2}$ and has the coordinates (x, e^{-x^2})

where $x > 0$ Point B has coordinates $(x, 0)$ If 'O' is the origin, then the

maximum area of $\triangle AOB$ is :

A. $\frac{1}{\sqrt{8}e}$

B. $\frac{1}{\sqrt{4}e}$

C. $\frac{1}{\sqrt{2}e}$

D. $\frac{1}{\sqrt{e}}$

Answer: A

 [Watch Video Solution](#)

11. If the area enclosed between the curves $y = ax^2$ and $x = ay^2$ ($a > 0$) is 1 square unit, then find the value of a .

A. $\frac{1}{\sqrt{3}}$

B. $\frac{1}{2}$

C. 1

D. $\frac{1}{3}$

Answer: D

 [Watch Video Solution](#)

12. Let $f(x) = x^3 - 3x^2 + 3x + 1$ and g be the inverse of it , then area bounded by the curve $y = g(x)$ wirth x-axis between $x = 1$ to $x = 2$ is (in square units):

A. $\frac{1}{2}$

B. $\frac{1}{4}$

C. $\frac{3}{4}$

D. 1

Answer: B



[Watch Video Solution](#)

13. Area bounded by $x^2y^2 + y^4 - x^2 - 5y^2 + 4 = 0$ is equal to :

A. $\frac{4\pi}{2} + \sqrt{2}$

B. $\frac{4\pi}{3} - \sqrt{2}$

C. $\frac{4\pi}{3} - \sqrt{2}$

D. none of these

Answer: C

 [Watch Video Solution](#)

14. Let $f(x): R^+ \rightarrow R^+$ is an invertible function such that $f'(x) > 0$ and $f(x) > 0 \forall x \in [1, 5]$. If $f(1) = 1$ and $f(5) = 5$ and area under the curve $y = f(x)$ on x-axis from $x = 1 \rightarrow x = 5$ is 8 sq. units, then area bounded by $y = f^{-1}(x)$ on x-axis from $x = 1 \rightarrow x = 5$ is

A. 12

B. 16

C. 18

D. 20

Answer: B

 [Watch Video Solution](#)

15. A circle centered at origin and having radius π units is divided by the curve $y = \sin x$ in two parts. Then area of the upper part equals to

: (a) $\frac{\pi^2}{2}$ (b) $\frac{\pi^3}{4}$ (c) $\frac{\pi^3}{2}$ (d) $\frac{\pi^3}{8}$

A. $\frac{\pi^2}{2}$

B. $\frac{\pi^3}{4}$

C. $\frac{\pi^3}{2}$

D. $\frac{\pi^3}{8}$

Answer: C

 [Watch Video Solution](#)

16. The area of the loop formed by $y^2 = x(1 - x^3)$ dx is:

A. $\int_0^1 \sqrt{x - x^4} dx$

$$\text{B. } 2 \int_0^1 \sqrt{x - x^4} dx$$

$$\text{C. } \int_{-1}^1 \sqrt{x - x^4} dx$$

$$\text{D. } 4 \int_0^{1/2} \sqrt{x - x^4} dx$$

Answer: B

 [Watch Video Solution](#)

17. If $f(x) = \min \left[x^2, \sin \frac{x}{2}, (x - 2\pi)^2 \right]$, the area bounded by the curve $y = f(x)$, x -axis, $x = 0$ and $x = 2\pi$ is given by

Note: x_1 is the point of intersection of the curves x^2 and $\sin \frac{x}{2}$, x_2 is the point of intersection of the curves $\sin \frac{x}{2}$ and $(x - 2\pi)^2$

A.

$$\int_0^{x_1} \left(\sin \frac{x}{2} \right) dx + \int_{x_1}^{\pi} x^2 dx + \int_{\pi}^{x_2} (x - 2\pi)^2 dx + \int_{x_2}^{2\pi} \left(\sin \frac{x}{2} \right) dx$$

B. $\int_0^{x_1} x^2 dx + \int_{x_1}^{x_3} \left(\sin \frac{x}{2} \right) dx + \int_{x_2}^{2\pi} (x - 2\pi)^2 dx,$ where

$$x_1 \in \left(0, \frac{\pi}{3} \right) \text{ and } x_2 \in \left(\frac{5\pi}{3}, 2\pi \right)$$

$$C. \int_0^{x_1} x^2 dx + \int_{x_1}^{x_2} \sin\left(\frac{x}{2}\right) dx + \int_{x_2}^{2\pi} (x - 2\pi)^2 dx,$$

where

$$x_1 \in \left(\frac{\pi}{3}, \frac{\pi}{2}\right) \text{ and } x_2 \in \left(\frac{3\pi}{2}, 2\pi\right)$$

$$D. \int_0^{x_1} x^2 dx + \int_{x_1}^{x_2} \sin\left(\frac{x}{2}\right) dx + \int_{x_2}^{2\pi} (x - 2\pi)^2 dx,$$

where

$$x_1 \in \left(\frac{\pi}{2}, \frac{2\pi}{3}\right) \text{ and } x_2 \in (\pi, 2\pi)$$

Answer: B



Watch Video Solution

18. The area enclosed between the curves

$$|x| + |y| \geq 2 \text{ and } y^2 = 4\left(1 - \frac{x^2}{9}\right) \text{ is :}$$

A. $(6\pi - 4)$ sq. units

B. $(6\pi - 8)$ se. units

C. $(3\pi - 4)$ se. units


D. $(3\pi - 2)$ sq. units

Answer: B

 Watch Video Solution

Exercise One Or More Than One Answer Is Are Correct

1. Let $f(x)$ be a polynomial function of degree 3 where $a < b < c$ and $f(a) = f(b) = f(c)$. If the graph of $f(x)$ is as shown, which of the following statements are INCORRECT? (Where $c > |a|$)



A. $\int_a^c f(x)dx = \int_b^c f(x)dx + \int_a^b f(x)dx$

B. $\int_a^c f(x)dx < a$

$$C. \int_a^b f(x)dx < \int_c^b f(x)dx$$

$$D. \frac{1}{b-a} \int_a^b f(x)dx > \frac{1}{c-b} \int_b^c f(x)dx$$

Answer: B::C::D

 Watch Video Solution

2. $T_n = \sum_{r=2n}^{3n-1} \frac{r}{r^2 + n^2}, S_n = \sum_{r=2n+1}^{3n} \frac{r}{r^2 + n^2},$ then

$\forall n \in \{1, 2, 3, \dots\}:$

A. $T_n > \frac{1}{2} \ln 2$

B. $S_n < \frac{1}{2} \ln 2$

C. $T_n < \frac{1}{2} \ln 2$

D. $S_n > \frac{1}{2} \ln 2$

Answer: A::B

 Watch Video Solution

3. If a curve $y = a\sqrt{x} + bx$ passes through point $(1, 2)$ and the area bounded by curve, line $x = 4$ and x-axis is 8, then : (a) $a = 3$ (b) $b = 3$
(c) $a = -1$ (d) $b = -1$

A. $a = 3$

B. $b = 3$

C. $a = -1$

D. $b = -1$

Answer: A::D



[Watch Video Solution](#)

4. Area enclosed by the curves $y = x^2 + 1$ and a normal drawn to it with gradient -1, is equal to:

A. $\frac{2}{3}$

B. $\frac{1}{3}$

C. $\frac{3}{4}$

D. $\frac{4}{3}$

Answer: D



Watch Video Solution

Exercise Comprehension Type Problems

1. Let $f: A \rightarrow B$ $f(x) = \frac{x + a}{bx^2 + cx + 2}$, where A represent domain set and B represent range set of function $f(x)$ $a, b, c \in R$, $f(-1) = 0$ and $y = 1$ is an asymptote of $y = f(x)$ and $y = g(x)$ is the inverse of $f(x)$.

$g(0)$ is equal to :

A. -1

B. -3

C. $-\frac{5}{2}$

D. $-\frac{3}{2}$

Answer: A



Watch Video Solution

2. Let $f: A \rightarrow B$ $f(x) = \frac{x+a}{bx^2+cx+2}$, where A represent domain set and B represent range set of function $f(x)$ $a, b, c \in R$, $f(-1) = 0$ and $y = 1$ is an asymptote of $y = f(x)$ and $y = g(x)$ is the inverse of $f(x)$.

Area of region enclosed by asymptotes of curves $y = f(x)$ and $y = g(x)$ is:

- A. $2\sqrt{5} + \ln\left(\frac{3-\sqrt{5}}{5+\sqrt{5}}\right)$
- B. $2\sqrt{5} + 2\ln\left(\frac{3+\sqrt{5}}{3-\sqrt{5}}\right)$
- C. $3\sqrt{5} + 4\ln\left(\frac{3-\sqrt{5}}{3+\sqrt{5}}\right)$
- D. $3\sqrt{5} + 2\ln\left(\frac{3-\sqrt{5}}{3+\sqrt{5}}\right)$

Answer: D

 [Watch Video Solution](#)

3. Let $f: A \rightarrow B$ $f(x) = \frac{x + a}{bx^2 + cx + 2}$, where A represent domain set and B represent range set of function $f(x)$ $a, b, c \in R$, $f(-1) = 0$ and $y = 1$ is an asymptote of $y = f(x)$ and $y = g(x)$ is the inverse of $f(x)$.

Area of region enclosed by asymptotes of curves $y = f(x)$ and $y = g(x)$ is:

A. 4

B. 9

C. 12

D. 25

Answer: B

 [Watch Video Solution](#)

4. For $j = 0, 1, 2, \dots, n$ let S_j be the area of region bounded by the x-axis and the curve $ye^x = \sin x$ for $j\pi \leq x \leq (j+1)\pi$

The value of $\sum_{j=0}^{\infty} S_j$ equals to :

A. $\frac{e^x(1+e^x)}{2(e^\pi-1)}$

B. $\frac{1+e^\pi}{2(e^\pi-1)}$

C. $\frac{1+e^\pi}{e^\pi-1}$

D. $\frac{e^\pi(1+e^\pi)}{(e^\pi-1)}$

Answer: B



[Watch Video Solution](#)

5. For $j = 0, 1, 2, \dots, n$ let S_j be the area of region bounded by the x-axis and the curve $ye^x = \sin x$ for $j\pi \leq x \leq (j+1)\pi$

The ratio $\frac{S_{2009}}{S_{2010}}$ equals :

A. e^{-x}

B. (e^x)

C. $\frac{1}{2}e^x$

D. $2e^\pi$

Answer: B



Watch Video Solution

6. For $j = 0, 1, 2, \dots, n$ let S_j be the area of region bounded by the x-axis and the curve $ye^x = \sin x$ for $j\pi \leq x \leq (j+1)\pi$

The value of $\sum_{j=0}^{\infty} S_j$ equals to :

A. $\frac{e^x(1+e^x)}{2(e^\pi-1)}$

B. $\frac{1+e^\pi}{2(e^\pi-1)}$

C. $\frac{1+e^\pi}{e^\pi-1}$

D. $\frac{e^\pi(1+e^\pi)}{(e^\pi-1)}$

Answer: B

 Watch Video Solution

Exercise Matching Type Problems

Column-I		Column-II	
(A)	Area of region formed by points (x, y) satisfying $[x]^2 = [y]^2$ for $0 \leq x \leq 4$ is equal to (where $[\]$ denotes greatest integer function)	(P)	48
(B)	The area of region formed by points (x, y) satisfying $x + y \leq 6$, $x^2 + y^2 \leq 6y$ and $y^2 \leq 8x$ is $\frac{k\pi - 2}{12}$, then $k =$	(Q)	27
(C)	The area in the first quadrant bounded by the curve $y = \sin x$ and the line $\frac{2y - 1}{\sqrt{2} - 1} = \frac{2}{\pi}(6x - \pi)$ is $\left[\frac{\sqrt{3} - \sqrt{2}}{2} - \frac{(\sqrt{2} + 1)\pi}{k} \right]$, then $k =$	(R)	7
(D)	If the area bounded by the graph of $y = xe^{-ax}$ ($a > 0$) and the abscissa axis is $\frac{1}{9}$ then the value of 'a' is equal to	(S)	4
		(T)	3

1.

 Watch Video Solution

Exercise Subjective Type Problems

1. Let f be a differentiable function satisfying the condition $f\left(\frac{x}{y}\right) = \frac{f(x)}{f(y)} (y \neq 0, f(y) \neq 0) \forall x, y \in R$ and $f'(1) = 2$. If the smaller area enclosed by $y = f(x)$, $x^2 + y^2 = 2$ is A , then find $[A]$, where $[.]$ represents the greatest integer function.

 [Watch Video Solution](#)

2. Let $f(x)$ be a function which satisfy the equation $f(xy) = f(9x) + f(y)$ for all $x > 0, y > 0$ such that $f'(1) = 2$. Let A be the area of the region bounded by the curves $y = f(x)$, $y = |x^3 - 6x^2 + 11x - 6|$ and $x = 0$, then find value of $\frac{28}{17}A$.

 [Watch Video Solution](#)

3. If the area bounded by circle $x^2 + y^2 = 4$, the parabola $y = x^2 + x + 1$ and the curve $y = \left[\sin^2 \frac{x}{4} + \cos \frac{x}{4}\right]$, (where $[.]$

denotes the greatest integer function) and x-axis is $\left(\sqrt{3} + \frac{2\pi}{3} - \frac{1}{k}\right)$,

then the numerical quantity is should be :

 [Watch Video Solution](#)

4. Let the function $f: [-4, 4] \rightarrow [-1, 1]$ be defined implicitly by the equation $x + 5y - y^5 = 0$ If the area of triangle formed by tangent and normal to $f(x)$ at $x = 0$ and the line $y = 5$ is A, find $\frac{A}{13}$.

 [Watch Video Solution](#)

5. Area of the region bounded by $[x]^2 = [y]^2$, if $x \in [1, 5]$, where $[]$ denotes the greatest integer function is:

 [Watch Video Solution](#)

6. Consider $y = x^2$ and $f(x)$ where $f(x)$, is a differentiable function satisfying

$$f(x+1) + f(z-1) = f(x+z) \forall x, z \in R \text{ and } f(0) = 0, f'(0) = 4.$$

If area bounded by curve $y = x^2$ and $y = f(x)$ is Δ , find the value of $\left(\frac{3}{16}, \Delta\right)$.

 [Watch Video Solution](#)

7. The least integer which is greater than or equal to the area of region in $x - y$ plane satisfying $x^6 - x^2 + y^2 \leq 0$ is:

 [Watch Video Solution](#)

8. The set of points (x,y) in the plane satisfying $x^{2/5} + |y| = 1$ form a curve enclosing a region of area $\frac{p}{q}$ square units, when p and q are relatively prime positive integers. Find $p - q$.

 [Watch Video Solution](#)

