



## MATHS

### BOOKS - VK JAISWAL ENGLISH

### COMPOUND ANGLES

#### Exercise 1 Single Choice Problems

1.  $\left(\cos^4 \frac{\pi}{24} - \sin^4 \frac{\pi}{24}\right)$  equals :

A.  $\frac{1}{\sqrt{2}}$

B.  $\frac{\sqrt{6} - \sqrt{2}}{4}$

C.  $\frac{\sqrt{6} + \sqrt{2}}{4}$

D.  $\frac{\sqrt{3} + 1}{2}$

**Answer: C**



Watch Video Solution

2. If  $a \sin x + b \cos(c + x) + b \cos(c - x) = \alpha$ ,  $\alpha > a$ , then the minimum value of  $|\cos c|$  is :

A.  $\sqrt{\frac{\alpha^2 - a^2}{b^2}}$

B.  $\sqrt{\frac{\alpha^2 - a^2}{2b^2}}$

C.  $\sqrt{\frac{\alpha^2 - a^2}{3b^2}}$

D.  $\sqrt{\frac{\alpha^2 - a^2}{4b^2}}$

**Answer: D**



**Watch Video Solution**

3. If all values of  $x \in (a, b)$  satisfy the inequality  $\tan x \tan 3x < -1$ ,  $x \in \left(0, \frac{\pi}{2}\right)$ , then the maximum value  $(b - a)$  is :

A.  $\frac{\pi}{12}$

B.  $\frac{\pi}{3}$

C.  $\frac{\pi}{6}$

D.  $\frac{\pi}{4}$

**Answer: A**



[Watch Video Solution](#)

4.  $\sum_{r=1}^8 \tan(rA)\tan((r+1)A)$  where  $A = 36^\circ$  is :

A.  $-10 - \tan A$

B.  $-10 + \tan A$

C. -10

D. -9

**Answer: D**



[Watch Video Solution](#)

5. Let  $f(x) = 2 \operatorname{cosec} 2x + \sec x + \operatorname{cosec} x$ . Then find the minimum value of  $f(x)$  or  $x \in \left(0, \frac{\pi}{2}\right)$ .

A.  $\frac{1}{\sqrt{2} - 1}$

B.  $\frac{2}{\sqrt{2} - 1}$

C.  $\frac{1}{\sqrt{2} + 1}$

D.  $\frac{2}{\sqrt{2} + 1}$

**Answer: B**



**Watch Video Solution**

6. The value of  $\operatorname{cosec} 10^\circ + \operatorname{cosec} 50^\circ - \operatorname{cosec} 70^\circ$  is \_\_\_\_\_

A. 4

B. 5

C. 6

D. 8

**Answer: C**



**Watch Video Solution**

7. If  $u = \sqrt{a^2 \cos^2 \theta + b^2 \sin^2 \theta} + \sqrt{a^2 \sin^2 \theta + b^2 \cos^2 \theta}$ , then the difference between the maximum and minimum values of  $u^2$  is given by :

(a)  $(a - b)^2$  (b)  $2\sqrt{a^2 + b^2}$  (c)  $(a + b)^2$  (d)  $2(a^2 + b^2)$

A.  $2(a^2 + b^2)$

B.  $2\sqrt{a^2 + b^2}$

C.  $(a + b)^2$

D.  $(a - b)^2$

**Answer: D**



**Watch Video Solution**

8. If  $u_n = \sin(n\theta)\sec^n \theta$ ,  $v_n = \cos(n\theta)\sec^n \theta$ ,  $n \in N$ ,  $n \neq 1$ , then

$$\frac{v_n - v_{n-1}}{v_{n-1}} + \frac{1}{n} \left( \frac{u_n}{v_n} \right) =$$

A.  $-\cos \theta + \frac{1}{n} \tan(n\theta)$

B.  $\cot \theta + \frac{1}{n} \tan(n\theta)$

C.  $\tan \theta + \frac{1}{n} \tan(n\theta)$

D.  $-\tan \theta + \frac{\tan(n\theta)}{n}$

**Answer: D**



**Watch Video Solution**

9.  $\alpha \cos^2 3\theta + \beta \cos^4 \theta = 16 \cos^6 \theta + 9 \cos^2 \theta$ . Find  $\alpha$  &  $\beta$  if its a identity

A.  $a = 1, b = 24$

B.  $a = 3, b = 24$

C.  $a = 4, b = 2$

D.  $a = 7, b = 18$

**Answer: A**



**Watch Video Solution**

10. Maximum value of  $\cos x(\sin x + \cos x)$  is equal to :

A.  $\sqrt{2}$

B. 2

C.  $\frac{\sqrt{2} + 1}{2}$

D.  $\sqrt{2} + 1$

**Answer: C**



**Watch Video Solution**

11. If  $\frac{\sin A}{\sin B} = \frac{\sqrt{3}}{2}$  and  $\frac{\cos A}{\cos B} = \frac{\sqrt{5}}{2}$ ,  $0 < A, B < \frac{\pi}{2}$ , then

$\tan A + \tan B$  is equal to

A.  $\sqrt{\frac{3}{5}}$

B.  $\sqrt{\frac{5}{3}}$

C.  $\frac{\sqrt{3} + \sqrt{5}}{\sqrt{5}}$

D.  $\frac{\sqrt{3} + \sqrt{5}}{\sqrt{3}}$

**Answer: C**



**Watch Video Solution**

12. Let  $0 \leq \alpha, \beta, \gamma, \delta \leq \pi$  where  $\beta$  and  $\gamma$  are not complementary such that

$$2 \cos \alpha, 6 \cos \beta + 7 \cos \gamma + 9 \cos \delta = 0$$

$$\text{and } 2 \sin \alpha - 6 \sin \beta + 7 \sin \gamma - 9 \sin \delta = 0$$

If  $\frac{\cos(\alpha + \delta)}{\cos(\beta + \gamma)} = \frac{m}{n}$  where  $m$  and  $n$  are relatively prime positive

numbers, then the value of  $(m + n)$  is equal to :



A. 11

B. 10

C. 9

D. 2

**Answer: B**



**Watch Video Solution**

13. If  $-\pi < \theta < -\frac{\pi}{2}$ , then  $\left| \sqrt{\frac{1 - \sin \theta}{1 + \sin \theta}} + \sqrt{\frac{1 + \sin \theta}{1 - \sin \theta}} \right|$  is equal to :

A.  $2 \sec \theta$

B.  $-2 \sec \theta$

C.  $2 \sec \frac{\theta}{2}$

D.  $-\sec \frac{\theta}{2}$

**Answer: B**



**Watch Video Solution**

14. If  $A = \sum_{r=1}^3 \cos \frac{2r\pi}{7}$  and  $B = \sum_{r=1}^3 \cos \frac{2^r\pi}{7}$ , then :

A.  $A + B = 0$

B.  $2A + B = 0$

C.  $A + 2B = 0$


D.  $A = B$

**Answer: D**



**Watch Video Solution**

15. In a  $\triangle PQR$  (as shown in figure) if  $x : y : z = 2 : 3 : 6$ , then the value of  $\angle QPR$  is :



- A.  $\frac{\pi}{6}$
- B.  $\frac{\pi}{4}$
- C.  $\frac{\pi}{3}$
- D.  $\frac{\pi}{2}$

**Answer: B**



**Watch Video Solution**

16. If  $A = \sum_{r=1}^3 \cos \frac{2r\pi}{7}$  and  $B = \sum_{r=1}^3 \cos \frac{2^r\pi}{7}$ , then :

A.  $A + B = 0$

B.  $2A + B = 0$

C.  $A + 2B = 0$

D.  $A - B = 0$

**Answer: D**

 [Watch Video Solution](#)

17. Let  $f(x) = \sin x + 2 \cos^2 x$ ,  $\frac{\pi}{6} \leq x \leq \frac{2\pi}{3}$ , then maximum value of  $f(x)$  is :

A. 1

B.  $\frac{3}{2}$

C. 2

D.  $\frac{5}{2}$

**Answer: C**



**Watch Video Solution**

18. In  $\Delta ABC$ ,  $\angle C = \frac{2\pi}{3}$  then the value of  $\cos^2 A + \cos^2 B - \cos A \cdot \cos B$  is equal to :

A.  $\frac{3}{4}$

B.  $\frac{3}{2}$

C.  $\frac{1}{2}$

D.  $\frac{1}{4}$

**Answer: A**



**Watch Video Solution**

19. The number of solutions of the equation

$$4\sin^2 x + \tan^2 x + \cot^2 x + \operatorname{cosec}^2 x = 6 \text{ in } [0, 2\pi]:$$

A. 1

B. 2

C. 3

D. 4

**Answer: D**



[Watch Video Solution](#)

20. If  $\sin A$ ,  $\cos A$  and  $\tan A$  are in G.P, the  $\cos^3 A + \cos^2 A$  is equal to :

A. 1

B. 2

C. 4

D. none

**Answer: A**



**Watch Video Solution**

**21.** Range of function  $f(x) = \sin\left(x + \frac{\pi}{6}\right) + \cos\left(x - \frac{\pi}{6}\right)$  is :

A.  $[-\sqrt{2}, \sqrt{2}]$

B.  $[-\sqrt{2}(\sqrt{3} + 1), \sqrt{2}(\sqrt{3} + 1)]$

C.  $\left[-\frac{\sqrt{3} + 1}{\sqrt{2}}, \frac{\sqrt{3} + 1}{\sqrt{2}}\right]$

D.  $\left[-\frac{\sqrt{3} - 1}{\sqrt{2}}, \frac{\sqrt{3} - 1}{\sqrt{2}}\right]$

**Answer: C**



**Watch Video Solution**

**22.** The value of

$\tan(\log_2 6) \cdot \tan(\log_2 3) \cdot \tan 1$  is always equal to :

A.  $\tan(\log_2 6) + \tan(\log_2 3) + \tan 1$

B.  $\tan(\log_2 6) - \tan(\log_2 3) - \tan 1$

C.  $\tan(\log_2 6) - \tan(\log_2 3) + \tan 1$

D.  $\tan(\log_2 6) + \tan(\log_2 3) - \tan 1$

**Answer: B**



**Watch Video Solution**

**23.** In a triangle ABC, side  $BC = 3$ ,  $AC = 4$  and  $AB = 5$ . The value of  $\sin A + \sin 2B + \sin 3C$  is equal to :

A.  $\frac{24}{25}$

B.  $\frac{14}{25}$

C.  $\frac{64}{25}$

D. none

**Answer: B**



 [Watch Video Solution](#)

24.

If

$$A + B + C = 180^\circ, \text{ then } \frac{\cos A \cos C + \cos(A + B)\cos(B + C)}{\cos A \sin C - \sin(A + B)\cos(B + C)}$$

simplifies to :

A.  $-\cot C$

B. 0

C.  $\tan C$

D.  $\cot C$

**Answer: D**

 [Watch Video Solution](#)

25. If  $\alpha + \gamma = 2\beta$  then the expression  $\frac{\sin \alpha - \sin \gamma}{\cos \gamma - \cos \alpha}$  simplifies to:

A.  $\tan \beta$

B.  $-\tan \beta$

C.  $\cot \beta$

D.  $-\cot \beta$

**Answer: C**



**Watch Video Solution**

26. The product  $\left(\cos \frac{x}{2}\right) \cdot \left(\cos \frac{x}{4}\right) \cdot \left(\cos \frac{x}{8}\right) \cdot \dots \cdot \left(\cos \frac{x}{256}\right)$  is equal to :

A.  $\frac{\sin x}{128 \sin \frac{x}{256}}$

B.  $\frac{\sin x}{256 \sin \frac{x}{256}}$

C.  $\frac{\sin x}{128 \sin \frac{x}{128}}$

D.  $\frac{\sin x}{512 \sin \frac{x}{512}}$

**Answer: B**



**Watch Video Solution**

27. The value of the expression

$\frac{\sin 7\alpha + 6 \sin 5\alpha + 17 \sin 3\alpha + 12 \sin \alpha}{\sin 6\alpha + 5 \sin 4\alpha + 12 \sin 2\alpha}$ , where  $\alpha = \frac{\pi}{5}$  is equal to :

A.  $\frac{\sqrt{5} - 1}{4}$

B.  $\frac{\sqrt{5} + 1}{4}$

C.  $\frac{\sqrt{5} + 1}{2}$

D.  $\frac{\sqrt{5} - 1}{2}$

**Answer: C**



[Watch Video Solution](#)

28. In a triangle ABC if  $\sum \tan^2 A = \sum \tan A \tan B$ , then largest angle of the triangle in radian will be :

A.  $\frac{2\pi}{3}$

B.  $\frac{\pi}{3}$

C.  $\frac{\pi}{2}$

D.  $\frac{3\pi}{4}$

**Answer: B**



**Watch Video Solution**

**29.** Which one of the following values is not the solution of the equation

$$\log_{|\sin x|}(|\cos x|) + \log_{|\cos x|}(|\sin x|) = 2$$

A.  $\frac{7\pi}{4}$

B.  $\frac{11\pi}{4}$

C.  $\frac{3\pi}{4}$

D.  $\frac{3\pi}{8}$

**Answer: D**



**Watch Video Solution**

30. If  $f(x) = \sin^6 x + \cos^6 x$ , then range of  $f(x)$  is  $\left[\frac{1}{4}, 1\right]$  (b)  $\left[\frac{1}{4}, \frac{3}{4}\right]$

(c)  $\left[\frac{3}{4}, 1\right]$  (d) none of these

A.  $\left[\frac{1}{4}, 1\right]$

B.  $\left[\frac{1}{4}, \frac{3}{4}\right]$

C.  $\left[\frac{3}{4}, 1\right]$

D.  $[1, 2]$

**Answer: A**



**Watch Video Solution**

31. If  $\frac{2 \sin \alpha}{\{1 + \cos \alpha + \sin \alpha\}} = y$ , then  $\frac{\{1 - \cos \alpha + \sin \alpha\}}{1 + \sin \alpha} =$

A.  $\frac{1}{y}$

B.  $y$

C.  $1 - y$

D.  $1 + y$

**Answer: B**

 [Watch Video Solution](#)

32. If  $\frac{\tan^3 A}{1 + \tan^2 A} + \frac{\cot^3 A}{1 + \cot^2 A} = p \sec A \operatorname{cosec} A + q \sin A \cos A$ , then :

A.  $p = 2, q = 1$

B.  $p = 1, q = 2$

C.  $p = 1, q = -2$

D.  $p = 2, q = -1$

**Answer: C**

 [Watch Video Solution](#)

33. If  $\theta$  lies in the second quadrant. Then the value of

$\sqrt{\frac{1 - \sin \theta}{1 + \sin \theta}} + \sqrt{\frac{1 + \sin \theta}{1 - \sin \theta}}$  is equal to :

A.  $2 \sec \theta$

B.  $-2 \sec \theta$

C.  $2 \operatorname{cosec} \theta$

D. 2

**Answer: B**

 [Watch Video Solution](#)

34. If  $y = (\sin \theta + \operatorname{cosec} \theta)^2 + (\cos \theta + \sec \theta)^2$ , then minimum value of  $y$  is :

A. 7

B. 8

C. 9

D. none of these

**Answer: C**

 [Watch Video Solution](#)

35. If  $\log_3 \sin x - \log_3 \cos x - \log_3(1 - \tan x) - \log_3(1 + \tan x) = -1$ ,  
then  $\tan 2x$  is equal to (wherever defined)

A. -2

B.  $\frac{3}{2}$

C.  $\frac{2}{3}$

D. 6

**Answer: C**

 [Watch Video Solution](#)

36. If  $\sin \theta + \operatorname{cosec} \theta = 2$ , then the value of  $\sin^8 \theta + \operatorname{cosec}^8 \theta$  is equal to :

A. 2

B.  $2^4$



C.  $2^8$

D. more than  $2^8$

**Answer: A**

 [Watch Video Solution](#)

37. If  $\tan^3 \theta + \cot^3 \theta = 52$ , then the value of  $\tan^2 \theta + \cot^2 \theta$  is equal to :

A. 14

B. 15

C. 16

D. 17

**Answer: A**

 [Watch Video Solution](#)

38. The maximum value of  $(\log)_{20}(3 \sin x - 4 \cos x + 15)$  – a. 1 b. 2 c. 3  
d. 4

A. 1

B. 2

C. 3

D. 4

**Answer: A**



[Watch Video Solution](#)

39. If  $x^2 + y^2 = 9$  and  $4a^2 + 9b^2 = 16$ , then maximum value of  $4a^2x^2 + 9b^2y^2 - 12abxy$  is :

A. 81

B. 100

C. 121

**Answer: D**

 [Watch Video Solution](#)

40. If  $A = \sqrt{\sin 2 - \sin \sqrt{3}}$ ,  $B = \sqrt{\cos 2 - \cos \sqrt{3}}$ , then which of the following statement is true ?

A. A and B both are real numbers and  $A > B$

B. A and B both are real numbers and  $A < B$

C. Exactly one of A and B is not real number

D. Both A and B are not real numbers

**Answer: D**

 [Watch Video Solution](#)

41. The number of real values of  $x$  such that

$$(2^x + 2^{-x} - 2 \cos x)(3^{x+\pi} + 3^{-x-\pi} + 2 \cos x)(5^{\pi-x} + 5^{x-\pi} - 2 \cos x) =$$

is :

- A. 1
- B. 2
- C. 3
- D. infinite

**Answer: B**



[Watch Video Solution](#)

42. The equation  $e^{\sin x} - e^{-\sin x} - 4 = 0$  has

- A. infinite number of real roots
- B. no real roots
- C. exactly one real root

D. exactly four real roots

**Answer: B**



**Watch Video Solution**

43. If  $\pi < \alpha < \frac{3\pi}{2}$ , then find the value of expression  $\sqrt{4\sin^4 \alpha + \sin^2 2\alpha} + 4\cos^2\left(\frac{\pi}{4} - \frac{\alpha}{2}\right)$ .

A.  $2 + 4\sin \alpha$

B.  $2 - 4\cos \alpha$

C. 2

D.  $2 - 4\sin \alpha$

**Answer: C**



**Watch Video Solution**

44.  $\left(\cos\frac{\pi}{12} - \sin\frac{\pi}{12}\right)\left(\tan\frac{\pi}{12} + \cot\frac{\pi}{12}\right) =$

A.  $\frac{1}{\sqrt{2}}$

B.  $4\sqrt{2}$

C.  $\sqrt{2}$

D.  $2\sqrt{2}$

**Answer: D**



**Watch Video Solution**

45.  $\tan(100^\circ) + \tan(125^\circ) + \tan(100^\circ)\tan(125^\circ) =$

A. 0

B.  $\frac{1}{2}$

C. -1

D. 1

**Answer: D**



**Watch Video Solution**

**46.** If  $\sin x + \sin^2 = 1$ , then  $\cos^8 x + 2 \cos^6 x + \cos^4 x =$

A. 2

B. 1

C. 3

D.  $\frac{1}{2}$

**Answer: B**



**Watch Video Solution**

**47.** The maximum value of  $\log_5(3x + 4y)$ , if  $x^2 + y^2 = 25$  is :

A. 1

B. 2

C. 3

D. 4

**Answer: B**



**Watch Video Solution**

**48.** The number of values of  $\theta$  between  $-\pi$  and  $\frac{3\pi}{2}$  that satisfies the equation  $5 \cos 2\theta + 2\cos^2 \frac{\theta}{2} + 1 = 0$  is :

A. 3

B. 4

C. 5

D. 6

**Answer: C**



**Watch Video Solution**



49. Given that  $\sin \beta = \frac{4}{5}$ ,  $0 < \beta < \pi$  and  $\tan \beta > 0$ , then  $((3 \sin(\alpha + \beta) - 4 \cos(\alpha + \beta))$  is equal to :

- A. 2
- B. 3
- C. 4
- D. 5

**Answer: D**

 [Watch Video Solution](#)

50. The maximum value of  $\sin\left(x + \frac{\pi}{5}\right) + \cos\left(x + \frac{\pi}{5}\right)$ , where  $x \in \left(0, \frac{\pi}{2}\right)$ , is attained at

- A.  $\frac{\pi}{12}$

B.  $\frac{\pi}{6}$

C.  $\frac{\pi}{3}$

D.  $\frac{\pi}{20}$

**Answer: A**



**Watch Video Solution**

51. The value of 'a' for which the equation  $\sin x(\sin x + \cos x) = a$  has a real solution are

A.  $1 - \sqrt{2} \leq a \leq 1 + \sqrt{2}$

B.  $2 - \sqrt{3} \leq a \leq 2 + \sqrt{3}$

C.  $0 \leq a \leq 2 + \sqrt{3}$

D.  $\frac{1 - \sqrt{2}}{2} \leq a \leq \frac{1 + \sqrt{2}}{2}$

**Answer: D**



**Watch Video Solution**

52. The value of  $\cos 12^\circ \cos 24^\circ \cos 36^\circ \cos 48^\circ \cos 60^\circ \cos 72^\circ \cos 84^\circ$  is :

A.  $\frac{1}{64}$

B.  $\frac{1}{128}$

C.  $\frac{1}{256}$

D.  $\frac{1}{512}$

**Answer: B**



[Watch Video Solution](#)

53. The ratio of the maximum value to minimum value of  $2 \cos^2 \theta + \cos \theta + 1$  is :

A. 32 : 7

B. 32 : 9

C. 4 : 1

D. 2:1

**Answer: A**



[Watch Video Solution](#)

54. If all values of  $x \in (a, b)$  satisfy the inequality  $\tan x \tan 3x < -1$ ,  $x \in \left(0, \frac{\pi}{2}\right)$ , then the maximum value (b - a) is :

A.  $\frac{\pi}{12}$

B.  $\frac{\pi}{3}$

C.  $\frac{\pi}{6}$

D.  $\frac{\pi}{4}$

**Answer: A**



[Watch Video Solution](#)

55. If a regular polygon of  $n$  sides has circum radius  $R$  and inradius  $r$  then each side of polygon is:

A.  $(R + r)\tan\left(\frac{\pi}{2n}\right)$

B.  $2(R + r)\tan\left(\frac{\pi}{2n}\right)$

C.  $(R + r)\sin\left(\frac{\pi}{2n}\right)$

D.  $2(R + r)\cot\left(\frac{\pi}{2n}\right)$

**Answer: B**



**Watch Video Solution**

56. Find the value of  $\cos 12^{\circ} + \cos 84^{\circ} + \cos 156^{\circ} + \cos 132^{\circ}$

A.  $\frac{1}{8}$

B.  $-\frac{1}{2}$

C. 1

D.  $\frac{1}{2}$

**Answer: B**

 [Watch Video Solution](#)

$$57. \frac{\sin \theta}{\cos(3\theta)} + \frac{\sin(3\theta)}{\cos(9\theta)} + \frac{\sin(9\theta)}{\cos(27\theta)} + \frac{\sin(27\theta)}{\cos(81\theta)} =$$

A.  $\frac{\sin(81\theta)}{2 \cos(80\theta) \cos \theta}$

B.  $\frac{\sin(80\theta)}{2 \cos(81\theta) \cos \theta}$

C.  $\frac{\sin(81\theta)}{\cos(80\theta) \cos \theta}$

D.  $\frac{\sin(80\theta)}{2 \cos(81\theta) \cos \theta}$

**Answer: B**

 [Watch Video Solution](#)

58. The value of  $\left(\sin \frac{\pi}{9}\right) \left(4 + \sec \frac{\pi}{9}\right)$  is :

A.  $\frac{1}{2}$

B.  $\sqrt{2}$

C. 1

D.  $\sqrt{3}$

**Answer: D**

 [Watch Video Solution](#)

59. If  $\frac{dy}{dx} = \sin\left(\frac{x\pi}{2}\right)\cos(x\pi)$ , then y is strictly increasing in :

A. (3, 4)

B.  $\left(\frac{5}{2}, \frac{7}{2}\right)$

C. (2, 3)

D.  $\left(\frac{1}{2}, \frac{3}{2}\right)$

**Answer: B**

 [Watch Video Solution](#)

60. Smallest positive value of  $\theta$  satisfying

$$8 \sin \theta \cos 2\theta \sin 3\theta \cos 4\theta = \cos 6\theta$$
 is

A.  $\frac{\pi}{18}$

B.  $\frac{\pi}{22}$

C.  $\frac{\pi}{24}$

D. none of these

**Answer: A**



[Watch Video Solution](#)

61. If an angle  $A$  of a triangle  $ABC$  is given by  $3 \tan A + 1 = 0$ , then  $\sin A$  and  $\cos A$  are the roots of the equation

A.  $10x^2 - 2\sqrt{10}x + 3 = 0$

B.  $10x^2 - 2\sqrt{10}x - 3 = 0$

C.  $10x^2 + 2\sqrt{10}x + 3 = 0$



D.  $10x^2 + 2\sqrt{10}x - 3 = 0$

**Answer: D**



**Watch Video Solution**

62. If  $\tan \theta = \frac{1}{\sqrt{7}}$ , find the value of  $\frac{\operatorname{cosec}^2 \theta - \sec^2 \theta}{\operatorname{cosec}^2 \theta + \sec^2 \theta}$ .

A.  $3/4$

B.  $1/2$

C.  $2$

D.  $5/4$

**Answer: A**



**Watch Video Solution**

63. If  $2 \cos \theta + \sin \theta = 1$  then  $7 \cos \theta + 6 \sin \theta$  is equal to

A. 1 or 2

B. 2 or 3

C. 2 or 4

D. 2 or 6

**Answer: D**



[Watch Video Solution](#)

**64.** If  $\sin \theta + \operatorname{cosec} \theta = 2$ , then the value of  $\sin^8 \theta + \operatorname{cosec}^8 \theta$  is equal to :

A. 2

B.  $2^4$

C.  $2^8$

D. more than  $2^8$

**Answer: A**



[Watch Video Solution](#)

65. If  $\tan^3 \theta + \cot^3 \theta = 52$ , then the value of  $\tan^2 \theta + \cot^2 \theta$  is equal to :

A. 14

B. 15

C. 16

D. 17

**Answer: A**



[Watch Video Solution](#)

66. If ABCD is a cyclic quadrilateral such that  $12 \tan A - 5 = 0$  and

$5 \cos B + 3 = 0$  then  $\tan C + \tan D$  is equal to :

A.  $\frac{21}{12}$

B.  $\frac{11}{12}$

C.  $-\frac{11}{12}$

D.  $-\frac{21}{12}$

**Answer: B**



**Watch Video Solution**

67. If  $\frac{\pi}{2} < \theta < \frac{3\pi}{2}$  then  $\sqrt{\tan^2 \theta - \sin^2 \theta}$  is equal to :

A.  $\tan \theta \sin \theta$

B.  $-\tan \theta \sin \theta$

C.  $\tan \theta - \sin \theta$

D.  $\sin \theta - \tan \theta$

**Answer: B**



**Watch Video Solution**

68. The value of  $\frac{\sin 10^\circ + \sin 20^\circ}{\cos 10^\circ + \cos 20^\circ}$  equals

A.  $2 + \sqrt{3}$

B.  $\sqrt{2} - 1$

C.  $2 - \sqrt{3}$

D.  $\sqrt{2} + 1$

**Answer: C**

 [Watch Video Solution](#)

69. The value of the expression  $\cos^6 \theta + \sin^6 \theta + 3 \sin^2 \theta \cos^2 \theta =$

A. 0

B. 1

C. 2

D. 3

**Answer: B**

 [Watch Video Solution](#)

70.  $\frac{\sin x + \cos x}{\sin x - \cos x} - \frac{\sec^2 x + 2}{\tan^2 x - 1} =$ , where  $x \in \left(0, \frac{\pi}{2}\right)$

A.  $\frac{1}{\tan x + 1}$

B.  $\frac{2}{1 + \tan x}$

C.  $\frac{2}{1 + \cot x}$

D.  $\frac{2}{1 - \tan x}$

**Answer: B**



**Watch Video Solution**

71. If  $\frac{\cot \alpha + \cot(270^\circ + \alpha)}{\cot \alpha - \cot(270^\circ + \alpha)} - 2 \cos(135^\circ + \alpha) \cos(315^\circ - \alpha) = \lambda$ ,

where  $\alpha \in \left(0, \frac{\pi}{2}\right)$ , then  $\lambda =$

A. 0

B. 1

C. 2

D. 4

**Answer: C**



**Watch Video Solution**

72. The expression  $\frac{\sin \alpha + \cos \alpha}{\cos \alpha - \sin \alpha} \tan\left(\frac{\pi}{4} + \alpha\right) + 1, \alpha \in \left(-\frac{\pi}{4}, \frac{\pi}{4}\right)$

simplifies to :

A.  $\operatorname{cosec}^2\left(\frac{\pi}{4} - \alpha\right)$

B.  $\sec^2\left(\frac{\pi}{4} - \alpha\right)$

C.  $\tan^2\left(\frac{\pi}{4} - \alpha\right)$

D.  $\cot^2\left(\frac{\pi}{4} - \alpha\right)$

**Answer: A**



**Watch Video Solution**

73. The value of expression  $\frac{\tan \alpha + \sin \alpha}{2\cos^2 \frac{\alpha}{2}}$  for  $\alpha = \frac{\pi}{4}$  is :

A. 4

B. 3

C. 2

D. 1

**Answer: D**



**Watch Video Solution**

74.  $\cos 2\alpha - \cos 3\alpha - \cos 4\alpha + \cos 5\alpha$  simplifies to :

A.  $-4\sin \frac{\alpha}{2} \sin \alpha \cos \frac{7\alpha}{2}$

B.  $4\sin \frac{\alpha}{2} \sin \alpha \cos \frac{7\alpha}{2}$

C.  $-4\sin \frac{\alpha}{2} \sin \frac{7\alpha}{2} \cos \alpha$

D.  $-4\sin \alpha \cos \frac{\alpha}{2} \sin \frac{7\alpha}{2}$



**Answer: A**



**Watch Video Solution**

75. If  $\tan \gamma = \sec \alpha \sec \beta + \tan \alpha \tan \beta$ , then  $\cos 2\gamma$  is necessarily

A. -1

B.  $\frac{1}{2}$

C.  $-\frac{1}{2}$

D. 0

**Answer: D**



**Watch Video Solution**

76. If  $\operatorname{cosec} x = \frac{2}{\sqrt{3}}$ ,  $\cot x = -\frac{1}{\sqrt{3}}$ ,  $x \in [0, 2\pi]$ , then

$\cos x + \cos 2x + \cos 3x + \dots + \cos 100x =$

A.  $\frac{1}{2}$

B.  $-\frac{1}{2}$

C.  $-\frac{\sqrt{3}}{2}$

D.  $\frac{\sqrt{3}}{2}$

**Answer: B**



**Watch Video Solution**

77. The value of  $\sum_{r=0}^{10} \cos^3 \frac{r\pi}{3}$  is equal to

A.  $-\frac{7}{8}$

B.  $-\frac{9}{8}$

C.  $-\frac{3}{8}$

D.  $-\frac{1}{8}$

**Answer: D**



**Watch Video Solution**

78. about to only mathematics

A. 1

B. 2

C.  $\sqrt{3}$

D.  $\frac{\sqrt{3}}{2}$

**Answer: A**



**Watch Video Solution**

79. If  $x, y \in R$  and satisfy  $(x + 5)^2 + (y - 12)^2 = 14^2$ , then the minimum value of  $x^2 + y^2$  is :

A. 2

B. 1

C.  $\sqrt{3}$

D.  $\sqrt{2}$

**Answer: B**

 [Watch Video Solution](#)

80. If  $\theta_1, \theta_2, \theta_3$  are three values lying in  $[0, 3\pi)$  for which  $\tan \theta = \lambda$ , then

the value of

$$\left| \tan\left(\frac{\theta_1}{3}\right)\tan\left(\frac{\theta_2}{3}\right) + \tan\left(\frac{\theta_2}{3}\right)\tan\left(\frac{\theta_3}{3}\right) + \tan\left(\frac{\theta_3}{3}\right)\tan\left(\frac{\theta_1}{3}\right) \right| \text{ is}$$

A. -3

B. -2

C. 2

D. 3

**Answer: A**

 [Watch Video Solution](#)

81.

If

$\tan \alpha = \frac{b}{a}$ ,  $a > b > 0$  and if  $0 < \alpha < \frac{\pi}{4}$ , then  $\sqrt{\frac{a+b}{a-b}} + \sqrt{\frac{a-b}{a+b}}$

is equal to :

- A.  $\frac{2 \sin \alpha}{\sqrt{2\alpha}}$
- B.  $\frac{2 \cos \alpha}{\sqrt{\cos 2\alpha}}$
- C.  $\frac{2 \sin \alpha}{\sqrt{\sin 2\alpha}}$
- D.  $\frac{2 \cos \alpha}{\sqrt{\sin 2\alpha}}$

**Answer: B**



[Watch Video Solution](#)

82. Minimum value of  $3 \sin \theta + 4 \cos \theta$  in the interval  $\left[0, \frac{\pi}{2}\right]$  is : (a) -5 (b)

3 (c) 4 (d)  $\frac{7}{\sqrt{2}}$

A. -5

B. 3

C. 4

D.  $\frac{7}{\sqrt{2}}$

**Answer: B**



**Watch Video Solution**

83. If  $f(n) = \prod_{r=1}^n \cos r$ ,  $n \in \mathbb{N}$ , then

A.  $|f(n)| > |f(n + 1)|$

B.  $f(5) > 0$

C.  $f(4) > 0$

D.  $|f(n)| < |f(n + 1)|$

**Answer: A**



**Watch Video Solution**

84. If  $\tan A + \sin A = p$  and  $\tan A - \sin A = q$ , then the value of  $\frac{(p^2 - q^2)^2}{pq}$  is :

A. 16

B. 22

C. 18

D. 42

Answer: A



Watch Video Solution

85.

Let

$$t_1 = (\sin \alpha)^{\cos \alpha}, t_2 = (\sin \alpha)^{\sin \alpha}, t_3 = (\cos \alpha)^{\cos \alpha}, t_4 = (\cos \alpha)^{\sin \alpha},$$

where  $\alpha \in \left(0, \frac{\pi}{4}\right)$ , then which of the following is correct

A.  $t_3 > t_1 > t_2$

B.  $t_4 > t_2 > t_1$

C.  $t_4 > t_1 > t_2$

D.  $t_1 > t_3 > t_2$

**Answer: B**



**Watch Video Solution**

86. If  $\cos A = 3/4$ , then  $32 \sin(A/2) \sin((5A)/2) = \text{-----}$  (A)  $\sqrt{11}$  (B)  $-\sqrt{11}$  (C) 11

(D) -11

A. 11

B. -11

C. 12

D. 4

**Answer: A**



**Watch Video Solution**



87.  $\cos(\alpha + \beta) + \sin(\alpha - \beta) = 0$  and  $\tan \beta = \frac{1}{2009}$  then value of  $\tan 3\alpha$

A. 2

B. 1

C. 3

D. 4

**Answer: B**



[Watch Video Solution](#)

88. If  $2^x = 3^y = 6^{-z}$ , the value of  $\frac{1}{x} + \frac{1}{y} + \frac{1}{z}$  is equal to :

A. 0

B. 1

C. 2

D. 3

Answer: A



Watch Video Solution

89. Let  $\alpha$  and  $\beta$  be such that  $\pi < \alpha - \beta < 3\pi$ , If  $\sin \alpha + \sin \beta = -\frac{21}{65}$  and  $\cos \alpha + \cos \beta = -\frac{27}{65}$ , then the value of  $\frac{\cos(\alpha - \beta)}{2}$  is (a)  $-\frac{3}{\sqrt{130}}$  (b)  $\frac{3}{\sqrt{130}}$  (c)  $\frac{6}{25}$  (d)  $\frac{6}{65}$

A.  $\frac{-3}{\sqrt{130}}$

B.  $\frac{3}{\sqrt{130}}$

C.  $\frac{6}{65}$

D.  $-\frac{6}{65}$

Answer: A



Watch Video Solution

90. If  $u = \sqrt{a^2 \cos^2 \theta + b^2 \sin^2 \theta} + \sqrt{a^2 \sin^2 \theta + b^2 \cos^2 \theta}$ , then the difference between the maximum and minimum values of  $u^2$  is given by :

(a)  $(a - b)^2$  (b)  $2\sqrt{a^2 + b^2}$  (c)  $(a + b)^2$  (d)  $2(a^2 + b^2)$

A.  $2(a^2 + b^2)$

B.  $(a + b)^2$

C.  $2\sqrt{a^2 + b^2}$

D.  $(a - b)^2$

**Answer: D**



**Watch Video Solution**

91. If  $P = (\tan(3^{n+1}\theta) - \tan \theta)$  and  $Q = \sum_{r=0}^n \frac{\sin(3^r \theta)}{\cos(3^{r+1} \theta)}$ , then

A.  $P = 2Q$

B.  $P = 3Q$

C.  $2P = Q$

D.  $3P = Q$

**Answer: A**



**Watch Video Solution**

92. If  $270^\circ < \theta < 360^\circ$ , then find  $\sqrt{2 + \sqrt{2(1 + \cos \theta)}}$

A.  $-2 \sin\left(\frac{\theta}{4}\right)$

B.  $2 \sin\left(\frac{\theta}{4}\right)$

C.  $\pm 2 \sin \frac{\theta}{4}$

D.  $2 \cos \frac{\theta}{4}$

**Answer: B**



**Watch Video Solution**

93. If  $y = (\sin x + \cos x) + (\sin 4x + \cos 4x)^2$ , then :

A.  $y > 0 \forall x \in R$

B.  $y \geq 0 \forall x \in R$

C.  $y < 2 + \sqrt{2} \forall x \in R$

D.  $y = 2 + \sqrt{2}$  for some  $x \in R$

**Answer: C**



**Watch Video Solution**

**94.**

**If**

$\cos x + \cos y + \cos z = \sin x + \sin y + \sin z = 0$  then  $\cos(x - y) =$

(a) 0 (b)  $-\frac{1}{2}$  (c) 2 (d) 1

A. 0

B.  $-\frac{1}{2}$

C. 2

D. 1

**Answer: B**



**Watch Video Solution**

95. The value of  $\operatorname{cosec}10^\circ + \operatorname{cosec}50^\circ - \operatorname{cosec}70^\circ$  is \_\_\_\_\_

A. 4

B. 5

C. 6

D. 8

**Answer: C**



**Watch Video Solution**

96. If  $270^\circ < \theta < 360^\circ$ , then find  $\sqrt{2 + \sqrt{2(1 + \cos \theta)}}$

A.  $-2 \sin\left(\frac{\theta}{4}\right)$

B.  $2 \sin\left(\frac{\theta}{4}\right)$

C.  $\pm 2 \sin \frac{\theta}{4}$

D.  $2 \cos \frac{\theta}{4}$

**Answer: B**



**Watch Video Solution**

### Exercise 2 One Or More Than One Answer Is Are Correct

1.  $\cot 12^\circ \cdot \cot 24^\circ \cdot \cot 28^\circ \cdot \cot 32^\circ \cdot \cot 48^\circ \cdot \cot 88^\circ = \dots\dots\dots$

A.  $\tan 45^\circ$

B. 2

C.  $2 \tan 15^\circ \cdot \tan 45^\circ \cdot \tan 75^\circ$

D.  $\tan 15^\circ \cdot \tan 45^\circ \cdot \tan 75^\circ$

**Answer: A:D**



Watch Video Solution

2. If the equation  $\cot^4 x - 2 \operatorname{cosec}^2 x + a^2 = 0$  has at least one solution, then the sum of all possible integral values of  $a$  is equal to 4 (b) 3 (c) 2 (d) 0

A. -1

B. 0

C. 1

D. 2

Answer: A::B::C



Watch Video Solution

3. Which of the following is/are true ?

A.  $\tan 1 > \tan^{-1} 1$



B.  $\sin 1 > \cos 1$

C.  $\tan 1 < \sin 1$

D.  $\cos(\cos 1) > \frac{1}{\sqrt{2}}$

**Answer: A::B::D**



**Watch Video Solution**

**4. Which of the following is/are true ?**

A.  $\log_{\sin 1} \tan 1$

B.  $\log_{\cos 1} (1 + \tan 3)$

C.  $\log_{\log_{10} 5} (\cos \theta + \sec \theta)$

D.  $\log_{\tan 15^\circ} (2 \sin 18^\circ)$

**Answer: B::D**



**Watch Video Solution**

5. If  $\sin \alpha + \cos \alpha = \frac{\sqrt{3} + 1}{2}$ ,  $0 < \alpha < 2\pi$ , then possible values  $\tan \frac{\alpha}{2}$  can take is/are :

A.  $2 - \sqrt{3}$

B.  $\frac{1}{\sqrt{3}}$

C. 1

D.  $\sqrt{3}$

**Answer: A:B**



**Watch Video Solution**

6. If  $3 \sin \beta = \sin(2\alpha + \beta)$ , then

A.  $(\cot \alpha + \cot(\alpha + \beta))(\cot \beta - 3 \cot(2\alpha - \beta)) = 6$

B.  $\sin \beta = \cos(\alpha + \beta) \sin \alpha$

C.  $\tan(\alpha + \beta) = 2 \tan \alpha$

D.  $2 \sin \beta = \sin(\alpha + \beta) \cos \alpha$

Answer: A::B::C::D



Watch Video Solution

7. If  $\sin(x + 20^\circ) = 2 \sin x \cos 40^\circ$ , where  $x \in (0, \frac{\pi}{2})$ , then which of the following hold(s) good?  $\cos 2x = \frac{1}{2}$  (b)  $\operatorname{cosec} 4x = 2 \frac{\sec x}{2} = \sqrt{6} - \sqrt{2}$   
(d)  $\frac{\tan x}{2} = (2 - \sqrt{3})$

A.  $\sec \frac{x}{2} = \sqrt{6} - \sqrt{2}$

B.  $\cot \frac{x}{2} = 2 + \sqrt{3}$

C.  $\tan 4x = \sqrt{3}$

D.  $\operatorname{cosec} 4x = 2$

Answer: A::B



Watch Video Solution

8. If  $\cos x + \cos y + \cos z = 0$  and  $\sin x + \sin y + \sin z = 0$ , then show that

$$\cos(x - y) + \cos(y - z) + \cos(z - x) = -\frac{3}{2}.$$

A.  $\cos x \cos y \cos z = 1$

B.  $\cos x + \cos y + \cos z = 0$

C.  $\sin x + \sin y + \sin z = 1$

D.  $\cos 3x + \cos 3y + \cos 3z = 12 \cos x \cos y \cos z$

**Answer: B::D**



**Watch Video Solution**

9. If  $0 < x < \frac{\pi}{2}$  and  $\sin^n x + \cos^n x \geq 1$ , then

A.  $[1, 2]$

B.  $[3, 4]$

C.  $[-\infty, 2]$

D.  $[-1, 1]$

**Answer: A::C::D**



**Watch Video Solution**

10.

If

$$x = \sin(\alpha - \beta) \cdot \sin(\gamma - \delta), y = \sin(\beta - \gamma) \cdot \sin(\alpha - \delta), z = \sin(\gamma - \alpha) \cdot \sin(\beta - \delta)$$

, then :

A.  $x + y + z = 0$

B.  $x^3 + y^3 + z^3 = 3xyz$

C.  $x + y - z = 0$

D.  $x^3 + y^3 - z^3 = 3xyz$

**Answer: A::B**



**Watch Video Solution**

11.

If

$$X = x \cos \theta - y \sin \theta, Y = x \sin \theta + y \cos \theta \text{ and } X^2 + 4XY + Y^2 = Ax^2 -$$

, then :

(where A and B are constants)

A.  $\theta = \frac{\pi}{6}$

B.  $\theta = \frac{\pi}{4}$

C.  $A = 3$

D.  $B = -1$

**Answer: B::C::D**



[Watch Video Solution](#)

12. If  $2a = 2\tan 10^\circ + \tan 50^\circ$ ,  $2b = \tan 20^\circ + \tan 50^\circ$

$2c = 2\tan 10^\circ + \tan 70^\circ$ ,  $2d = \tan 20^\circ + \tan 70^\circ$

Then which of the following is/are correct ?

A.  $a + d = b + c$

B.  $a + b = c$

C.  $a > b < c > d$

D.  $a < b < c < d$

**Answer: A::B::D**



**Watch Video Solution**

13. Which of the following real numbers when simplified are neither terminating nor repeating decimal ?

A.  $\sin 75^\circ \cdot \cos 75^\circ$

B.  $\log_2 28$

C.  $\log_3 5 \cdot \log_5 6$

D.  $8^{-(\log_{27} 3)}$

**Answer: B::C**



**Watch Video Solution**

14. If  $\alpha = \sin x \cos^3 x$  and  $\beta = \cos x \sin^3 x$ , then :

A.  $\alpha - \beta > 0$ , for all  $x$  in  $\left(0, \frac{\pi}{4}\right)$

B.  $\alpha - \beta < 0$ , for all  $x$  in  $\left(0, \frac{\pi}{4}\right)$

C.  $\alpha + \beta > 0$ , for all  $x$  in  $\left(0, \frac{\pi}{2}\right)$

D.  $\alpha + \beta < 0$ , for all  $x$  in  $\left(0, \frac{\pi}{2}\right)$

**Answer: A:C**



**Watch Video Solution**

15. If  $\frac{\pi}{2} < \theta < \pi$ , then possible answers of  $\sqrt{2 + \sqrt{2 + 2 \cos 4\theta}}$  is/are :

A.  $2 \cos \theta$

B.  $2 \sin \theta$

C.  $-2 \sin \theta$

D.  $-2 \cos \theta$



**Answer: B::D**



**Watch Video Solution**

**16.** If  $\cot^3 \alpha + \cot^2 \alpha + \cot \alpha = 1$  then which of the following is/are correct

A.  $\cos 2\alpha \tan \alpha = 1$

B.  $\cos 2\alpha \cdot \tan \alpha = -1$

C.  $\cos 2\alpha - \tan 2\alpha = -1$

D.  $\cos 2\alpha - \tan 2\alpha = 1$

**Answer: B::D**



**Watch Video Solution**

**17.** The value of  $x$  in  $\left(0, \frac{\pi}{2}\right)$  satisfying

$\frac{\sqrt{3}-1}{\sin x} + \frac{\sqrt{3}+1}{\cos x} = 4\sqrt{2}$  is/are  $\frac{\pi}{12}$  (b)  $\frac{5\pi}{12}$  (c)  $\frac{7\pi}{24}$  (d)  $\frac{11\pi}{36}$

A.  $\frac{\pi}{15}$

B.  $\frac{\pi}{12}$

C.  $\frac{11\pi}{36}$

D.  $\frac{3\pi}{10}$

**Answer: B::C**



**Watch Video Solution**

18. If  $\alpha > \frac{1}{\sin^6 x + \cos^6 x} \forall x \in R$ , then  $\alpha$  can be

A. 3

B. 4

C. 5

D. 6

**Answer: B::D**



**Watch Video Solution**

19. If  $x \in \left(0, \frac{\pi}{2}\right)$  and  $\sin x = \frac{3}{\sqrt{10}}$ ,

Let  $k = \log_{10} \sin x + \log_{10} \cos x + 2\log_{10} \cot x + \log_{10} \tan x$  then the value of  $k$  satisfies

A.  $k = 0$

B.  $k + 1 = 0$

C.  $k - 1 = 0$

D.  $k^2 - 1 = 0$

**Answer: B::D**



**Watch Video Solution**

20. If  $A, B, C$  are angles of a triangle  $ABC$  and  $\tan A \tan C = 3, \tan B \tan C = 6$  then which is (are) correct :

A.  $A = \frac{\pi}{4}$

B.  $\tan A \tan B = 2$

C.  $\frac{\tan A}{\tan C} = 3$

D.  $\tan B = 2 \tan A$

**Answer: A::B::D**



**Watch Video Solution**

21. The value of  $\frac{\sin x - \cos x}{\sin^3 x}$  is equal to :

A.  $\operatorname{cosec}^2 x (1 - \cot x)$

B.  $1 - \cot x + \cot^2 x - \cot^3 x$

C.  $\operatorname{cosec}^2 x - \cot x - \cot^3 x$

D.  $\frac{1 - \cot x}{\sin^2 x}$

**Answer: A::B::C::D**



**Watch Video Solution**

22. If  $f(x) = \sin^2 x + \sin^2\left(x + \frac{2\pi}{3}\right) + \sin^2\left(x + \frac{4\pi}{3}\right)$  then :

A.  $f\left(\frac{\pi}{15}\right) = \frac{3}{2}$

B.  $f\left(\frac{15}{\pi}\right) = \frac{2}{3}$

C.  $f\left(\frac{\pi}{10}\right) = \frac{3}{2}$

D.  $f\left(\frac{10}{\pi}\right) = \frac{2}{3}$

Answer: A:C



Watch Video Solution

23. The range of  $y = \frac{\sin 4x - \sin 2x}{\sin 4x + \sin 2x}$  satisfies

A.  $y \in \left(-\infty, \frac{1}{3}\right)$

B.  $y \in \left(\frac{1}{3}, 1\right)$

C.  $y \in (1, 3)$

D.  $y \in (3, \infty)$

Answer: A::D



Watch Video Solution

24.

If

$\sqrt{2} \cos A = \cos B + \cos^3 B$ , and  $\sqrt{2} \sin A = \sin B - \sin^3 B$  then  $\sin(A - B)$

$\pm 1$  (b)  $\pm \frac{1}{2}$  (c)  $\pm \frac{1}{3}$  (d)  $\pm \frac{1}{4}$

A.  $\frac{1}{2}$

B.  $\frac{1}{3}$

C.  $-\frac{1}{2}$

D.  $-\frac{1}{3}$

Answer: B::D



Watch Video Solution

25. If  $\alpha > \frac{1}{\sin^6 x + \cos^6 x} \forall x \in R$ , then  $\alpha$  can be

A. 3

B. 4

C. 5

D. 6

**Answer: C::D**



**Watch Video Solution**

**26.** If  $\cot^3 \alpha + \cot^2 \alpha + \cot \alpha = 1$  then which of the following is/are correct

A.  $\cos 2\alpha \tan \alpha = 1$

B.  $\cos 2\alpha \cdot \tan \alpha = -1$

C.  $\cos 2\alpha - \tan 2\alpha = -1$

D.  $\cos 2\alpha - \tan 2\alpha = 1$

**Answer: B::D**

 [Watch Video Solution](#)

### Exercise 3 Comprehension Type Problems

1. Let  $l = \sin \theta$ ,  $m = \cos \theta$  and  $n = \tan \theta$ .

If  $\theta = 5$  radian, then :

A.  $l > m$

B.  $l < m$

C.  $l = m$

D. none of these

**Answer: B**

 [Watch Video Solution](#)

2. Let  $l = \sin \theta$ ,  $m = \cos \theta$  and  $n = \tan \theta$ .

Q. If  $\theta = -1042^\circ$ , then : (a)  $n > 1$  (b)  $n < 1$  (c)  $n = 1$  (d) nothing can be



said

A.  $n > 1$

B.  $n < 1$

C.  $n = 1$

D. nothing can be said

**Answer: B**



[Watch Video Solution](#)

3. Let  $l = \sin \theta$ ,  $m = \cos \theta$  and  $n = \tan \theta$ .

Q. If  $\theta = 7$  radian, then : (a)  $l + m > 0$  (b)  $l + m < 0$  (c)  $l + m = 0$  (d)

nothing can be said

A.  $l + m > 0$

B.  $l + m < 0$

C.  $l + m = 0$

D. nothing can be said

**Answer: A**



[Watch Video Solution](#)

4. Let  $a, b, c$  are respectively the sines and  $p, q, r$  are respectively the cosines of  $\alpha, \alpha + \frac{2\pi}{3}$  and  $\alpha + \frac{4\pi}{3}$ , then :

Q. The value of  $(a + b + c)$  is :

A. 0

B.  $\frac{3}{4}$

C. 1

D. none of these

**Answer: A**



[Watch Video Solution](#)

5. Let  $a, b, c$  are respectively the sines and  $p, q, r$  are respectively the cosines of  $\alpha, \alpha + \frac{2\pi}{3}$  and  $\alpha + \frac{4\pi}{3}$ , then :

Q. The value of  $(ab + bc + ca)$  is :

A. 0

B.  $-\frac{3}{4}$

C.  $-\frac{1}{2}$

D. -1

**Answer: B**



[Watch Video Solution](#)

6. Let  $a, b, c$  are respectively the sines and  $p, q, r$  are respectively the cosines of  $\alpha, \alpha + \frac{2\pi}{3}$  and  $\alpha + \frac{4\pi}{3}$ , then :

Q. The value of  $(qc - rb)$  is :

A. 0

B.  $-\frac{\sqrt{3}}{2}$

C.  $\frac{\sqrt{3}}{2}$

D. depends on  $\alpha$

**Answer: C**



**Watch Video Solution**

7. Consider a right angle triangle ABC right angle at B such that  $AC = \sqrt{8 + 4\sqrt{3}}$  and  $AB = 1$ . A line through vertex A meet BC at D such that  $AD = BC$ . An arc DE of radius AD is drawn from vertex A to meet AC at E and another arc DF of radius CD is drawn from vertex C to meet AC at F. On the basis of above information, answer the following questions.

Q.  $\sqrt{\tan A + \cot C}$  is equal to :

A.  $\sqrt{3}$

B. 1

C.  $2 + \sqrt{3}$

D.  $\sqrt{3} + 1$

**Answer: D**



**Watch Video Solution**

8. Consider a right angle triangle ABC right angle at B such that  $AC = \sqrt{8 + 4\sqrt{3}}$  and  $AB = 1$ . A line through vertex A meet BC at D such that  $AD = DC$ . An arc DE of radius AD is drawn from vertex A to meet AC at E and another arc DF of radius CD is drawn from vertex C to meet AC at F. On the basis of above information, answer the following questions.

Q.  $\log AE \left( \frac{AE + CF}{CD} \right)$  is equal to :

A.  $\sqrt{2}$

B. 1

C. 0

D. -1

**Answer: B**



[Watch Video Solution](#)

9. In  $\Delta ABC$ , if  $\cot \theta = \cot A + \cot B + \cot C$ , prove that

$$\sin^3 \theta = \sin(A - \theta)\sin(B - \theta)\sin(C - \theta)$$

A.  $60^\circ$

B.  $25^\circ$

C.  $35^\circ$

D.  $45^\circ$

**Answer: B**



[Watch Video Solution](#)

10. In  $\Delta ABC$ , if  $\cot \theta = \cot A + \cot B + \cot C$ , prove that

$$\sin^3 \theta = \sin(A - \theta)\sin(B - \theta)\sin(C - \theta)$$

A.  $\tan^3 \theta$

B.  $\cot^3 \theta$

C.  $\sin^3 \theta$

D.  $\cos^3 \theta$

**Answer: C**

 [Watch Video Solution](#)

11. Consider the function  $f(x) = \frac{\sqrt{1 + \cos x} + \sqrt{1 - \cos x}}{\sqrt{1 + \cos x} - \sqrt{1 - \cos x}}$  then

Q. If  $x \in (\pi, 2\pi)$  then  $f(x)$  is

A.  $\cot\left(\frac{\pi}{2} + \frac{x}{2}\right)$

B.  $\tan\left(\frac{\pi}{4} + \frac{x}{2}\right)$

C.  $\cot\left(\frac{\pi}{4} - \frac{x}{2}\right)$

D.  $\tan\left(\frac{\pi}{4} - \frac{x}{2}\right)$

**Answer: D**



Watch Video Solution

12. If the function  $f(x) = \frac{\sqrt{1 + \cos x} + \sqrt{1 - \cos x}}{\sqrt{1 + \cos x} - \sqrt{1 - \cos x}}$  if the value of  $f\left(\frac{\pi}{3}\right) = a + b\sqrt{c}$  then  $a + b + c =$

A. 4

B. 5

C. 6

D. 7

Answer: C



Watch Video Solution

Exercise 4 Matching Type Problems



Column-I		Column-II	
(A)	The value of $\frac{\cos 68^\circ}{\sin 56^\circ \sin 34^\circ \tan 22^\circ}$ equals to	(P)	16
(B)	The value of $(\cos 65^\circ + \sqrt{3} \sin 5^\circ + \cos 5^\circ)^2 = \lambda \cos^2 25^\circ$ ; then value of $\lambda$ be	(Q)	3

1.

 [Watch Video Solution](#)

### Exercise 5 Subjective Type Problems

1. Let  $P = \frac{\sin 80^\circ \sin 65^\circ \sin 35^\circ}{\sin 20^\circ + \sin 50^\circ + \sin 110^\circ}$ , then the value of  $24P$  is :

 [Watch Video Solution](#)

2. The value of expression  $(1 - \cot 23^\circ)(1 - \cot 22^\circ)$  is equal to :

 [Watch Video Solution](#)

3. If  $\tan A$  and  $\tan B$  the roots of the quadratic equation,  
 $4x^2 - 7x + 1 = 0$  then evaluate

$$4\sin^2(A + B) - 7\sin(A + B) \cdot \cos(A + B) + \cos^2(A + B).$$



[Watch Video Solution](#)

4.  $A_1A_2A_3 \dots A_{18}$  is a regular 18 sided polygon. B is an external point such that  $A_1A_2B$  is an equilateral triangle. If  $A_{18}A_1$  and  $A_1B$  are adjacent sides of a regular n sided polygon, then  $n =$



[Watch Video Solution](#)

5. If  $10\sin^4 \alpha + 15\cos^4 \alpha = 6$  and the value of  $9\operatorname{cosec}^4 \alpha + 8\sec^4 \alpha$  is S,  
then find the value of  $\frac{S}{25}$ .



[Watch Video Solution](#)

6. The value of  $\left(1 + \tan \frac{3\pi}{8} \tan \frac{\pi}{8}\right) + \left(1 + \tan \frac{5\pi}{8} \tan \frac{3\pi}{8}\right) + \left(1 + \tan \frac{7\pi}{8} \tan \frac{5\pi}{8}\right) + \left(1 + \tan \frac{9\pi}{8} \tan \frac{7\pi}{8}\right)$  is

 [Watch Video Solution](#)

7. If  $\alpha = \frac{\pi}{7}$  then find the value of  $\left(\frac{1}{\cos \alpha} + \frac{2 \cos \alpha}{\cos 2\alpha}\right)$

 [Watch Video Solution](#)

8. Given the for  $a, b, c, d \in R$ , if a

$\sec(200^\circ) - c \tan(200^\circ) = d$  and  $b \sec(200^\circ) + d \tan(200^\circ) = c$ ,

then find the value of  $\left(\frac{a^2 + b^2 + c^2 + d^2}{bd - ac}\right) \sin 20^\circ$ .

 [Watch Video Solution](#)

9. The expression  $2 \cos \frac{\pi}{17} \cdot \cos \frac{9\pi}{17} + \cos \frac{7\pi}{17} + \cos \frac{9\pi}{17}$  simplifies to an integer P. Find the value of P.



Watch Video Solution

10. If the expression  $\frac{\sin \theta \sin 2\theta + \sin 3\theta \sin 6\theta + \sin 4\theta \sin 13\theta}{\sin \theta \cos 2\theta + \sin 3\theta \cos 6\theta + \sin 4\theta \cos 13\theta} = \tan k\theta$ , where  $k \in \mathbb{N}$ .

Find the value of  $k$ .



Watch Video Solution

11. Let  $a = \sin 10^\circ$ ,  $b = \sin 50^\circ$ ,  $c = \sin 70^\circ$ , then  $8abc \left( \frac{a+b}{c} \right) \left( \frac{1}{a} + \frac{1}{b} - \frac{1}{c} \right)$

is equal to



Watch Video Solution

12. If  $\sin^3 \theta + \sin^3 \left( \theta + \frac{2\pi}{3} \right) + \sin^3 \left( \theta + \frac{4\pi}{3} \right) = a \sin b\theta$ . Find the value of  $\left| \frac{b}{a} \right|$ .



Watch Video Solution

13. If  $\sum_{r=1}^n \left( \frac{\tan 2^{r-1}}{\cos 2^r} \right) = \tan p^n - \tan q$ , then find the value of  $(p + q)$ .

 [Watch Video Solution](#)

14. If  $x = \sec \theta - \tan \theta$  and  $y = \operatorname{cosec} \theta + \cot \theta$ , then  $y - x - xy =$

 [Watch Video Solution](#)

15. Prove that:  $\cos 18^\circ - \sin 18^\circ = \sqrt{2} \sin 27^\circ$

 [Watch Video Solution](#)

16. The value of  $3(\sin 1 - \cos 1)^4 + 6(\sin 1 + \cos 1)^2 + 4(\sin^6 1 + \cos^6 1)$  is equal to

 [Watch Video Solution](#)

17. If  $x = \alpha$  satisfy the equation  $3^{\sin 2x + 2 \cos^2 x} + 3^{1 - \sin 2x + 2 \sin^2 x} = 28$ ,

then  $(\sin 2\alpha - \cos 2\alpha)^2 + 8 \sin 4\alpha$  is equal to :

 [Watch Video Solution](#)

18. If  $y = (\sin \theta + \operatorname{cosec} \theta)^2 + (\cos \theta + \sec \theta)^2$ , then minimum value of  $y$  is

:

 [Watch Video Solution](#)

19. If  $\tan 20^\circ + \tan 40^\circ + \tan 80^\circ - \tan 60^\circ = \lambda \sin 40^\circ$ , then  $\frac{\lambda}{4}$  is equal to

 [Watch Video Solution](#)

20. If  $K^\circ$  lies between  $360^\circ$  and  $540^\circ$  and  $K^\circ$  satisfies the equation

$1 + \cos 10x \cos 6x = 2 \cos^2 8x + \sin^2 8x$ , then  $\frac{K}{10} =$

 [Watch Video Solution](#)

[Watch Video Solution](#)

21. If  $\cos 20^\circ + 2 \sin^2 55^\circ = 1 + \sqrt{2} \sin K^\circ$ ,  $K$  in  $(0, 90^\circ)$ , then  $K =$  \_\_\_\_\_.

[Watch Video Solution](#)

22. If the value of  $\cos\left(\frac{2\pi}{7}\right) + \cos\left(\frac{4\pi}{7}\right) + \cos\left(\frac{6\pi}{7}\right) + \cos\left(\frac{7\pi}{7}\right) = -\frac{l}{2}$  Find the value of  $l$

[Watch Video Solution](#)

23. Let  $\alpha$  be the smallest integral value of  $x, x > 0$  such that  $\tan 19x = \frac{\cos 96^\circ + \sin 96^\circ}{\cos 96^\circ - \sin 96^\circ}$ . The last digit of  $\alpha$  is :

[Watch Video Solution](#)

24. Find the value of the expression  $\frac{\sin 20^\circ (4\cos 20^\circ + 1)}{\cos 20^\circ \cos 30^\circ}$

 [Watch Video Solution](#)

25. If the value of  $\cos\left(\frac{2\pi}{7}\right) + \cos\left(\frac{4\pi}{7}\right) + \cos\left(\frac{6\pi}{7}\right) + \cos\left(\frac{7\pi}{7}\right) = -\frac{l}{2}$  Find the value of  $l$

 [Watch Video Solution](#)

26. If  $\cos A = \frac{3}{4}$  and  $k \sin\left(\frac{A}{2}\right) \sin\left(\frac{5A}{2}\right) = \frac{11}{8}$ . Find  $k$ .

 [Watch Video Solution](#)

27. Find the least value of the expression  $3\sin^2 x + 4\cos^2 x$ .

 [Watch Video Solution](#)



28. If  $\tan \alpha$  and  $\tan \beta$  are the roots of equation  $x^2 - 12x - 3 = 0$ , then the value of  $\sin^2(\alpha + \beta) + 2 \sin(\alpha + \beta)\cos(\alpha + \beta) + 5 \cos^2(\alpha + \beta)$  is :

 [Watch Video Solution](#)

29. The value of  $\frac{\cos 24^\circ}{2 \tan 33^\circ \sin^2 57^\circ} + \frac{\sin 162^\circ}{\sin 18^\circ - \cos 18^\circ \tan 9^\circ} + \cos 162^\circ$  is equal to :

 [Watch Video Solution](#)

30. Find the value of  $\tan \theta(1 + \sec 2\theta)(1 + \sec 4\theta)(1 + \sec 8\theta)$ , when  $\theta = \frac{\pi}{32}$

 [Watch Video Solution](#)

31. If  $\lambda$  be the minimum value of  $y = (\sin x + \operatorname{cosec} x)^2 + (\cos x + \sec x)^2 + (\tan x + \cot x)^2$  where

$x \in R$ . Find  $\lambda - 6$ .



**Watch Video Solution**