

MATHS

BOOKS - VK JAISWAL ENGLISH

ELLIPSE

Exercise 1 Single Choice Problems

1. If CF be the perpendicular from the centre C of the ellipse $\frac{x^2}{12} + \frac{y^2}{8} = 1$, on the tangent at any point P and G is the point where the

normal at P meets the major axis, then the value of $(CF \cdot PG)$ equals to :

A. 5

B. 6

C. 8

D. None of these

Answer: C



Watch Video Solution

2. The minimum length of intercept on any tangent to the ellipse $\frac{x^2}{4} + \frac{y^2}{9} = 1$ cut by the circle $x^2 + y^2 = 25$ is :

A. 8

B. 9

C. 2

D. 11

Answer: A



Watch Video Solution

3. Find a point on the curve $x^2 + 2y^2 = 6$ whose distance from the line $x+y=7$, is minimum.

A. (2, 3)

B. (2, 1)

C. (1, 0)

D. None of these

Answer: B



Watch Video Solution

4. If lines $2x + 3y = 10$ and $2x - 3y = 10$ are tangents at the extremities of a latus rectum of an ellipse, whose centre is origin, then the length of the latus rectum is :

A. $\frac{110}{27}$

B. $\frac{98}{27}$

C. $\frac{100}{27}$

D. $\frac{120}{27}$

Answer: C



Watch Video Solution

5. Prove that the area bounded by the circle $x^2 + y^2 = a^2$ and the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ is equal to the area of another ellipse having semi-axis $a - b$ and b , $a > b$.

A. $a + b$ and b

B. $a - b$ and a

C. a and b

D. None of these

Answer: B



Watch Video Solution

6. If F_1 and F_2 are the feet of the perpendiculars from the foci S_1 and S_2 of the ellipse $\frac{x^2}{25} + \frac{y^2}{16} = 1$ on the tangent at any point P on the ellipse, then prove that $S_1F_1 + S_2F_2 \geq 8$.

A. $S_1F_1 + S_2F_2 \geq 2$

B. $S_1F_1 + S_2F_2 \geq 3$

$$C. S_1 F_1 + S_2 F_2 \geq 6$$

$$D. S_1 F_1 + S_2 F_2 \geq 8$$

Answer: D



Watch Video Solution

7. Consider the ellipse

$$\frac{x^2}{f(k^2 + 2k + 5)} + \frac{y^2}{f(k + 11)} = 1. \text{ If } f(x) \text{ is}$$

a positive decreasing function, then the set

of values of k for which the major axis is the x-

axis is $(-3, 2)$. the set of values of k for

which the major axis is the y-axis is $(-\infty, 2)$.

the set of values of k for which the major axis

is the y-axis is $(-\infty, -3) \cup (2, \infty)$ the set

of values of k for which the major axis is the y-

axis is $(-3, -\infty,)$

A. $k \in (-7, -5)$

B. $k \in (-5, -3)$

C. $k \in (-3, 2)$

D. None of these

Answer: C



8. If area of the ellipse $\frac{x^2}{16} + \frac{y^2}{b^2} = 1$ inscribed in a square of side length $5\sqrt{2}$ is A, then $\frac{A}{\pi}$ equals to :

A. 12

B. 10

C. 8

D. 11

Answer: A



Watch Video Solution

9. Any chord of the conic $x^2 + y^2 + xy = 1$ passing through origin is bisected at a point (p, q) , then $(p + q + 12)$ equals to :

A. 13

B. 14

C. 11

D. 12

Answer: D



Watch Video Solution

10. Tangents are drawn from the point $(4, 2)$ to the curve $x^2 + 9y^2 = 9$, the tangent of angle between the tangents :

A. $\frac{3\sqrt{3}}{5\sqrt{17}}$

B. $\frac{\sqrt{43}}{10}$

C. $\frac{\sqrt{43}}{5}$

D. $\sqrt{\frac{3}{17}}$

Answer: C



Watch Video Solution

Exercise 2 Comprehension Type Problems

1. An ellipse has semi-major of length 2 and semi-minor axis of length 1. It slides between the coordinate axes in the first quadrant while maintaining contact with both x-axis and y-axis. The locus of the centre of the ellipse is

A. $x^2 + y^2 = 3$

B. $x^2 + y^2 = 5$

$$C. (x - 2)^2 + (y - 1)^2 = 5$$

$$D. (x - 2)^2 + (y - 1)^2 = 3$$

Answer: B



Watch Video Solution

2. An ellipse has semi-major axis of length 2 and semi-minor axis of length 1. It slides between the coordinate axes in the first quadrant while maintaining contact with both x-axis and y-axis. The locus of the centre of the ellipse is

$$\text{A. } x^2 + y^2 + \frac{1}{x^2} + \frac{1}{y^2} = 16$$

$$\text{B. } x^2 + y^2 + \frac{1}{x^2} - \frac{1}{y^2} = 2\sqrt{3} + 4$$

$$\text{C. } x^2 + y^2 - \frac{1}{x^2} - \frac{1}{y^2} = 2\sqrt{3} + 4$$

$$\text{D. } x^2 - y^2 + \frac{1}{x^2} - \frac{1}{y^2} = 2\sqrt{3} + 4$$

Answer: A



Watch Video Solution

3. A coplanar beam of light emerging from a point source has the equation

$\lambda x - y + 2(1 + a\lambda) = 0, \lambda \in R$. The rays of

the beam strike an elliptical surface and get reflected. The reflected rays form another convergent beam having equation $\mu x - y + 2(1 - \mu) = 0$, $\mu \in R$. Further, it is found that the foot of the perpendicular from the point (2,2) upon any tangent to the ellipse lies on the circle $x^2 + y^2 - 4y - 5 = 0$

The eccentricity of the ellipse of is equal to

- A. $\frac{1}{3}$
- B. $\frac{1}{\sqrt{3}}$
- C. $\frac{2}{3}$

D. $\frac{1}{2}$

Answer: C



Watch Video Solution

4. A coplanar beam of light emerging from a point source has the equation $\lambda x - y + 2(1 + a\lambda) = 0$, $\lambda \in R$. The rays of the beam strike an elliptical surface and get reflected. The reflected rays form another convergent beam having equation

$\mu x - y + 2(1 - \mu) = 0, \mu \in R$. Further, it is found that the foot of the perpendicular from the point (2,2) upon any tangent to the ellipse lies on the circle $x^2 + y^2 - 4y - 5 = 0$

The eccentricity of the ellipse of is equal to

A. $4\sqrt{5}$

B. $\sqrt{5}$

C. $3\sqrt{5}$

D. $2\sqrt{5}$

Answer: D



5. A coplanar beam of light emerging from a point source has the equation $\lambda x - y + 2(1 + a\lambda) = 0, \lambda \in R$. The rays of the beam strike an elliptical surface and get reflected. The reflected rays form another convergent beam having equation $\mu x - y + 2(1 - \mu) = 0, \mu \in R$. Further, it is found that the foot of the perpendicular from the point (2,2) upon any tangent to the ellipse

lies on the circle $x^2 + y^2 - 4y - 5 = 0$

The eccentricity of the ellipse of is equal to

A. 6

B. 3

C. $\sqrt{5}$

D. $2\sqrt{5}$

Answer: A



Watch Video Solution

1. For the ellipse $\frac{x^2}{9} + \frac{y^2}{4} = 1$. Let O be centre and S and S' be the foci. For any point P on the ellipse the value of $PS \cdot PS' \cdot d^2$ (where d is the distance of O from the tangent at P) is equal to



[Watch Video Solution](#)

2. Number of perpendicular tangents that can be drawn on the ellipse $\frac{x^2}{16} + \frac{y^2}{25} = 1$ from point (6, 7) is



Watch Video Solution