



MATHS

BOOKS - ARIHANT MATHS (ENGLISH)

PAIR OF STRAIGHT LINES

Example

1. Find the joint equation of lines $y = x$ and $y = -x$.

 [Watch Video Solution](#)

2. Find the separate equation of lines represented by the equation

$$x^2 - 6xy + 8y^2 = 0$$

 [Watch Video Solution](#)

3. Find the condition that the slope of one of the lines represented by $ax^2 + 2hxy + by^2 = 0$ should be n times the slope of the other .

 [Watch Video Solution](#)

4. If the slope of one of the lines represented by $ax^2 + 2hxy + by^2 = 0$ be the n th power of the other, prove that , $(ab^n)^{\frac{1}{n+1}} + (a^n b)^{\frac{1}{n+1}} + 2h = 0$.

 [Watch Video Solution](#)

5. Find the product of the perpendiculars drawn from the point (x_1, y_1) on the lines $ax^2 + 2hxy + by^2 = 0$

 [Watch Video Solution](#)

6. Find the condition that the one of the lines given by $ax^2 + 2hxy + by^2 = 0$

may be perpendicular to one of the lines given by

$$a'x^2 + 2h'xy + b'y^2 = 0$$

 [Watch Video Solution](#)

7. Show that the area of the triangle formed by the lines

$$ax^2 + 2hxy + by^2 = 0 \text{ and } lx+my+n=0$$

$$\text{is } \frac{n^2 \sqrt{(h^2 - ab)}}{|(am^2 - 2hlm + bl^2)|}$$

 [Watch Video Solution](#)

8. Show that the area of the triangle formed by the lines

$$ax^2 + 2hxy + by^2 = 0 \text{ and } lx+my+n=0$$

$$\text{is } \frac{n^2 \sqrt{(h^2 - ab)}}{|(am^2 - 2hlm + bl^2)|}$$

 [Watch Video Solution](#)

9. Show that the two straight lines

$$x^2(\tan^2 \theta + \cos^2 \theta) - 2xy \tan \theta + y^2 \sin^2 \theta = 0$$

Make with the axis of x angles such that the difference of their tangents is 2 .

 [Watch Video Solution](#)

10. The angle between the lines $(x^2 + y^2) \sin^2 \alpha = (x \cos \beta - y \sin \beta)^2$ is

 [Watch Video Solution](#)

11. Show that the angle between the lines given by

$$(a + 2hm + bm^2)x^2 + 2\{(b - a)m - (m^2 - 1)h\}xy + (am^2 - 2hm + b)$$

is the same whatever be the value of m ,.

 [Watch Video Solution](#)

12. Show that the straight lines $x^2 + 4xy + y^2 = 0$ and the line $x-y=4$ form an equilateral triangle .

 [Watch Video Solution](#)

13. If two of the three lines represented by $ax^3 + bx^2y + cxy^2 + dy^3 = 0$ may be at right angles then

 [Watch Video Solution](#)

14. Find the equation of the bisectors of the angle between the lines represented by $3x^2 - 5xy + 4y^2 = 0$

 [Watch Video Solution](#)

15. The lines $y = mx$ bisects the angle between the lines $ax^2 + 2hxy + by^2 = 0$ if

 [Watch Video Solution](#)

16. If the pair of straight lines $x^2 - 2pxy - y^2 = 0$ and $x^2 - 2qxy - y^2 = 0$ are such that each pair bisects the angle between the other pair, then prove that $pq = -1$.

 [Watch Video Solution](#)

17. If the lines given by $ax^2 + 2hxy + by^2 = 0$ are equally inclined to the lines given by $ax^2 + 2hxy + by^2 + \lambda(x^2 + y^2) = 0$, then

 [Watch Video Solution](#)

18. Show that the pair of lines given by $a^2x^2 + 2h(a + b)xy + b^2y^2 = 0$ is equally inclined to the pair given by $ax^2 + 2hxy + by = 0$.

 [Watch Video Solution](#)

19. If the lines represented by $x^2 - 2pxy - y^2 = 0$ are rotated about the origin through an angle θ , one clockwise direction and other in anti-clockwise direction, then the equation of the bisectors of the angle between the lines in the new position is

 [Watch Video Solution](#)

20. For what value of λ does the equation $12x^2 - 10xy + 2y^2 + 11x - 5y + \lambda = 0$ represent a pair of straight lines? Find their equations and the angle between them.

 [Watch Video Solution](#)

21. Prove that the equation $8x^2 + 8xy + 2y^2 + 26x + 13y + 15 = 0$ represents a pair of parallel straight lines. Also find the perpendicular distance between them.

 [Watch Video Solution](#)

22. Find the combined equation of the straight lines passing through the point (1,1) and parallel to the lines represented by the equation .

$$x^2 - 5xy + 4y^2 + x + 2y - 2 = 0 .$$

 [Watch Video Solution](#)

23. Find the point of intersection of lines represented by

$$2x^2 - 7xy - 4y^2 - x + 22y - 10 = 0$$

 [Watch Video Solution](#)

24. Find the new equation of curve

$$12x^2 + 7xy - 12y^2 - 17x - 31y - 7 = 0$$
 after removing the first degree

terms.

 [Watch Video Solution](#)

25. Prove that the angle between the lines joining the origin to the points of intersection of the straight line $y = 3x + 2$ with the curve $x^2 + 2xy + 3y^2 + 4x + 8y - 11 = 0$ is $\tan^{-1}\left(\frac{2\sqrt{2}}{3}\right)$

 [Watch Video Solution](#)

26. Find the equation to the pair of straight lines joining the origin to the intersections of the straight line $y = mx + c$ and the curve $x^2 + y^2 = a^2$. Prove that they are at right angles if $2c^2 = a^2(1 + m^2)$.

 [Watch Video Solution](#)

27. Prove that the pair of lines joining the origin to the intersection of the curve $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ and the line $lx + my + n = 0$ are coincident, if $a^2l^2 + b^2m^2 = n^2$

 [Watch Video Solution](#)

28. The pair of lines joining origin to the points of intersection of, the two curves $ax^2 + 2hxy + by^2 + 2gx = 0$ and $a'x^2 + 2h'xy + b'y^2 + 2g'x = 0$ will be at right angles, if

 [Watch Video Solution](#)

29. If the pairs of lines $x^2 + 2xy + ay^2 = 0$ and $ax^2 + 2xy + y^2 = 0$ have exactly one line in common, then the joint equation of the other two lines is given by $3x^2 + 8xy - 3y^2 = 0$ $3x^2 + 10xy + 3y^2 = 0$ $y^2 + 2xy - 3x^2 = 0$ $x^2 + 2xy - 3y^2 = 0$

A. $3x^2 + 8xy - 3y^2 = 0$

B. $3x^2 + 10xy + 3y^2 = 0$

C. $x^2 + 2xy - 3y^2 = 0$

D. $3x^2 + 2xy - y^2 = 0$

Answer: b

 [Watch Video Solution](#)

30. The combined equation of the lines l_1 and l_2 is $2x^2 + 6xy + y^2 = 0$ and that of the lines m_1 and m_2 is $4x^2 + 18xy + y^2 = 0$. If the angle between l_1 and m_2 is α then the angle between l_2 and m_1 will be $\frac{\pi}{2} - \alpha$

(b) 2α $\frac{\pi}{4} + \alpha$ (d) α

A. $\frac{\pi}{2} - \alpha$

B. $\frac{\pi}{4} + \alpha$

C. α

D. 2α

Answer: c



Watch Video Solution

31. If the pair of lines $\sqrt{3}x^2 - 4xy + \sqrt{3}y^2 = 0$ is rotated about the origin by $\pi/6$ in the anticlockwise sense, then find the equation of the pair of lines in the new position.

A. $x^2 - \sqrt{3}xy = 0$

B. $y^2 - \sqrt{3}xy = 0$

C. $\sqrt{3}x^2 - xy = 0$

D. $\sqrt{3}y^2 - xy = 0$

Answer: c



Watch Video Solution

32. If the pair of lines $ax^2 - 2xy + by^2 = 0$ and $bx^2 - 2xy + ay^2 = 0$

be such that each pair bisects the angle between the other pair, then $|a - b|$ equals to

A. 1

B. 2

C. 3

D. 4

Answer: b



Watch Video Solution

33. The equation of line which is parallel to the line common to the pair of lines given by $3x^2 + xy - 4y^2 = 0$ and $6x^2 + 11xy + 4y^2 = 0$ and at a distance of 2 units from it is

A. $3x - 4y = -10$

B. $x - y = 2$

C. $3x + 4y = 10$

D. $2x + y = -2$

Answer: c



Watch Video Solution

34. The lines joining the origin to the point of intersection of $3x^2 + mxy - 4x + 1 = 0$ and $2x + y - 1 = 0$ are at right angles. Then which of the following is a possible value of m ? -4 (b) 4 (c) 7 (d) 3

A. $g^2 + f^2 = c$

B. $g^2 - f^2 = c$

C. $g^2 - f^2 = 2c$

D. $g^2 + f^2 = c^2$

Answer: c



Watch Video Solution

35. The lines joining the origin to the point of intersection of $3x^2 + mxy - 4x + 1 = 0$ and $2x + y - 1 = 0$ are at right angles. Then which of the following is a possible value of m ?

A. -4

B. 3

C. 4

D. 7

Answer: (a,b,c,d)



Watch Video Solution

36. The lines $(lx + my)^2 - 3(mx - ly)^2 = 0$ and $lx + my + n = 0$ forms

A. an isosecles triangle

B. a right angled triangle

C. an equilateral triangle

D. None of these

Answer: (a,c)



Watch Video Solution

37. If the equation $ax^2 - 6xy + y^2 + 2bx + 2cy + d = 0$ represents a pair of lines whose slopes are m and m^2 , then value (s) of a is /are

A. -27

B. -8

C. 8

D. 27

Answer: (a,c)



[Watch Video Solution](#)

38. Consider the equation of a pair of straight lines as

$$\lambda xy - 8x + 9y - 12 = 0$$

A. 0

B. 2

C. 4

D. 6

Answer: d



Watch Video Solution

39. The point of intersection of lines is (α, β) , then the equation whose roots are α, β , is

A. $4x^2 + x - 8 = 0$

B. $6x^2 + x - 12 = 0$

C. $4x^2 - x - 8 = 0$

D. $6x^2 - x - 12 = 0$

Answer: b



Watch Video Solution

40. If the sum of the slopes of the lines given by $x^2 - 2cxy - 7y^2 = 0$ is four times their product, then find the value of c .

 [Watch Video Solution](#)

41. If one of the lines given by $6x^2 - xy + 4cy^2 = 0$ is $3x + 4y = 0$, then value of $|c|$ is

 [Watch Video Solution](#)

42. Find the slope of tangent to the curve if $ax^2 + 2hxy + by^2 = 0$

 [Watch Video Solution](#)

43. Statement I . The combined equation of l_1, l_2 is $3x^2 + 6xy + 2y^2 = 0$ and that of m_1, m_2 is $5x^2 + 18xy + 2y^2 = 0$. If angle between l_1, m_2 is θ , then angle between l_2, m_1 is θ .

Statement II . If the pairs of lines $l_1l_2 = 0, m_1m_2 = 0$ are equally inclined that angle between l_1 and $m_2 =$ angle between l_2 and m_1 .

 [Watch Video Solution](#)

44. Statement I . The equation $2x^2 - 3xy - 2y^2 + 5x - 5y + 3 = 0$ represents a pair of perpendicular straight lines.

Statement II A pair of lines given by $ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0$ are perpendicular if $a + b = 0$

 [Watch Video Solution](#)

45. If the lines represented by $2x^2 - 5xy + 2y^2 = 0$ be the sides of a parallelogram and the line $5x + 2y = 1$ be one of its diagonal. Find the equation of the other diagonal, and area of the parallelogram .

 [Watch Video Solution](#)

46. Prove that the equation $(a + 2h + b)x^2 - 2(a - b)xy + (a - 2h + b)y^2 = 0$ represents a pair of lines each inclined at an angle of 45° to one or other of the lines given by $ax^2 + 2hxy + by^2 = 0$

 [Watch Video Solution](#)

47. If $u \equiv ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0$

represents a pair of straight lines, prove that the equation of the third pair of straight lines passing through the points where these meet the axes is $ax^2 - 2hxy + by^2 + 2gx + 2fy + c + 4fgxy = 0$.

 [Watch Video Solution](#)

48. If the equation $ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0$ represents a pair of parallel lines, prove that

$$h = \sqrt{ab} \text{ and } g\sqrt{b} = f\sqrt{a} \text{ or } (h = -\sqrt{ab} \text{ and } g\sqrt{b} = -f\sqrt{a}).$$

$$\text{The distance between them is } 2\sqrt{\left(\frac{g^2 - ac}{a(a+b)}\right)}.$$

 [Watch Video Solution](#)

49. Find $\frac{dy}{dx}$ if $x - 3y = x^4$

 [Watch Video Solution](#)

50. A point moves so that the distance between the foot of perpendiculars from it on the lines $ax^2 + 2hxy + by^2 = 0$ is a constant $2d$. Show that the equation to its locus is $(x^2 + y^2)(h^2 - ab) = d^2\{(a - b)^2 + 4h^2\}$.

 [Watch Video Solution](#)

51. Show that if two of the lines $ax^3 + bx^2y + cxy^2 + dy^3 = 0$ ($a \neq 0$) make complementary angles with X-axis in anti-clockwise sense, then a(a-

$$c)+d(b-d)=0 .$$



[Watch Video Solution](#)

52. Show that the equation $a(x^4 + y^4) - 4bxy(x^2 - y^2) + 6cx^2y^2 = 0$ represents two pairs of lines at right angles and that if $2b^2 = a^2 + 3ac$, the two pairs will coincide.



[Watch Video Solution](#)

53. if one of the lines given by the equation $ax^2 + 2hxy + by^2 = 0$ coincides with one of the lines given by $a'x^2 + 2h'xy + b'y^2 = 0$ and the other lines represented by them be perpendicular , then .

$$\frac{ha'b'}{b' - a'} = \frac{h'ab}{b - a} = \frac{1}{2}\sqrt{(-aa'f)}$$



[Watch Video Solution](#)

1. The lines given by the equation $(2y^2 + 3xy - 2x^2)(x + y - 1) = 0$ form a triangle which is

- A. equilateral
- B. isosceles
- C. right angled
- D. obtuse angled

Answer: C



[Watch Video Solution](#)

2. Area of the triangle formed by the lines $y^2 - 9xy + 18x^2 = 0$ and $y = 9$ is

- A. $27/4$
- B. 0
- C. $9/4$

D. 27

Answer: A



[Watch Video Solution](#)

3. The equation $3x^2 + 2hxy + 3y^2 = 0$ represents a pair of straight lines passing through the origin . The two lines are

- A. real and distinct , if $h^2 > 3$
- B. real and distinct , if $h^2 > 9$
- C. real and coincident , if $h^2 = 3$
- D. real and coincident , if $h^2 > 3$

Answer: B



[Watch Video Solution](#)

4. If one of the lines of the pair $ax^2 + 2hxy + by^2 = 0$ bisects the angle between the positive direction of the axes. Then find the relation for a , b and h .

A. $a + b = 2|h|$

B. $a + b = -2h$

C. $a - b = 2|h|$

D. $(a - b)^2 = 4h^2$

Answer: B



[Watch Video Solution](#)

5. If the slope of the line given by $a^2x^2 + 2hxy + b^2y^2 = 0$ be three times of the other, then h is equal to

A. (a) $2\sqrt{3}ab$

B. (b) $-2\sqrt{3}ab$

C. (c) $\frac{2}{\sqrt{3}}ab$

D. (d) $-\frac{2}{\sqrt{3}}ab$

Answer: C::D



Watch Video Solution

6. Find the separate equation of two straight lines whose joint equation is $ab(x^2 - y^2) + (a^2 - b^2)xy = 0$



Watch Video Solution

7. Find the coordinates of the centroid of the triangle whose sides are $12x^2 - 20xy + 7y^2 = 0$ and $2x - 3y + 4 = 0$



Watch Video Solution

8. If the lines $ax^2 + 2hxy + by^2 = 0$ be two sides of a parallelogram and the line $lx+my=1$ be one of its diagonal, show that the equation of the other diagonal is $y(bl-hm)=x(am-hl)$.



Watch Video Solution

9. Find the condition that one of the lines given by $ax^2 + 2hxy + by^2 = 0$ may coincide with one of the lines given by $a'x^2 + 2h'xy + b'y^2 = 0$



Watch Video Solution

Exercise For Session 2

1. The angle between the pair of straight lines $y^2 \sin^2 \theta - xy \sin^2 \theta + x^2 (\cos^2 \theta - 1) = 0$ is

A. $\frac{\pi}{4}$

B. $\frac{\pi}{2}$

C. $\frac{\pi}{3}$

D. $\frac{2\pi}{3}$

Answer: B



Watch Video Solution

2. The angle between the lines $ay^2 - (1 + \lambda^2)xy - ax^2 = 0$ is same as the angle between the line:

A. (a) $5x^2 + 2xy - 3y^2 = 0$

B. (b) $x^2 - 2xy - 3y^2 = 0$

C. (c) $x^2 - y^2 = 100$

D. (d) $xy=0$

Answer: C::D



Watch Video Solution

3. Which of the following pair of straight lines intersect at right angles ?

A. $2x^2 = y(x + 2y)$

B. $(x + y)^2 = x(y + 3x)$

C. $2y(x + y) = xy$

D. $y = + 2x$

Answer: A



[Watch Video Solution](#)

4. if $h^2 = ab$, then the lines represented by $ax^2 + 2hxy + by^2 = 0$ are

A. Parallel

B. perpendicular

C. coincident

D. None of these

Answer: C



Watch Video Solution

5. Equation $ax^3 - 9x^2y - xy^2 + 4y^3 = 0$ represents three straight lines.

If the two of the lines are perpendicular, then a is equal to

A. -5

B. 5

C. -4

D. 4

Answer: B::C



Watch Video Solution

6. Find the angle between the lines whose joint equation is

$$2x^2 - 3xy + y^2 = 0$$



Watch Video Solution

7. Show that the lines

$$(1 - \cos \theta \tan \alpha)y^2 - (2 \cos \theta + \sin^2 \theta \tan \alpha)xy + \cos \theta(\cos \theta + \tan \alpha)x^2 = 0$$

include an angle α between them .



Watch Video Solution

8. Find the angle between the lines represented by the equation

$$x^2 - 2pxy + y^2 = 0$$



Watch Video Solution

9. Show that the lines $x^2 - 4xy + y^2 = 0$ and $x + y = 1$ form an equilateral triangle and find its area.



Watch Video Solution

10. Find $\frac{dy}{dx}$ if $ax^2 + 2hxy + by^2 = 0$



Watch Video Solution

Exercise For Session 3

1. If the coordinate axes are the bisectors of the angles between the pair of lines $ax^2 + 2hxy + by^2 = 0$, then

A. (a) $a=b$

B. (b) $h=0$

C. (c) $a^2 = b = 0$

D. (d) $a + b^2 = 0$

Answer: B



Watch Video Solution

2. The equation of the bisectors of angle between the lines $x^2 - 4xy + y^2 = 0$ is

 [Watch Video Solution](#)

3. If one of the lines of $my^2 + (1 - m^2)xy - mx^2 = 0$ is a bisector of the angle between lines $xy=0$, then $\cos^{-1}(m)$ is

A. 0

B. $\pi/2$

C. π

D. $3\pi/2$

Answer: A:C

 [Watch Video Solution](#)

4. The bisectors of the angles between the lines $(ax + by)^2 = c(bx - ay)^2, c > 0$ are respectively parallel and perpendicular to the line

A. $bx - ay + \mu = 0$

B. $ax + by + \lambda = 0$

C. $ax = by + v = 0$

D. $bx + ay + \tau = 0$

Answer: B



[Watch Video Solution](#)

5. If the pairs of straight lines $ax^2 + 2hxy - ay^2 = 0$ and $bx^2 + 2gxy - by^2 = 0$ be such that each bisects the angles between the other, then



[Watch Video Solution](#)

6. Prove that the lines $2x^2 + 6xy + y^2 = 0$ are equally inclined to the lines $4x^2 + 18xy + y^2 = 0$



[Watch Video Solution](#)

7. Show that the equation of the pair of lines bisecting the angles between the pair of bisectors of the angles between the pair of lines $ax^2 + 2hxy + by^2 = 0$ is $(a - b)(x^2 - y^2) + 4hxy = 0$



[Watch Video Solution](#)

8. Prove that the bisectors of the angle between the lines $ax^2 + acxy + cy^2 = 0$ and $\left(3 + \frac{1}{c}\right)x^2 + xy + \left(3 + \frac{1}{a}\right)y^2 = 0$ are always the same .



[Watch Video Solution](#)

9. The lines represented by $x^2 + 2\lambda xy + 2y^2 = 0$ and the lines represented by $(1 + \lambda)x^2 - 8xy + y^2 = 0$ are equally inclined, then $\lambda =$

 [Watch Video Solution](#)

Exercise For Session 4

1. Prove that the equation $3y^2 - 8xy - 3x^2 - 29x + 3y - 18 = 0$ represents two straight lines. Find also their point of intersection and the angle between them.

- A. $\left(1, \frac{1}{2}\right)$
- B. $\left(1, -\frac{1}{2}\right)$
- C. $\left(-\frac{3}{2}, \frac{5}{2}\right)$
- D. $\left(-\frac{3}{2}, -\frac{5}{2}\right)$

Answer: D



Watch Video Solution

2. If the angle between the two lines represented by $2x^2 + 5xy + 3y^2 + 6x + 7y + 4 = 0$ is $\tan^{-1}(m)$, then find the value of m .

A. $-\frac{1}{5}$

B. $\frac{1}{5}$

C. $-\frac{3}{5}$

D. $\frac{3}{5}$

Answer: B



Watch Video Solution

3. The equation of second degree $x^2 + 2\sqrt{2}xy + 2y^2 + 4x + 4\sqrt{2}y + 1 = 0$ represents a pair of straight lines. The distance between them is

A. 2

B. $2\sqrt{3}$

C. 4

D. $4\sqrt{3}$

Answer: A



[Watch Video Solution](#)

4. Find the area of the parallelogram formed by the lines

$$2x^2 + 5xy + 3y^2 = 0 \text{ and } 2x^2 + 5xy + 3y^2 + 3x + 4y + 1 = 0$$



[Watch Video Solution](#)

5. Find the locus of the incentre of the triangle formed by

$$xy - 4x - 4y + 16 = 0 \text{ and } x + y = a (a > 4, a \neq \sqrt{2} \text{ and } a \text{ is the parameter})$$



[Watch Video Solution](#)

6. If the equation $2hxy + 2gx + 2fy + c = 0$ represents two straight lines, then show that they form a rectangle of area $\frac{|fg|}{h^2}$ with the coordinate axes.

 [Watch Video Solution](#)

7. Find the area of the triangle formed by the lines represented by $ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0$ and axis of x .

 [Watch Video Solution](#)

8. Find the combined equation of the straight lines passing through the point $(1,1)$ and parallel to the lines represented by the equation $x^2 - 5xy + 4y^2 + x + 2y - 2 = 0$.

 [Watch Video Solution](#)

Exercise For Session 5

1. If the straight lines joining origin to the points of intersection of the line $x+y=1$ with the curve $x^2 + y^2 + x - 2y - m = 0$ are perpendicular to each other, then the value of m should be

A. $-\frac{1}{2}$

B. 0

C. $\frac{1}{2}$

D. 1

Answer: A



[Watch Video Solution](#)

2. The angle between the pair of straight lines formed by joining the points of intersection of $x^2 + y^2 = 4$ and $y = 3x + c$ to the origin is a right angle. Then c^2 is equal to

A. -1

B. 6

C. 13

D. 20

Answer: A



Watch Video Solution

3. If θ is an angle by which axes are rotated about origin and equation

$$ax^2 + 2hxy + by^2 = 0$$

does not contain xy term in the new system, then prove that

$$\tan 2\theta = \frac{2h}{a - b}.$$

A. $\frac{(a - b)}{2h}$

B. $\frac{2h}{(a + b)}$

C. $\frac{(a + b)}{2h}$

D. $\frac{2h}{(a - b)}$

Answer: A



Watch Video Solution

4. The lines joining the origin to the points of intersection of $2x^2 + 3xy - 4x + 1 = 0$ and $3x + y = .1$ given by

A. $x^2 - y^2 - 5xy = 0$

B. $x^2 - y^2 + 5xy = 0$

C. $x^2 + y^2 - 5xy = 0$

D. $x^2 + y^2 + 5xy = 0$

Answer: A



Watch Video Solution

5. The equation of the line joining the origin to the point of intersection of the lines $2x^2 + xy - y^2 + 5x - y + 2 = 0$ is

A. $x+y=0$

B. $x-y=0$

C. $x-2y=0$

D. $2x+y=0$

Answer: A

 [Watch Video Solution](#)

6. The lines joining the origin to the points of intersection of the line $3x-2y-1$ and the curve $3x^2 + 5xy - 3y^2 + 2x + 3y = 0$, are

 [Watch Video Solution](#)

7. If the straight lines joining the origin and the points of intersection of $y = mx + 1$ and $x^2 + y^2 = 1$ are perpendicular to each other, then find the value of m .

 [Watch Video Solution](#)

8. Prove that the straight lines joining the origin to the point of intersection of the straight line $hx + ky = 2hk$ and the curve $(x - k)^2 + (y - h)^2 = c^2$ are perpendicular to each other if $h^2 + k^2 = c^2$.



[Watch Video Solution](#)

9. Show that for all values of λ , the lines joining the origin to the points common to $x^2 + 2hxy - y^2 + gx + fy = 0$ and $fx - gy = \lambda$ are at right angles .



[Watch Video Solution](#)

10. Find the equations of the straight lines joining the origin to the points of intersection of $x^2 + y^2 - 4x - 2y = 4$ and $x^2 + y^2 - 2x - 4y = 4$.



Exercise Single Option Correct Type Questions

1. If the sum of the slopes of the lines given by $4x^2 + 2\lambda xy - 7y^2 = 0$ is equal to the product of the slope, then λ is equal to

- A. a) -4
- B. b) -2
- C. c) 2
- D. d) 4

Answer: B

2. The equation $3ax^2 + 9xy + (a^2 - 2)y^2 = 0$ represents two perpendicular straight lines for

- A. a) only one value of a
- B. b) for all values of a
- C. c) for only two values of a
- D. d) for no value of a

Answer: C

 [Watch Video Solution](#)

3. The image of the pair of lines represented by $ax^2 + 2hxy + by^2 = 0$ by the line mirror $y = 0$ is

- A. $ax^2 + 2hxy + by^2 = 0$
- B. $bx^2 - 2hxy + ay^2 = 0$
- C. $bx^2 + 2hxy + ay^2 = 0$
- D. $ax^2 - 2hxy + by^2 = 0$

Answer: D

 [Watch Video Solution](#)

4. Number of points lying on the line $7x + 4y + 2 = 0$ which is equidistant from the lines $15x^2 + 56xy + 48y^2 = 0$ is

A. 0

B. 1

C. 2

D. 4

Answer: C

 [Watch Video Solution](#)

5. Orthocentre of the triangle formed by the lines $xy - 3x - 5y + 15 = 0$ and $3x + 5y = 15$ is

A. (-5,-3)

B. (5,3)

C. (-3,-5)

D. (3,5)

Answer: B



[Watch Video Solution](#)

6. Two of the straight lines given by $3x^3 + 3x^2y - 3xy^2 + dy^3 = 0$ are at right angles , if d equal to

A. -4

B. -3

C. -2

D. -1

Answer: B



[Watch Video Solution](#)

7. Two lines are given by $(x - 2y)^2 + k(x - 2y) = 0$. The value of k , so that the distance between them is 3, is:

- A. (a) $\sqrt{5}$
- B. (b) $2\sqrt{5}$
- C. (c) $3\sqrt{5}$
- D. (d) $4\sqrt{5}$

Answer: C



[Watch Video Solution](#)

8. The point of intersection of the two lines given by $2x^2 - 5xy + 2y^2 - 3x + 3y + 1 = 0$ is

- A. A. (-2,2)
- B. B. (-3,3)

C. C. (3,3)

D. D. (2,2)

Answer: C



[Watch Video Solution](#)

9. Let $0 < p < q$ and $a \neq 0$ such that the equation $px^2 + 4\lambda xy + qy^2 + 4a(x + y + 1) = 0$ represents a pair of straight lines, then a can lie in the interval

A. $\alpha \leq p \leq \beta$

B. $p \leq \alpha$

C. $p \leq \alpha$ or $p \geq \beta$

D.

Answer: D



[Watch Video Solution](#)

10. If the equation of the pair of straight lines passing through the point $(1, 1)$, one making an angle θ with the positive direction of the x-axis and the other making the same angle with the positive direction of the y-axis, is $x^2 - (a + 2)xy + y^2 + a(x + y - 1) = 0, a \neq 2$, then the value of $\sin 2\theta$ is $a - 2$ (b) $a + 2$ (c) $2(a + 2)$ (d) $\frac{2}{a}$

A. $a-2$

B. $a+2$

C. $\frac{2}{(a + 2)}$

D. $\frac{2}{a}$

Answer: C



Watch Video Solution

Exercise More Than One Correct Option Type Questions

1. The equation of image of pair of lines $y = |x - 1|$ with respect to y-axis is :

A. $y = |x + 1|$

B. $y = |x - 1| + 3$

C. $x^2 - y^2 + 2x + 1 = 0$

D. $x^2 - y^2 + 2x - 1 = 0$

Answer: A::C



[Watch Video Solution](#)

2. If the equation $ax^2 + by^2 + cx + cy = 0$ represents a pair of straight lines , then

A. a) $a + b = 0$

B. b) $c = 0$

C. c) $a + c = 0$

D. d) $c(a + b) = 0$

Answer: A::B::D



Watch Video Solution

3. If $x^2 + \alpha y^2 + 2\beta y = a^2$ represents a pair of perpendicular straight lines , then

A. $\alpha = 1, \beta = a$

B. $\alpha = 1, \beta = -a$

C. $\alpha = -1, \beta = -a$

D. $\alpha = -1, \beta = a$

Answer: C::D



Watch Video Solution

4. If the pair of lines $ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0$ intersect on the y-axis, then prove that $2fgh = bg^2 + ch^2$.

A. $f^2 = bc$

B. $abc = 2fgh$

C. $bg^2 \neq ch^2$

D. $2fgh = bg^2 + ch^2$

Answer: A::D



[Watch Video Solution](#)

5. Two pairs of straight lines have the equations $y^2 + xy - 12x^2 = 0$ and $ax^2 + 2hxy + by^2 = 0$. One line will be common among them if

A. $a = -3(2h + 3b)$

B. $a = 8(h - 2b)$

C. $a=2(b+h)$

D. $a=-3(b+h)$

Answer: A::B



Watch Video Solution

6. The three sides of a triangle are given by $(x^2 - y^2)(2x + 3y - 6) = 0$.

If the points $(-2,a)$ lies inside and $(b,1)$ lies outside the triangle, then

A. $2 < a < \frac{10}{3}$

B. $-2 < a < \frac{10}{3}$

C. $-1 < b < \frac{9}{2}$

D. $-1 < b < 1$

A. $2 < a < \frac{10}{3}$

B. $-2 < a < \frac{10}{3}$

C. $-1 < b < \frac{9}{2}$

D. $-1 < b < 1$

Answer: A::D



Watch Video Solution

Exercise Passage Based Questions

1. Consider the equation of a pair of straight lines as

$$x^2 - 3xy + \lambda y^2 + 3x = 5y + 2 = 0$$

The value of λ is

A. 1

B. 2

C. 3

D. 4

Answer: B



Watch Video Solution

2. Consider the equation of a pair of straight lines as

$$x^2 - 3xy + \lambda y^2 + 3x - 5y + 2 = 0$$

The point of intersection of line is (α, β) , then the value of $\alpha^2 + \beta^2$ is

A. 2

B. 5

C. 10

D. 17

Answer: C

[Watch Video Solution](#)

3. Consider the equation of a pair of straight lines as

$$x^2 - 3xy + \lambda y^2 + 3x - 5y + 2 = 0$$

The angle between the lines is θ then the value of $\cos 2\theta$ is

A. $\frac{1}{3}$

B. $\frac{2}{3}$

C. $\frac{3}{5}$

D. $\frac{4}{5}$

Answer: D



Watch Video Solution

4. Let $f_1(x, y) \equiv ax^2 + 2hxy + by^2 = 0$ and let $f_{i+1}(x, y) = 0$ denote the equation of the bisectors of $f_i(x, y) = 0$ for all $i=1,2,3,\dots$

$f_3(x, y) = 0$ is

A. (a) $hx^2 - (a - b)xy - hy^2 = 0$

B. (b) $(a - b)x^2 + 4hxy - (a - b)y^2 = 0$

C. (c) $ax^2 + 2hxy + by^2 = 0$

D. (d) None of the above

Answer: B



Watch Video Solution

5. Let $f_1(x, y) \equiv ax^2 + 2hxy + by^2 = 0$ and let $f_{i+1}(x, y) = 0$ denote the equation of the bisectors of $f_i(x, y) = 0$ for all $i=1,2,3,\dots$

$f_3(x, y) = 0$ is

A. $f_1(x, y) = 0$

B. $f_2(x, y) = 0$

C. $hx^2 - (a - b)xy - hy^2 = 0$

D. None of the above

Answer: A



Watch Video Solution

6. Let $f_1(x, y) \equiv ax^2 + 2hxy + by^2 = 0$ and let $f_{i+1}(x, y) = 0$ denote the equation of the bisectors of $f_i(x, y) = 0$ for all $i=1,2,3,\dots$

The value of $\sum_{n=2}^5 \frac{f_{n+2}(x, y)}{f_n(x, y)}$ is

A. 14

B. 4

C. 54

D. 6

Answer: B



[Watch Video Solution](#)

7. Consider a pair of perpendicular straight lines

$2x^2 + 3xy + by^2 - 11x + 13y + c = 0$ The value of c is

A. -2

B. 2

C. -3

D. 3

Answer: A



Watch Video Solution

8. Find $\frac{dy}{dx}$ if $2x^2 + 3xy + by - 11x + 13y + c = 0$



Watch Video Solution

9. Consider a pair of perpendicular straight lines

$$2x^2 + 3xy + by - 11x + 13y + c = 0$$

The value of c is

A. 2

B. 3

C. 4

D. 5

Answer: C

 [Watch Video Solution](#)

Exercise Single Integer Answer Type Questions

1. If the lines joining the origin to the intersection of the line $y=nx+2$ and the curve $x^2 + y^2 = 1$ are at right angles, then the value of n^2 is

 [Watch Video Solution](#)

2. Area of the triangle formed by the line $x + y = 3$ and angle bisectors of the pair of straight lines $x^2 - y^2 + 2y = 1$ is $2sq\text{units}$ b. $4sq\text{units}$ c. $6sq\text{units}$ d. $8sq\text{units}$

 [Watch Video Solution](#)

Exercise Statement I And Ii Type Questions

1. Statement I. The four straight lines given by

$6x^2 + 5xy - 6y^2 = 0$ and $6x^2 + 5xy - 6y^2 - x + 5y - 1 = 0$ are the sides of a square .

Statement II . The lines represented by general equation of second degree $ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0$ are perpendicular if $a+b=0$.

- A. A. Statement I is true, Statement II is true , Statement II is a correct explanation for Statement I
- B. B. Statement I is true , Statement II is true , Statement II is not a correct explanation for statement I
- C. C. Statement I is true , Statement II is false
- D. D. Statement I is false , Statement II is true

Answer: b



[Watch Video Solution](#)

2. Statement I. Two of the straight lines represented by $dx^3 + cx^2y + bxy^2 + ay^3 = 0$ will be at right angles if $d^2 + bd + bc + a^2 = 0$

Statement II. Product of the slopes of two perpendicular line is -1

- A. Statement I is true, Statement II is true , Statement II is a correct explanation for Statement I
- B. Statement I is true , Statement II is true , Statement II is not a correct explanation for statement I
- C. Statement I is true , Statement II is false
- D. Statement I is false , Statement II is true

Answer: b



Watch Video Solution

3. Statement I. if $\alpha\beta = -1$ then the pair of straight lines $x^2 - 2\alpha xy - y^2 = 0$ and $y^2 + 2\beta xy - x^2 = 0$ are the angle bisector of each other.

Statement II. Pair of angle bisector lines of the pair of lines $ax^2 + 2hxy + by^2 = 0$ is $(x^2 - y^2) = (a - b)xy$.

A. Statement I is true, Statement II is true, Statement II is a correct explanation for Statement I

B. Statement I is true, Statement II is true, Statement II is not a correct explanation for statement I

C. Statement I is true, Statement II is false

D. Statement I is false, Statement II is true

Answer: a

 [Watch Video Solution](#)

4. Statement 1 : If $-2h = a + b$, then one line of the pair of lines $ax^2 + 2hxy + by^2 = 0$ bisects the angle between the coordinate axes in the positive quadrant. Statement 2 : If $ax + y(2h + a) = 0$ is a factor of $ax^2 + 2hxy + by^2 = 0$, then $b + 2h + a = 0$

- A. Statement I is true, Statement II is true , Statement II is a correct explanation for Statement I
- B. Statement I is true , Statement II is true , Statement II is not a correct explanation for statement I
- C. Statement I is true , Statement II is false
- D. Statement I is false , Statement II is true

Answer: b



[Watch Video Solution](#)

Exercise Subjective Type Questions

1. The straight lines represented by $(y - mx)^2 = a^2(1 + m^2)$ and $(y - nx)^2 = a^2(1 + n^2)$ form a rectangle (b) rhombus trapezium (d) none of these

 [Watch Video Solution](#)

2. Prove that the equation $m(x^3 - 3xy^2) + y^3 - 3x^2y = 0$ represents three straight lines equally inclined to each other.

 [Watch Video Solution](#)

3. Show that straight lines $(A^2 - 3B^2)x^2 + 8ABxy + (B^2 - 3A^2)y^2 = 0$ form with the line $Ax + By + C = 0$ an equilateral triangle of area $\frac{C^2}{\sqrt{3}(A^2 + B^2)}$.

 [Watch Video Solution](#)

4. Find $\frac{dy}{dx}$ if $x \cos x = 2 \sin y$



Watch Video Solution

5. Find $\frac{dy}{dx}$ if $y = ax^2 + 2hxy + by^2$



Watch Video Solution

6. Find $\frac{dy}{dx}$ if $ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0$



Watch Video Solution

Exercise Questions Asked In Previous 13 Years Exam

1. If the pair of lines $ax^2 + 2(a + b)xy + by^2 = 0$ lie along diameters of a circle and divide the circle into four sectors such that the area of one of the sectors is thrice the area of another sector then

A. $3a^2 + 2ab + 3b^2 = 0$

B. $3a^2 + 10ab + 3b^2 = 0$

C. $3a^2 - 2ab + 3b^2 = 0$

D. $3a^2 - 10ab + 3b^2 = 0$

Answer: A



[Watch Video Solution](#)

2. If one of the lines of $my^2 + (1 - m^2)xy - mx^2 = 0$ is a bisector of the angle between the lines $xy = 0$, then m is

A. $-\frac{1}{2}$

B. -2

C. 1

D. 2

Answer: C



[Watch Video Solution](#)

