



## CHEMISTRY

### BOOKS - PATHFINDER CHEMISTRY (BENGALI ENGLISH)

## SOLUTIONS

#### Question Bank

1. Calculate the mass percentage of benzene ( $C_6H_6$ ) and carbon tetrachloride ( $CCl_4$ ) if 22g of benzene is dissolved in 122g of carbon tetrachloride.



Watch Video Solution

2. Calculate the mole fraction of benzene in solution containing 30% by mass in carbon tetrachloride.

 [Watch Video Solution](#)

3. Calculate the molarity and normality of a solution containing 5g  $Na_2CO_3$  in 250 mL solution.

 [Watch Video Solution](#)

4. Calculate the mass of urea ( $NH_2CONH_2$ ) required in making 2.5 kg of 0.25 molal aqueous solution.

 [Watch Video Solution](#)

5. Henry's law constant for  $CO_2$  in water is  $1.67 \times 10^8$  Pa at 298 K. Calculate the quantity of  $CO_2$  in 500 mL of soda water packed under 2.5 atm  $CO_2$  pressure at 298 K.

 [Watch Video Solution](#)

6. Vapour pressure of chloroform ( $CHCl_3$ ) and dichloromethane ( $CH_2Cl_2$ ) at 298 K are 200 mm Hg and 415 mm Hg respectively.

Calculate the vapour pressure of the solution prepared by mixing 25.5 g of  $CHCl_3$  and 40 g of  $CH_2Cl_2$  at 298 K and,

 [Watch Video Solution](#)

7. A solution containing 30 g of non-volatile solute exactly in 90 g of water has a vapour of 2.8 kPa at 298 K. Further, 18g of water is then added to the solution and the new vapour pressure becomes 2.9 kPa at 298 K.

Calculate,

molar mass of the solute

 [Watch Video Solution](#)

8. A solution containing 30 g of non-volatile solute exactly in 90 g of water has a vapour of 2.8 kPa at 298 K. Further, 18g of water is then added to the solution and the new vapour pressure becomes 2.9 kPa at 298 K.

Calculate,

molar mass of the solute

 [Watch Video Solution](#)

9. A mixture of chloroform and acetone shows negative deviation from Raoult's law. Explain it

 [Watch Video Solution](#)

10. The vapour pressure of pure liquids A and B are 450 and 700mm Hg respectively, at 350K. Find out the composition of the liquid mixture if total vapour pressure is 600 mm Hg. Also find the composition of the vapour phase.

 [Watch Video Solution](#)

11. The vapour pressure of pure benzene at a certain temperature is 0.850 bar. A non-volatile, non-electrolyte solid weighing 0.5 g when added to 39.0 g of benzene (molar mass  $78 \text{ g mol}^{-1}$ ), vapour pressure of the solution, then, is 0.845 bar. What is the molar mass of the solid substance ?

 [Watch Video Solution](#)

12. Vapour pressure of pure water at 298 K is 23.8 mm Hg. 50 g of urea ( $\text{NH}_2\text{CONH}_2$ ) is dissolved in 850 g of water. Calculate the vapour pressure of water for the solution and its relative lowering.

 [Watch Video Solution](#)

13. Boiling point of water at 750 mm Hg is  $99.63^\circ \text{C}$ . How much sucrose is to be added to 500 g of water such that it boils at  $100^\circ \text{C}$  ?

 [Watch Video Solution](#)

14. Calculate the mass of ascorbic acid (Vitamin C,  $C_6H_8O_6$ ) to be dissolved in 75 g of acetic acid to lower its melting by  $1.5^\circ C$ .  $K_f = 3.9Kkgmol^{-1}$ .

 [Watch Video Solution](#)

15. 0.6 mL of acetic acid ( $CH_3COOH$ ), having density  $1.06gmL^{-1}$ , is dissolved in 1 litre of water. The depression in freezing point observed for this strength of acid was  $0.0205^\circ C$ . Calculate the van't Hoff factor and the dissociation constant of acid

 [Watch Video Solution](#)

16. 2g of benzoic acid ( $C_6H_5COOH$ ) dissolved in 25 g of benzene shows a depression in freezing point equal to 1.62 K. Molal depression constant for benzene is  $4.9kkgmol^{-1}$ . What is the percentage association of acid if it forms dimer in solution ?

 [Watch Video Solution](#)

17. Calculating the osmotic pressure in pascals exerted by a solution prepared by dissolving 1.0 g of a polymer of molar mass 185,000 in 450 mL of water at  $37^{\circ}C$ .

 [Watch Video Solution](#)

18. The vapour pressure of a given liquid will decrease if :

- A. surface area of liquid is decreased
- B. the volume of liquid in the container is decreased
- C. the volume of the vapour phase is increased
- D. the temperature is decreased

**Answer:**

 [Watch Video Solution](#)

19. Raoult's law is obeyed by each constituent of a binary liquid solution when :

- A. the force of attraction between like molecules are greater than those between unlike molecules
- B. the force of attraction between like molecules are smaller than those between unlike molecules
- C. the force of attraction between like molecules are identical with those between unlike molecules
- D. the volume occupied by unlike molecules are different

**Answer:**



[Watch Video Solution](#)

20. For a dilute solution, Raoult's law states that :



- A. the lowering of vapour pressure is equal to the mole fraction of solute
- B. the relative lowering of vapour pressure is equal to the mole fraction of solute.
- C. the relative lowering of vapour pressure is proportional to the amount of solute in solution
- D. the vapour pressure of the solution is equal to the mole fraction of solvent

**Answer:**



**Watch Video Solution**

21. 6.0 g of urea (molecular weight=60) was dissolved in 9.9 moles of water. If the vapour pressure of pure water is  $P^\circ$ , the vapour pressure of solution is :

A.  $0.10P^\circ$

B.  $1.10P^\circ$

C.  $0.90P^\circ$

D.  $0.99P^\circ$

**Answer:**

 [Watch Video Solution](#)

**22.** The vapour pressure of an aqueous solution of sucrose at 373 K is found to be 750 mm Hg. The molality of the solution at the same temperature will be :

A. 0.26

B. 0.73

C. 0.74

D. 0.039

**Answer:**



[Watch Video Solution](#)

23. If two liquid A and B form minimum boiling azeotrope at same specific composition then

- A. A-B interactions are stronger than those between A-A or B-B
- B. Vapour pressure of solution increases because more number of molecules of liquid A and B can escape from the solution.
- C. Vapour pressure of solution decreases because less number of molecules of only one of the liquids escape from the solution
- D. A-B interaction are weaker than those between A-A or B-B

**Answer:**



[Watch Video Solution](#)

24. On adding acetone to methanol some of the hydrogen bonds between methanol molecules break. Hence

A. at specific composition, methanol-acetone mixture will form minimum boiling azeotrope and will show positive deviation from Raoult's law

B. at specific composition, methanol-acetone mixture will form maximum boiling azeotrope and will show positive deviation from Raoult's law

C. at specific composition, methanol-acetone mixture will form minimum boiling azeotrope and will show negative deviation from Raoult's law

D. at specific composition, methanol-acetone mixture will form maximum boiling azeotrope and will show negative deviation from Raoult's law

**Answer:**



Watch Video Solution

25. Which of the following solutions can be separated into its pure components by fractional distillation ?

A. Benzene-toluene

B. Water- $HNO_3$

C. Water-HCl

D. Water- $C_2H_5OH$

Answer:



Watch Video Solution

26. A solution is made by mixing the two components. It is said to be ideal if

A.  $\Delta V = 0$ ,  $\Delta H = +V_e$ , Raoult's law is obeyed

B.  $\Delta V = +V_e$ ,  $\Delta H = -V_e$ , Raoult's law is obeyed

C.  $\Delta V = 0$ ,  $\Delta H = 0$ , Raoult's law is obeyed

D.  $\Delta V = 0$ ,  $\Delta H = -V_e$ , Raoult's law is obeyed

**Answer:**

 [Watch Video Solution](#)

**27. Select correct statements):**

A. Heat of evaporation for a pure solvent and for a solution are similar because similar intermolecular forces between solvent molecules must be overcome in both cases.

B. Entropy change between solution and vapour is smaller than the entropy change between pure solvent and vapour.

C. Boiling point of the solution is large than that of the pure solvent.

D. All are correct statements

**Answer:**

 [Watch Video Solution](#)

28. One molal solution of a carboxylic acid in benzene shows the elevation of boiling point of 1.518 K. The degree of association for dimerization of the acid in benzene is ( $K_b$  for benzene =  $2.53 \text{ K kg mol}^{-1}$ ):

A. 60 %

B. 70 %

C. 75 %

D. 80 %

**Answer:**

 [Watch Video Solution](#)

29. The boiling point elevation constant for toluene is  $3.32 \text{ K kg mol}^{-1}$ . The normal boiling point of toluene is  $110.7^\circ\text{C}$ . The enthalpy of evaporation of toluene would be nearly :

A.  $17.0 \text{ kJ mol}^{-1}$

B.  $34.0 \text{ kJ mol}^{-1}$

C.  $51.0 \text{ kJ mol}^{-1}$

D.  $68.0 \text{ kJ mol}^{-1}$

**Answer:**

 [Watch Video Solution](#)

30. Two solutions of glucose have osmotic pressure 1.5 and 2.5 atm. respectively. 1 L of first solution is mixed with 2 L of second solution. The osmotic pressure of the resultant solution will be :

A. 1.62 atm



B. 6.12 atm

C. 1.26 atm

D. 2.16 atm

**Answer:**

 [Watch Video Solution](#)

31. Calculating the depression in the freezing point of water when 10g of  $CH_3CH_2CHClCOOH$  is added to 250g of water.  $K_a=1.4 \times 10^{-3}$ ,  $k_f=1.86 \text{ K kg mol}^{-1}$

A. 0.56

B. 0.84

C. 0.65

D. 0.27

**Answer:**

32. Following are equimolal aqueous solutions

Urea

KCl

$MgCl_2$

$Na_3PO_4$  Arrange them in order of increasing osmotic pressure

A.  $A < B < C < D$

B.  $D < A < B < C$

C.  $B < A < C < D$

D.  $C < A < B < D$

Answer:

33. 18 g glucose ( $C_6H_{12}O_6$ ) is added to 178.2 g water. The vapor pressure of water (in torr) for this aqueous solution is.

- A. 759
- B. 7.6
- C. 76
- D. 752.4

**Answer: D**

 [Watch Video Solution](#)

34. The van't Hoff factor ( $i$ ) for a dilute aqueous solution of the strong electrolyte barium hydroxide is

- A. 3
- B. 0
- C. 1

D. 2

**Answer: A**



[Watch Video Solution](#)

**35.** Which one of the following is incorrect for ideal solution ?

A.  $\Delta G_{mix} = 0$

B.  $\Delta H_{mix} = 0$

C.  $\Delta U_{mix} = 0$

D.  $\Delta P_{obs} - P_{calculated\ by\ Raoult's\ law} = 0$

**Answer: A**



[Watch Video Solution](#)

36. Which of the following statements about the composition of the vapour over an ideal 1:1 molar mixture of benzene and toluene is correct ? Assume that that the temperature is constant at  $25^{\circ}C$ . (Given, Vapour pressure Data at  $25^{\circ}C$ , benzene = 12.8 KPa, toluene = 3.85 KPa)

- A. The vapour will contain equal amounts of benzene and toluene.
- B. Not enough information is given to make a prediction.
- C. The vapour will contain a higher percentage of benzene.
- D. The vapour will contain a higher percentage of toluene.

**Answer: C**

 [Watch Video Solution](#)

37. At  $100^{\circ}C$  the vapour pressure of a solution of 6.5 g of a solute in 100 g water is 732 mm. If  $K_b = 0.52$ , the boiling point of this solution will be.

- A.  $102^{\circ}C$

B.  $103^{\circ} C$

C.  $101^{\circ} C$

D.  $100^{\circ} C$

**Answer: C**

 [Watch Video Solution](#)

**38.** Consider the following liquid - vapour equilibrium Liquid  $\rightleftharpoons$  Vapour

Which of the following relations is correct ?

A.  $\frac{d/nP}{dT^2} = \frac{-\Delta H}{T^2}$

B.  $\frac{d/nP}{dT} = \frac{\Delta H}{R} T^2$

C.  $\frac{d/nG}{dT^2} = \frac{\Delta H}{R} T^2$

D.  $\frac{d/nP}{dT} = \frac{-\Delta H}{RT}$

**Answer: B**

 [Watch Video Solution](#)

39. If  $P^0$  and  $P$  are the vapour pressure of the pure solvent and solution and  $n_1$  and  $n_2$  are the moles of solute and solvent respectively in the solution then the correct relation between  $P$  and  $P^0$  is

A.  $P^0 = P \left[ \frac{n_1}{n_1 + n_2} \right]$

B.  $P^0 = P \left[ \frac{n_2}{n_1 + n_2} \right]$

C.  $P = P^0 \left[ \frac{n_2}{n_1 + n_2} \right]$

D.  $P = P^0 \left[ \frac{n_1}{n_1 + n_2} \right]$

**Answer: C**

 [Watch Video Solution](#)

40. The vapour pressure of acetone at  $20^\circ C$  is 185 torr. When 1.2 g of a non-volatile solute was dissolved in 100 g of acetone at  $20^\circ C$  its vapour pressure was 183 torr. The mass ( $gmol^{-1}$ ) of the substance is.

A. 64

B. 128

C. 488

D. 32

**Answer: A**

 [Watch Video Solution](#)

**41.** Of the following 0.10 m aqueous solutions, which one will exhibit the largest freezing point depression ?

A. KCl

B.  $C_6H_{12}O_6$

C.  $Al_2(SO_4)_3$

D.  $K_2SO_4$

**Answer: C**





Watch Video Solution

42. A solution of 1.25 g of 'P' in 50g of water lowers freezing point by  $0.3^{\circ}C$ . Molar mass of 'P' is 94.  $K_f(\text{water})$  is  $1.86 \text{ K kg mol}^{-1}$ . The degree of association of 'P' in water is [Assume dimerisation of P]

A. 80 %

B. 60 %

C. 65 %

D. 75 %

Answer: A



Watch Video Solution

43.  $25\text{cm}^3$  of oxalic acid completely neutralised 0.064g of sodium hydroxide. Molarity of the oxalic acid solution is

A. 0.064

B. 0.045

C. 0.015

D. 0.032

**Answer: D**



**Watch Video Solution**

**44.** 5.5 mg of nitrogen gas dissolves in 180 g of water at 273K and 1 atm pressure due to nitrogen gas. The mole fraction of nitrogen in 180g of water at 5 atm nitrogen pressure is approximately.

A.  $1 \times 10^{-6}$

B.  $1 \times 10^{-5}$

C.  $1 \times 10^{-3}$

D.  $1 \times 10^{-4}$

**Answer: D**

 [Watch Video Solution](#)

**45.** What will be the value of molality for an aqueous solution of 10 % w/w NaOH? (Na = 23, O = 16, H = 1)

A. 5

B. 2.778

C. 10

D. 2.5

**Answer: B**

 [Watch Video Solution](#)

**46.** If 10ml of 0. 1M aqueous solution of NaCl is divided into 1000 drops of equal volume, what will be the concentration of one drop ?

A. 0.10M

B. 0.01M

C. 0.001M

D. 0.0001M

**Answer: A**



**Watch Video Solution**

**47.** What is the molality of a solution containing 200 mg of urea (molar mass  $60\text{g mol}^{-1}$ ) dissolved in 40g of water?

A. 0.0825

B. 0.825

C. 0.498

D. 0.0013

**Answer: A**



Watch Video Solution

48. The molarity of a solution obtained by mixing 750 mL of 0.5(M) HCl with 250 mL of 2(M) HCl will be

A. 0.875 M

B. 1

C. 1.75 M

D. 0.0975 M

Answer: A





Watch Video Solution

49. Identify the mixture that shows positive deviation from Raoult's law.

A.  $CHCl_3 + (CH_3)_2CO$

B.  $(CH_3)_2CO + C_6H_5NH$



**Answer: D**



**Watch Video Solution**

50. The temperature at which the vapour pressure of a liquid becomes equal to external (atmospheric) pressure is its

A. melting point

B. sublimation point

C. inversion point

D. BOILING POINT

**Answer: D**



**Watch Video Solution**

51. An aqueous solution freezes at  $-0.186^{\circ}\text{C}$  then elevation in boiling point is ( $K_b = 0.512$ ,  $K_f = 1.86$ )

- A.  $0.0512^{\circ}\text{C}$
- B.  $100.0512^{\circ}\text{C}$
- C.  $-0.0512^{\circ}\text{C}$
- D. None of these

**Answer: A**



[Watch Video Solution](#)

52. Which one of the following solutions exhibits the maximum elevation in boiling point ?

- A.  $0.1\text{ m NaCl}$
- B.  $0.1\text{ m FeCl}_3$
- C.  $0.1\text{ m CaCl}_2$

D.  $0.1mBaCl_2$

**Answer: B**

 [Watch Video Solution](#)

53. Which colligative property is most suitable for the determination of molecular masses of polymer?

- A. elevation in boiling point
- B. osmotic pressure
- C. optical activity
- D. relative lowering of vapour pressure

**Answer: C**

 [Watch Video Solution](#)



54. Which one of the statements given below concerning properties of solution, describes a colligative effect ?

A. Boiling point of pure water decreases by the addition of ethanol.

B. Vapour pressure of pure water decreases by the addition of nitric acid

C. Vapour pressure of pure benzene decreases by the addition of naphthalene.

D. Vapour pressure of pure benzene increases by the addition of toluene.

**Answer: C**



**Watch Video Solution**

55.  $6.02 \times 10^{20}$  molecules of urea are present in 100 mL of its solution.

The concentration of solution is

A. 0.02 M

B. 0.01 M

C. 0.001 M

D. 0.1 M

**Answer: B**

 [Watch Video Solution](#)

**56.** How many gram of concentrated nitric acid solution should be used  
be prepare 250 mL of 2.0 M  $\text{HNO}_3$  ?

A. 45.0 g conc.  $\text{HNO}_3$

B. 90.0 g conc.  $\text{HNO}_3$

C. 70.0 g conc.  $\text{HNO}_3$

D. 50.0 g conc.  $\text{HNO}_3$

**Answer: A**



[Watch Video Solution](#)

57. Distribution law was given by

A. Ostwald

B. Nernst

C. Henry

D. van't Hoff

**Answer: B**



[Watch Video Solution](#)

58. Which of the following is not a colligative property ?

A. Depression in freezing point

B. Osmotic pressure

C. Elevation of boiling point

D. Increase in freezing point

**Answer: D**

 [Watch Video Solution](#)

59. The osmotic pressure of 12 % solution of cane sugar at  $17^{\circ}C$  is

A. 2.42 atm

B. 4.33 atm

C. 8.35 atm

D. 16.30atm

**Answer: C**

 [Watch Video Solution](#)

60. The normality of orthophosphoric acid having purity of 70 % by weight and specific gravity 1.54 is

- A. 11 N
- B. 22 N
- C. 33 N
- D. 44 N

**Answer: C**

 [Watch Video Solution](#)

61. The molarity of  $H_2SO_4$  solution, which has a density 1.84 g/cc at  $35^\circ C$  and contains 98 % by weight is

- A. 1.84 M
- B. 18.4 M
- C. 20.6 M

D. 24.5 M

**Answer: B**



[Watch Video Solution](#)

62. For a dilute solution containing 2.5 g of a non-volatile non-electrolyte solute in 100 g of water, the elevation in boiling point at 1 atm pressure is  $2^\circ C$ . Assuming concentration of solute is much lower than the concentration of solvent, the vapour pressure (mm of Hg) of the solution is ( $take\ k_b = 0.76\ K\ kg\ mol^{-1}$ ).

A. 726

B. 740

C. 736

D. 718

**Answer: A**



[Watch Video Solution](#)

63.  $k_f$  for water is  $1.86 \text{ k kg mol}^{-1}$ . If your automobile radiator holds 1.0 kg of water, how many grams of ethylene glycol ( $C_2H_6O_2$ ) must you add to get the freezing point of the solution lowered to  $-2.8^\circ C$ ?

A. 72 g

B. 93 g

C. 39 g

D. 27 g

Answer: B



Watch Video Solution

64. The density of a 3M sodium thiosulphate ( $Na_2S_2O_3$ ) solution is 1.25 g/mL. Calculate the percent by weight of sodium thiosulphate.

A. 12.65 %

B. 37.92 %

C. 0.87 %

D. 63.21 %

**Answer: B**



**Watch Video Solution**

**65.** Two solutions of HCl, A and B have concentrations of 0.5 N and 0.1 M respectively. The volume of solutions A and B required to make 2 L of 0.2 N HCl are

A. 0.5 L of A + 1.5 L of B

B. 1.5 L of A + 0.5 L of B

C. 1.0 L of A + 1.0 L of B

D. 0.75 L of A + 1.25 L of B

**Answer: A**



 Watch Video Solution

66. The highest osmotic pressure corresponds to the following solution

- A.  $\frac{M}{10}$  urea
- B.  $\frac{M}{10}$  glucose
- C.  $\frac{M}{10}$  HCl
- D.  $\frac{M}{10}$   $BaCl_2$

Answer: D

 Watch Video Solution

67. The correct order of increasing boiling points of the following aqueous solutions

(I) 0.0001 M NaCl (II) 0.000 1 M Urea (III) 0.001 M  $MgCl_2$  (IV) 0.01 M NaCl

- A.  $I < II < III < IV$

B.  $IV < III < II < I$

C.  $II < I < III < IV$

D.  $III < II < IV < I$

**Answer: C**

 [Watch Video Solution](#)

**68.** 58.4 g of NaCl and 180 g of glucose were separately dissolved in 1000 mL of water. Identify the correct statement regarding the elevation of boiling point (b.p.) of the resulting solutions.

A. NaCl solution will show higher elevation of boiling point

B. Glucose solution will show higher elevation of boiling point

C. Both the solutions will show equal elevation of boiling point

D. The boiling point elevation be shown by neither of the solutions

**Answer: A**

 [Watch Video Solution](#)

69. Mixture of chloroform and acetone shows a negative deviation from Raoult's law. Explain.

- A. Acetone-benzene
- B. Acetone-ethanol
- C. Benzene-methanol
- D. Acetone-chloroform

**Answer: D**

 [Watch Video Solution](#)

70. The relative lowering of vapour pressure of a dilute solution is equal to mole fraction of solute present in the solution. Which of the following low is state this ?

- A. Henry low
- B. Avogadro's law
- C. Raoult's law
- D. Law of definite proportion

**Answer: C**

 [Watch Video Solution](#)

71. The mass of a non-volatile solute of molar mass  $40 \text{ g mol}^{-1}$  that should be dissolved in 114 g of octane to lower its vapour pressure by 20 % , is

- A. 11.4 g
- B. 9.8 g
- C. 12.8 g
- D. 10 g

**Answer: D**

 [Watch Video Solution](#)

72. The relative lowering of vapour pressure of an aqueous solution containing a non-volatile solute is 0.0125. The molality of the solution is

A. 0.7

B. 0.3

C. 0.125

D. 0.07

**Answer: A**

 [Watch Video Solution](#)

73. Which of the following will represent the normality of 10%  $\left(\frac{w}{V}\right)$  acetic acid

A. 1.66 N

B. 16.6 N

C. 3.32 N

D. 33.2 N

**Answer: A**



[Watch Video Solution](#)

**74.** Milk is an emulsion. Identify the dispersed phase and the emulsifier in milk.



[Watch Video Solution](#)

**75.** In which case van't Hoff factor are equal ?

A. 0.5

B. 1

C. 2

D. 6

**Answer: C**



**Watch Video Solution**

76.  $P_A$  and  $P_B$  are the vapour pressure of pure liquid components, A and B, respectively of an ideal binary solution. If  $X_A$  represents the mole fraction of component A, the total pressure of the solution will be

A.  $P_A + X_A(P_B - P_A)$

B.  $P_A + X_A(P_A - P_B)$

C.  $P_B + X_A(P_B - P_A)$

D.  $P_B + X_A(P_A - P_B)$

**Answer: D**



**Watch Video Solution**

77. Vapour pressure of chloroform ( $CHCl_3$ ) and dichloromethane ( $CH_2Cl_2$ ) at 298 K are 200 mm Hg and 415 mm Hg respectively.

Calculate the vapour pressure of the solution prepared by mixing 25.5 g of  $CHCl_3$  and 40 g of  $CH_2Cl_2$  at 298 K and,

- A. 17.9 mmHg
- B. 615.0 mmHg
- C. 347.9 mmHg
- D. 90.63 mmHg

**Answer: D**

 [Watch Video Solution](#)

78. A solution has higher osmotic pressure than its standard solution. Which of the following term will be used for this solution ?



- A. Isotonic
- B. Hypertonic
- C. Dilute
- D. Hypotonic

**Answer: B**

 [Watch Video Solution](#)

79. The vapour pressure lowering caused by the addition of 100 g of sucrose (molecular mass = 342) to 1000 g of water, if the vapour pressure of pure water at  $25^{\circ}C$  is 23.8 mm Hg, is

 [Watch Video Solution](#)

80. Molarity of a given orthophosphoric acid solution is 3 M. Its normality is

A. 9 N

B. 0.3 N

C. 3 N

D. 1 N

**Answer: A**



**Watch Video Solution**

**81.** 138 g of ethyl alcohol is mixed with 72 g of water. The ratio of mole fraction of alcohol to water is

A. 3 : 4

B. 1 : 2

C. 1 : 4

D. 1 : 1

**Answer: A**



Watch Video Solution

82. In which case Raoult's law is not applicable ?

- A. 1 M NaCl
- B. 1 M urea
- C. 1M glucose
- D. 1 M sucrose

Answer: A



Watch Video Solution

83. The freezing point of one molal NaCl solution assuming NaCl to be 100 % dissociated in water is (molal depression constant is 1.86)

- A.  $-2.72^{\circ}C$
- B.  $-3.72^{\circ}C$

C.  $2.72^{\circ}C$

D.  $3.72^{\circ}C$

**Answer: B**



[Watch Video Solution](#)

**84.** The vapour pressure of water depends upon

A. Surface area of container

B. Volume of container

C. Temperature

D. All

**Answer:**



[Watch Video Solution](#)

85. A vessel has nitrogen gas and water vapour in equilibrium with liquid water at a total pressure of 1 atm. The partial pressure of water vapours is 0.3 atm. The volume of this vessel is reduced to one third of the original volume, at the same temperature, then total pressure of the system is: (Neglect volume occupied by liquid water)

A. 3.0 atm

B. 1 atm

C. 3.33 atm

D. 2.4 atm

**Answer:**



[Watch Video Solution](#)

86. For a dilute solution, Raoult's law states that :

- A. The lowering of vapour-pressure is equal to the mole-fraction of solute.
- B. The relative lowering of vapour-pressure is equal to the mole-fraction of solute.
- C. The relative lowering of vapour-pressure is proportional to the amount of the solution is equal to the mole-fraction of solvent.
- D. The vapour-pressure of the solution is equal to the mole-fraction of solvent.

**Answer:**

 [Watch Video Solution](#)

**87.** The mathematical expression of Raoult's law is:

Where  $n$  is no. of moles of solute &  $N$  is no. of moles of solvent

$$A. \frac{P^\circ - P}{P^\circ} = n/N$$

$$B. \frac{P^\circ - P}{P^\circ} = N/n$$

$$C. \frac{P^\circ - P}{P} = n/N$$

$$D. \frac{P^\circ - P}{P^\circ} = n \times N$$

**Answer:**



**Watch Video Solution**

**88.** Total vapour pressure of mixture of 1 mol of volatile component A ( $P_A^\circ = 100\text{mmHg}$ ) and 3 mol of volatile component B ( $P_B^\circ = 60\text{mmHg}$ ) is 75 mm. For such case

- A. there is positive deviation from Raoult's law
- B. boiling point has been lowered
- C. force of attraction between A and B is smaller than that between A and A or between B and B
- D. all the above statement are correct

**Answer:**



[Watch Video Solution](#)

89. Two liquids A and B have vapour pressure in the ratio  $P_A^\circ : P_B^\circ = 2:3$  at a certain temperature. Assume A and B form an ideal solution and the ratio of mole fractions of A to B in the vapour phase is 1:3, then the mole fraction of A in the solution at the same temperature is :

A. 1/3

B. 2/3

C. 1/4

D. 3/4

**Answer:**



[Watch Video Solution](#)



90. Which of the following azeotropic solutions has the b.p. less than b.p. of the constituents A and B ?

A.  $CHCl_3$  and  $CH_3COCH_3$

B.  $CS_2$  and  $CH_3COCH_3$

C.  $CH_3CH_2OH$  and  $CH_3COCH_3$

D.  $CH_3CHO$  and  $CS_2$

Answer:



Watch Video Solution

91. The mass of glucose that should be dissolved in 100 g of water in order to produce same lowering of vapour pressure as is produced by dissolving 1 g of urea in 50 g of water is

A. 1g

B. 2g

C. 6g

D. 12g

**Answer:**



[Watch Video Solution](#)

**92.** The vapour pressure of water at room temperature is lowered by 5 % by dissolving a solute in it, then the approximate molality of solution is

A. 2

B. 1

C. 4

D. 3

**Answer:**



[Watch Video Solution](#)

93. The vapour pressure of a dilute aqueous of glucose is 750 mm of mercury at 373 K. The mole fraction of solute is

A. 44470

B. 1/7.6

C. 1/35

D. 1/76

**Answer:**

 [Watch Video Solution](#)

94. Relative decrease in vapour pressure of an aqueous solution containing 2 moles  $[Cu(NH_3)_3Cl]$  Cl in 3 moles  $H_2O$  is 0.50. On reaction with  $AgNO_3$  this solution will form (assuming no change in degree of ionisation of substance on adding  $AgNO_3$ ).

A. 1mol AgCl

B. 0.25mol AgCl

C. 0.5 mol AgCl

D. 0.40 mol AgCl

**Answer:**



[Watch Video Solution](#)

95. A solution of one mole of benzoic acid in 15 moles of benzene produces a relative lowering in vapour pressure of  $\frac{1}{31}$ . The mol. wt. of benzoic acid is

A. 122

B. 244

C. 61

D. 189

**Answer:**



Watch Video Solution

96. When mercuric iodide is added to the aqueous solution of potassium iodide

- A. freezing point is raised
- B. freezing point is lowered
- C. freezing point is does not change
- D. boiling point is does not change

**Answer:**



Watch Video Solution

97. 50g of antifreeze (ethylene glycol) is added to 200 g water. What amount of ice will separate out at  $-9.3^{\circ}C$ ?

- A. 42 mg

B. 42 g

C. 38.71 g

D. 38.71 mg

**Answer:**



[Watch Video Solution](#)

**98.** Which of the following 0.1 M aqueous solution will have the lowest freezing point ?

A. Potassium sulphate

B. Sodium chloride

C. Urea

D. Glucose

**Answer:**



[Watch Video Solution](#)

99. The freezing point of equimolal aqueous solutions will be highest for

A.  $C_6H_5NH_3Cl$ (aniline hydrochloride)

B.  $Ca(NO_3)_2$

C.  $La(NO_3)_3$

D.  $C_6H_{12}O_6$

**Answer:**



[Watch Video Solution](#)

100. 15 g of a solute in 100 g water makes a solution freeze at  $-1^\circ C$ . 30 g of a solute in 100 g of water will give a depression in freezing point equal to :

A.  $-2^\circ C$

B.  $0.5^\circ C$

C.  $2^{\circ}C$

D.  $1^{\circ}C$


**Answer:**




**Watch Video Solution**

**101.** Which of the following has been arranged in order of decreasing freezing point ?


A.



B.




C.





D.



**Answer:**

 [Watch Video Solution](#)

**102.** The molecular weight of benzoic acid in benzene as determined by depression in freezing point method corresponds to

- A. ionization of benzoic acid
- B. dimerization of benzoic acid
- C. trimerization of benzoic acid
- D. solvation of benzoic acid

**Answer:**

 [Watch Video Solution](#)

103. The fraction of phenol dimerised in benzene if 20 g of phenol in 1 kg benzene exhibits a freezing point depression of 0.69 K. ( $K_f$ benzene) =5.12 kkg/mol),(MW of phenol=94)

A. 0.74

B. 0.37

C. 0.46

D. 0.64

Answer:

 [Watch Video Solution](#)

104. When 20 g of naphthoic acid ( $C_{11}H_8O_2$ ) is dissolved in 50 g of benzene ( $K_f = 1.72Kkgmol^{-1}$ ), a freezing point depression of 2 K is observed. The van't Hoff factor(i) is

A. 0.5

B. 1

C. 2

D. 3

**Answer:**



[Watch Video Solution](#)

**105.** Pressure cooker reduces cooking time because

A. the heat is more evenly distributed inside the cooker

B. a large flame is used

C. boiling point of water is elevated

D. whole matter is converted into steam

**Answer:**



[Watch Video Solution](#)

106.  $y$  g of non-volatile solute of molar mass  $M$  is dissolved in 250 g benzene. If  $K_b$  is molal elevation constant, the value of  $\Delta T$  is given by

A.  $\frac{4M}{k_b y}$

B.  $\frac{4k_b y}{M}$

C.  $k_b \frac{y}{4M}$



D.  $k_b \frac{y}{M}$



**Answer:**



**Watch Video Solution**

107. A complex of iron and cyanide ions is 100 % ionised at 1m (molal). If its elevation in b.p. is  $2.08^\circ C$ . Then the complex is ( $K_b = 0.52 K mol^{-1} kg$ ):





**Answer:**

 [Watch Video Solution](#)

**108.** The elevation in boiling point of a solution of 13.44 g of  $CuCl_2$  in 1 Kg of water using the following information will be:

(Molecular weight of  $CuCl_2 = 134.4$  and  $K_b = 0.52Kmolal^{-1}$ )

A. 0.16

B. 0.05

C. 0.1

D. 0.2

**Answer:**

 [Watch Video Solution](#)

**109.** A solution of glucose ( $C_6H_{12}O_6$ ) is isotonic with 4 g of urea ( $NH_2 - CO - NH_2$ ) per liter of solution. The concentration of glucose is

- A. 4 g/l
- B. 8 g/l
- C. 12 g/l
- D. 14 g/l

**Answer:**

 [Watch Video Solution](#)

**110.** Osmotic pressure of 30 % solution of glucose is 1.20 atm and that of 3.42 % solution of cane sugar is 2.5 atm. The osmotic pressure of the mixture containing equal volumes of the two solutions will be

A. 2.5 atm

B. 3.7 atm

C. 1.85 atm

D. 1.3 atm

**Answer:**

 [Watch Video Solution](#)

111.  $0.004MNa_2SO_4$  is isotonic with  $0.01M$  glucose. Degree of ionisation of  $Na_2SO_4$  is :

A. 75 %

B. 50 %

C. 25 %

D. 85 %

**Answer:**



[Watch Video Solution](#)

**112.** We have 100 mL of 0.1 M KCl solution. To make it 0.2 M,

- A. evaporate 50 mL water
- B. evaporate 50 mL solution
- C. add 0.01 mol KCl
- D. both (1) and (3)

**Answer:**



[Watch Video Solution](#)

**113.** A molal solution is one that contains one mole of a solute in :

- A. 1000 g of the solvent
- B. 1 L of the solvent



C. 1 L of the solution

D. 22.4 L of the solution

**Answer:**

 [Watch Video Solution](#)

**114.** At higher altitudes, water boils at temperature  $> 100^{\circ}C$  because

A. temperature of higher altitudes is low

B. atmospheric pressure is low

C. the proportion of heavy water increases

D. atmospheric pressure becomes more.

**Answer:**

 [Watch Video Solution](#)

115. The solubility of a specific non-volatile salt is 4 g in 100 g of water at  $25^{\circ}\text{C}$ . If 2.0 g, 4.0 g and 6.0 g of the salt added to 100 g of water at  $25^{\circ}\text{C}$ , in system X, Y and Z. The vapour pressure would be in the order :

A.  $X < Y < Z$

B.  $X > Y > Z$

C.  $Z > X = Y$

D.  $X > Y = Z$

Answer: D



Watch Video Solution

116. According to Henry's law, the solubility of a gas in a given volume of liquid increases with increase in

A. Temperature

B. Pressure

C. BOTH (1) & (2)

D. None of these

**Answer: B**

 [Watch Video Solution](#)

117. Calculate the concentration of  $CO_2$  in a soft drink that is bottled with a partial pressure of  $CO_2$  of 4 atm over the liquid at  $25^\circ C$ . The Henry's Law constant for  $CO_2$  in water is  $3.1 \times 10^{-2}$  mol/litre atm at  $25^\circ C$ .

A. 1.24 mol/lit.

B. 0.24 mol/lit

C. 0.124 mol/lit

D. 12.4 mol/lit


**Answer: C**




118. Which of the following curves represents the Henry's law ?




A.



B.



C.



D.

Answer: A

119. An ideal solution has two components A and B. A is More volatile than B, i.e.,  $p_A^0 > p_B^0$  and also  $p_A^0 > P_{\text{total}}$  . If  $x_A$  and  $y_A$  are mole fractions of components A in liquid and vapour phases, then:

A.  $x_A = y_A$

B.  $x_A > y_A$

C.  $x_A < y_A$

D. Data insufficient

Answer: C



Watch Video Solution

120. Two liquids A and B form an ideal solution. At 300 k, the vapour pressure of a solution containing 1 mole of A and 3 moles of B is 550 mm of Hg. At the same temperature, if one mole of B is added to this solution, the vapour pressure of the solution increases by 10 mm of Hg. Determine the vapour pressure of A and B in their pure states.

A. 400, 600

B. 500, 500

C. 600, 400

D. None of these

**Answer: A**



**Watch Video Solution**

**121.** Water and chlorobenzene are immiscible liquids. Their mixture boils at  $90^\circ C$  under a reduced pressure of  $7.82 \times 10^4$  Pa. The vapour pressure of pure water at  $90^\circ C$  is  $7.03 \times 10^4$  Pa. On weight percent basis, chlorobenzene in the distillate is equal to (mol. wi. of chlorobenzene is  $112.5 \text{ gmol}^{-1}$ )

A. 50

B. 60

C. 41

D. 80

**Answer: C**

 [Watch Video Solution](#)

**122.** Two liquids having vapour pressures  $p_1^0$  and  $p_2^0$  in pure state in the ratio of 2:1 are mixed in the molar ratio of 1:2. The ratio of their moles in the vapour state would be:

A. 1:1

B. 1:2

C. 2:1

D. 3:2

**Answer: A**

 [Watch Video Solution](#)

**123.** For a binary ideal liquid solution, the total pressure of the solution is given as :

A.  $P_{\text{total}} = p_A^0 + (p_A^0 - p_B^0)x_B$

B.  $P_{\text{total}} = p_B^0 + (p_A^0 - p_B^0)x_A$

C.  $P_{\text{total}} = p_B^0 + (p_B^0 - p_A^0)x_A$

D.  $P_{\text{total}} = p_B^0 + (p_B^0 - p_A^0)x_B$

**Answer: B**



**Watch Video Solution**

**124.** The correct order for the bond dissociation enthalpy of Halogen molecules:

A. (A) Bromine > Iodine > Fluorine > Chlorine

B. (B) Fluorine > Chlorine > Bromine > Iodine

C. (C) Iodine > Bromine > Chlorine > Fluorine



D. (D) Chlorine > Bromine > Fluorine > Iodine

**Answer: A**



[Watch Video Solution](#)

**125.** When non ideal solution was prepared by mixing 30 mL chloroform and 50 mL acetone, the volume of mixture will be

A.  $> 80mL$

B.  $= 80mL$

C.  $< 80mL$

D.  $\geq 80mL$

**Answer: C**



[Watch Video Solution](#)

**126.** Two liquids A and B form an ideal solution. What is the vapour pressure of solution containing 2 moles of A and 3 moles of B at 300 K?

[Given : At 300 K, vapour pressure of pure liquid A

$(p_A^0) = 100$  torr and vapour pressure of pure liquid B

$(p_B^0) = 300$  torr]

- A. 200 torr
- B. 140 torr
- C. 180 torr
- D. None of these

**Answer: D**

 [Watch Video Solution](#)

**127.** The vapour pressure of pure liquid A is 70 torr at  $27^\circ C$ . It forms an ideal solution with another liquid B. The mole fraction of B is 0.2 and

total vapour pressure of the solution is 84 torr at  $27^{\circ}C$ . The vapour pressure of pure liquid B at  $27^{\circ}C$  is

- A. 140 torr
- B. 50 torr
- C. 14 torr
- D. 70 torr

**Answer: A**

 [Watch Video Solution](#)

**128.** At  $40^{\circ}C$ , the vapour pressure of pure liquids, benzene and toluene, are 160 mm Hg and 60 mm Hg respectively. At the same temperature, the vapour pressure of an equimolar solution of the two liquids, assuming the ideal solution should be

- A. 140 mmHg
- B. 110 mm Hg

C. 220 mm Hg

D. 100 mm Hg

**Answer: B**

 [Watch Video Solution](#)

**129.** At  $40^\circ C$  the vapour pressures in torr, of methyl alcohol ethyl alcohol solutions is represented by the equation.  $P = 119X_A + 135$  where  $X_A$  is mole-fraction of methyl alcohol, then the value of  $\lim_{X_A \rightarrow 1} \frac{P}{X_A}$  is :

A. 254 torr

B. 135 torr

C. 119 torr

D. 140 torr

**Answer: A**

 [Watch Video Solution](#)

**130.** Two liquids A and B form an ideal solution. The ratio of vapour pressures of pure liquids A and B are 3: 5. If the mole fraction of A in the liquid solution at equilibrium with vapours is 0.4, then the mole-fraction of B in the liquid is

A. 0.3

B. 0.6

C. 4

D. 4.3

**Answer: B**



[Watch Video Solution](#)

**131.** Which will form maximum boiling point azeotrope ?

A.  $C_6H_6 + C_6H_5CH_3$  solution

B.  $C_2H_5OH + H_2O$  solution

C.  $HNO_3 + H_2O$  solution

D. Both (2) and (3)

**Answer: C**

 [Watch Video Solution](#)

**132.** The relative lowering of vapour pressure is equal to the ratio between the number of

A. solute molecules to the solvent molecules

B. solute molecules to the total molecules in solution

C. solvent molecules to the total molecules in the solution

D. solvent molecules to the total number of ions in solution

**Answer: B**

 [Watch Video Solution](#)

**133.** The vapour pressure of a solvent is found to be 400 mm Hg at 300 K. When a certain amount of a nonvolatile solute is added to 50 mol of the solvent its vapour pressure decreases to 360 mm Hg. The amount of solute added is

- A. 2.41 mol
- B. 3.42 mol
- C. 4.54 mol
- D. 5.55 mol

**Answer: D**



[Watch Video Solution](#)

**134.** If relative decrease in vapour pressure is 0.4 for a solution containing 1 mol NaCl in 3 mol of  $H_2O$ , then % ionization of NaCl is

A. 60 %

B. 80 %

C. 40 %

D. 100 %

**Answer: A**

 [Watch Video Solution](#)

**135.** When a solution containing non-volatile solute freezes, which equilibrium would exist ?

A. solid solvent  $\rightleftharpoons$  liquid solvent

B. solid solute  $\rightleftharpoons$  liquid solution

C. solid solute  $\rightleftharpoons$  liquid solvent

D. solid solvent  $\rightleftharpoons$  liquid solution

**Answer: D**





Watch Video Solution

**136.** 2.56 g of sulphur in 100 g of  $CS_2$  has depression in freezing point of  $0.010^\circ$  .  $K_f = 0.1^\circ kg(mol)^{-1}$  . Hence atomicity of sulphur in the solution is

A. 2

B. 4

C. 6

D. 8

**Answer: D**



Watch Video Solution

**137.** The temperature of a city was found to be  $-9.30^\circ C$ . A car was used, whose radiator was filled with 5 L of water. What quantity of anti freezing

agent ethylene glycol were added to water of radiator in order to use the car for travelling ? ( $K_f$  of water  $1.86 \text{ K kg mol}^{-1}$ )

A. 3200 g

B. 1670 g




C. 1550 g

D. 2100 g

**Answer: C**

 [Watch Video Solution](#)

**138.** The freezing point of equimolal aqueous solutions will be highest for



**Answer: D**

 [Watch Video Solution](#)

**139.** KBr is 80 % dissociated in aqueous solution. The freezing point of a 0.5 molal solution is  $K_f = 1.86 K kg mol^{-1}$

A. 271.5 K

B. 273 K

C. 277 K

D. 218 K

**Answer: A**

 [Watch Video Solution](#)

**140.** A 0.2 molal aqueous solution of weak acid HX is 20 % ionized. The freezing point of solution is ( $K_f = 1.86$ )

A.  $-0.45^{\circ}C$

B.  $-0.9^{\circ}C$

C.  $-0.31^{\circ}C$

D.  $-0.53^{\circ}C$

**Answer: A**

 [Watch Video Solution](#)

**141.** Which of the following aqueous solutions has minimum freezing point ?

A. 0.01 m NaCl

B. 0.005m  $MgCl_2$

C. 0.005 m  $C_2H_5OH$

D. 0.005m  $MgSO_4$

**Answer: A**



Watch Video Solution

142. A .001 molal solution of a complex  $[MA_8]$  in water has the freezing point of  $-0.0054^\circ C$ . Assuming 100 % ionization of the complex salt and  $K_f$  for  $H_2O$   $1.86 K m^{-1}$ , write the correct representation for the complex

- A.  $[MA_8]$
- B.  $[MA_7]A$
- C.  $[MA_6]A_2$
- D.  $[MA_5]A_3$

Answer: C



Watch Video Solution

143. Ratio of  $\frac{\Delta T_b}{K_b}$  of 10 g  $AB_2$  and 14 g  $A_2B$  per 100 g of solvent in their respective, solution ( $AB_2$  and  $A_2B$  both are non-electrolytes) is 1 mol/kg in both cases. Hence, atomic wt. of A and B are respectively

A. 100, 40

B. 60, 20

C. 20, 60

D. None of these

**Answer: B**

 [Watch Video Solution](#)

**144.** The elevation in boiling point, when 13.44 g. of freshly prepared  $CuCl_2$ , are added to 1 kg of water is [some useful data,  $K_b(H_2O) = 0.52$  kg K mol<sup>-1</sup> and M.W of  $CuCl_2 = 134.4g$ ]

A. 0.05

B. 0.1

C. 0.16

D. 0.21

**Answer: A**



**Watch Video Solution**

**145.** A water sample contains 9.5 %  $MgCl_2$  and 11.7 %  $NaCl$  (by weight).

Assuming 80 % ionisation of each salt boiling point of water will be (

$K_b = 0.52$ )

A.  $110.01^\circ C$

B. 377.16 K

C. 277.25 K

D.  $102.5^\circ C$

**Answer: B**



**Watch Video Solution**

146. Consider equimolal aqueous solutions of  $NaHSO_4$  and  $NaCl$  with  $\Delta T_b$  and  $\Delta 'T_b$ , as their respective boiling point elevations. The value of  $\lim_{m \rightarrow 0} \frac{\Delta T_b}{\Delta 'T_b}$  will be

- A. 1
- B. 1.5
- C. 3.5
- D. 44257

**Answer: B**



[Watch Video Solution](#)

147. Blood cells retain their normal shape in solutions which are

- A. hypotonic to blood
- B. isotonic to blood
- C. hypertonic to blood



D. equi normal to blood

**Answer: B**

 [Watch Video Solution](#)

**148.** The osmotic pressures of equimolar solutions of urea,  $BaCl_2$  and  $AlCl_3$  will be in the order :

A.  $AlCl_3 > BaCl_2 > \text{urea}$

B.  $BaCl_2 > AlCl_3 > \text{urea}$

C.  $\text{urea} > BaCl_2 > AlCl_3$

D.  $BaCl_2 > \text{urea} > AlCl_3$

**Answer: A**

 [Watch Video Solution](#)

149. Which of the following pair of solutions can be expected to be isotonic at the same temperature ?

- A. 0.1 M urea and 0.1 M NaCl
- B. 0.1M  $NaCl$  and 0.1M  $Na_2SO_4$
- C. 0.1M  $Ca(NO_3)_2$  and 0.1M  $Na_2SO_4$
- D. 0.1 M glucose and 0.2 M  $MgCl_2$

Answer: C

 [Watch Video Solution](#)

150. Sulphur-Dioxide is passed through acidified Potassium-Dichromate solution. Write the reaction with observation.

 [Watch Video Solution](#)

**151.** In the phenomenon of osmosis through the semipermeable membrane

- A. solvent molecules pass from solution to solvent
- B. solvent molecules pass from solvent to solution
- C. solute molecules pass from solution to solvent
- D. solute molecules pass from solvent to solution

**Answer: B**



[Watch Video Solution](#)

**152.** Consider following solution

- (i) 1M aqueous glucose solution
- (ii) 1M aqueous sodium chloride solution
- (iii) 1 M aqueous ammonium phosphate solution
- (iv) 1M benzoic acid in benzene

Select correct statements for the above solution

A. All are isotonic solution

B. (ii) is hypotonic with (i), (ii) and (iv)

C. (iv) is hypertonic with (i), (ii) and (iii)

D. (ii) is hypotonic with (iii) but hypertonic with (i) and (iv)

**Answer: A**

 [Watch Video Solution](#)

**153.** Osmotic pressure of blood is  $7.40 \text{ atm}$  at  $27^\circ \text{C}$ . Number of mol of glucose to be used per L for an intravenous injection that is to have the same osmotic pressure as blood is

A. 0.3

B. 0.2

C. 0.1

D. 0.4

**Answer: A**



[Watch Video Solution](#)

**154.** Which has maximum osmotic pressure at temperature T

- A. 100 mL of 1 M urea solution
- B. 300 mL of 1 M glucose solution
- C. mixture of 100 ml. of 1 M urea solution and 300 mL of 1 M glucose solution
- D. all are isotonic

**Answer: D**



[Watch Video Solution](#)

**155.** The relationship between osmotic pressures ( $\pi_1$ ,  $\pi_2$  and  $\pi_3$ ) at a definite temperature when 1 g glucose, 1 g urea and 1 g sucrose are

dissolved in 1 litre of water is

(assume  $i = 1$  for all)

A.  $\pi_1 > \pi_2 > \pi_3$

B.  $\pi_3 > \pi_1 > \pi_2$

C.  $\pi_2 > \pi_1 > \pi_3$

D.  $\pi_2 > \pi_3 > \pi_2$

**Answer: C**

 [Watch Video Solution](#)

**156.** Which of the following would have the highest osmotic pressure

(Assume that all salts are 90 % dissociated) ?

A. Decimolar aluminium sulphate

B. Decimolar barium sulphate

C. Decimolar sodium sulphate

D. A solution obtained by mixing equal volume of (2) and (3) and filtering

**Answer: A**

 [Watch Video Solution](#)





157. 2 millimolar solution of sodium ferrocyanide is 60% dissociated at  $27^{\circ}C$  Osmotic pressure of the solution is

- A. 2.14 atm
- B. 1.02 atm
- C. 0.167 atm
- D. 0.0234 atm

**Answer: C**

 [Watch Video Solution](#)

158. A complex is written as  $M(en)_y \cdot xBr$ . Its 0.05 molar solution shows 2.46 atm osmotic pressure at  $27^\circ C$ . Assuming 100% ionization and coordination number of metal (III) is six, complex may be



Answer: A



Watch Video Solution

159. If  $M_{\text{normal}}$  is the normal molecular mass and  $\alpha$  is the degree of ionization of  $K_3[Fe(CN)_6]$ , then the abnormal molecular mass of the complex in the solution will be

A.  $M_{\text{normal}}(1 + 2\alpha)^{-1}$



B.  $M_{\text{normal}}(1 + 3\alpha)^{-1}$

C.  $M_{\text{normal}}(1 + \alpha)^{-1}$

D. equal to  $M_{\text{normal}}$

**Answer: B**

 [Watch Video Solution](#)

**160.** The van't Hoff factor  $i$  for an infinitely dilute solution of  $\text{NaHSO}_4$  is

A. 44228

B. 44256

C. 3

D. 2

**Answer: C**

 [Watch Video Solution](#)

161. One mole of a solute A is dissolved in a given volume of a solvent.

The association of the solute take place as follows,  $nA \rightleftharpoons A_n$

If  $\alpha$  is the degree of association of A, the van't Hoff factor  $i$  is expressed as

A.  $i = 1 - \alpha$

B.  $i = 1 + \frac{\alpha}{n}$

C.  $i = (1 - \alpha + \alpha/n)/1$

D.  $i = 1$

**Answer: C**



**Watch Video Solution**

162. The values of observed and calculated molecular mass of  $Ca(NO_3)_2$  are 65.4 and  $164 \text{ g mol}^{-1}$  respectively. The degree of dissociation of  $Ca(NO_3)_2$  will be

A. 75 %

B. 50 %

C. 98 %

D. 80 %

**Answer: C**



**Watch Video Solution**

**163.** Acetic acid dimerises in benzene solution, The Van't Hoff factor for the dimerisation of acetic acid is 0.8. The % of dimerisation of acetic acid is

A. 20 %

B. 40 %

C. 60 %

D. 80 %

**Answer: B**



[Watch Video Solution](#)

**164.** The value of observed and calculated molecular weights of silver nitrate are 92.64 and 170 respectively. The degree of dissociation of silver nitrate is

- A. 60 %
- B. 83.5 %
- C. 46.7 %
- D. 60.23 %

**Answer: B**



[Watch Video Solution](#)

165. When some NaCl was dissolved in water, the freezing point depression was numerically equal to twice the molal depression constant. The relative lowering of vapour pressure of the solution is

A. 0.117

B. 0.036

C. 0.0585

D. none

**Answer: B**



[Watch Video Solution](#)

166.  $\Delta T_b$  or  $\Delta T_f$  for ethanoic acid in benzene will be (i) of normal value and molar mass will be (ii) of expected value.

Here (i) and (ii) refer to

A. (i) = half, (ii) = half

B. (i) = twice, (II) = half

C. (i) = half, (II) = twice

D. (i) = twice, (II) = twice

**Answer: C**

 [Watch Video Solution](#)

**167.** The b.p. of  $0.2 \text{ mol kg}^{-1}$  solution of  $x$  in  $H_2O$  is greater than equimolal solution for  $y$  in  $H_2O$ . Which one of the following statements is true ?

A.  $y$  is undergoing dissociation in  $H_2O$  while  $x$  undergoes no change

B.  $x$  is undergoing dissociation in  $H_2O$  while  $y$  undergoes no change

C. molecular mass of  $x$  is greater than molecular mass of  $y$

D. molecular mass of  $x$  is less than the molecular mass of  $y$

**Answer: B**

 [Watch Video Solution](#)

**168.** Acetic acid exists in benzene solution in the dimeric form. In an actual experiment the vant Hoff factor was found to be 0.52. Then the degree of association of acetic acid is

- A. 0.48
- B. 0.88
- C. 0.96
- D. 0.52

**Answer: C**

 [Watch Video Solution](#)

**169.** The mole fraction of a solute in a binary solution is 0.1. At 298 K, molarity of this solution is same as its molality. Density of this solution at

298 K is  $2.0 \text{ g cm}^{-3}$ . The ratio of molecular weights of the solute and the solvent ( $M_{\text{solute}} / M_{\text{solvent}}$ ) is

 [Watch Video Solution](#)

**170.** Identify the mixture that shows positive deviation from Raoult's law.

- A. carbon tetrachloride + methanol
- B. carbon disulphide + acetone
- C. benzene + toluene
- D. phenol + aniline

**Answer:**

 [Watch Video Solution](#)

**171.** If the freezing point of a 0.01 molal aqueous solution of a cobalt (III) chloride-ammonia complex (which behaves as a strong electrolyte) is -



$0.0558^\circ C$ , the number of chloride(s) in the coordination sphere of the complex is  $[K_f \text{ of water} = 1.86 K kg mol^{-1}]$

 [Watch Video Solution](#)

**172.** A compound  $H_2X$  with molar weight of 80 g is dissolved in a solvent having density of  $0.4 \text{ g ml}^{-1}$ . Assuming no change in volume upon dissolution, the molality of a 3.2 molar solution is

 [Watch Video Solution](#)

**173.** Benzene and naphthalene form an ideal solution at room temperature. For this process, the true statement(s) is/are

- A.  $\Delta G$  is positive
- B.  $\Delta S_{system}$  is positive
- C.  $\Delta S_{surroundings} = 0$
- D.  $\Delta H = 0$

**Answer:**



[Watch Video Solution](#)

**174.** 29.2 % (wiW) HCl stock solution has density of  $1.25\text{gmL}^{-1}$ . The molecular weight of HCl is  $36.5\text{gmol}^{-1}$ . The volume (mL) of stock solution required to prepare a 200 mL solution 0.4 M HCl is

- A. 5.0 mL
- B. 6.0 mL
- C. 8.0 mL
- D. 15.0 mL

**Answer:**



[Watch Video Solution](#)

**175.** In which case van't Hoff factor are equal ?

A.  $KCl$ , 50 % ionised

B.  $K_2SO_4$ , 40 % ionised

C.  $FeCl_3$ , 30 % ionised

D.  $SnCl_4$ , 20 % ionised

**Answer:**

 [Watch Video Solution](#)

**176.** The total vapour pressure of a binary solution is given by

$$P = (100x_A + 260x_B)mmHg$$

where  $x_A$  and  $x_B$  are the mole fraction of components A and B. This indicates that the :

A. vapour pressure of solution is less than the pure B component

B. vapour pressure of solution is more than that of pure A component

C. vapour pressure of pure A is 100 mm Hg and that of pure B is 260

mm Hg

D. the vapour pressure of pure A and B are 260 mm Hg and 100 mm Hg respectively

**Answer:**

 [Watch Video Solution](#)

177. Which has the equal elevation of boiling point ?

A. 0.1 M  $Na_2SO_4$

B. 0.1 M  $C_6H_{12}O_6$  (glucose)

C. 0.1 M  $MgCl_2$

D. 0.1 M  $Al(NO_3)_3$

**Answer:**

 [Watch Video Solution](#)

178. Which of the following is correct for an ideal solution ?

A. Raoult's law is obeyed for entire concentration range and temperatures

B.  $\Delta H_{mix} = 0$

C.  $\Delta V_{mix} = 0$

D.  $\Delta S_{mix} = 0$

**Answer:**



[Watch Video Solution](#)

179. Which of the following will form non-ideal solution ?

A.  $C_2H_5OH$  and water

B.  $HNO_3$  and water

C.  $CHCl_3$  and  $CH_3COCH_3$

D.  $C_6H_6$  and  $C_6H_5CH_3$

**Answer:**

 [Watch Video Solution](#)

**180.** Which of the following is/are correct for an ideal binary solution of two volatile liquids (eg. benzene & toluene) ?

- A. its vapour is always richer in the more volatile component  
(compared to the liquid)
- B. The liquid will gradually become richer in the less volatile component if such a mixture is boiled (distilled)
- C. The  $P_T$  (ie. the total pressure) above the solution will be the sum of the vapour pressures of the two pure components.
- D. The boiling point of the solution will be less than the boiling of the two components.

**Answer:**

 [Watch Video Solution](#)

**181.** Draw and state correct Geometry and hybridization for  $\text{XeF}_4$

 [Watch Video Solution](#)

**182.** In which of the following pairs of solutions will the values of the colligative properties be the same?

A.  $0.05M\text{K}_4[\text{Fe}(\text{CN})_6]$  and  $0.10M\text{FeSO}_4$

B.  $0.10M\text{K}_4[\text{Fe}(\text{CN})_6]$  and  $0.10M\text{FeSO}_4(\text{NH}_4)_2\text{SO}_4 \cdot 6\text{H}_2\text{O}$

C.  $0.20M\text{NaCl}$  and  $0.10M\text{BaCl}_2$

D.  $0.05M\text{FeSO}_4(\text{NH}_4)_2\text{SO}_4 \cdot 6\text{H}_2\text{O}$  and  $0.05M\text{KCl} \cdot \text{MgCl}_2 \cdot 6\text{H}_2\text{O}$

**Answer:**

 [Watch Video Solution](#)

**183.** Why will copper and  $\text{HCl(aq)}$  not evolve Hydrogen gas?

 [Watch Video Solution](#)

**184.** Match Column - I with Column - II



 [View Text Solution](#)

**185.** Match Column - I with Column - II



 [View Text Solution](#)

**186.** Match Column - I with Column - II







 [View Text Solution](#)

**187.** Match Column - I with Column - II



 [View Text Solution](#)

**188.** What is the maximum value of van't Hoff factor for  $AlCl_3$  ?

 [Watch Video Solution](#)

**189.** Phenol ( $C_6H_5OH$ ) is found to exist as polymer. If there is 100 % polymerisation, determine number of phenol molecules polymerised in aqueous solution if 9.4 g of phenol solution in 100 g water freezes at  $-0.93^\circ C$ .

$$K_f H_2O = 1.86^\circ C mol^{-1} kg$$



Watch Video Solution

190. The freezing point depression of a  $1.00 \times 10^{-3}$  m solution of  $K_x[Fe(CN)_6]$  is  $7.10 \times 10^{-3}$  K. Determine  $x$  given

$K_f = 1.86 K kg mol^{-1}$  for  $H_2O$ . Assume 100 % dissociation.

$K_f = 1.86 K kg mol^{-1}$  for  $H_2O$



Watch Video Solution

191. The concentration of pollutant in ppm (w/w). that has been measured at 450 mg per 150 kg of sample is :



Watch Video Solution

192. An aqueous solution of 0.01 M  $CH_3COOH$  has Van't Hoff factor 1.01.

If  $pH = -\log[H^+]$ , pH of 0.01 M  $CH_3COOH$  solution would be



Watch Video Solution

**193.** Insulin  $(C_2H_{10}O_5)_n$  is dissolved in a suitable solvent and the osmotic pressure  $\pi$  of the solution of various concentration ( $\text{in kg/m}^3$ ) is measured at  $20^\circ\text{C}$ . The slope of a plot of  $\pi$  against  $c$  is found to be  $8.134 \times 10^{-3}$  (SI units) The molecular weight of the insulin

 [Watch Video Solution](#)

**194.** Relative decrease in vapour pressure of an aqueous NaCl is 0.181. Number of moles of NaCl present in 180g of  $H_2O$  is (consider 100% dissociation of NaCl)

 [Watch Video Solution](#)

**195.** One would expect proton to have very large:

A. (a) charge

B. (b) ionization potential

C. (c) hydration energy

D. (d) radius

**Answer:**

 [Watch Video Solution](#)

**196.** Which of the following statements is true for all the alkali metals?

A. (a) Their nitrates decomposes on heating to give the corresponding nitrites and oxygen.

B. (b) Their chlorides are deliquescent and crystallise as hydrates.

C. (c) They react with water to form hydroxide and hydrogen.

D. (d) They readily react with halogens to form ionic halides.

**Answer:**

 [Watch Video Solution](#)

197. What is the vapour pressure of pure water at  $100^{\circ}C$  temperature?

 [Watch Video Solution](#)

198. A solution is prepared by dissolving 10 g of nonvolatile solute in 180 g of  $H_2O$ . If the relative lowering of vapour pressure is 0.005, find the mol. wt. of the solute.

 [Watch Video Solution](#)

199. A very small amount of a nonvolatile solute (that does not dissociate) is dissolved in  $56.8\text{ cm}^3$  of benzene (density  $0.889\text{ g cm}^{-3}$ ). At room temperature, vapour pressure of this solution is 98.88 mm Hg while that of benzene is 100 mm Hg. Find the molality of this solution. If the freezing temperature of this solution is 0.73 degree lower than that of benzene, what is the value of molar freezing point depression constant of benzene ?

 [Watch Video Solution](#)

 Watch Video Solution

**200.** A solution of .73 g of 'A' in 100 cc of water is found to be isotonic with a 3.42 % (wt./vol.) solution of sucrose ( $C_{12}H_{22}O_{11}$ ). Calculate molecular weight of A. ( $C_{12}H_{22}O_{11} = 342$ ). Assume no dissociation/association.



Watch Video Solution

**201.** A solute 'A' undergoes association in a medium as  $nA \rightleftharpoons A_n$

If the degree of association is ' $\alpha$ ', then Van't Hoff factor may be given as

:



Watch Video Solution

**202.** Ethylene glycol is used as antifreezing agent to prevent the freezing of water in car radiator in cold climate. Calculate the amount of ethylene

glycol to be added to 5 kg of water to prevent it from freezing at  $-6^{\circ}C$ .

Given  $K_f$  for water =  $1.85 \text{ kg mol}^{-1}$ .

 [Watch Video Solution](#)

**203.** What is the weight of 1 mole of a solute, 0.132 g of which in 29.7 g benzene gave a freezing point depression of  $0.108^{\circ}C$ ? ( $K_f$  for benzene = 5.12)

 [Watch Video Solution](#)

**204.** A solution containing 28 g phosphorus in 315 g  $CS_2$  (b. pt.  $46.3^{\circ}C$ ) boils at  $47.98^{\circ}C$ .  $K_b$  for  $CS_2$  is  $2.34 \text{ K mol}^{-1} \text{ kg}$ . Calculate m.wt. of phosphorus and deduce its molecular formula. Assume its complete association.

 [Watch Video Solution](#)

205. A binary liquid (AB) shows positive deviation from Raoult's law when

A.  $P_A > P_A^0 X_A^{liq}$  and  $P_B > P_B^0 X_B^{liq}$

B. Intermolecular forces :  $A - A, B - B > A - B$

C.  $\Delta V_{mix} > 0$

D.  $\Delta H_{mix} > 0$

Answer:



Watch Video Solution

206. Consider 0.1 M solutions of two solutes X and Y. The solute X behaves as univalent electrolyte, will the solute Y dimerises in solution.

Selected correct statement

(s) regarding these solutions:

A. The boiling point of solution of 'X' will be higher than that of 'Y'



- B. The osmotic pressure of solution of 'Y' will be higher than that of 'X'.
- C. The freezing point of solution of 'X' will be lower than that of 'Y'.
- D. The relative lowering of vapour pressure of both the solution will be the same

**Answer:**

 [Watch Video Solution](#)

**207.** Which of the following statements is (are) incorrect ?

- A. 0.1 M KCL solution will have the same osmotic pressure as 0.1 M glucose solution.
- B. 0.1 M KCl solution will have the same boiling point as 0.1 M urea solution.
- C. 0.1 m glucose and 0.1 m urea isotonic.

D. 0.1 m  $MgCl_2$  solution will have less relative lowering of vapour pressure than 0.1 m NaCl.

**Answer:**

 [Watch Video Solution](#)

**208.** Consider following solutions :

0.1 m  $C_6H_5NH_3^+ Cl^-$  : 0.1 m Glucose , 0.1 m  $Na_2C_2O_4 \cdot 10H_2O$

- A. the solution with highest boiling point is 0.1 m  $Na_2C_2O_4 \cdot 10H_2O$
- B. the solution with highest freezing point 0.1 m glucose
- C. 0.1 m  $C_6H_5NH_3Cl$  and 0.1 m NaCl will have the same osmotic pressure
- D. 0.1 m glucose solution will have the lowest osmotic pressure

**Answer:**

 [Watch Video Solution](#)

**209.** The azeotropic solutions of two miscible liquids

- A. can be separated by simple distillation
- B. may show positive or negative deviation from Raoult's Law
- C. are supersaturated solution
- D. behave like a single component and boil at a constant temperature

**Answer:**

 [Watch Video Solution](#)

**210.** For a nonvolatile solute which statement (s) is/are correct ?

- A. vap. pressure of solution = vap. pressure of pure solvent
- B. vap. pressure of solute is zero
- C. vap. pressure of solution = vap. pressure of solvent in solution

D. vap. pressure of solution  $>$  vap. pressure of pure solvent

**Answer:**

 [Watch Video Solution](#)

**211.** In the depression of freezing point experiment it is found that

- A. the vap. pressure of the solution is less than that of pure solvent
- B. the vap. pressure of solution is more than that of pure solvent
- C. only solute molecules solidify at freezing point
- D. only solvent molecules solidify at the freezing point

**Answer:**

 [Watch Video Solution](#)

**212.** Addition of non-volatile solute to a solvent always increases the colligative properties such as osmotic pressure.  $\Delta P$ .  $\Delta T_b$  and  $\Delta T_f$ . All these colligative properties are directly proportional to molality if solutions are dilute. The increases in colligative properties on addition of non-volatile solute is due to increases in number of solute particles.

For different aqueous solutions of 0.1 N NaCl, 0.1 N urea, 0.1 N  $Na_2SO_4$  and 0.1 N  $Na_3PO_4$  solution at  $27^\circ C$ , the correct statements are :

1. The order of osmotic pressure is  $NaCl = Na_2SO_4 = Na_3PO_4 = urea$
2.  $\pi = \frac{\Delta T_b}{k_b} \times RT$  for urea solution
3. Addition of salt on ice increases its melting point
4. Addition of salt on ice bring in melting of ice earlier

A. 2,3,4

B. 2,4

C. 1,2,3

D. 3,4

**Answer: B**

**213.** Addition of non-volatile solute to a solvent always increases the colligative properties such as osmotic pressure.  $\Delta P$ .  $\Delta T_b$  and  $\Delta T_f$ . All these colligative properties are directly proportional to molality if solutions are dilute. The increases in colligative properties on addition of non-volatile solute is due to increases in number of solute particles.

For different aqueous solutions of 0.1 N NaCl, 0.1 N urea, 0.1 N  $Na_2SO_4$  and 0.1 N  $Na_3PO_4$  solution at  $27^\circ C$ , the correct statements are :

1. The order of osmotic pressure is  $NaCl = Na_2SO_4 = Na_3PO_4 = urea$
2.  $\pi = \frac{\Delta T_b}{k_b} \times RT$  for urea solution
3. Addition of salt on ice increases its melting point
4. Addition of salt on ice bring in melting of ice earlier

A. 1,2,3

B. 1,2,3,5

C. 2,4,5

D. 1,4,5

**Answer: A**

 [Watch Video Solution](#)

**214.** An aqueous solution of 0.1 molal concentration of sucrose should have freezing point ( $k_f = 1.86 \text{ kmol}^{-1} \text{ kg}$ )

A.  $0.186^\circ \text{C}$

B.  $1.86^\circ \text{C}$

C.  $-1.86^\circ \text{C}$

D.  $-0.186^\circ \text{C}$

**Answer: D**

 [Watch Video Solution](#)

**215.** When 250 mg of eugenol is added to 100 g of camphor ( $k_f = 39.7 \text{ kmolality}^{-1}$ ), it lowered the freezing point by  $0.62^\circ \text{C}$ . The

molar mass of eugenol is :

A.  $1.6 \times 10^2 \text{ g/mol}$

B.  $1.6 \times 10^4 \text{ g/mol}$

C.  $1.6 \times 10^3 \text{ g/mol}$

D. 200 g/mol

**Answer: A**



[Watch Video Solution](#)

**216.** Match column - I with Column - II



[View Text Solution](#)

**217.** Match column - I with Column - II





 [View Text Solution](#)

218. Match column - I with Column - II



 [View Text Solution](#)

219. The vapour pressure of two pure liquid A and B are 5 and 10 torr respectively. Calculate the total pressure of the solution (in torr) obtained by mixing 2 moles of A and 3 moles of B.

 [Watch Video Solution](#)

220. The vapour pressure of a liquid solution containing A and B is 99 torr. Calculate mole % of B in vapour phase.

(Given :  $P_A^0 = 100$  torr ,  $P_B^0 = 80$  torr)

 [Watch Video Solution](#)

**221.** Calculate elevation in boiling point for 2 molal aqueous solution of glucose.

$$(Given: k_b(H_2O) = 0.5Kkg. mol^{-1})$$

 [Watch Video Solution](#)

**222.** A solution containing 500 g of a protein per litre is isotonic with a solution containing 3.42 g sucrose per litre. The molecular mass of protein is  $5 \times 10^x$  hence x is

 [Watch Video Solution](#)

**223.** An aqueous solution of urea has a freezing point of  $-0.515^\circ C$ . Predict the osmotic pressure (in atm) of the same solution at  $37^\circ C$ .

$$k_f = 1.86kkg(mol)^{-1}$$

 [Watch Video Solution](#)

**224.** Calculate the concentration of NaOH solution in g/mL which has the same normality as that of a solution of HCl of concentration 0.0365 g/mL.

 [Watch Video Solution](#)

**225.** 10 g of solute A and 20 g of solute B both are dissolved in 500 mL of water. The solution has the same osmotic pressure as 6.67 g of A and 30 g of B are dissolved in the same volume of water at the same temperature. What is the ratio of molar masses of A and B ?

 [Watch Video Solution](#)

**226.** Arrange the osmotic pressure in increasing order for the following (assume salts are 100 % dissociated).

I. 0.M urea

II. 0.1 M NaCl

III. 0.1MNa<sub>2</sub>SO<sub>4</sub>

IV. 0.1MNa<sub>3</sub>PO<sub>4</sub>

 [Watch Video Solution](#)

227. If equal volumes of two isotonic solutions are mixed then calculate the osmotic pressure of the net resultant solution obtained.

 [Watch Video Solution](#)

228. Dry air was successively passed through a solution of 5 g solute in 80 g water & then through pure water. The loss in weight of solution was 2.5 g & that of pure water was 0.04 g. What is the mol. wt. of solute ?

 [Watch Video Solution](#)

229. A beaker containing 20 g sugar in 100 g water and another containing 10 g sugar in 100 g water are placed under a bell-jar and allowed to stand until equilibrium is reached. How much water will be transferred from one beaker to another ?

 [Watch Video Solution](#)

 [Watch Video Solution](#)

**230.** A solution containing 0.5 g of naphthalene in 50 g  $CCl_4$  yield a boiling point elevation of 0.4 k, while a solution of 0.6 g of an unknown solute in the same mass of the solvent gives a boiling point elevation of 0.65 k. Find the molar mass of the unknown solute.

 [Watch Video Solution](#)

**231.** Calculate the freezing point of a solution of a non-volatile solute in a unknown solvent of molar mass 30 g/mol having mole fraction of solvent equal to 0.8. Given that latent heat of fusion of solid solvent = 2.7 kcal  $mol^{-1}k^{-1}$ , freezing point of solvent =  $27^\circ C$  and  $R = 2 \text{ cal } mol^{-1}k^{-1}$

 [Watch Video Solution](#)

**232.** Two liquids A and B form an ideal solution. At 300 K, the vapour pressure of a solution containing 1 mole of A and 3 moles of B is 550 mm of Hg. At the same temperature, if one mole of B is added to this solution, the vapour pressure of the solution increases by 10 mm of Hg. Determine the vapour pressure of A and B in their pure states.

 [Watch Video Solution](#)

**233.** Vapour pressure of  $C_6H_6$  and  $C_7H_8$  mixture at  $50^\circ C$  is given  $P$  (mm Hg) =  $180X_B + 90$ , where  $X_B$  is the mole fraction of  $C_6H_6$ . A solution is prepared by mixing 936 g benzene and 736 g toluene and if vapours over this solution are removed and condensed into liquid and again brought to the temperature  $50^\circ C$ , what would be mole fraction of  $C_6H_6$  in the vapour state. (At. wt. of C 12, H = 1)

 [Watch Video Solution](#)

**234.** At a constant temperature liquid 'A' has vapour pressure of 170 mm Hg and liquid 'B' has vapour pressure of 280 mm Hg. A solution of the two at the same temperature in which mole fraction of A is 0.7 has a total vapour pressure of 376 mm. Identify whether the solution process is Endothermic or Exothermic ?



[Watch Video Solution](#)

**235.** This question has statement I and II. Of the four choices given after the statements, choose the one that best describes the two statements.

Statement - I : The difference in the boiling points of equimolar solution of HCl and HF decreases as their molarity is decreased.

Statement - II : The extent of dissociation decreases steadily with increasing dilution.

A. Statement - I is true, Statement - II is true. Statement - II is a correct explanation of Statement - I

- B. Statement - I is true, Statement - II is true. Statement - II is not correct explanation of Statement - I
- C. Statement - I is true, Statement - II is false.
- D. Statement - I is false, Statement - II is true.

**Answer: C**

 [Watch Video Solution](#)

**236.** This question has statement I and II. Of the four choices given after the statements, choose the one that best describes the two statements.

Statement - I : When 'a' mL of a 0.1 molal urea solution is mixed with another 'b' mL of 0.1 molal glucose solution, the boiling point of the solution is no different from the boiling points of the samples prior to mixing but if a mL of molal urea is mixed with 'b' mL of 0.1 molal HF the boiling point is different from the separate samples

Statement -II : HF is an electrolyte (weak) whereas glucose is a non electrolyte.



- A. Statement - I is true, Statement - II is true. Statement - II is a correct explanation of Statement - I
- B. Statement - I is true, Statement - II is true. Statement - II is not correct explanation of Statement - I
- C. Statement - I is true, Statement - II is false.
- D. Statement - I is false, Statement - II is true.

**Answer: A**



**Watch Video Solution**

**237.** This question has statement I and II. Of the four choices given after the statements, choose the one that best describes the two statements.

Statement - I : The freezing point of water is depressed by the addition of glucose.

Statement -II : Entropy of solution is less than entropy of pure solvent.

- A. Statement - I is true, Statement - II is true. Statement - II is a correct explanation of Statement - I
- B. Statement - I is true, Statement - II is true. Statement - II is not correct explanation of Statement - I
- C. Statement - I is true, Statement - II is false.
- D. Statement - I is false, Statement - II is true.

**Answer: C**

 [Watch Video Solution](#)

**238.** This question has statement I and II. Of the four choices given after the statements, choose the one that best describes the two statements.

Statement - I : In binary ideal solution more volatile component boils at less temperature and less volatile component boils at high temperature than their normal boiling point of pure component.

Statement -II : For any composition, mixture boils at temperature in between the their normal boiling point temperature of pure component.

- A. Statement - I is true, Statement - II is true. Statement - II is a correct explanation of Statement - I
- B. Statement - I is true, Statement - II is true. Statement - II is not correct explanation of Statement - I
- C. Statement - I is true, Statement - II is false.
- D. Statement - I is false, Statement - II is true.

**Answer: D**



**Watch Video Solution**

**239.** This question has statement I and II. Of the four choices given after the statements, choose the one that best describes the two statements.

Statement - I : The boiling point of 0.1 M urea solution is less than that of 0.1 M KCl solution.

Statement -II : Elevation of boiling point is directly proportional to the number of moles of non-volatile solute particles present in the solution.

- A. Statement - I is true, Statement - II is true. Statement - II is a correct explanation of Statement - I
- B. Statement - I is true, Statement - II is true. Statement - II is not correct explanation of Statement - I
- C. Statement - I is true, Statement - II is false.
- D. Statement - I is false, Statement - II is true.

**Answer: A**



[Watch Video Solution](#)

**240.** This question has statement I and II. Of the four choices given after the statements, choose the one that best describes the two statements.

Statement - I : The observed molar mass of acetic acid in benzene is more than the normal molar mass of the acetic acid

Statement -II : Molecules of acetic acid dimerise in benzene due to hydrogen bonding.

- A. Statement - I is true, Statement - II is true. Statement - II is a correct explanation of Statement - I
- B. Statement - I is true, Statement - II is true. Statement - II is not correct explanation of Statement - I
- C. Statement - I is true, Statement - II is false.
- D. Statement - I is false, Statement - II is true.

**Answer: A**



[Watch Video Solution](#)

**241.** This question has statement I and II. Of the four choices given after the statements, choose the one that best describes the two statements.

Statement - I : Molarity and molality of 111 mg  $CaCl_2$  in 1 kg  $H_2O$  is nearly same.

Statement -II : In dilute solution molarity and molality are approximately equal.

- A. Statement - I is true, Statement - II is true. Statement - II is a correct explanation of Statement - I
- B. Statement - I is true, Statement - II is true. Statement - II is not correct explanation of Statement - I
- C. Statement - I is true, Statement - II is false.
- D. Statement - I is false, Statement - II is true.

**Answer: A**



[Watch Video Solution](#)

**242.** This question has statement I and II. Of the four choices given after the statements, choose the one that best describes the two statements.

Statement - I : At ice  $\rightleftharpoons$  water equilibrium, on increasing the pressure freezing point of water decreases.

Statement -II : Ice have low density so equilibrium shift in forward direction when pressure is increased.

- A. Statement - I is true, Statement - II is true. Statement - II is a correct explanation of Statement - I
- B. Statement - I is true, Statement - II is true. Statement - II is not correct explanation of Statement - I
- C. Statement - I is true, Statement - II is false.
- D. Statement - I is false, Statement - II is true.

**Answer: A**



**Watch Video Solution**

**243.** This question has statement I and II. Of the four choices given after the statements, choose the one that best describes the two statements.

Statement - I : An increase in surface area increases the the rate of evaporation.

Statement -II : Stronger the intermolecular attraction forces, faster is the rate of evaporation at a given temperature.

- A. Statement - I is true, Statement - II is true. Statement - II is a correct explanation of Statement - I
- B. Statement - I is true, Statement - II is true. Statement - II is not correct explanation of Statement - I
- C. Statement - I is true, Statement - II is false.
- D. Statement - I is false, Statement - II is true.

**Answer: B**



[Watch Video Solution](#)

**244.** This question has statement I and II. Of the four choices given after the statements, choose the one that best describes the two statements.

Statement - I : An ideal solution obeys Raoult's law.

Statement -II : In an ideal solution, solute - solute as well as solvent - solvent, interactions are similar to solute solvent interactions.



- A. Statement - I is true, Statement - II is true. Statement - II is a correct explanation of Statement - I
- B. Statement - I is true, Statement - II is true. Statement - II is not correct explanation of Statement - I
- C. Statement - I is true, Statement - II is false.
- D. Statement - I is false, Statement - II is true.

**Answer: A**



[Watch Video Solution](#)

**245.** This question has statement I and II. Of the four choices given after the statements, choose the one that best describes the two statements.

Statement - I : If a liquid solute more volatile than the solvent is added to the solvent, the vapour pressure of the solution is greater than vapour pressure of pure solvent.

Statement -II : Vapour pressure of solution is equal to vapour pressure of solvent.

- A. Statement - I is true, Statement - II is true. Statement - II is a correct explanation of Statement - I
- B. Statement - I is true, Statement - II is true. Statement - II is not correct explanation of Statement - I
- C. Statement - I is true, Statement - II is false.
- D. Statement - I is false, Statement - II is true.

**Answer: C**



**Watch Video Solution**

**246.** This question has statement I and II. Of the four choices given after the statements, choose the one that best describes the two statements.

Statement - I :  $\Delta V_m^i x$  and  $\Delta S_m^i x$  for an ideal solution is zero.

Statement -II : A... B interaction in an ideal solution are same as between A...A and B...B.

- A. Statement - I is true, Statement - II is true. Statement - II is a correct explanation of Statement - I
- B. Statement - I is true, Statement - II is true. Statement - II is not correct explanation of Statement - I
- C. Statement - I is true, Statement - II is false.
- D. Statement - I is false, Statement - II is true.

**Answer: D**

 [Watch Video Solution](#)

**247.** This question has statement I and II. Of the four choices given after the statements, choose the one that best describes the two statements.

Statement - I : Elevation in boiling point will be high if the molal elevation constant of the liquid is high.

Statement - II : Elevation in boiling point is a colligative property.

- A. Statement - I is true, Statement - II is true. Statement - II is a correct explanation of Statement - I
- B. Statement - I is true, Statement - II is true. Statement - II is not correct explanation of Statement - I
- C. Statement - I is true, Statement - II is false.
- D. Statement - I is false, Statement - II is true.

**Answer: B**

 [Watch Video Solution](#)

**248.** This question has statement I and II. Of the four choices given after the statements, choose the one that best describes the two statements.

Statement - I : Addition of ethylene glycol to water lowers the freezing point of water, therefore, used as antifreeze substance.

Statement -II : Ethylene glycol is soluble in water.

- A. Statement - I is true, Statement - II is true. Statement - II is a correct explanation of Statement - I
- B. Statement - I is true, Statement - II is true. Statement - II is not correct explanation of Statement - I
- C. Statement - I is true, Statement - II is false.
- D. Statement - I is false, Statement - II is true.

**Answer: B**

 [Watch Video Solution](#)

**249.** This question has statement I and II. Of the four choices given after the statements, choose the one that best describes the two statements.

Statement - I : Osmotic pressure is a colligative property.

Statement -II : Osmotic pressure developed in a column due to osmosis.

- A. Statement - I is true, Statement - II is true. Statement - II is a correct explanation of Statement - I

- B. Statement - I is true, Statement - II is true. Statement - II is not correct explanation of Statement - I
- C. Statement - I is true, Statement - II is false.
- D. Statement - I is false, Statement - II is true.

**Answer: B**

 [Watch Video Solution](#)

**250.** This question has statement I and II. Of the four choices given after the statements, choose the one that best describes the two statements.

Statement - I : Osmosis involves movement of solvent molecules from its higher concentration to its lower concentration.

Statement -II : Solutions having the same osmotic pressure are called isotonic solutions.

- A. Statement - I is true, Statement - II is true. Statement - II is a correct explanation of Statement - I

- B. Statement - I is true, Statement - II is true. Statement - II is not correct explanation of Statement - I
- C. Statement - I is true, Statement - II is false.
- D. Statement - I is false, Statement - II is true.

**Answer: D**

 [Watch Video Solution](#)

## Question Bank

1.  $pH$  of a 0.1 M monobasic acid is measured to be 2. Its osmotic pressure at a given temperature  $TK$  is

- A. 0.1 RT
- B. 0.11 RT
- C. 1.1 RT

D. 0.01 RT

**Answer:**



**Watch Video Solution**