



PHYSICS

BOOKS - MTG PHYSICS (BENGALI ENGLISH)

QUESTION PAPER 2014

Physics Category I

1. A drop of some liquid of volume 0.04cm^3 is placed on the surface of a glass slide. Then

another glass slide is placed on it in such a way that the liquid forms a thin layer of area 20cm^2 between the surfaces of the two slides. To separate the slides a force of 16×10^5 dyne has to be applied normal to the surfaces. The surface tension of the liquid is (in dyne-cm^{-1})

A. 60

B. 0

C. 80

D. 90

Answer:



Watch Video Solution

2. A very small circular loop of radius a is initially (at $t = 0$) coplanar and concentric with a much larger fixed circular loop of radius b . A constant current I flows in the larger loop. The smaller loop is rotated with a constant angular speed ω about the common diameter. The emf induced in the smaller loop as a function of time t is

$$\text{A. } \pi \frac{a^2 \mu_0 I}{2b} \omega \cos(\omega t)$$

B. $\frac{\pi a^2 \mu_0 I}{2b} \omega \sin(\omega^2 t^2)$

C. $\frac{\pi a^2 \mu_0 I}{2b} \omega \sin(\omega t)$

D. $\frac{\pi a^2 \mu_0 I}{2b} \omega \sin^2(\omega t)$

Answer:



Watch Video Solution

3. An electron in a circular orbit of radius .05 nm performs 10^{16} revolutions per second. The magnetic moment due to this rotation of electron is (in Am^2)

A. 2.16×10^{-23}

B. 3.21×10^{-22}

C. 3.21×10^{-24}

D. 1.26×10^{-23}

Answer:



Watch Video Solution

4. Consider three vectors

$$\vec{A} = \hat{i} + \hat{j} - 2\hat{k}, \vec{B} = \hat{i} - \hat{j} + \hat{k} \quad \text{and}$$

$\vec{C} = 2\hat{i} - 3\hat{j} + 4\hat{k}$.A vector X of the form

$\alpha\vec{A} + \beta\vec{B}$ (α and β are numbers) is perpendicular to \vec{C} . The ratio of α and β is

A. 1:1

B. 2:1

C. -1:1

D. 3:1

Answer:



Watch Video Solution

5. A galvanometer having internal resistance 10Ω requires 0.01 A for a full scale deflection. To convert this galvanometer to a voltmeter of full-scale deflection at 120 V , we need to connect a resistance of

- A. 11990Ω in series
- B. 11990Ω in parallel
- C. 12010Ω in series
- D. 12010Ω in parallel

Answer:



Watch Video Solution

6. Three capacitors, $3\mu F$, $6\mu F$ and $6\mu F$ are connected in series to a source of 120 V. The potential difference, in volts, across the $3\mu F$ capacitor will be

A. 24

B. 30

C. 40

D. 60

Answer:



Watch Video Solution

7. An infinite sheet carrying a uniform surface charge density σ lies on the xy -plane. The work done to carry a charge q from the point $\vec{A} = a(\hat{i} + 2\hat{j} + 3\hat{k})$ to the point $\vec{B} = \alpha(\hat{i} - 2\hat{j} + 6\hat{k})$ (where a is a constant with the dimension of length and ϵ_0 is the permittivity of free space) is

A. $\frac{3\sigma a q}{2\epsilon_0}$

B. $\frac{2\sigma a q}{\epsilon_0}$

C. $\frac{5\sigma a q}{2\epsilon_0}$

D. $\frac{3\sigma a q}{\epsilon_0}$

Answer:



[Watch Video Solution](#)

8. The intensity of magnetization of a bar magnet is $5.0 \times 10^4 \text{ Am}^{-1}$. The magnetic length and the area of cross section of the

magnet are 12cm and 1cm^2 respectively. The magnitude of magnetic moment of this bar magnet is (in Si unit)

A. 0.6

B. 1.3

C. 1.24

D. 2.4

Answer:



Watch Video Solution

9. A luminous object is separated from a screen by distance d . A convex lens is placed between the object and the screen such that it forms a distinct image on the screen. The maximum possible focal length of this convex lens is

A. $4d$

B. $2d$

C. $d/2$


D. $d/4$

Answer:



Watch Video Solution

10. The output of the logic circuit given below is



A. $\bar{A} + B$

B. \bar{A}

C. $\overline{(\bar{A} + B)} \cdot \bar{A}$

$$D. \overline{(\overline{A} + B)}. A$$

Answer:



Watch Video Solution

11. One mole of an ideal monoatomic gas is heated at a constant pressure from $0^\circ C$ to $100^\circ C$. Then the change in the internal energy of the gas is (Given $R = 8.32 J mol^{-1} k^{-1}$)

A. $0.83 \times 10^3 J$

B. $4.6 \times 10^3 J$

C. $2.98 \times 10^3 J$

D. $1.25 \times 10^3 J$

Answer:



Watch Video Solution

12. A whistle whose air column is open at both ends has a fundamental frequency of 5100 Hz. If the speed of sound in air is $340m s^{-1}$, the length of the whistle, in cm, is

A. $5/3$

B. $10/3$

C. 5

D. $20/3$

Answer:



Watch Video Solution

13. A particle moves with constant acceleration along a straight line starting from rest. The percentage increase in its displacement

during the 4th second compared to that in the 3 second is

A. 0.33

B. 0.4

C. 0.66

D. 0.7

Answer:



Watch Video Solution

14. An artificial satellite moves in a circular orbit around the earth. Total energy of the satellite is given by E . The potential energy of the satellite is

A. $-2E$

B. $2E$

C. $2E/3$

D. $-2E/3$

Answer:



Watch Video Solution

15. In which of the following phenomena, the heat waves travel along straight lines with the speed of light?

- A. conduction
- B. forced convection
- C. natural convection
- D. thermal radiation

Answer:





16. A proton of mass m and charge q is moving in a plane with kinetic energy E . If there exists a uniform magnetic field B , perpendicular to the plane of the motion, the proton will move in a circular path of radius

A. $\frac{2Em}{qB}$

B. $\frac{\sqrt{2Em}}{qB}$

C. $\frac{\sqrt{Em}}{2qB}$

D. $\sqrt{\frac{2Eq}{mB}}$

Answer:



Watch Video Solution

17. The ionization energy of hydrogen is 13.6 eV. The energy of the photon released when an electron jumps from the first excited state ($n = 2$) to the ground state of a hydrogen atom is

A. 3.4 eV

B. 4.53 eV

C. 10.2 eV

D. 13.6 eV

Answer:



Watch Video Solution

18. When a particle executing SHM oscillates with a frequency ν , then the kinetic energy of the particle

- A. changes periodically with a frequency of v
- B. changes periodically with a frequency of $2v$
- C. changes periodically with a frequency of $v/2$
- D. remains constant

Answer:



Watch Video Solution

19. A parallel plate capacitor is charged and then disconnected from the charging battery. If the plates are now moved farther apart by pulling at them by means of insulating handles, then

A. the energy stored in the capacitor decreases

B. the capacitance of the capacitor increases

C. the charge on the capacitor decreases

D. the voltage across the capacitor
increases

Answer:



Watch Video Solution

20. If the bandgap between valence band and conduction band in a material is 5.0 eV, then the material is

A. semiconductor

B. good conductor

C. superconductor

D. insulator

Answer:



Watch Video Solution

21. Two coherent monochromatic beams of intensities I and $4I$ respectively are superposed. The maximum and minimum intensities in the resulting pattern are

A. $5I$ and $3I$

B. $9I$ and $3I$

C. $4I$ and I

D. $9I$ and I

Answer:



Watch Video Solution

22. A uniform solid spherical ball is rolling down a smooth inclined plane from a height h . The velocity attained by the ball when it

reaches the bottom of the inclined plane is v . If the ball is now thrown vertically upwards with the same velocity v , the maximum height to which the ball will rise is

A. $5h / 8$

B. $3h / 5$

C. $5h / 7$

D. $7h / 9$

Answer:



Watch Video Solution

23. A cricket ball thrown across a field is at heights h_1 and h_2 from the point of projection at times t_1 and t_2 respectively after the throw. The ball is caught by a fielder at the same height as that of projection. The time of flight of the ball in this journey is

A. $\frac{h_1 t_2^2 - h_2 t_1^2}{h_1 t_2 - h_2 t_1}$

B. $\frac{h_1 t_1^2 + h_2 t_2^2}{h_1 t_2 - h_2 t_1}$

C. $\frac{h_1 t_2^2 + h_2 t_1^2}{h_1 t_2 - h_2 t_1}$

D. $\frac{h_1 t_1^2 - h_2 t_2^2}{h_1 t_2 - h_2 t_1}$

Answer:



Watch Video Solution

24. A small metal sphere of radius a is falling with a velocity v through a vertical column of a viscous liquid. If the coefficient of viscosity of the liquid is η , then the sphere encounters an opposing force of

A. $6\pi\eta a^2 v$

B. $\frac{6\eta v}{\pi a}$

C. $6\pi\eta av$

D. $\frac{\pi\eta v}{6a^3}$

Answer:



Watch Video Solution

25. In which of the following pairs, the two physical quantities have different dimensions?

A. Planck's constant and angular momentum

B. Impulse and linear momentum

C. Moment of inertia and moment of a
force

D. Energy and torque

Answer:



Watch Video Solution

26. If n denotes a positive integer, h the Planck's constant, q the charge and B the

magnetic field, then the quantity $\left(\frac{nh}{2\pi qB}\right)$ has the dimension of

A. area

B. length

C. speed

D. acceleration

Answer:



Watch Video Solution

27. In a transistor output characteristics commonly used in common emitter configuration, the base current I_B the collector current I_C and the collector-emitter voltage V_{CE} have values of the following orders of magnitude in the active region

A. I_B and I_C both are in μA , and V_{CE} in

Volts

B. I_B is in μA and I_C is in mA and V_{CE} in

Volts

C. I_B is in mA and I_C is in μA and V_{CE} in mV

D. I_B is in mA and I_C is in mA and V_{CE} in mV


Answer:



Watch Video Solution

28. In the circuit shown assume the diode to be ideal. When V_i increases from 2 V to 6 V,

the change in the current is (in mA)



A. zero

B. 20

C. $80/3$

D. 40

Answer:



Watch Video Solution

29. A scientist proposes a new temperature scale in which the ice point is 25 X(X is the new unit of temperature) and the steam point is 305 X. The specific heat capacity of water in this new scale is (in $Jkg^{-1}X^{-1}$)

A. 4.2×10^3

B. 3.0×10^3

C. 1.2×10^3

D. 1.5×10^3

Answer:






Watch Video Solution

30. One mole of a van der Waals' gas obeying the equation

$$\left(P + \frac{a}{V^2}\right)(V - b) = RT$$

undergoes the quasi-static cyclic process which is shown in the P-V diagram. The net

heat absorbed by the gas in this process is



A. $\frac{1}{2}(P_1 - P_2)(V_1 - V_2)$

B. $\frac{1}{2}(P_1 + P_2)(V_1 - V_2)$

C. $\frac{1}{2} \left(P_1 + \frac{a}{V_1^2} - P_2 - \frac{a}{V_2^2} \right) (V_1 - V_2)$

D. $\frac{1}{2} \left(P_1 + \frac{a}{V_1^2} + P_2 - \frac{a}{V_2^2} \right) (V_1 - V_2)$

Answer:



Watch Video Solution

31. The displacement of a particle in a periodic motion is given by $y = 4 \cos^2\left(\frac{t}{2}\right) \sin(1000t)$.

This displacement may be considered as the result of superposition of n no. of independent harmonic oscillations. Here n is

A. 1

B. 2

C. 3

D. 4

Answer:



Watch Video Solution

32. Consider two concentric spherical metal shells of radii r_1 and r_2 ($r_2 > r_1$). If the outer shell has a charge q and the inner one is grounded, the charge on the inner shell is (A) q (B) zero

A. $\frac{-r_2}{r_1}q$

B. zero

C. $\frac{-r_1}{r_2}q$

D. $-q$

Answer:



Watch Video Solution

33. Four cells, each of emf E and internal resistance r , are connected in series across an external resistance R . By mistake one of the

cells is connected in reverse. Then the current in the external circuit is

A. $\frac{2E}{4r + R}$

B. $\frac{3E}{4r + R}$

C. $\frac{3E}{3r + R}$

D. $\frac{2E}{3r + R}$

Answer:



Watch Video Solution

34. Consider a blackbody radiation in a cubical box at absolute temperature T . If the length of each side of the box is doubled and the temperature of the walls of the box and that of the radiation is halved, then the total energy

A. halves

B. doubles

C. quadruples

D. remains the same

Answer:



Watch Video Solution

35. A wooden block is floating on water kept in a beaker. 40% of the block is above the water surface. Now the beaker is kept inside a lift that starts going upward with acceleration equal to $g/2$. The block will then

A. sink

B. float with 10% above the water surface

C. float with 40% above the water surface

D. float with 70% above the water surface

Answer:



Watch Video Solution

36. To determine the coefficient of friction between a rough surface and a block, the surface is kept inclined at 45° and the block is released from rest. The block takes a time t in moving a distance d . The rough surface is then

replaced by a smooth surface and the same experiment is repeated. The block now takes a time $t/2$ in moving down the same distance d .

The coefficient of friction is

A. $\frac{3}{4}$

B. $\frac{5}{4}$

C. $\frac{1}{2}$

D. $1/\sqrt{2}$

Answer:



Watch Video Solution

37. A smooth massless string passes over a smooth fixed pulley. Two masses m_1 and m_2 ($m_1 > m_2$) are tied at the two ends of the string. The masses are allowed to move under gravity starting from rest. The total external force acting on the two masses is

A. $(m_1 + m_2)g$

B. $\frac{(m_1 - m_2)^2}{m_1 + m_2}g$

C. $(m_1 + m_2)g$

$$D. \frac{(m_1 + m_2)^2}{m_1 - m_2} g$$

Answer:



Watch Video Solution

38. Same quantity of ice is filled in each of the two metal containers P and Q having the same size, shape and wall thickness but made of different materials. The containers are kept in identical surroundings. The ice in P melts completely in time t_1 whereas that in Q takes a

time t_2 . The ratio of thermal conductivities of the materials of P and Q is

A. $t_2 : t_1$

B. $t_1 : t_2$

C. $t_1^2 : t_2^2$

D. $t_2^2 : t_1^2$

Answer:



Watch Video Solution

39. A car is moving with a speed of 72km-hour^{-1} towards a roadside source that emits sound at a frequency of 850 Hz. The car driver listens to the sound while approaching the source and again while moving away from the source after crossing it. If the velocity of sound is 340ms^{-1} , the difference of the two frequencies, the driver hears is

A. 50 Hz

B. 85 Hz

C. 100 Hz

D. 150Hz

Answer:



Watch Video Solution

40. For the radioactive nuclei that undergo either α or β decay, which one of the following cannot occur?

A. isobar of original nucleus is produced

B. isotope of the original nucleus is produced

C. nuclei with higher atomic number than that of the original nucleus is produced

D. nuclei with lower atomic number than that of the original nucleus is produced

Answer:



Watch Video Solution

41. A particle is moving uniformly in a circular path of radius r . When it moves through an angular displacement θ , then the magnitude of the corresponding linear displacement will be

A. $2r \cos\left(\frac{\theta}{2}\right)$

B. $2r \cot\left(\frac{\theta}{2}\right)$

C. $2r \tan\left(\frac{\theta}{2}\right)$

D. $2r \sin\left(\frac{\theta}{2}\right)$

Answer:



Watch Video Solution

42. The intermediate image formed by the objective of a compound microscope is

- A. real, inverted and magnified
- B. real, erect and magnified
- C. virtual, erect and magnified
- D. virtual, inverted and magnified

Answer:



43. A uniform rod is suspended horizontally from its mid-point. A piece of metal whose weight is W is suspended at a distance l from the mid-point. Another weight W_1 is suspended on the other side at a distance l_1 from the mid-point to bring the rod to a horizontal position. When W is completely immersed in water, W_1 needs to be kept at a distance l_2 from the mid-point to get the rod

back into horizontal position. The specific gravity of the metal piece is

A. $\frac{w}{w_1}$

B. $\frac{wl_1}{wl - w_1l_2}$

C. $\frac{l_1}{l_1 - l_2}$

D. $\frac{l_1}{l_2}$

Answer:



Watch Video Solution

44. The energy of gamma ray photon is E_Y and that of an X-ray photon is E_X . If the visible light photon has an energy of E_V , then we can say that

A. $E_X > E_Y > E_V$

B. $E_Y > E_V > E_X$

C. $E_Y > E_X > E_V$

D. $E_X > E_V > E_Y$

Answer:



Watch Video Solution

45. A metal rod is fixed rigidly at two ends so as to prevent its thermal expansion. If L , α and Y respectively denote the length of the rod, coefficient of linear thermal expansion and Young's modulus of its material, then for an increase in temperature of the rod by ΔT , the longitudinal stress developed in the rod is

A. inversely proportional to α

B. inversely proportional to Y

C. directly proportional to $\frac{\Delta T}{Y}$

D. independent of L

Answer:



Watch Video Solution

Physics Category Ii

1. To determine the composition of a bimetallic alloy, a sample is first weighed in air and then in water. These weights are found to be w_1

and w_2 respectively. If the densities of the two constituent metals are ρ_1 and ρ_2 respectively, then the weight of the first metal in the sample is (where ρ_w is the density of water)

A. $\frac{\rho_1}{\rho_w(\rho_2 - \rho_1)} [w_1(\rho_2 - \rho_w) - w_2\rho_2]$

B. $\frac{\rho_1}{\rho_w(\rho_2 + \rho_1)} [w_1(\rho_2 - \rho_w) + w_2\rho_2]$

C. $\frac{\rho_1}{\rho_w(\rho_2 - \rho_1)} [w_1(\rho_2 + \rho_w) - w_2\rho_2]$

D. $\frac{\rho_1}{\rho_w(\rho_2 - \rho_1)} [w_1(\rho_1 - \rho_w) - w_2\rho_1]$

Answer:



Watch Video Solution

2. The de Broglie wavelength of an electron is the same as that of a 50 keV X-ray photon. The ratio of the energy of the photon to the kinetic energy of the electron is the energy equivalent of electron mass is 0.5 MeV)

A. 1 : 50

B. 1 : 20

C. 20 : 1

D. 50 : 1

Answer:



Watch Video Solution

3. Sound waves are passing through two routes- one in straight path and the other along a semicircular path of radius r , and are again combined into one pipe and superposed as shown in the figure. If the velocity of sound waves in the pipe is v , then frequencies of resultant waves of maximum amplitude will be

integral multiples of



A. $\frac{v}{r(\pi - 2)}$

B. $\frac{v}{r(\pi - 1)}$

C. $\frac{2v}{r(\pi - 1)}$

D. $\frac{v}{r(\pi + 1)}$

Answer:



Watch Video Solution

4. A solid uniform sphere resting on a rough horizontal plane is given a horizontal Impulse directed through its center so that it starts sliding with an initial velocity v_0 . When it finally starts rolling without slipping the speed of its center is

A. $\frac{2}{7}v_0$

B. $\frac{3}{7}v_0$

C. $\frac{5}{7}v_0$

D. $\frac{6}{7}v_0$


Answer:



Watch Video Solution

5. A glass slab consists of thin uniform layers of progressively decreasing refractive indices n (see figure) such that the n of any layer is $n_0 - m\Delta n$. Here n_0 and Δn denote the n of 0^{th} layer and the difference in n between any two consecutive layers, respectively. The integer $m = 0,1,2,3\dots$ denotes the numbers of the successive layers. A ray of light from the

0^{th} layer enters the 1^{st} layer at an angle of incidence of 30° . After undergoing the m^{th} refraction, the ray emerges parallel to the interface. If $\mu = 1.5$ and $\Delta\mu = 0.015$, the value of m is



- A. 20
- B. 30

C. 40

D. 50


Answer:



Watch Video Solution

6. Three identical square plates rotate about the axes shown in the figure in such a way their kinetic energies are equal. Each of the rotation axes passes through the centre square. Then the ratio of angular speeds

$\omega_1 : \omega_2 : \omega_3$ is



A. $1 : 1 : 1$

B. $\sqrt{2} : \sqrt{2} : 1$

C. $1 : \sqrt{2} : 1$


D. $1 : 2 : \sqrt{2}$

Answer:



Watch Video Solution

7. A long conducting wire carrying a current I is bent at 120° (see figure). The magnetic field B at a point P on the right bisector of bending angle at a distance d from the bend is (μ_0 is the permeability of free space)



A. $\frac{3\mu_0 I}{2\pi d}$

B. $\frac{\mu_0 I}{2\pi d}$

C. $\frac{\mu_0 I}{\sqrt{3}\pi d}$

D. $\frac{\sqrt{3}\mu_0 I}{2\pi d}$


Answer:



Watch Video Solution

8. A circuit consists of three batteries of emf $E_1 = 1V$, $E_2 = 2V$ and $E_3 = 3V$ and internal resistances 1Ω , 2Ω and 1Ω respectively which are connected in parallel as shown in the figure. The potential difference

between points P and Q is



A. $1.0V$

B. $2.0V$

C. $2.2V$

D. $3.0V$

Answer:



9. A 10 Watt electric heater is used to heat a container filled with 0.5 kg of water. It is found that the temperature of water and the container rises by 3° K in 15 minutes. The container is then emptied, dried and filled with 2 kg of oil. The same heater now raises the temperature of container-oil system by 2° K in 20 minutes. Assuming that there is no heat loss in the process and the specific heat of

water as $4200 \text{ J kg}^{-1} \text{ K}^{-1}$, the specific heat of oil in the same unit is equal to

A. 1.50×10^3

B. 2.55×10^3

C. 3.00×10^3

D. 5.10×10^3

Answer:



Watch Video Solution

10. An object is placed 30 cm away from a convex lens of focal length 10 cm and a sharp image is formed on a screen. Now a concave lens is placed in contact with the convex lens. The screen now has to be moved by 45 cm to get a sharp image again. The magnitude of focal length of the concave lens is (in cm)

A. 72

B. 60

C. 36

D. 20

Answer:




Watch Video Solution

Physics Category Iii

1. Half of the space between the plates of a parallel-plate capacitor is filled with a dielectric material of dielectric constant K . The remaining half contains air as shown in the

figure. The capacitor is now given a charge Q .

Then



A. electric field in the dielectric filled region

is higher than that in the air-filled region

B. on the two halves of the bottom plate

the charge densities are unequal

C. charge on the half of the top plate

above the air-filled part is $\frac{Q}{K + 1}$

D. capacitance of the capacitor shown

above is $(1 + K) \frac{C_0}{2}$, where is the

capacitance of the same capacitor with

the dielectric removed.

Answer:




Watch Video Solution

2. A heating element of resistance r is fitted inside an adiabatic cylinder which carries a frictionless piston of mass m and cross-section

A as shown in diagram. The cylinder contains one mole of an ideal diatomic gas. The current flows through the element such that the temperature rises with time t as $\Delta T = \alpha t + \frac{1}{2}\beta t^2$ (α and β are constants), while pressure remains constant. The atmospheric pressure above the piston is P_0 .

Then



the rate of increase in internal energy is

$$\frac{5}{2}R(\alpha + \beta t)$$

the current flowing in the element is

$$\sqrt{\frac{5}{2r}}(\alpha + \beta t)$$

the piston moves upwards with constant

acceleration


the piston moves upwards with constant speed



Watch Video Solution

3. A stream of electrons and protons are directed towards a narrow slit in a screen (see figure). The intervening region has a uniform electric field (vertically downwards) and a uniform magnetic field \vec{B} (out of the plane of

the figure) as shown. Then



A. electrons and protons with speed $\frac{|\vec{E}|}{|\vec{B}|}$

will pass through the slit.

B. protons with speed $\frac{|\vec{E}|}{|\vec{B}|}$ will pass

through the slit, electrons of the same speed will not

C. neither electrons nor protons will go through the slit irrespective of their speed.

D. electrons will always be deflected upwards irrespective of their speed.

Answer:



Watch Video Solution