

BIOLOGY

BOOKS - TRUEMAN BIOLOGY

Plant Kingdom

Multiple Choice Question

1. A full mature plant body, undifferentiated into root, stem and leaf is called.

A. thallus

B. coenocytic

- C. hyphae
- D. callus



Watch Video Solution

- 2. Algae are included in
 - A. Thallophytes
 - B. spermatophytes
 - C. Embryophytes
 - D. Tracheophytes

Answer: A



- 3. Algae are characterised by
 - A. thalloid plant body
 - B. ph/ycobilins
 - C. unicellular sex organs
 - D. mechanical tissue.



Watch Video Solution

4. Algae resemble fungi in the presence of similar

B. cell wall
C. similar reserve food
D. similar mode of nutrition
Answer: A
Watch Video Solution
5. Green algae are ancestors of angiosperms/land plants
because
A. both have vascular bundles
B. both have starch as reserve food

A. reproductive structure

- C. both have jacketed sex organs
- D. all of the above

Answer: B



- **6.** Classification of Algae into 11 classes was made by Fritsch on the basis of flagellation, pigmentation and type of reserve food. Out of this, the main criteria used in algal classification grouping of algae is
 - A. chemical composition of cell wall
 - B. type of pigmentation
 - C. nature of food storage

D. shape and colony formation.

Answer: B



Watch Video Solution

7. Study of Algae is called

A. phycology

B. phytology

C. mycology

D. phenology

Answer: A



8. Choose the correct statement.

A. Algae show embryo stage

B. Algae show only haplontic life cycle

C. Algae are plants as they possess cell wall, chl a and starch food.

D. all of the above

Answer: C



- **9.** All algae possess
 - A. chl a and b
 - B. chl a, carotenes and phycobilins
 - C. chl b and carotenes
 - D. chl a and carotene

Answer: D



Watch Video Solution

10. Largest a cellular, green, marine algae ,popularly called umbrella plant is

A. Ulva B. Acetabularia C. Spirogyra D. Volvox **Answer: B Watch Video Solution** 11. Kelps are A. fresh water algae

B. marine green algae

C. large marine parenchymatous brown algae

D. large marine parenchymatous and algae.

Answer: C



Watch Video Solution

12. Red eye spot (stigma) is meant for

A. movement

B. vision

C. photoreception

D. photosynthesis

Answer: C



13. Red snow is caused by

A. stigma

B. chloroplast

C. hypnospore

D. aplanospore

Answer: C



Watch Video Solution

14. The diploid stage is represented by one celled zygote only in

B. Spirogyra C. both 1 and 2 D. Funaria **Answer: C Watch Video Solution 15.** Meiosis in spirogyra/Chlamydomonas / Ulothrix occurs A. as the zygospore germinates B. during conjugation

A. Chlamydomonas

- C. during palmella formation
- D. during formation of gametes



Watch Video Solution

- **16.** The product of conjugation of Spirogyra is called
 - A. zoospore
 - B. zygospore
 - C. endospore
 - D. akinete

Answer: B

17. Asexual spores with flagella in algae are called

- A. zoospore
- B. zygospores
- C. aplanospores
- D. hypnospores

Answer: A



18. In spriogyra lateral conjugation takes place in the cells of

- A. two adjacent cells of same filament
- B. two cells of different filaments
- C. among three filaments
- D. within one cell of a filament

Answer: B



Watch Video Solution

19. What is the shape of chloroplast in Spirogyra?

A. spiral, bank like B. cup like C. stellate like D. U-like **Answer: A Watch Video Solution** 20. Gametes of Spirogyra are A. biflagellated B. uniflagellated C. non-flagellated

D. diploid

Answer: C



- 21. Choose the correct statement in Spirogyra.
 - A. Filaments showing scalariform conjugation are always dioecious
 - B. Filaments showing lateral conjugation are always monoecious
 - C. Filaments showing lateral conjugation may be monoecious

D. Asexual reproduction occurs by zosspores.

Answer: B



Watch Video Solution

22. Agar-agar is obtained from

A. red algae

B. green algae

C. kelps

D. yellow algae

Answer: A



23. Red algae show maximum photosynthesis in blue green light. Their main pigment to trap this light is

- A. chlorophyll
- B. phycoerythrin
- C. chl a and xanthophyll
- D. carotenes.

Answer: B



- A. phycocyanin
- B. phycoerythrin
- C. fucoxanthin
- D. chlorophyll b

Answer: C



- **25.** Food reserve in Rhodophyta is :-
 - A. floridean starch
 - B. laminarian starch
 - C. animal starch

D. none of these

Answer: A



Watch Video Solution

26. Reserve food in Phaeophyceae is

- A. laminarin and manitol
- B. glycogen and sorbitol
- C. floridean starch
- D. glucose and starch.

Answer: A



27. Sieved septa/trumpet hyphae occur in

- A. kelps
- B. All green algae
- C. All red algae
- D. Marine algae.

Answer: A



Watch Video Solution

28. Agar, Alginic acid, Carrageenan and Funori in sea weeds are

B. 2.phycocolloids
C. 3.acids
D. 4.fats
Answer: B
Watch Video Solution
29. Red algae are similar to blue-green algae in possession of
A. similar reserve food
B. nucleus

A. 1.proteins

- C. phycobilins
- D. gas-vascuoles

Answer: C



Watch Video Solution

30. Red algae are able to grow deep in sea as

- A. 1.they can trap blue green light of short wave length in deep layer of water
- B. 2.they have chlorophyll b to trap blue green light in deep layer of water
- C. 3.Both 1 and 2

D. 4.both wrong.

Answer: A



Watch Video Solution

31. Irish Moss is

- A. Chondrus (a red alga)
- B. Ulva (green alga)
- C. Porphyra (red alga)
- D. Gelidium (red alga)

Answer: A



32. A colourless parasitic red algae is

- A. Harveyella
- B. Batrachospermum
- C. Porphyra (red alga)
- D. Cephaleuros

Answer: A



Watch Video Solution

33. Red rust of tea is caused by

- A. Cephaleuros parasiticus
- B. Puccinia graminis
- C. Harveyella
- D. Chlamydomonas gametos



- **34.** A student collected an alga and found that its cells contain both chl a and chl d as well as phycoerythrin but no chl b and flagella. The alga belongs to
- 1. Rhodophyceae
- 2. Phaeophyceae

- 3. Bacillariophyceae
- 4. Chlorophyceae
 - A. Rhodophyceae
 - B. Phaeophyceae
 - C. Bacillariophyceae
 - D. Chlorophyceae



- 35. Allophycocyanin is found in
- 1. Chlorophyceae
- 2. Rhodophyceae

- 3. Both 1 and 2
- 4. All marine algae
 - A. Chlorophyceae
 - B. Rhodophyceae
 - C. Both 1 and 2
 - D. All marine algae

Answer: B



- 36. The algal forms are mostly marine in
- 1. Chlorophyceae
- 2. Phaeophyceae

3. Ascomycetes 4. All of these A. Chlorophyceae B. Phaeophyceae C. Ascomycetes D. All of these **Answer: B Watch Video Solution** 37. Algae and fungi are characterised by the possession of

- A. 1.mitospores B. 2.chloroplast C. 3.multicellular jacketed sex organs D. 4.unicellular jacketed sex organs **Answer: A Watch Video Solution**
- **38.** What is common in Thallphytes, Bryophytes and pteridophytes?
 - A. Dependence on water
 - B. Presence of conductive system

- C. Presence of cones
- D. Absence of vascular tissue.



Watch Video Solution

- **39.** Most of the green algae are
 - A. fresh water
 - B. marine
 - C. terrestrial
 - D. epiphytic/Epizoic

Answer: A

40. Chlorophyll common between phaeophyceae and bacillariophyceae but absent in rhodophyceae is

- A. chlorophyll a
- B. chlorophyll b
- C. chlorophyll c
- D. chlorophyll e

Answer: C



A. chlorophyll a
B. xanthophyll
C. carotene
D. phycobilins
Answer: D
Watch Video Solution
42. The structure in algae helping in perennation to tide over drought are

41. The pigment found only in some algae

- A. zoospores, hypnospores and aplanospores B. zoospores, akinetes, zygospores C. aplanospores, hypnospores and zygospores. D. none of the above **Answer: C Watch Video Solution**
- **43.** A singel thylakoid per granunis found in the chloroplast of
 - A. red algae
 - B. green algae

- C. brwon algae
- D. yellow algae



Watch Video Solution

- **44.** The evolutionary sequence is
 - A. Thallophyta-Bryophyta-Pteridophyta -Phanerogams
 - B. Protophyta-Phanerogams-Crypto-gams-Monocots
 - C. Archegoniatae-Embryophyte
 - D. Archegoniatae-Embryophyte-Phan-erogams-

Monocots



Watch Video Solution

45. Sexual reproduction where a smaller and motile male gamete fuses with larger and motile female gamete known as

- A. isogamy
- B. anisogamy
- C. oogamy
- D. heterogamy

Answer: B

- **46.** Life cycle in Chlamydomonas / Ulothrix/ Spirogya is
 - A. haplontic
 - B. haplobiontic
 - C. diplontic
 - D. diplobiontic



Watch Video Solution

47. Shape of Spirogyra chloroplast is

A. scalariform conjugation
B. lateral conjugation
C. direct conjugation
D. parthenogenesis
Answer: A
Watch Video Solution
48. Which is known as Pond Scum/mermaids tresses?
A. Ulothrix
B. Nostoc
C. Spirogyra

D. Anabaena

Answer: C



Watch Video Solution

- 49. The gametangia and sporangia of Ulothrix are
 - A. jacketed and multicellular
 - B. jacketed and unicellular
 - C. nonjacketed and multicellular
 - D. non-jacketed and unicellular

Answer: D



50. Alginic acid is present in the cell wall of

A. bacillariophyceae

B. Euglena

C. Laminaria

D. Diatoms

Answer: C



Watch Video Solution

51. In phaeophyceae, thylakoids are found in group os

- A. 3's
- B. 4's
- C. 5's
- D. 6's



Watch Video Solution

52. Match the following

`{:("Phaeophyceae","fucin"),("Cyanophyceae","r-phycoerythrin"),("Rhodophyceae","c-phycocyanin"),

A. 1-A,2-B,3-C,4-D

("Diatoms", "Chrysolaminarin"):}

- B. 1-A,2-C,3-B,4-D
- C. 1-C,2-B,3-A,4-D
- D. 1-B,2-A,3-D,4-C

Answer: B



Watch Video Solution

53. Plants lacking seed and vascular tissue but forming spores and embryo are

- A. bryophytes
- B. pteridophytes
- C. angiosperms

D. gymnosperms

Answer: A



Watch Video Solution

54. Bryophytes grow in moist, humid places as

A. they have no cuticle

B. require water for fertilization

C. they do not have roots

D. all of the above

Answer: D



55. Bryophytes don't attain much height because

A. they do not have vascular supply

B. they do not have roots and mechanical tissue

C. they require water for transport of sperms

D. all of the above.

Answer: D



Watch Video Solution

56. Bryophytes can be distinguished from algae/fungi/thallophytes because they have

- A. thallus, haploid, gametophytic body
- B. chloroplast
- C. no conducting tissue
- D. sterile jacket of cells around multicellular sex organs.

Answer: D



Watch Video Solution

57. Bryophytes show

- A. 1.asexual reproduction & zygotic meiosis
- B. 2.asexual reproduction & sporic meiosis

- C. 3.no asexual reproduction but sporic meiosis
- D. 4.gametophytic dominance and sporic meiosis

Answer: D



Watch Video Solution

58. Rhizoids differ from roots as

- A. rhizoids are always unicellular but roots are multicellular
- B. rhizoids arise in gametophytes but roots in sporophytes

C. roots are branched but rhizoids are always unbranched.

D. all of the above.

Answer: B



Watch Video Solution

59. Predominant and largest gametophyte is of

- 1. Selaginella
- 2. Pinus
- 3. Moss
- 4. Rice

A. Selaginella

B. Pinus C. Moss D. Rice **Answer: C Watch Video Solution 60.** Which one of the following is true moss 1. Cord moss 2. Club moss 3. Irish moss

4. All of the above.

A. Cord moss

- B. Club moss
- C. Irish moss
- D. All of the above.



- **61.** The spores produced in capsule in moss, germinate to form haploid structure called
- 1. leafy gametophyte
- 2. protonema
- 3. prothallus
- 4. peristome

A. leafy gametophyte B. protonema C. prothallus D. peristome **Answer: B Watch Video Solution** 62. Sex organ of Funaria are A. projected and sessile B. projected and stalked C. embedded and stalked D. embedded and sessile

Answer: B



Watch Video Solution

63. Funaria is included in bryophytes because

A. it is rootless

B. it is without vascular supply

C. its sporophyte is parasite and attached to gametophyte

D. it grows near water.

Answer: C

64. Sporophyte of Funaria is

- A. Parasite on gametophyte as it is non green
- B. semiparasite on gametophyte as it is green
- C. independent of gametophyte as it is green
- D. none of the above

Answer: B



A. sucrose B. glucose C. malic acid D. citric acid **Answer: A Watch Video Solution** 66. Polytrichum has A. heart shaped prothallus B. foot, seta and capsule C. vascular bundles

D. mitospores

Answer: B



Watch Video Solution

- **67.** The sporophyte of Funaria begins development within
 - A. 1.antheridium
 - B. 2.archegonium
 - C. 3.capsule
 - D. 4.protonema

Answer: B



68. Bryophytes show

- A. oogamous
- B. isogamous
- C. anisogamous
- D. hologamy

Answer: A



Watch Video Solution

69. Bryophytes called amphibians of plant king-dom as

- A. they inhabits damp places
- B. they have tracheids
- C. their sex organs are multicellular and jacketed
- D. their reproductive phase requires water.

Answer: D



- 70. Bryophytes differ from pteridophytes in
 - A. having mitospores
 - B. the absence of vascular tissue system
 - C. lacking embryo stage

D. all of the stage

Answer: B



Watch Video Solution

71. Match List I with List II and pick up the correct choice

from the under assigned codes

List I

- (a) Thallophyta (c) Bryophyta
- (b) Embryophyta (d) Tracheophyta

List II

- (I) Embryophyta excluding Tracheophyta
- (II) Bryophyta and Tracheophyta
- (III) Phycophyta and Mycophyta
- (IV) Bryophyta and Pteridophyta
- (V) Pteridophyta and Spermatophyta

A.
$$rac{a}{IV} rac{b}{II} rac{c}{V} rac{d}{I}$$

Answer: B



Watch Video Solution

72. A liver wort is

- A. a parasite causing infection of liver
- B. a kind of virus infecting the liver
- C. a flowering plant for treating liver disorders

D. a plant without differentiation into roots, stem and leaves.

Answer: D



Watch Video Solution

73. Branched rhizoids and leafy gametophytes are characteristics of

- 1. all bryophytes
- 2. some bryophytes like mosses
- 3. algae
- 4. liverworts

A. all bryophytes

- B. some bryophytes like mosses
- C. algae
- D. liverworts.

Answer: B



- **74.** In the life cycle of Funaria, spores are beginning of the generation
- 1. gametophyte
- 2. sporophyte
- 3. peristome
- 4. both 1 and 2

- A. gametophyte
- B. sporophyte
- C. peristome
- D. both 1 and 2



- 75. Sphagnum is also called 'Peat Moss' because it
- 1. it grown in acidic marshes (bogs) and helps in peat formation
- 2. it is found in peat

- 3. it retains water
- 4. it is fossilized quickly

A. it grown in acidic marshes (bogs) and helps in peat

formation

- B. it is found in peat
- C. it retains water
- D. it is fossilized quickly.

Answer: A



Watch Video Solution

76. Moss plant is a

- A. gametophyte
- B. sporophyte
- C. sometimes gametophyte and sometimes sporophyte
- D. predominantly gametophyte with sporophyte attached to it.

Answer: D



Watch Video Solution

77. Bryophytes are close to pteridophytes in having

A. multicellualr jacketed sex organs

- B. flagellated oosphere
- C. mitospores in life cycle
- D. all of the above



Watch Video Solution

78. To which would you assign a plant that has xylem and phloem, produces meiospores, embryo but lacks seeds and flowers?

- A. Bryophytes
- B. Pteridophytes

- C. Tracheophytes
- D. Spermatophytes.

Answer: B



Watch Video Solution

79. Vascular cryptogams and botanical snakes of plant kingdom are

- A. pteridophytes
- B. tracheophytes
- C. angiosperms
- D. spermatophytes



Watch Video Solution

80. Tracheophyte include

- A. bryophytes, pteridophytes and spermato-phytes
- B. bryophytes,pteridophytes,gymnoperms and angiosperms
- C. pteridophytes and spermatophytes (gymnosperms and monocotyledons)
- D. pteridophytes and spermatophytes (gym-nosperms and angiosperms)

Answer: D



Watch Video Solution

81. Gametophytic and sporophytic phases are independent in

- A. bryophyte
- B. pteridophyte
- C. gymnosperms
- D. all of the above.

Answer: B



82. Which one of the following is considered important in the development of seed habit

Or

Seloginella has the character of evolutionary importance.

That character is

A. seed habit and heterosporous nature

B. vivipar

C. rhizophore

D. spikes.

Answer: A



83. Heterosporous pteridophytes produce

- A. dioecious gametophytes
- B. monoecious gametophytes
- C. homothallic gametophytes
- D. hermaphrodite gametophytes

Answer: A



- 84. Rhizophore in Selaginella is
- 1. root
- 2. stem

3. leaf 4. organ sui generis A. root stem B. leaf C. organ sui generis D. **Answer: D Watch Video Solution** 85. Adiantum, Pteris/ferns are 1. heterosporous 2. homosporous

3. homogametic 4. isomerous A. heterosporous B. homosporous C. homogametic D. isomerous **Answer: B Watch Video Solution** 86. The main plant body/dominant plant in life cycle of a fern/Pleridophytes is

A. gametophyte (x) B. sporophyte (2x) C. apogamous D. aposporous **Answer: B Watch Video Solution** 87. Stem in ferns is underground, monopodial root stock and called A. rhizome B. corm

- C. bulb
- D. sucker

Answer: A



Watch Video Solution

- **88.** The leaves of fern plants are called
 - A. fronds with reticulate venation
 - B. haploid
 - C. pinnately compound and grow by apical meristem
 - D. all of the above.

Answer: C

- 89. Young fern leaves and rhizome are protected by :-
 - A. fronds with reticulate venation
 - B. leaf bases
 - C. ramenta
 - D. root stock

Answer: C



A. 1.Adiantum B. 2.Cheilanthis C. 3.Pteris D. 4. Dryopteris **Answer: A Watch Video Solution** 91. Sporangia bearing leaves in Pteridophytes are called A. 1.sporophylls B. 2.megaphylls C. 3.fronds

D. 4.sorus

Answer: A



Watch Video Solution

92. Fern sperms (antherozoids) are

A. multicillated or multi flagellated and spirally coiled

B. biciliated and coiled

C. uninucleated and uniflagellated

D. multiciliated sickle shaped

Answer: A



93. What represents the gametophytic generation in pteridophytes?

- A. Main plant body
- B. Heart shaped prothallus
- C. Indusium
- D. Stomium

Answer: B



94. A collection of sporangia attached to placenta and covered over by indusium is known as

- A. sorus
- B. sporophyll
- C. ramenta
- D. spike

Answer: A



Watch Video Solution

95. Archegonium of Funaria secretes mucilage rich in

B. malic acid
C. glucose
D. citric acid
Answer: B
Watch Video Solution
96. A thick, multicelluar covering called indusium in ferns over sori is meant for
A. protection
B. producing spores

A. sucrose

- C. help in dispersal
- D. has no role

Answer: A



Watch Video Solution

97. Antheridium in ferns is

- A. multicellular and jacekted
- B. stalked, hemispherical and borne ventrally in posterior part between rhizoids
- C. its jacket has 32-48 cells
- D. all of the above.

Answer: A



Watch Video Solution

98. Life cycle in ferns is

- A. haplontic
- B. diplohaplontic
- C. haplo-haplontic
- D. diplobionntic

Answer: B



A. Ferns					
B. Grasses					
C. Gymnosperms					
D. Algae.					
Answer: A					
Watch Video Solution					
100. Spike moss is					

99. First land plant was represented by

- B. Funaria
- C. Lycopodium
- D. Adiantum

Answer: A



Watch Video Solution

101. Ferns are characterised by

- A. cambium
- B. xylem vessles
- C. young leaves with circinate ptyxis and presence of

ramenta

D. all of the above.

Answer: C



Watch Video Solution

102. Ferns resemble with mosses in one respect

- A. 1.both are embryophytes
- B. 2.both are tracheophytes
- C. 3.both are sporophytes
- D. 4.both are aquatic

Answer: A



103	Heteromor	nhic :	alternation	of gen	eration	ic	found	in
105.	петегоппог	priica	aiteinatioi	i oi gen	eration	12	iouna	ш

- A. Spirogyra
- B. Mucor
- C. Selaginella
- D. All of these.

Answer: C



Watch Video Solution

104. In ferns, circinate ptyxis (vernation) is defined as

- A. arrangement of leaf gaps in stems
- B. coiling of young leaves
- C. arrangement of sori on leaves
- D. attachement of ramenta on young parts.

Answer: B



Watch Video Solution

105. If sperms of moss and fern are put together near the archegonium of fern, only the sperms of fem find entry into archeogonium, the reason being that

A. sperms of moss are killed by larger sperms of fern

- B. archegonia of fern secretes a toxoid to kill moss of fern.
- C. archegonia of fern secrete mucilage rich in malic acid to attract sperms of fern only
- D. sperms of moss are less motile and find difficulty in entering archegonium.

Answer: C



Watch Video Solution

106. What may be the possible advantage occurring out of the presence of antheridia and archegonia on the underside of a fern prothallus?

- A. They are protected from direct rays of the sun.
- B. Capillary water accumulates on the underside of prothallus between its lower surface and the soil surface sex organs projecting in this water can be readily fertilised by the cilliated sperms which are chemotactically attracted by the archegonia
- C. The sex organs remain protected from wind
- D. Nutrients manufactured by the green prothallus can radily seep downwards to the sex organs due to the action of gravity.

Answer: B



107. Walking fern is named so as

- A. 1.it knows walking
- B. 2.it is dispersed through walking of animals in forests
- C. 3.its spores are able to move with wind
- D. 4.it spreads and propagates vegetatively by its leaf tips.

Answer: D



108. A water fern capable of fixing atmospheric nitrogen and used as biofertilizer.

- A. 1.Azolla
- B. 2.Nostoc
- C. 3.Adiantum
- D. 4.Spirulina

Answer: A



Watch Video Solution

109. Which group of gymnosperms is close to angiosperms?

A. Gnetales				
B. Ginkgoales				
C. Cycadales				
D. Confierales				
Answer: A				
Watch Video Solution				
110. Gymnosperms are different from angiosperms in				
A. 1.absence of seeds				
B. 2.absence of ovary				
C. 3.absence of ovule				

D. 4.absence of sieve cells.

Answer: B



Watch Video Solution

111. Which one constitutes the dominant vegetation in colder regions?

- A. Dicots
- B. Gymnosperms
- C. Monocots
- D. legumes

Answer: B

112. Which one of the following has not changed for the last several thousand years

- A. Pinus excelsa
- B. Ginkgo biloba
- C. Welwitschia
- D. Sequola

Answer: B



113. Largest ovules, largest male and female gametes and tallest trees are found among

- A. 1.angiosperms
- B. 2.tree ferns and some monocots
- C. 3.gymnosperms
- D. 4.dicots

Answer: C



Watch Video Solution

114. Siphonogamy in traceophytes

- A. eliminates dependence on water
- B. brings pollen grains together
- C. carries spores
- D. protects embryo

Answer: A



Watch Video Solution

115. Cycas is a

- A. living fossil
- B. fossil
- C. endangered species

D. exotic species.

Answer: A



Watch Video Solution

116. The cortex and pith of stem Cycas serves as a source of 'Sago' which is a

- A. proteins
- B. cellulose
- C. starch
- D. mixture of starch and protein.

Answer: C

117. Cycas has the largest

- A. ovule
- B. egg
- C. sperm
- D. all of the above.

Answer: C



A. 1.hermaphrodite B. 2.dioecious C. 3.monoecious D. 4.none of these **Answer: B Watch Video Solution** 119. Organised female cone is absent in A. 1.Ephedra B. 2.Pinus C. 3.Cycas

D. 4. None of these

Answer: C



Watch Video Solution

120. Cycas differs from Pteris in having

A. 1.vessels and tracheids

B. 2.motile sperms

C. 3.pollen tube

D. 4.archegonia

Answer: B



121. Coralloid roots of Cycas are useful in

A. absorption of water

B. fixation

C. absorption and fixation of nitrogen

D. respiration from air.

Answer: C



Watch Video Solution

122. Coralloid roots of Cycas possess a symbiotic alga

B. Anabaena C. Oscillatoria D. Chlorella **Answer: B Watch Video Solution** 123. Phloem of gymnosperms differ from angiosperms in A. having no companion cells B. having no sieve cells C. having phloem fibre

A. Aulosira

D. having phloem parenchyma

Answer: A



Watch Video Solution

124. Foliage leaves (needles) in Pinus are brone by

A. dwarf shoots

B. long shoots

C. both 1 and 2

D. female strobilus

Answer: A



125. At the time of dehiscence winged pollen grain of Pinus is

- A. 1)1 celled
- B. 2)4 celled
- C. 3)3 celled
- D. 4)2 celled

Answer: B



Watch Video Solution

126. Simplest and highly reduced archegonium is found in

A. 1.Pinus
B. 2.Fern
C. 3.Bryophytes
D. 4.Liver worts/mosses
Answer: A
Watch Video Solution
127. Microsporangia of Cycas occur over microsporophyll
A. abaxial
B. adaxial
C. lateral

D. margin.

Answer: A



Watch Video Solution

128. Number of generations present in Pinus seed are

A. 2

B. 3

C. 1

D. 4

Answer: B



129. Ploidy in wing of pollen grain and wing of seed of Pinus is

- A. x in pollen grain and 2x in seed
- B. 2x in pollens and 2x in seed
- C. being an outgrowth, chromosomes are absent
- D. 1/2 x in wing of pollens and polyploidy in seed.

Answer: A



130. The phenomenon of sulphur shower in pine forest is due to

- A. dispersal of winged pollens of Pinus
- B. dispersal of winged seeds of Pinus
- C. bursting of thrid year female cone of Pinus
- D. none of the above

Answer: A



Watch Video Solution

131. Which is homologous?

- A. Leaves of Moss and Selaginella
- B. Roots of ferns and moss
- C. Endosperm of Pinus and prothallus of Adianlum
- D. Pinus endosperm & endosperm of Maize



Watch Video Solution

132. In pinus

- A. seeds and ovules are winged
- B. endosperm is triploid
- C. fruits and flowers absent and seeds are naked

D. all of the above

Answer: C



Watch Video Solution

133. Chilgoza' is a fruit, obtained from a gymnosperm which is

- A. Pinus Roxburghii
- B. Pinus geradiana
- C. Cycas revoluta
- D. Abies balsamiana

Answer: B

134. Ephedrine obtained from the stem of Ephedra is given to cure

- A. gastric disorders
- B. respiratory disorders
- C. arhritis
- D. all of the above.

Answer: B



135. No Gymnosperms is

- A. 1.annual and herbaceous
- B. 2.perennial and herbaceous
- C. 3.xerophyte and woody
- D. 4.tree and shrubby

Answer: A



Watch Video Solution

136. Vessels are absent in the xylem of

A. angiosperms

- B. monocots
- C. gymnosperms excluding gnetales
- D. pteridophytes excluding ferns.



- **137.** One of the main evolutionary features of the alternation of generations from algae to flowering plants is.
 - A. gradual elaboration of gametophyte
 - B. gradual elaboration of sporophyte

- C. elinination of sporophytic tissue
- D. eliminination of gametophyte.

Answer: B



Watch Video Solution

138. In which of the following the angiosperms resemble the Gymnosperms

- A. nature of endosperm
- B. presence of vessels
- C. siphonogamy
- D. double fertilization.



139. In gymnosperms and angiosperms the pollen tube carries the male gamete to the site of fertilization and thus does not require water for fertilization. It is referred as

- A. porogamy
- B. siphonogamy
- C. mesogamy
- D. syngamy.

Answer: B



Watch Video Solution

- **140.** which one of the following statements is correct?
 - A. 1. Vascular cryptogams are all heterosporous
 - B. 2.vascular cryptogams are all homosporous
 - C. 3.Archegoniates are all homosporous
 - D. 4. Spermatophytes are all heterosporous.

Answer: D



141. Cycas has 2 cotyledons yet it is classified as Gymnosperm and not as dicot plant because

- A. 1.it resembles date palm/monocot in appearance.
- B. 2.it has compound leaves
- C. 3.its ovules are naked
- D. 4.it bears megasporophylls

Answer: C



- 142. In Cycas pollen grains shed at
- a. 2 celled stage

- b. 3 celled stage
- c. 4 celled stage
- d. 5 celled stage
 - A. 2 celled stage
 - B. 3 celled stage
 - C. 4 celled stage
 - D. 5 celled stage

Answer: B



Watch Video Solution

143. What is true for Cycas?

- A. 1.Cycas is dioecious
- B. 2.Ovules without integument
- C. 3.Largest female cone
- D. 4.All of the above.

Answer: A



- **144.** Which is incorrect about Cycas
 - A. 1.Its xylem has vessels
 - B. 2.Its young leaves show circinate vernation
 - C. 3.It does not have organised female flower

D. 4.motile sperms

Answer: A



- 145. Cycas reproduces vegetatively by
- a. sporophylis
- b. bulbils
- c. fragmentation
- d. rhizome
 - A. sporophylis
 - B. bulbils
 - C. fragmentation

D. rhizome

Answer: B



- 146. Which one is common between Funaria and Pinus?
- 1. No fruits are produced
- 2. No seeds are produced
- 3. Pollen tube is formed
- 4. Antheridia and archegonia are present
 - A. No fruits are produced
 - B. No seeds are produced
 - C. Pollen tube is formed

D. Antheridia and archegonia are present

Answer: A



- **147.** Among the following, which does not belong to sporophyte generation in Pinus?
- 1. Long shoot
- 2. Dwarf shoot
- 3. roots are branched but rhizoids are always unbranched
- 4. Endosperm
 - A. Long shoot
 - B. Dwarf shoot

C. roots are branched but rhizoids are always unbranched.

D. Endosperm

Answer: D



Watch Video Solution

148. Needle in Pinus represents

A. root

B. scale leaf

C. follage leaf

D. spur



Watch Video Solution

149. The male cone of Pinus is formed of

Or

In pinus male cone bears is large number of

A. antheridia

B. megasporophylls/green

C. microsporophylls

D. ligules

Answer: C

150. Seeds of Pinus are

- A. adaxial, endospermic and polycotyledonous
- B. abaxial, monocotyledonous and endospermic
- C. hypogeal, endospermic and monocot
- D. monocotyledonous,epigeal.

Answer: A



Watch Video Solution

151. Which of the following is not a correct match?

- A. Maiden hari fern: Ginkgo
- B. Bog Moss : Sphagnum
- C. Cord/green moss: Funaria
- D. Walking fern: Adiantum

Answer: A



- **152.** Following are are given some trends in the evolution of plants
 - A. origin of vascular system
 - B. origin of rhizoids

- C. origin of seeds
- D. origin of flowers.



Watch Video Solution

153. Characteristic of Angiosperms which distinguish them from gymnosperms

- A. presence of fruits and flowers
- B. double fertilization and triploid endosperm formed after double fertilization
- C. companion cells in phloem and vessels in xylem

D. all of the above.

Answer: D



Watch Video Solution

154. Beginning with germination of a moss spore, what is the sequence of structures that develop after germination?

I. embryo II. Gametes III. Sporophyte IV. Potonema V. gametophore

A. IV,V,III,I,II

B. V,IV,III,II,I

C. III,IV,V,II,I

D. IV,V,II,I,III

Answer: D



Watch Video Solution

155. In which of the following the angiosperms resemble the Gymnosperms

- A. presence of companion cells
- B. type of fertilization
- C. presence of ovules
- D. nature of endosperm

Answer: C

156. An example of a marine angiosperm thriving in shallow seas is

- A. Zostera
- B. Wolffia
- C. Pistia
- D. Rhizophora

Answer: A



157. Evolutionary important character of Selaginella is

- A. Haplontic life cycle
- B. Fee-living gametophyte
- C. Dependent sporophyte
- D. Heterospory

Answer: D



Watch Video Solution

158. Which of the following is grouped under phanerogams?

A. Algae show embryo stage B. Bryophytes C. Gymnosperms D. Pteridophytes **Answer: C Watch Video Solution**

- **159.** The stamen in angiosperms is homologous to which part in gymnosperm and pteridophytes?
 - A. Microsporangium
 - B. Micorosporophyll

C. Megasporophyll D. Male gametophyte **Answer: B Watch Video Solution** 160. The megasporophyll of vascular plants is analogous to which structure in angiosperms A. stamen B. ovule C. carpel D. leaf



Watch Video Solution

161. To which group of plants does the Banyan tree belong

- A. angiosperms
- B. Gymnosperms
- C. cryptogams
- D. phaeophyta

Answer: A



162. Mark the national tree

- A. Mangifera indica (Mango tree)
- B. Ficus benghalensis (banyan tree)
- C. Ficus religiosa (pipal tree)
- D. Azadirachta indica (Neem tree)

Answer: B



Watch Video Solution

163. Of the four widely known systems of classification one remains less phylogenetic and more natural, which is

- A. Engler & Prantl
- B. Benthan & Hooker
- C. Rendle
- D. Hutchinson

Answer: B



- 164. The book Genera plantarum was written by
 - A. Benthan and Hooker
 - B. Hutchinson

- C. Rendle
- D. Engler and Prantl

Answer: A



Watch Video Solution

165. Dryopteris differs from Funaria in having

- A. an independent gametophyte
- B. an independent sporophyte
- C. swimming antherozoids
- D. archegonia

Answer: B

166. Angiosperms have dominated the land flora primarily by their

- A. nature of self pollination
- B. property of producing large number of seeds
- C. domestication by man
- D. power of adaptability in diverse habitat.

Answer: D



167. In which of the following group, all the organisms are homosporous


- A. Funaria, Salvinia, Dryopteris
- B. Azolla, Selaginella, salvinia
- C. Salvinia, Cedrus, Funaria
- D. lycopodium, equisetum, phylloglossum

Answer: D



Watch Video Solution

168. See the given life cycle pattern and choose the option which it correctly represents.



- A. Haplontic
- B. Diplontic
- C. Haplo-diplontic
- D. 1 or 3

Answer: B



169. Read the following statements carefully

(i) Protonema is found in mosses and is absent in liverworts.

(ii) In Marchantia, the sporophyte is divided into the foot, seta and capsule

(iii) Archegonia are partially embedded in pteridophytes
(iv) Gymnosperms do not possess vessels in xylem except
in some gnetophytes.

A. i,ii,iii

B. ii,iv


C. i,iii,iv

D. All are correct.




Watch Video Solution

170. Find the correct match which identifies these diagrams







A. $\frac{A}{\text{Chara}}$ B C D Chara Sphagnum Salvinia Ginkgo

- B. $\frac{A}{\text{Ginkgo}}$ $\frac{B}{\text{Chara}}$ $\frac{C}{\text{Sphagnum}}$ $\frac{D}{\text{Salvinia}}$
- c. $\frac{A}{\text{Sphagnum}}$ $\frac{B}{\text{Ginkgo}}$ $\frac{C}{\text{Chara}}$ $\frac{D}{\text{Salvinia}}$
- D. $\frac{A}{\text{Ginkgo}}$ $\frac{B}{\text{Sphagnum}}$ $\frac{C}{\text{Salvinia}}$ $\frac{D}{\text{Chara}}$



171. The algal group which most biologists believe to have led to evolution of land plants is

- A. phaeophyta
- B. Rhodophyta
- C. Chlorophyta

D. Chrysophyta

Answer: C



Watch Video Solution

172. Phycocolloids are found in the cell walls of

- A. Diatoms
- B. Spirogyra
- C. Red and brown algae
- D. All algae

Answer: C



173. Which one of the following is not characteristic of all divisions of vascular plants?

- A. An alternation of generations
- B. The development of seeds
- C. Differentiation into roots, stems and leaves
- D. Xylem and phloem for transporting materials between roots and leaves.

Answer: B



174. Red algae differs from green and brown algae in


- A. No chlorophyll a
- B. No differentiated cells
- C. No phycocyanin within their cells
- D. No flagellated stages in their life cycles

Answer: D



Watch Video Solution

175. Given is a picture of bryophyte. The correct ploidy levels of the indicated structures are



A. 1)1:2n,2:n,3:n

B. 2)1:n,2:n,3:n

C. 3)1:2n,2:2n,3:2n

D. 4)1:2n,2:n,3:2n

Answer: A



- 176. Bryophytes resemble algae in the following aspects -
 - A. Differentiation of the plant body into root,stem and heterotrophic mode of nutrition
 - B. Thallus-like plant body, lack of vascular tissue: absence of root: and autotrophic mode of nutrition
 - C. Thallus-like plant body, presence of roots, and autotrophic mode of nutrition
 - D. Filamentous body: presence of vascular tissue: and autotrophic mode of nutrition

Answer: B



177. Funaria differs from Pteridium in the absence of

- A. Root
- B. Seed
- C. Archegonia
- D. Embryo

Answer: A



Watch Video Solution

178. Prothallus is

- A. Gametophyte,monoecious,autotroph present in pteridophytes
- B. Gametophyte,monoecious,autotroph found in bryophytes
- C. Sporophyte,dioecious,heterotroph found in bryophytes
- D. Gametophyte,dioecious,heterotroph present in pteridophytes

Answer: A



179. In gymnosperms how many male gametes are produced by each pollen grain

- A. 4
- B. 3
- C. 2
- D. 1

Answer: C



Watch Video Solution

180. Which arrangements of the organisms represents a rank ordering (based on size/importance) from dominant

gametophyte on the left to dominant sporophyte on the right?

A. Polytrichum,Cycas,Equisetum

B. Lycopodium,Zostera,Riccia

C. Sphagnum,Pteridium,Cedrus

D. Pinus, Selaginella, Marchantia

Answer: C



Watch Video Solution

181. Red algae do not have

A. zoospores, hypnospores and aplanospores

- B. Cellulose in cell wall
- C. Floridean starch
- D. Sexual reproduction.

Answer: A



Watch Video Solution

182. Select the wrong statement

- A. Gymnosperm do not have antheridia
- B. Cycas and ferns have multiflagellate gametes
- C. Selaginella is a heterosporous non seed plant
- D. Pinus tree is dioecious



Watch Video Solution

183. Ovule of Pinus can be called

- A. Female gametophyte
- B. Male gametophyte
- C. Megasporangium
- D. Integumented megasoporangium

Answer: D



184	Conifers	differ	from	σταςςρς	in	the
104.	Conners	umer	11 0111	grasses	Ш	uie

- A. Production of seeds from ovules
- B. Lack of xylem tracheids
- C. Absence of pollen tubes
- D. Formation of endosperm before fertilization.



Watch Video Solution

185. "Ordines Anomali" of Benthan and Hooker includes

- A. Seed plants showing abnormal forms of growth and development
- B. Plants represented only in fossil state.
- C. Plants described in the literature but which

 Bentham and Hooker did not see in original
- D. A few orders which could not be placed satisfactorily in the classification.



Watch Video Solution

186. Which of the following is a wrong combination?

- A. Haploid endosperm, archegonia present but antheridium absent-Gymnosperms
- B. Triploid endosperm and 7-celled female gametophyte called embryo sac-Angiosperms
- C. Embryo stage absent,reproduction by mitospores-Algae
- D. Gametophyte independent, biflagellate gametes and reproduction by accessory spores-Mosses.



187. Green algae are also included among the items carried by astronauts in a space ship Which of the following are the reasons for this?

1. These being very simple plants, are amenable to a variety of experiments. 2. These may provide nourishment to the persons on the space ship. 3. These act as a source of oxygen.

Select the correct answer using the codes given below.

- A. 1,2 and 3
- B. 1 and 3
- C. 1 and 2
- D. 2 and 3

Answer: A



188. Consider the following pigments 1. Chlorophyll a 2.

Chlorophyll b 3. Chlorophyll c 4. Carotenes 5. Biliproteins

6. Xanthophylls

The characteristic photosynthetic pigments present in Chlorophyceae would include

- A. 1,2,3 and 4
- B. 1,3,4, and 5
- C. 2,3,5 and 6
- D. 1,2,4 and 6



- **189.** Which one of the following pairs is not correctly matched?
 - A. Red Sea: Trichodesmium erythreum a cyanobacterium
 - B. Red Tides: Rhodymenia, a red algae
 - C. Red Snow: Chalmydomonas nivalis, a green algae
 - D. Red Rust of Tea: Cephaleuros virescens, a green algae

Answer: B



Watch Video Solution

190. Consider the following statements

1. In pteridophytes, the phloem lacks companion cells. 2.

The xylem lacks vessels in majority of pteridophytes. 3.

Heterosporous pteridophytes have monoecious

gametophytes. 4. Water is essential for fertilization in

pteridophytes.

Which of these statements are correct?

- A. 1,2,3 and 4
- B. 2,3 and 4

- C. 1 and 3
- D. 1,2 and 4



- **191.** Which one of the following statements is correct?
 - A. Leaves of conifers are perennial
 - B. All vascular plants have Sieve Tube Members
 - C. The red and edible flesh of strawberry is the carpel
 - tissue
 - D. Conifers have motile gametes.

Answer: A



Watch Video Solution

192. In pinus

- A. Megaspore, endosperm and pollen
- B. Microspore, megaspore and endosperm
- C. Megaspore, integument and root
- D. Microspore, leaf and endosperm

Answer: B



193. Read the following statements carefully

(i) Protonema is found in mosses and is absent in liverworts.

(ii) In Marchantia,the sporophyte is divided into the foot, seta and capsule

(iii) Archegonia are partially embedded in pteridophytes
(iv) Gymnosperms do not possess vessels in xylem except
in some gnetophytes.

A. 1 and 2

B. 3 and 4

C. 1,2 and 3

D. 2 and 3

Answer: A

194. Consider the following statemetns

1. The thylakoids of blue-green algae are arranged singly in the stroma of chloroplasts. 2. In blue-green algae, the thylkoids are not only the sites for photosynthesis but also for respiration

Which of the statements given above is/are correct?

- A. 1 only
- B. 2 only
- C. both 1 and 2
- D. Neither 1 nor 2

Answer: B



- 195. In gymnosperms the pollen chamber represents
 - A. A cavity in the ovule in which pollen grains are stored after pollination
 - B. An opening in the megagametophyte through which the pollen tube approaches the egg
 - C. The microsporangium in which pollen grains develop

D. A cell in the pollen grain in which the sperms are formed.

Answer: A



Watch Video Solution

196. Bryophytes can be compared with amphibians since both are

(ii) Autotrophic (ii) Dependent on water for reproduction (iii) Without endoskeleton (iv) Devoid of impervious body surface (v) With separate sexes

A. ii,iv

B. i,ii,iv

C. ii,iii,iv

D. i,iii,iv

Answer: A



Watch Video Solution

197. In which of the following groups all the organisms contain chlorophyll c

A. Ectocarpus, Fucus, Sargassum, Laminaria

B. Macrocystis, Batrachospermum, Polysliphonia, Characium

C. Chlamydomonas, Asterocystis, Ulothrix, Gelidium

D. Harveyella, Chondrus, Volvox, Chlorella

Answer: A



Watch Video Solution

198. In which of the following group all the orgaisms contain phycobilins?

- A. 1. Anabaena, Gelidium, Nereocystis, Zoochlorella
- B. 2.Nostoc, Porphyra, Chondrus, Batrachospermum
- C. 3.Oscillatoria, Sargassum, Volvox, Fucus
- D. 4. None of the above.

Answer: B



199. In which of the following groups all the organisms lack motile stages in the life cycle?

- 1. Chlamydomonas, Ulothrix, Sargassum
- 2. Chondrus, Gelidium, Batrachospermum
- 3. Chlorella, Polysiphonia, Volvox
- 4. All of the above
 - A. Chlamydomonas, Ulothrix, Sargassum
 - B. Chondrus, Gelidium, Batrachospermum
 - C. Chlorella, Polysiphonia, Volvox
 - D. All of the above

Answer: B



Match video Solution

200. Go through the following statements

- (i) In chlorophyceae the flagella are 2 in number, unequal in size and lateral in position
- (ii) In phaeophyceae flagella are 2-8 in number, equal in

size and apical in position (iii) Chlorella and Spirullina re rich in proteins and are used as food supplements even

by space travellers. (iv) The leaves in pteridophyta are microphylls as in Selaginella or macrophylls as in ferns.

Which of these are correct

- a. i,ii,iii
- b. ii,iii
- c. i,ii,iv
- d. iii,iv

- A. i,ii,iii
- B. ii,iii
- C. i,ii,iv
- D. iii,iv



- 201. In a moss the sporophyte
- 1. arises from a spore produced from the gametophyte
- 2. manufactures food for itself, as well as for the gametophyte

- 3. is partially parasitic on the gametophyte
- 4. Produces gametes that give rise to the gametophyte
 - A. arises from a spore produced from the gametophyte
 - B. manufactures food for itself, as well as for the gametophyte
 - C. is partially parasitic on the gametophyte
 - D. Produces gametes that give rise to the gametophyte

Answer: C



202. Which of these is mismatched

- A. 1.phaneros-visible
- B. 2.kryptos-concealed
- C. 3.gymno-naked
- D. 4.thallus-diploid

Answer: D



Watch Video Solution

203. Which of the following is a vascular cryptogam

A. 1.Marchantia

- B. 2.Cedrus
- C. 3.Equisetum
- D. 4.Ginkgo

Answer: C



Watch Video Solution

204. Which one of the following plants is monoecious

- A. Cycas
- B. Papaya
- C. Marchantia
- D. Pinus



Watch Video Solution

205. Which one of the following has haplontic life cycle

- A. Ustilago
- B. Wheat
- C. Funaria
- D. Polytrichum

Answer: A



A. Fucus
B. Gracillaria
C. Chara
D. Porphyra
Answer: A Watch Video Solution
207. Young leaves of Cycas show
A. Simple venation

206. Mannitol (sugar alcohol) is the stored food in

- B. Circinate venation
- C. Alternate arrangement
- D. Opposite arrangement

Answer: B



Watch Video Solution

208. Algae have cell wall made up of

- A. Cellulose, hemicellulose and pectins
- B. Cellulose, galactans and mannans
- C. Hemicellulose, pectins and proteins
- D. Pectins, cellulose and proteins

Answer: B



209. Which of the following algal groups has no motile stage

- 1. Brown
- 2. Yellow
- 3. Red
- 4. Green
 - A. Brown
 - B. Yellow
 - C. Red

D. Green

Answer: C



Watch Video Solution

210. Sphagnum is an example of

- 1. Moss
- 2. Pteridophyte
- 3. Algae
- 4. Gymnosperm
 - A. Moss
 - B. Pteridophyte
 - C. Algae

D. Gymnosperm

Answer: A



Watch Video Solution

211. Consider the following four statements whether they are correct or wrong.

- A. The sporophyte in liverworts is more elaborate than that in mosses
- B. Salvinia is heterosporous
- C. The life-cycle in all seed-bearing plants is diplontic

D. In Pinus male and female cones are borne on different trees

Answer: B



Watch Video Solution

212. Selaginella and Salvinia are considered to represent a significant step toward evolution of seed habit because

A. Female gametophyte is free and gets dispersed like seeds

B. Female gametophyte lacks archegonia


- C. Megaspores possess endosperm and embryo surrounded by seed coat
- D. Embryo develops in female gametophyte which is retained on parent sporophyte

Answer: D



Watch Video Solution

213. Examine the figure given below and select the right option given all the four parts (a,b,c,d) correctly



identified.

A.

B. Archegoniphore Female thallus Bud Foot

C. $\frac{a}{\text{Seta}} \frac{b}{\text{Sporophyte}} \frac{c}{\text{Protonema}} \frac{d}{\text{Rhizoids}}$

D. $\frac{a}{\text{Anteridiophore}}$ $\frac{b}{\text{Male thallus}}$ $\frac{c}{\text{Globule}}$ Roots

Answer: A



Watch Video Solution

214. Archegoniophore is present in

- A. 1.Marchantia
- B. 2.Chara
- C. 3.Adiantum
- D. 4.Funaria

Answer: A



215. Compared with the gametophytes of the bryophytes, the gametophytes of vascular plants tend to be:-

- A. smaller but to have larger sex organs
- B. larger but to have smaller sex organs
- C. larger and to have larger sex organs
- D. smaller and to have smaller sex organs

Answer: D



Watch Video Solution

216. The gametophyte is not an independent, free-living generation in

B. Adiantum C. Marchantia D. Pinus **Answer: D Watch Video Solution** 217. Asexual reproduction in Liverworts takes place by the formation of specialized structures called A. Gemmae B. zoospores, akinetes, zygospores

A. Polytrichum, Cycas, Equisetum

- C. Sporangia
- D. Microspores

Answer: A



Watch Video Solution

218. Pinus belongs to the class

- A. gnetopsida
- B. cycadopsida
- C. coniferopsida
- D. sphenopsida

Answer: C

- **219.** Read the following five statements (A-D) and answer as asked next to them
- (A) In Equisetum the female gametophyte is retained on the parent sporophyte
- (B) In ginkgo male gametophyte is not independent
- (C) Sexual reproduction in Volvox is isogamous
- (E) The spores of slime moulds lack cell walls

How many of the above statements are correct

- A. Four
- B. One
- C. Two

D. Three

Answer: B



Watch Video Solution

220. Which one of the following pairs is wrongly matched?

- A. viroids-RNA
- B. Mustard-Synergids
- C. Ginkgo-Archegonia
- D. Salvinia-Prothallus

Answer: D

221. How many organism in the list given below are autotrophs Lactobacillus, Nostoc, Chara, Nitrosomonas, Nitrobacter, Streptomuces, Sacharomyces, Trypanosoma, Porphyra Wolfia

- A. Six
- B. Three
- C. Four
- D. Five

Answer: A



222. Cycas and Adiantum resemble each other in having

- A. Motile sperms
- B. Cambium
- C. Vessles
- D. Seeds

Answer: A



Watch Video Solution

223. Which one of the following is common to multicellular fungi, filamentous algae and protonema of

mosses
A. Members of kingdom plantae
B. Mode of Nutrition
C. Multiplication by fragmentation
D. Diplontic life cycle

Answer: C



Watch Video Solution

224. Which one of the following is a correct statement

A. In gymnosperms female gametophyte is free-living

- B. Anteridiophores and archegoniophores are present in pteridophytes
- C. Origin of seed habit can be traced in pteridophytes
- D. Pteridophyte gametophyte has a protonemal and leafy stage.

Answer: C



- 225. Sex-organs of pteridophytes are
 - A. 1.Multicellular and jacketed
 - B. 2.Unicellular and jacketed

- C. 3.Unicellular and non jacketed
- D. 4.Multicellular and non jacketed

Answer: A



Watch Video Solution

226. The chloroplasts of algae usually lack

- A. Quantasomes
- B. Lamellae
- C. Pigments reticulum
- D. Grana

Answer: D

227. First amphibian plants of the plant kingdom are

- A. Thallophytes
- B. Bryophytes
- C. pteridophytes
- D. Gymnosperms

Answer: B



A. macrophylls B. microphylls C. sporophylls D. megasporophylls **Answer: A Watch Video Solution** 229. The tallest tree species of the gymnosperms is A. Cycas B. Pinus C. Sequoia

D. None of the Above

Answer: C



Watch Video Solution

230. Besides paddy fields, cyanobacteria are also found inside vegetative part of:

- A. 1.Equisetum
- B. 2.Psilotum
- C. 3.Pinus
- D. 4.Cycas

Answer: D

231. Select the wrong statement

- A. 1.In Oomycetes female gamete is smaller and motile, while male gamete is larger and non-motile
- B. 2.Chlamydomonas exhibits both isogamy and anisogamy and Fucus shows oogamy
- C. 3.Isogametes are similar in structure, function and behaviour
- D. 4. Anisogametes differ either in structure, function or behaviour.

Answer: A



232. Which of the following represent maximum number of species among global biodiversity

- 1. Fungi
- 2. Mosses and Ferns
- 3. Algae
- 4. Lichens
 - A. Fungi
 - B. Mosses and Ferns
 - C. Algae

D. Lichens

Answer: A



Watch Video Solution

233. Male gametophyte with least number of cells is present in

- A. Pinus
- B. Pteris
- C. Funaria
- D. Lilium

Answer: D

234. Which one of the following shows isogamy with non-flagellated gametes

- A. Spirogyra
- B. Sargassum
- C. Ectocarpus
- D. Ulothrix

Answer: A



235. An alga which can be employed as food for humna being is

- A. polysiphonia
- B. Ulothrix
- C. Chlorella
- D. Spirogyra

Answer: C



Watch Video Solution

236. Which of the following is responsible for peat formation?

B. Marchantia C. Riccia D. Funaria **Answer: A Watch Video Solution** 237. In which of the following, gametophyte is not independent in free living? A. Marchantia B. Pteris

A. Sphagnum

- C. Pinus
- D. Funaria

Answer: C



- **238.** Read the following five statements (A to E) and select the option with all correct statement
- (A) Mosses and Lichens are the first organisms to colonise a bare rock.
- (B) Selaginella is a homospourous pteridophyte.
- (C) Coralloid roots in Cycas have VAM.
- (D) Main plant body in bryophytes is gametophytic whereas in pteridophytes it is sporophytic.

(E) In Gymnoperms, male and female gametophytes are present within sporangia located on sporophyte.

- A. B,C and D
 - B. A,D and E
- C. B,C and E
- D. A,C and D

Answer: B



- 239. Male gametes are flagellated in
- 1. Anabaena
- 2. Ectocarpus

- 3. Spirogyra
- 4. Polysiphonia
 - A. Anabaena
 - B. Ectocarpus
 - C. Spirogyra
 - D. Polysiphonia

Answer: B



- **240.** Which one of the following statements is wrong?
- 1. Agar-agar is obtained from Gelidium and Gracilaria
- 2. Chlorella and Spirullina are used as space food

- 3. Mannitol is stored food in Rhodophyceae
- 4. Algin and carrageenan are products of algae.
 - A. Agar-agar is obtained from Gelidium and Gracilaria
 - B. Chlorella and Spirullina are used as space food
 - C. Mannitol is stored food in Rhodophyceae
 - D. Algin and carrogen are products of algae.

Answer: C



Watch Video Solution

241. Which one is a wrong statement?

- A. Archegonia are found in Bryophyta, Pteridophyta and Gymnosperms
- B. Mucor has biflagellate zoospores
- C. Haploid endosperm is typical feature of gymnosperms
- D. Brown algae have chlorophyll a and c,and fucoxanthin

Answer: B



242. Male gametophyte in angiosperms produces:

- 1. two sperms and a vegetative cell
- 2. single sperm and a vegetative cell
- 3. single sperm and two vegetative cells
- 4. three sperms
 - A. two sperms and a vegetative cell
 - B. single sperm and a vegetative cell
 - C. single sperm and two vegetative cells
 - D. three sperms

Answer: A



243. Select the correct statement

- A. Salvinia, Ginkgo and Pinus all are gymnosperms
- B. Sequoia is one of the tallest trees
- C. The leaves of gymnosperms are not well adapted to extremes of climate
- D. Gymnosperms are both homosporous and heterosporous

Answer: B



244. In bryophytes and and pteridophytes, transport of male gametes requires

- A. Insects
- B. Birds
- C. Water
- D. Wind

Answer: C



Watch Video Solution

245. Conifers are adapted to tolerate extreme environmental conditions because of

- A. 1.Broad hardy leaves
- B. 2.Superficial stomata
- C. 3. Thick cuticle
- D. 4.Presence of vessels.

Answer: C



Watch Video Solution

246. Which one of the following statements is wrong?

A. Algae increase the level of dissovledf oxygen in the immediate environment.

- B. Algin is obtained from red algae, and carrageenin from brown algae.
- C. Agar-agar is obtained from Gelidium and Gracilaria
- D. Laminaria and Sargassum are used as food.

Answer: B



- **247.** Zygotic meiosis is characteristic of
 - A. Marchantia
 - B. Fucus
 - C. Funaria

D. Chlamydomonas

Answer: D



Watch Video Solution

248. An example of colonial algae is

A. 1.Chlorella

B. 2.Volvox

C. 3. Ulothrix

D. 4.Spirogyra

Answer: B



249. Select the mismatch

- A. Pinus-Dioecious
- **B.** Cycas-Dioecious
- C. Salvinia -Heterosporous
- D. Equisetum-Homosporous

Answer: A



Watch Video Solution

250. Life cycle of Ectocarpus and Fucus respectivley are

- A. haplontic, diplontic
- B. diplontic, haplodiplontic
- C. haplodiplontic, diplontic
- D. haplodiplontic, haplontic

Answer: C



Watch Video Solution

251. Which one is wrongly matched

- A. Unicellular organism -Chlorella
- B. Gemma cups-Marchantia
- C. Biflagellate Zoospores-Brown algae

D. Uniflagellate gametes-Polysiphonia

Answer: D



Watch Video Solution

252. Which of the following statements is correct?

- 1. Stems are usually unbranched in both Cycas and Cedrus.
- 2. Horsetails are gymnosperms.
- 3. Selaginella is heterosporous, while salvinia is homosporous.
- 4. Ovules are not enclosed by ovary wall in gymnosperms.

- A. Stems are usually unbranched in both Cycas and Cedrus.
- B. Horsetails are gymnosperms.
- C. Selaginella is heterosporous, while salvinia is homosporous.
- D. Ovules are not enclosed by ovary wall in gymnosperms.

Answer: D



Watch Video Solution

Assertion And Reason

- 1. [A]: Flagella found in green algae are of tinsel type.
- [R]: Flagella in algae are whiplash type.
 - A. If both A and R are true and R is the correct explanation of A
 - B. If both A and R are true but R is not the correct explanation of A
 - C. If A is true and R is false
 - D. If both A and R are false

Answer:



2. [A]: The mosses form dense growth.

[R]: They produce a large number of spores.

A. If both A and R are true and R is the correct explanation of A

B. If both A and R are true but R is not the correct explanation of A

C. If A is true and R is false

D. If both A and R are false

Answer:



3. [A] : Pteridophytes and Gymnosperms are included in Archegoniate .

 $\left[R \right]$: They have archegonia as female reproductive organ .

A. 1)If both A and R are true and R is the correct explanation of A

B. 2)If both A and R are true but R is not the correct explanation of A

C. 3)If A is true and R is false

D. 4)If both A and R are false

Answer:



4. [A]: Pinus bears resin canals.

[R]: Resin canals are found in cortex and bounded by a glandular epithelial layer that secretes resin.

A. If both A and R are true and R is the correct explanation of A

B. If both A and R are true but R is not the correct explanation of A

C. If A is true and R is false

D. If both A and R are false

Answer:



- 5. [A]: In terms, prothallus is club shaped.
- [R] : Prothallus is derived from club shaped zoosporangia

•

- A. If both A and R are true and R is the correct explanation of A
- B. If both A and R are true but R is not the correct explanation of A
- C. If A is true and R is false
- D. If both A and R are false

Answer:



6. [A] : Either megasporophylls or microsporophylls are found in Gymnosperms .

[R]: Both mega and microsporophylls are common in Gymnosperms . Megasporophylls are smaller than microsporophylls .

A. If both A and R are true and R is the correct explanation of A

B. If both A and R are true but R is not the correct explanation of A

C. If A is true and R is false

D. If both A and R are false

Answer:

7. [A]: Rhizophore is considered to be stem structure.

[R]: Rhizophore develops from angle meristem present between the two branches of stem.

A. If both A and R are true and R is the correct explanation of A

B. If both A and R are true but R is not the correct explanation of A

C. If A is true and R is false

D. If both A and R are false

Answer:

8. [A] : In Selaginella/Pteris reduction division occurs during microspore formation only .

[R]: It was first shown by Zacharis and Strasburger in 1963.

A. If both A and R are true and R is the correct explanation of A

B. If both A and R are true but R is not the correct explanation of A

C. If A is true and R is false

D. If both A and R are false

Answer:



9. [A] : Neottia and Monotrapa are saprophytic angiosperms .

[R]: The association of a fungus with roots of higher plants is mycorrhiza.

A. If both A and R are true and R is the correct explanation of A

B. If both A and R are true but R is not the correct explanation of A

C. If A is true and R is false

D. If both A and R are false

Answer:



Watch Video Solution

10. [A] : Fossil plants serve as an index for paleoclimates .

[R] : Because plant adaptations to varied environmental conditions are well known .

A. If both A and R are true and R is the correct explanation of A

B. If both A and R are true but R is not the correct explanation of A

- C. If A is true and R is false
- D. If both A and R are false

Answer:



Watch Video Solution

11. [A] : Coralloid root of Cycas is confined to bacterial zone.

[R]: Nucleus is absent in young sieve cells.

A. If both A and R are true and R is the correct explanation of A

B. If both A and R are true but R is not the correct explanation of A

C. If A is true and R is false

D. If both A and R are false

Answer:



Watch Video Solution

12. [A] : Heterospory has led to the reduction and specialization in spores .

[R]: Incipient heterospory has been found to occur in the fern Platyzoma microphylla.

A. If both A and R are true and R is the correct explanation of A

B. If both A and R are true but R is not the correct explanation of A

C. If A is true and R is false

D. If both A and R are false

Answer:



Watch Video Solution

13. [A]: In Pinus only the dwarf roots become mycorrhizal

[R] : Seeds of Pinus are polycotyledonous .

A. If both A and R are true and R is the correct explanation of A

B. If both A and R are true but R is not the correct explanation of A

C. If A is true and R is false

D. If both A and R are false

Answer:



Watch Video Solution

14. [A] : The mosses like liverworts , do not exhibit alternation of generation .

[R] : The adult gametophyte is conspicuous leafy green , photosynthetic plant in ferns .

A. If both A and R are true and R is the correct explanation of A

B. If both A and R are true but R is not the correct explanation of A

C. If A is true and R is false

D. If both A and R are false

Answer:



15. [A]: Mosses do not seem to resist infestations by most fungi and are often seriously damaged by insects.

[R]: They do not possess antibiotic material.

A. If both A and R are true and R is the correct explanation of A

B. If both A and R are true but R is not the correct explanation of A

C. If A is true and R is false

D. If both A and R are false

Answer:



16. [A]: Monotrapa is a parasitic plant.

[R]: It does not show proper mycorrhizal association.

A. If both A and R are true and R is the correct explanation of A

B. If both A and R are true but R is not the correct explanation of A

C. If A is true and R is false

D. If both A and R are false

Answer:



17. [A]: Some of the Sea kelps are used as fertilizers.

[R]: They are important source of minerals.

A. 1)If both A and R are true and R is the correct explanation of A

B. 2)If both A and R are true but R is not the correct explanation of A

C. 3)If A is true and R is false

D. 4)If both A and R are false

Answer:



18. [A]: Angiosperms and gymnosperms both are flowering plants .

[R]: They both form covered seeds.

A. 1)If both A and R are true and R is the correct explanation of A

B. 2)If both A and R are true but R is not the correct explanation of A

C. 3)If A is true and R is false

D. 4)If both A and R are false

Answer:



