



# BIOLOGY

## BOOKS - TRUEMAN BIOLOGY

### HUMAN REPRODUCTION

#### Multiple Choice Questions

1. Which of the following is secondary sex organ?

A. Beard

B. Uterus

C. Ovary

D. Broad hips

**Answer: B**



**Watch Video Solution**

**2. Scrotal sacs of man is connected with the abdominal cavity by-**

A. epididymis

B. spermatic canal

C. inguinal canal

D. haversian canal

**Answer: C**



**Watch Video Solution**

**3.** In mammals, failure of testes to descend into the scrotum is known as

A. castration

B. impotency

C. paedogenesis

D. cryptorchidism

**Answer: D**



**Watch Video Solution**

**4. Mesorchium in frog refers to**

A. capsule in testis

B. capsule in ovary

C. a peritoneal fold that covers testis

D.

**Answer: C**



**Watch Video Solution**

**5. Tunica albuginea is the covering around**

A. testes

B. kidneys

C. uterus

D. epididymis

**Answer: A**



**Watch Video Solution**

6. Supporting cells found in between the germinal epithelium is called:

A. Phagocytes

B. Sertoli cells

C. Leydig cells

D. granular cells

**Answer: B**



**Watch Video Solution**

7. Sertoli cells are found in testis. These cells are

A. nurse cells

B. reproductive cells

C. receptor cells

D. none of these

**Answer: A**



**Watch Video Solution**

**8. Rete testis opens to**

A. urethra

B. vasa efferentia

C. bidder's canal



D. caput epididymis

**Answer: B**



**Watch Video Solution**

**9. Major part of semen is secreted by:**

A. Seminal vesicle

B. prostate gland

C. Cowper's gland

D. bartholin's gland

**Answer: A**



**Watch Video Solution**

**10.** The common duct formed by the union of vas deferens and duct from seminal vesical is :

- A. urethra
- B. stenson's duct
- C. spermatic duct
- D. ejaculatory duct

**Answer: D**



**Watch Video Solution**

**11.** Which gland in mammel makes alkaline secretion for lubrication?

A. Testis

B. Pineal body

C. Cervical glands

D. Cowper's gland

**Answer: D**



**Watch Video Solution**

**12. Seminal fluid contains the secretion of:**

A. follicles, uterus and prostate gland

B. prostate, cowper's and bartholin's gland

C. seminal vesicle, uterus and prostate  
gland

D. seminal vesicle, prostate and Cowper's gland

**Answer: D**



**Watch Video Solution**

**13.** Which one is unpaired gland in male reproductive system?

A. Seminal vesicle

B. Cowper's gland

C. Prostate gland

D. Lacrimal gland

**Answer: C**



**Watch Video Solution**

**14.** Which of the following sugars in semen is a source of energy for the spermatozoa?

A. sucrose

B. fructose

C. glucose

D. galactose

**Answer: B**



**Watch Video Solution**

**15.** Sugar fructose is present in the secretion of :

A. seminal vesicles

B. Cowper's gland

C. peneal gland

D. bartholin's gland

**Answer: A**



**Watch Video Solution**

**16.** At what speed a human sperm moves in the female genital tract?

A. 3mm/min

B. 10 mm/min



C. 15mm/min

D. 20 mm/min

**Answer: A**



**Watch Video Solution**

**17.** Which is the correct sequence of layers in the mammalian egg from outside to inside?

A. Zona pellucida, corona radiata, plasma membrane

B. Corona radiata, zona pellucida, plasma membrane

C. Plasma membrane, zona pellucida, corona radiata

D. None of the above

**Answer: B**



**Watch Video Solution**

**18.** Bartholin's glands of female correspond to which gland in male?

- A. Rectal glands
- B. inguinal glands
- C. prostate glands
- D. Cowper's glands

**Answer: D**



**Watch Video Solution**

19. labium majora of a female mammalis homologous to

- A. scrotal sac
- B. prostate gland
- C. epididymis
- D. seminal vesicle

**Answer: A**



**Watch Video Solution**

20. Which of the following hormones is active during proliferative phase of menstrual cycle?

- A. Estrogen
- B. Progesterone
- C. Testosterone
- D. All of these

**Answer: A**



**Watch Video Solution**

21. Progesterone hormone is active during

- A. follicular phase
- B. secretory phase
- C. menstrual phase
- D. proliferative phase

**Answer: B**



**Watch Video Solution**

22. Cessation of menstrual cycle in women is called

A. menarche

B. menopause

C. impotency

D. puberty

**Answer: B**



**Watch Video Solution**

23. At menopause there is rise in urinary excretion of

A. FSH

B. STH

C. Oxytocin

D. Oestrogen

**Answer: A**



**Watch Video Solution**



24. In spermatogenesis, the phase of maturation involves

A. The growth of spermatogonia into primary spermatocyte

B. The formation of spermatogonia from gonocytes through mitosis

C. The formation of spermatids from primary spermatocytes through meiosis

D. The formation of spermatogonia from the spermatocytes through meiosis

**Answer: C**



**Watch Video Solution**

**25.** Which of the following is haploid in nature?

A. Spermatids

B. Spermatogonia

C. Primary spermatocytes

D. Tertiary spermatocytes

**Answer: B**



**Watch Video Solution**

**26.** The male gamete is human is

A. Ovum

B. Ovary

C. Sperm

D. Testis

**Answer: B**



**Watch Video Solution**

**27. Acrosome of sperm is formed from**

A. nucleus of spermatid

B. centrosome of spermatid

C. mitochondria of spermatid

D. golgi complex of spermatid

**Answer: D**



**Watch Video Solution**

**28.** How many centrioles are normally present in a sperm?

A. One

B. Two

C. Many

D. None of these

**Answer: B**



**Watch Video Solution**

**29.** Which organelle is absent in human sperm?

A. ER

B. Nucleus

C. Centriole

D. Mitochondria

**Answer: A**



**Watch Video Solution**

**30.** Middle piece of sperm contains

A. mitochondria and golgi body

B. centriole and golgi body

C. axial filament and golgi body

D. mitochondria and axial filament

**Answer: D**



Watch Video Solution

31. Oogonium is

A. haploid

B. diploid

C. triploid

D. euploid

**Answer: B**



Watch Video Solution



32. The number of chromosomes in a mature gamete gets halved during

A. Meiosis II

B. Formation of first polar body

C. Formation of second polar body

D. Division of secondary oocyte and secondary spermatocyte

**Answer: B**



**Watch Video Solution**

**33.** In which phase of cell division is oocyte arrested ?

A. Interphase

B. Prophase I

C. Anaphase II

D. Both prophase I and II

**Answer: B**



**Watch Video Solution**

34. 100 eggs and 100 sperms can be produced from \_\_\_\_\_ and \_\_\_\_\_ meiotic division respectively.

A. 25,25

B. 100,25

C. 100,100

D. 25,100

**Answer: B**



**Watch Video Solution**

**35.** One million oocytes and one million secondary spermatocytes will give:

- A. 2 million ova 1 million sperms
- B. 2 million ova and 2 million sperms
- C. 1 million ova and 2 million sperms
- D. 1 million ova and 1 million sperms

**Answer: C**



**Watch Video Solution**

**36.** Which statement is true?

A. At the onset of menopause, the human female stops producing FSH & LH

B. Primary oocytes are produced by the human female throughout adolescence

C. Oocytes produced by the females are stored in the seminiferous tubules

D. At birth, the human female has produced all the oocytes she will ever

produce

**Answer: D**



**Watch Video Solution**

**37. Cytoplasm of ovum does not contain:**

- A. Ribosomes
- B. Mitochondria
- C. Golgi bodies
- D. Centrosomes

**Answer: D**



**Watch Video Solution**

**38.** During a women's life time, she produces about:

- A. 40-50 eggs
- B. 300-350 eggs
- C. 400-500 eggs
- D. 750-850 eggs

**Answer: C**



**Watch Video Solution**

**39. Capacitation occurs in**

A. epididymis

B. vas efferens

C. vas deferens

D. female genital tract

**Answer: D**





Watch Video Solution

40. Match the following with correct combination

- |   |               |                            |
|---|---------------|----------------------------|
| A | hyaluronidase | 1. acrosomal reaction      |
| B | corpus luteum | 2. morphogenetic movements |
| C | gastrulation  | 3. progesterone            |
| D | capacitation  | 4. mammary gland           |
| E | colostrum     | 5. sperm activation        |

A. A-5,B-2,C-4,D-1,E-3

B. A-1,B-3,C-2,D-5,E-4

C. A-3,B-2,C-5,D-4,E-1

D. A-1,B-2,C-3,D-4,E-5

**Answer: B**



**Watch Video Solution**

**41.** Find out the correct sequence in embryonic development of animal:

A. cleavage, zygote, fertilization, morula,  
glastula, gastrula

B. Fertilization, zygote, cleavage, morula,  
blastula, gastrula

C. Fertilization, cleavage, morula, zygote,  
blastula, gastrula

D. Fertilization, zygote, blastula, morula,  
cleavage, gastrula

**Answer: B**



**Watch Video Solution**

**42.** What is true about cleavage in fertilized egg in humans?

A. It is meroblastic

B. It is identical to normal mitosis

C. It starts when the egg reaches uterus

D. It starts while the egg is in fallopian

**Answer: D**



**Watch Video Solution**

**43.** Cleavage in mammals is:

A. Discoidal

B. superficial

C. equal holoblastic

D. unequal holoblastic

**Answer: C**



**Watch Video Solution**

**44.** Zona pellucida disintegrates just:

A. just after fertilization

B. before fertilization

C. before cleavage

D. after completion of cleavage

**Answer: D**



**Watch Video Solution**

**45.** The portion of the endometrium that covers the embryo and located between the embryo and the uterine cavity is the:

- A. decidua basalis
- B. decidua umbilicus
- C. decidua capsularis
- D. decidua functionalis

**Answer: C**



**Watch Video Solution**

**46.** Human embryo will be called as a 'foetus' after:

- A. two months
- B. six months
- C. four months
- D. seven months

**Answer: A**



**Watch Video Solution**

**47. Study the following:**

A. Testosterone influences the male secondary sexual characters



B. Gestation period in rabbit is approximately

276 days

C. Bulbo-urethral glands secrete a vaginal

lubricant

D. Placenta secretes estrogen

The correct answer is:

A. C and D

B. A and B

C. A and D

D. B, C and D

**Answer: C**



Watch Video Solution

**48.** In oogenesis, haploid egg is fertilized by sperm at which stage?

A. Ovum

B. Oogonium

C. Primary oocyte

D. Secondary oocyte

**Answer: D**



**49.**  $2n=6$  in a primary spermatocyte which is in metaphase of first meiotic division. What shall be the total number of chromatids in each of the secondary spermatocyte?

- A. 6
- B. 8
- C. 24
- D. 32

**Answer: A**



**Watch Video Solution**

**50.** Some important events in the human female reproductive cycle are given below.

Arrange the events in a proper sequence.

A- Secretion of FSH, B - Growth of corpus luteum,

C- Growth of the follicle and oogenesis, D- Ovulation

E - Sudden increase in the levels of LH.

A. ADCEB

B. BACDE

C. ACEDB

D. CADBE

**Answer: C**



**Watch Video Solution**

**51. The 16 cells stage of the human embryo is**

A. smaller than the fertilized egg

B. same size as the fertilized egg

C. two times of the size of the fertilized  
egg

D. four times the size of the fertilized egg

**Answer: B**



**Watch Video Solution**

**52.** The principal tail piece of human sperm shows the microtubular arrangement of

A.  $7+2$

B.  $9+2$

C.  $11+2$

D.  $13+2$

**Answer: B**



**Watch Video Solution**

**53.** Eggs of placental mammals/human egg are expected to be

A. alecithal

B. polylecithal

C. telolecithal

D. mesolecithal

**Answer: A**



**Watch Video Solution**

**54.** The eggs of some mammals have more yolk. They are



A. eutherians

B. prototherians

C. metatherians

D. aquatic mammals

**Answer: B**



**Watch Video Solution**

**55.** The primary egg membrane of mammalian egg is termed as -

A. Zona pellucida

B. corona radiata

C. shell

D. All of these

**Answer: A**



**Watch Video Solution**

**56.** After a sperm has penetrated an ovum, entry of further sperm is prevented by

A. condensation of yolk

B. formation of pigment coat

C. development of vitelline membrane

D. Development of fertilization membrane

**Answer: D**



**Watch Video Solution**

**57.** Fusion of male and female pronuclei of two conjugate Paramecium is known as

A. apomixis

B. capacitation

C. acrosome reaction

D. amphimixis

**Answer: D**



**Watch Video Solution**

**58.** Cleavage is a unique form of mitotic cell division in that

A. there is no growth of cells

B. the nucleus does not particular

C. no spindle develops to drag  
chromosomes

D. The plasma membranes of daughter  
cells do not separate

**Answer: A**



**Watch Video Solution**

**59.** During cleavage, the cell division is very rapid. The daughter cells do not undergo any growth and the cells thus become gradually smaller in volume. Hence

A. the embryo becomes haploid

B. the embryo grows in volume

C. the embryo becomes smaller in volume

D. there is no increase in the volume of the embryo

**Answer: D**



**Watch Video Solution**

**60.** Which one of the following is incorrect?

A. fertilization follows capacitation

B. cleavage of fertilized ovum results in  
blastula

C. fusiion of sperm and ovum occurs in  
fallopian tube

D. cleavage leads to increase in the mass of  
protoplasm

**Answer: D**



**Watch Video Solution**

**61.** In embryo, cleavage brings about

- A. Increased size
- B. increased DNA content
- C. change in shape and size



D. increased mass of protoplasm

**Answer: B**



**Watch Video Solution**

**62.** A zygote is completely divided into two by a cleavage furrow. The cleavage type is

A. radial

B. equatorial

C. holoblastic

D. meroblastic

**Answer: C**



**Watch Video Solution**

**63.** Solid ball like structure formed after completion of cleavage is

A. morula

B. foetus

C. gastrula

D. blastocyst

**Answer: A**



**Watch Video Solution**

**64.** A morula can be differentiated from blastula in

A. absence of yolk

B. presence of cavity

C. absence of cavity

D. presence of more yolk

**Answer: C**



**Watch Video Solution**

**65.** Morphogenetic movements convert hollow spherical blastula into

A. morula

B. gastrula

C. foetus

D. embryonic disc

**Answer: B**



**Watch Video Solution**

**66.** The best definition of the process of gastrulation is that it is a process where the

A. blastocoel is formed

B. zygote gets converted to larva

C. cells move the occupy their definite position

D. simple layeres blastula becomes two layeres

**Answer: C**



**Watch Video Solution**

**67. Notochord develops from**

A. ectoderm

B. endoderm

C. mesoderm

D. All of these

**Answer: C**



**Watch Video Solution**

**68.** Extra embryonic membrane of the mammals embryo are derived from

A. trophoblast

B. follicle cells

C. endodermal cells

D. inner cell mass

**Answer: A**



**Watch Video Solution**

**69.** Mammalian foetus is directly surrounded  
by

A. yolk sac cavity



B. amniotic cavity

C. allantoic cavity

D. primary digestive cavity

**Answer: B**



**Watch Video Solution**

**70.** The fluid released from the vagina just prior to childbirth is

A. amniotic fluid

B. baby's accumulated urine

C. mother's plasma from umbilical cord

D. baby's plasma form its foetal circulation

**Answer: A**



**Watch Video Solution**

**71.** Blood flowing through umbilical cord of mammalian embryo is

A. 100% foetal

B. 100% maternal

C. 50% maternal and 50% foetal

D. 75% maternal and 25% foetal

**Answer: A**



**Watch Video Solution**

**72.** Drugs causing embryo malformations during pregnancy are called

A. nicotine

B. sedatives

C. teratogens

D. tranquillizer

**Answer: C**



**Watch Video Solution**

**73.** The chemical substances released by activated spermatozoa that acts on the ground substances of the follicle cells is known as

A. relaxin

B. teratogen

C. progesterone

D. hyaluronidase

**Answer: D**



**Watch Video Solution**

**74.** Choose the incorrect statement from the following:

A. In birds and mammals internal fertilisation takes place

B. colostrum contains antibodies and nutrients

C. Polyspermy in mammals is prevented by the chemical changes in the sperm surface

D. In the human female implantation occurs almost seven days after fertilisation.

**Answer: C**



**Watch Video Solution**

**75.** Spot the odd one out from the following structures with reference to the male reproductive system :

- A. Rete testis
- B. Epididymis
- C. Vasa efferentia
- D. Isthmus

**Answer: D**



**Watch Video Solution**

**76.** Seminal plasma, the fluid part of semen, is contributed by.

(i) Seminal vesicle (ii) Prostate (iii) Urethra (iv)

Bulbourethral gland

A. Bulbourethral gland

B. i,ii and iv

C. ii,iii and iv



D. *i* and iv

**Answer: B**



**Watch Video Solution**

**77.** Spermiation is the process of the release of sperms from:

A. Seminiferous tubules

B. Vas deferens

C. Epididymis

D. prostate gland

**Answer: A**



**Watch Video Solution**

**78.** Mature Graafian follicle is generally present in the ovary of a healthy human female around  
stet:

A. 5-8 day of menstrual cycle

B. 11-17 day of menstrual cycle

C. 18-23 day of menstrual cycle

D. 24-28 day of menstrual cycle

**Answer: B**



**Watch Video Solution**

**79.** Acrosomal reaction of the sperm occurs

due to :

A. its contact with zona pellucida of the

ova

B. reactions within the uterine environment of the female

C. Reactions within the epididymal environment of the male

D. androgens produced in the uterus

**Answer: A**



**Watch Video Solution**

**80.** The immature male germ cell undergoes division to produce sperms by the process of spermatogenesis. Choose the correct one with reference to above.

A. Spermatogonia have 46 chromosomes and always undergo meiotic cell division

B. Primary spermatocytes divide by mitotic cell division

C. Secondary spermatocytes have 23 chromosomes and undergo second

meiotic division

D. Spermatozoa are transformed into spermatids

**Answer: C**



**Watch Video Solution**

**81.** Match between the following representing parts of the sperm and their functions and

choose the correct option.

Col. A	Col. B
(a) Head	(i) Enzymes
(b) Middle piece	(ii) Sperm motility
(c) Acrosome	(iii) Energy
(d) Tail	(iv) Genetic material

A. A-ii,B-iv,C-i,D-iii

B. A-iv,B-iii,C-i,D-ii

C. A-iv,B-i,C-ii,D-iii

D. A-ii,B-i,C-iii,D-iv

**Answer: B**



**Watch Video Solution**

82. Match the following and choose the correct options:

- |                     |   |
|---------------------|---|
| (a) Trophoblast     | (i) Embedding of blastocyst in the endometrium              |
| (b) Cleavage        | (ii) Group of cells that would differentiate as embryo      |
| (c) Inner cell mass | (iii) Outer layer of blastocyst attached to the endometrium |
| (d) Implantation    | (iv) Mitotic division of zygote                             |

A. A-ii,B-*i*,C-iii,D-iv

B. A-iii,B-iv,C-ii,D-*i*

C. A-*i*,B-iv,C-ii,D-iii

D. A-ii,B-iv,C-iii,D-*i*

**Answer: B**





Watch Video Solution

**83.** The vas deferens receives duct from the seminal vesicle and opens into urethra as :

- A. epididymis
- B. ejaculatory duct
- C. efferent ductule
- D. ureter

**Answer: B**



**84.** Morula is a developmental stage:

- A. between the zygote and blastocyst
- B. between the blastocyst and gastrula
- C. after the implantation
- D. between implantation and parturition

**Answer: A**



**85.** The cellular cover of the ovum at ovulation is

A. corona radiata

B. zona radiata

C. zona pellucida

D. chorion

**Answer: A**



**Watch Video Solution**

**86.** Identify the odd one from the following:

A. Labia minora

B. Fimbriae

C. Infundibulum

D. Isthmus

**Answer: A**



**Watch Video Solution**

**87.** Which of the following statements about the female reproductive system are true (+) or false (-)?

1. Both FSH and LH are necessary for ovulation to take place.

2. Oestrogen tends to inhibit the production of FSH by the anterior pituitary gland.

3. Fertilization of the ovum by the spermatozoon normally takes place in the uterus.

4. Progesterone production is largely under the control of LH

5. Throughout the part of the menstrual cycle that follows ovulation, there is a slight rise in body temperature.

A. + ± + +

B. - - + + +

C. - ± ±

D. + ± - -

**Answer: A**



**Watch Video Solution**

**88.** Go through the following statements

(i) The secondary oocyte undergoes meiosis II which proceeds only till metaphase until a sperm enters it.

(ii) Ovulation occurs about 36-38 hours after the start of LH surge at midcycle.

(iii) In humans, it takes about 74 hours to form a mature sperm from a primitive germ cells.

(iv). About 70% of the human ejaculate is contributed by bulbourethral glands

Which of these are correct?

A. (i) & (iii)

B. (ii) & (iii)

C. (i) & (ii)

D. all are correct

**Answer: C**



**Watch Video Solution**

**89.** What would happen if vasa deferentia of man are cut?

A. Sperms are non-nucleate



B. spermatogenesis does not occur

C. Semen is without sperms

D. Sperms are nonmotile

**Answer: C**



**Watch Video Solution**

**90.** Go through the following statement

(i) Relaxin produced by the ovary, facilitates delivery of the foetus by softening the connective tissue of pubic symphysis and

relaxing the pelvic ligaments and joints.

(ii) By the end of fifth month of pregnancy, the foetus develops limbs and digits.

(iii) Thalidomide is a teratogenic drug which causes a condition called phocomelia.

(iv). Although the levels of prolactin are high during pregnancy, milk secretion does not occur because of the high oestrogen and progesterone levels which make the breast unresponsive to the prolactin.

Which of these are correct?

A. i,ii and iii

B. *i*, iii and iv

C. ii, iii and iv

D. all are correct

**Answer: B**



**Watch Video Solution**

**91.** identify X and Y in the diagram related to human reproductive system. Choose the

correct option:-



A. X-Seminal vesicle Y-Epididymis

B. X-prostate Y-seminal

C. X-prostate Y-vasdeferens


D. X-Bulbourethral gland Y-Seminal vesicle

**Answer: B**



**Watch Video Solution**

**92.** Three cells are labelled A,B and C. select the option which correctly tells the cells which produce androgen.



A. Only A

B. Only B

C. A and C



D. All three

**Answer: A**



**Watch Video Solution**

**93.** Given is the stage of a growing embryo and different regions of the fallopian tube marked 1,2,3,4.



Where do you think the given stage of embryo will be seen in the fallopian tube

A. 4

B. 3

C. 2

D. 1

**Answer: A**



**Watch Video Solution**

**94.** Find the correct match regarding human foetal development

A. Month of Pregnancy- End of 4<sup>th</sup> month,

Event occurring-Eyelashes appear



B. Month of pregnancy-End of 3<sup>rd</sup> month,

Event occurring-Movements of foetus

C. Month of pregnancy-End of 5<sup>th</sup> month,

Event occurring-Hair on head

D. Month of pregnancy-End of 2<sup>nd</sup> month-

Event occurring-organ system develop

**Answer: C**



**Watch Video Solution**

**95.** Trace a sperm cell from the structure where it is produced to fertilization of the egg

1. Seminiferous tubules

2. Vas deferens

3. Uterus

4. Fallopian tube

5. Vagina

6. Epididymis

7. Urethra

A. 6,1,2,7,5,3,4

B. 1,6,2,7,5,3,4

C. 1,6,2,7,5,4,3

D. 1,2,6,7,5,3,4

**Answer: B**



**Watch Video Solution**

**96.** How are the time of ovulation and the onset of menstruation related in the human menstrual cycle?

- A. Both are triggered by high luteinizing hormone "spikes" (sharp increase in concentration).
- B. Ovulation occurs approximately 7 days after the first day of menstruation
- C. Ovulation occurs approximately 14 days before the first day of menstruation.
- D. All of the above

**Answer: C**



**Watch Video Solution**

97. Tick the correct statement

A. Proximal and distal centrioles in the middle piece of sperm help anchor the flagellum

B. whereas proximal centriole is non functional, distal centriole acts as basal body for the flagellum of sperm

C. The functions of proximal and distal centrioles are not known

D. Proximal centriole forms the spindle fibres during cleavage and the distal centriole anchors the flagellum of sperm

**Answer: D**



**Watch Video Solution**

**98.** The intra-testicular genital duct system does not involve

A. Rete testis

B. Ductuli efference


C. Tubuli recti

D. Ductus deference

**Answer: D**



**Watch Video Solution**



99.

In the above diagram identify X, Y and Z with the sequence of number of appropriate hormones given below:-

(i) progesteron

(ii) LH

(iii). HCG

(iv). Estradiol



(v). FSH

(vi). GnRH

A. (iii), (iv), (ii)

B. (vi), (ii), (v)

C. (i),(iv), (vi)

D. (vi), (v), (ii)

**Answer: B**



**Watch Video Solution**

**100.** Which of the following hormones is not secreted by human placenta?

A. Relaxin

B. Lactogen

C. Gonadotropin

D. Oxytocin

**Answer: D**



**Watch Video Solution**

**101.** In humans, the first polar body formed during oogenesis has

(i) 46 chromosomes

(ii) 23 chromosomes

(iii) 46 chromatids

(iv). 23 chromatids

A. (ii) and (iv)

B. (ii) and (iii)

C. (i) and (iii)

D. (i) and (iv)

**Answer: B**



**Watch Video Solution**

**102.** Go through the following statements

- (i) Androgens are produced by the interstitial cells
- (ii) Sertoli cells give rise to germ cells
- (iii). Secretions of male accessory glands constitute the seminal plasma which is rich in fructose, calcium and certain enzymes
- (iv) The presence or absence of hymen is a very

reliable indicator of virginity.

Which of these are correct?

A. *i* & *iii*

B. *iii* & *iv*

C. *i*, *iii* & *iv*

D. all are correct

**Answer: A**



**Watch Video Solution**

**103.** Go through the following statements:-

(i). Primary spermatocytes undergo mitotic divisions to produce secondary spermatocytes

(ii). Sperms released from the seminiferous tubules are fully mature and motile

(iii) The head of sperm possesses many mitochondria which produce energy for the movement of tail

(iv) The human male ejaculates about 20-30 million sperms during a coitus

which of these are correct?

A. (i), (ii) & (iii)

B. (ii), (iii) & (iv)

C. (i), (iii) & (iv)

D. All are wrong

**Answer: D**



**Watch Video Solution**

**104.** Go through the following statements

(i) Mammary glands are modified sweat glands and each mammary gland consists of 15-25

lobules of the compound tubulo-alveolar type.

(ii) The tertiary follicle is characterised by a fluid filled cavity called antrum

(iii) Both LH and FSH attain a peak level towards the middle of the menstrual cycle

(iv). In oogenesis, both first and second meiotic divisions are unequal

Which of these are correct?

A. (ii) & (iii)

B. (i), (iii) & (iv)

C. (i),(ii) & (iv)



D. All are correct

**Answer: D**



**Watch Video Solution**

**105.** Go through the following statement:-

(i) In both pre-pubertal and post-meno-pausal females, there are low levels of female sex hormones and high levels of gonadotrophins.

(ii) There is no bleeding in an oestrous cycle as the broken endometrium is absorbed.

(iii) Oestrogen is mainly secreted by the granulosa cells and progesterone mainly by the theca cells

(iv) Some of the menopausal symptoms can be reversed HRT wherein a small dose of gonadotrophins is given to the patient.

Which of these are correct?

A. (ii) & (iii)

B. (i), (ii), (iv)

C. (i), (ii) & (iii)

D. All are correct

**Answer: A**



**Watch Video Solution**

**106.** Go through the following statements

- (i) HCG from placenta stimulates the sertoli cells of the male foetus to produce testosterone and is thus indirectly involved in the development of male external genitalia
- (ii) Sertoli cells secrete a protein called inhibin, which suppresses FSH synthesis.
- (iii) Humans have haemo-endothelial type of

placenta.

(iv) oxytocin stimulates the placenta to secrete prostaglandins which in turn stimulate more contractions of uterus.

Which of these correct?

A. (i) & (iv)

B. (ii) & (iv)

C. (i), (ii) & (iv)

D. All are correct

**Answer: B**



**107.** Match list-I (cell type) with list-II (characteristic) and select the correct answer using the codes given below the lists

List-I	List-II
A. Spermatogonium	1. Does not divide.
B. Spermatocyte	2. Divides mitotically
C. Spermatid	3. Divides meiotically
D. Sertoli cell	4. Divides both mitotically and meiotically
	5. Nourishes other cell types

A. A-5,B-3,C-1,D-4

B. A-1,B-3,C-4,D-5

C. A-3,B-2,C-4,D-1

D. A-2,B-3,C-1,D-5

**Answer: D**



**Watch Video Solution**

**108.** Both corpus lutea and macula lutea are

A. found in human ovaries

B. a source of hormones

C. characterized by a yellow colour

D. Contributory I maintaining pregnancy

**Answer: C**



**Watch Video Solution**

**109.** The phase of menstrual cycle in humans that lasts for 7-8 days, is

A. Follicular phase

B. Ovulatory phase

C. Luteal phase

## D. Menstruation

**Answer: A**



**Watch Video Solution**

**110.** Which one of the following statements with regard to embryonic development in humans is correct?

A. Cleavage divisions bring about considerable increase in the mass of



protoplasm.

B. In the second cleavage division, one of the two blastomeres usually divides a little sooner than the second

C. With more cleavage divisions, the resultant blastomeres become larger and larger.

D. Cleavage division results in a hollow ball of cells called morula

**Answer: B**



Watch Video Solution

**111.** Assertion: Holoblastic cleavage with almost equal sized blastomeres is a characteristic of placental animals.

Reason: Eggs of most mammals, including humans, are of centrolecithal type.

A. If both assertion and reason are true  
and the reason is the correct

explanation of the assertion, then mark

a.

B. if both assertion and reason are true but the reason is not the correct explanation of assertion, then mark b.

C. if assertion is true statement but reason is false, then mark c.

D. If both assertion and reason are false statements, then mark d.

**Answer: C**



Watch Video Solution

112. Which one of the following events is correctly matched with the time period in a normal menstrual cycle?

- A. Release of egg, 5<sup>th</sup> day
- B. Endometrium regenerates: 5-10 days
- C. Endometrium secretes nutrients for implantation: 11-18 days
- D. Rise in progesterone level: 1-15 days

**Answer: B**



**Watch Video Solution**

**113.** Which of the following has the longest gestation period:-

A. Man

B. Cat

C. Dog

D. Elephant

**Answer: D**



**Watch Video Solution**

**114.** Go through the following diagram carefully which of the following option shows the correct labelling and events occurring in that organ:-

A. (i) Ovary-Follicle rupture, beginning of

$1^{st}$  and  $2^{nd}$  meiotic division.

Completion of  $2^{nd}$  meiotic division

(iii) Uterus-Blastocyst formation,  
implantation.

B. (i) Ovary-Follicle rupture, completion of  
 $2^{nd}$  meiotic division

(ii) Fallopian tube-Fertilisation,  
Blastocystformation.

(iii) Uterus-Implantation

C. (i) Ovary-Follicle maturation, Beginning  
of  $1^{st}$  and  $2^{nd}$  meiotic division

(ii) Fallopian tube-Fertilisation,

Completion of 2<sup>nd</sup> meiotic division

(iii) Uterus-Beginning of cleavage,  
Implantation.

D. (i) Ovary-Follicle maturation completion  
of 2<sup>nd</sup> meiotic division.

(ii) Fallopian tube-Beginning of cleavage,  
Implantation. ItBrgt (iii) Uterus-  
Fertilisation, blastocyst formation

**Answer: A**



**Watch Video Solution**



**115.** Secretion of GnRH would lead to

A. secretion of testosterone leading to mammary gland development

B. release of prolactin leading to milk production in mammary glands

C. secretion of LH and FSH leading to follicle development

D. All of the above

**Answer: C**



**Watch Video Solution**

**116.** Find the odd one out

A. HIV

B. Trichomoniasis

C. Gonorrhoea

D. typhoid

**Answer: D**



Watch Video Solution

**117.** Assertion: If scrotal sacs removed, testosterone is still found in the body.

Reason: Small amount of androgens is secreted by the adrenal cortex.

- A. If both assertion and reason are true and the reason is the correct explanation of the assertion, then mark
- a.

B. if both assertion and reason are true but the reason is not the correct explanation of assertion, then mark b.

C. if assertion is true statement but reason is false, then mark c.

D. If both assertion and reason are false statements, then mark d.

**Answer: A**



**Watch Video Solution**

**118.** The entry of additional sperms into the ovum is prevented because of:-

- A. release of secretions from the acrosome
- B. changes in the zonapellucida
- C. changes in the corona radiata
- D. contractions and secretions of the fallopian tube

**Answer: B**



**Watch Video Solution**

**119.** Ovulation in the human female normally takes place during the menstrual cycle

A. at the beginning of the proliferative phase

B. at the end of the proliferative phase

C. at the mid secretory phase

D. just before the end of the secretory phase

**Answer: B**





[Watch Video Solution](#)

**120.** Which part of ovary in mammals acts as an endocrine gland after ovulation ?

- A. Stroma
- B. Germinal epithelium
- C. Vitelline membrane
- D. Graffian follicle

**Answer: D**



[Watch Video Solution](#)

**121.** In the human female , menstruation can be referred by the administration of :-

A. combination of FSH and LH

B. combination of estrogen and progesterone

C. FSH only

D. LH only

**Answer: B**





**122.** Compared to a bull, a bullock is docile because of

A. higher levels of cortisone

B. lower levels of blood testosterone

C. lower levels of adrenalin/noradrenalin in its blood

D. higher levels of thyroxin

**Answer: B**



Watch Video Solution

**123.** In humans, at the end of the first meiotic division, the male germ cells differentiate into the

- A. spermatozonia
- B. Primary spermatocytes
- C. secondary spermatocytes
- D. spermatids

**Answer: C**



Watch Video Solution

**124.** Which extra-embryonic membrane in human prevents desiccation of the embryo inside the uterus?

A. Amnion

B. Chorion

C. Allantois

D. Yolk sac

**Answer: A**



Watch Video Solution

**125.** Which one of the following statement is incorrect about menstruation ?

- A. The beginning of the cycle of menstruation is called menarche.
- B. During normal menstruation about 40 ml blood is lost.
- C. The menstrual fluid can easily clot.

D. At menopause | the female, there is especially abrupt increase in gonadotropic hormones

**Answer: C**



**Watch Video Solution**

**126.** In human adult females, oxytocin

A. causes strong uterine contractions during parturition

B. is secreted by anterior pituitary

C. stimulates growth of mammary glands

D. stimulates pituitary to secrete  
vasopressin

**Answer: A**



**Watch Video Solution**

**127.** Which one of the following is the correct matching of the events occurring during menstrual cycle?

A. Development of corpus luteum-secretory phase and increased secretion of progesterone.

B. Menstruation-Breakdown of myometrium and ovum not fertilized.

C. Ovulation-LH and FSH attain peak level and sharp fall in the secretion of progesterone.

D. Proliferative phase- Rapid regeneration of myometrium and maturation of

graafian follicle.

**Answer: A**



**Watch Video Solution**

**128.** The correct sequence of spermatogenetic stages leading to the formation of sperms in a mature human testis is

A. Spermatid-spermatocyte-  
spermatogonia-sperms



B. Spermatogonia-spermatid-spermatocyte-  
sperms

C. spermatocyte-spermatogonia-spermatid-  
sperms

D. Spermatogonia-spermatocyte-spermatid-  
sperms

**Answer: D**



**Watch Video Solution**

**129.** Foetal ejection reflex in human female is induced by:

- A. fully developed foetus and placenta
- B. differentiation of mammary glands
- C. pressure exerted by amniotic fluid
- D. release of oxytocin from pituitary

**Answer: A**



**Watch Video Solution**

**130.** In a regularly cycling human female, which can be the root cause of menstrual failure?

A. Maintenance of high concentration of sex-hormones in the blood stream

B. Retention of well-developed corpus luteum

C. Fertilisation of the ovum

D. Maintenance of the hypertrophical endometrial lining


**Answer: C**



Watch Video Solution

**131.** Given below is a diagrammatic sketch of a portion of human male reproductive system.

Select the correct set of the names of the parts labelled A,B,C,D:-



A. A-Vasdeferens, B-Seminal vesicle, C-Bulbourethral gland, D-Prostate

B. A,Ureter, B-Seminal vesicle, C-Prostate, D-Bulbourethral gland

C. A-Ureter, B-Prostate, C-Seminal vesicle, D-Bulbourethral gland.

D. A-Vasdeferens, B-Seminal vesicle, C-prostate, D-Bulbourethral gland

**Answer: D**



**Watch Video Solution**

**132.** A change in the amount of yolk and its distribution in the egg will affect :-

- A. number of blastomeres produced
- B. fertilization
- C. Formation of zygote
- D. pattern of cleavage

**Answer: D**



**Watch Video Solution**

**133.** Seminal plasma in humans is rich in

A. glucose and certain enzymes but has no calcium

B. fructose and certain enzymes but poor in calcium

C. fructose, calcium and certain enzymes

D. fructose and calcium but has no enzymes

**Answer: C**

---



Watch Video Solution

**134.** Vasa efferentia are the ductules leading from:

- A. epididymis to urethra
- B. testicular lobules to rete testis
- C. rete testis to vas deferens
- D. vas deferens to epididymis

**Answer: C**



Watch Video Solution



**135.** The first movements of the fetus and appearance of hair on its head are usually observed during which month of pregnancy?

- A. Third month
- B. fourth month
- C. fifth month
- D. sixth month

**Answer: C**



**136.** Which is correct about human sperm

A. Acrosome serves no particular function

B. Acrosome has a conical pointed structure used for piercing and penetrating the egg resulting in fertilization

C. The sperm lysins in the acrosome dissolve the egg envelope facilitating fertilization

D. Acrosome serves as a sensory structure

leading the sperm towards the ovum

**Answer: C**



**Watch Video Solution**

**137.** Sertoli cells are found :

A. pancreas and secrete cholecystokinin

B. ovaries and secrete progesterone

C. adrenal cortex and secrete adrenaline

D. seminiferous tubules and provide nutrition to germ cells

**Answer: D**



**Watch Video Solution**

**138.** The part of Fallopian tube closest to the ovary is:

A. ampulla

B. isthmus

C. infundibulum

D. cervix

**Answer: C**



**Watch Video Solution**

**139.** The second maturation division of the mammalian ovum occurs

A. in the graafian follicle following the first maturation division

- B. shortly after ovulation before the ovum makes entry into the fallopian tube
- C. until after the ovum has been penetrated by a sperm
- D. until the nucleus of the sperm has fused with that of the ovum

**Answer: C**



**Watch Video Solution**

**140.** Which one of the following statements about morula humans is correct?

A. It has more cytoplasm and more DNA than an uncleaved zygote

B. It has almost equal quantity of cytoplasm as an uncleaved zygote but much more DNA

C. It has far less cytoplasm as less DNA than in an uncleaved zygote

D. It has more or less equal quantity of cytoplasm and DNA as in uncleaved zygote

**Answer: B**



**Watch Video Solution**

**141.** If for some reason, the vasa efferentia in the human reproductive system get blocked, the gametes will not be transported from



A. testes to epididymis

B. epididymis to vas deference

C. ovary to uterus

D. vagina to uterus

**Answer: A**



**Watch Video Solution**

**142.** The testes in humans are situated outside the scrotum. The pupose served is for


- A. maintaining the scrotal temperature lower than the internal body temperature
- B. escaping any possible compression by the vescceral organs
- C. providing more space for the growth of epididymis
- D. providing a secodary sexual feature for exhibiting the male sex.

**Answer: A**



Watch Video Solution

143. The figure given below depicts a diagrammatic sectional view of the female reproduction system of humans. Which one set of three parts out of I-VI have been correctly identified?



A. (II) Endometrium, (III) Infundibulum, (IV)

fimbriae

B. (III) infundibulum, (IV) fimbriae, (V) cervix

C. (IV) oviducal funnel, (V) uterus, (VI) cervix

D. (I) perimetrium, (II) Myometrium, (III)

Fallopian

**Answer: B**



**Watch Video Solution**

**144.** What is correct to say about the hormone action in humans ?

A. Secretion of thymosins is stimulated with aging

B. In females, FSH first binds with specific receptors on ovarian cell membrane

C. FSH stimulates the secretion of estrogen and progesterone

D. Glucagon is secreted by  $\hat{\alpha}$ -cells of islets of langerhans and stimulates glycogenolysis

**Answer: B**



**Watch Video Solution**

**145.** In a normal pregnant woman, the amount of total gonadotropin activity was assessed. The result expected was

A. High level of circulating HCG to stimulate  
endometrial thickening

B. High levels of FSH and LH in uterus to  
stimulate endometrial thickening

C. High level of circulating HCG to  
stimulate estrogen and progesterone  
synthesis

D. High level of circulating FSH and LH in  
the uterus to stimulate implantation of  
the embryo

**Answer: C**



**Watch Video Solution**

**146.** Signals for parturition originate from

- A. Oxytocin released from maternal pituitary
- B. placenta only
- C. Fully developed foetus only



D. Both placenta as well as fully developed foetus

**Answer: D**



**Watch Video Solution**

**147.** The leydig cells as found in the human body are the secretory source of

A. intestinal mucus

B. glucagon

C. androgens

D. progesterone

**Answer: C**



**Watch Video Solution**

**148.** Which one of the following statements is not true with respect to viability of mammalian sperm?

- A. Survival of sperm depends on the pH of the medium and is more active in alkaline medium
- B. Viability of sperm is determined by its motility
- C. sperms must be concentrated in a thick suspension
- D. Sperm is viable for only up to 24 hours

**Answer: D**



**Watch Video Solution**

**149.** Signals from the fully developed foetus and placenta ultimately lead to parturition which requires the release of

- A. Estrogen from placenta
- B. oxytocin from maternal pituitary
- C. oxytocin from foetal pituitary
- D. Relaxin from placenta

**Answer: B**



**Watch Video Solution**

**150.** In human female the blastocyst:

A. Forms placenta even before

implantation

B. gets implanted into uterus 3 days after

ovulation

C. Gets nutrition from uterine endometrial

secretion only after implantation

D. gets implanted in endometrium by the trophoblast cells

**Answer: D**



**Watch Video Solution**

**151.** What happens during fertilization in humans after many sperms reach close to the ovum?

- A. Secretions of acrosome helps one sperm enter cytoplasm of ovum through zonapellucida
- B. All sperms except the one nearest to the ovum lose their tails
- C. Cells of corona radiata trap all the sperms except one
- D. Only two sperms nearest the ovum penetrate zonapellucida

**Answer: A**



Watch Video Solution

**152.** About which day in a normal human menstrual cycle does rapid secretion of LH (popularly called LH-surge) normally occurs?

A. 14th day

B. 20th day

C. 5th day

D. 11th day

**Answer: A**





Watch Video Solution

**153.** The secretory phase in the human menstrual cycle is also called:

A. luteal phase and lasts for about 13 days

follicular phase and lasts for about 13 days

B. luteal phase and lasts for about 6 days

C. follicular phase and lasting for about 6 days

D. Follicular phase lasts for about 13 days

**Answer: A**



**Watch Video Solution**

**154.** Identify the human development stage shown below as well as the related right place of its occurrence in a normal pregnant woman, and select the right option for the two

together.



**Options :**

<b>Developmental stage</b>	<b>Site of occurrence</b>
(1) Blastocyst	Uterine wall
(2) 8-celled morula	Starting point of Fallopian tube
(3) Late morula	Middle part of Fallopian tube
(4) Blastula	End part of Fallopian tube

A. Developmental stage-Blastocyst, Site of occurrence-Uterine wall

- B. Developmental stage-8celled morula,  
Site of occurrence-Starting point of  
fallopian tube
- C. Developmental stage-Late morula, Site of  
occurrence-Middle part of Fallopian tube
- D. Developmental stage-Blastula, Site of  
occurrence-End part of fallopian tube

**Answer: A**



**Watch Video Solution**

**155.** Bartholin's glands are situated:

A. On the sides of head of some amphibians

B. At the reduced tail end of birds

C. On either side of vagina in humans

D. On either side of vas deferens in humans

**Answer: C**



**Watch Video Solution**

**156.** Placenta in human beings is formed by :

A. amnion

B. chorion

C. allantois

D. chorion and allantois

**Answer: B**



**Watch Video Solution**

157. Which gland secretes alkaline mucus in urethra to neutralise the acidity of urine?

- A. Prostrate gland
- B. Cowper's gland
- C. Seminal vesicles
- D. preputial glands

**Answer: B**



**Watch Video Solution**

**158.** A temporary endocrine gland in humans is

A. Islets of langerhans

B. Pineal body

C. Corpus luteum

D. corpora allata

**Answer: C**



**Watch Video Solution**

**159.** Braxton Hicks contractions occur during



- A. passage of food through alimentary canal
- B. peristaltic movements
- C. pregnancy
- D. lactation

**Answer: C**



**Watch Video Solution**

**160.** Correct sequence of human embryonic development is

A. Blastocoel-gastrocoel-neural crest-

notochord

B. gastrocoal-blastocoel-notochord-neural

crest

C. gastrocoel-blastocoel-neural crest-

notochord

D. blastocoel-neural

crest-gastrocoel-

notochord

**Answer: A**



**Watch Video Solution**

**161.** Onset of menstrual cycle in female anthropoid primates is

A. Puberty

B. Menarche

C. Menopause

D. Menstruation

**Answer: B**



**Watch Video Solution**

**162.** Thick yellow, high protein fluid produced by mammary glands of a woman during first 2-3 days after child birth is

A. Meconium

B. Hymen

C. Cumulus oophorus

D. Colostrum

**Answer: D**



**Watch Video Solution**

**163.** Which hormones is produced in women during pregnancy?

A. human chorionic gonadotropin (hcG)

B. human placental lactogen (hpL)

C. Relaxin

D. All of above

**Answer: D**



**Watch Video Solution**

**164.** Menstrual phase is followed by

A. luteal phase and lasts for about 13 days

follicular phase and lasts for about 13

days

B. follicular phase

C. Fertilization, cleavage, morula, zygote,

blastula, gastrula

D. Implantation

**Answer: B**



**Watch Video Solution**

**165.** Which layer of uterus undergoes cyclic changes during menstrual cycle

A. Perimetrium

B. Myometrium

C. Endometrium

D. All the above

**Answer: C**



**Watch Video Solution**



**166.** Which type of germ cells contain 23 chromosomes

- A. Spermatogonia
- B. Secondary spermatocytes
- C. Primary spermatocytes
- D. None of the above

**Answer: B**



**Watch Video Solution**

**167.** In human females, menstrual cycle stops around 50 years of age. It is termed as

A. Menarche

B. Diapause

C. Menopause

D. None of the above

**Answer: C**



**Watch Video Solution**

**168.** Which hormone is mainly secreted by corpus luteum ?

A. Luteinizing hormone

B. Estrogen

C. Follicle stimulating hormone

D. Progesterone

**Answer: D**



**Watch Video Solution**

**169.** The internal cavity commonly formed by cell division prior to gastrulation is the

A. Enteron

B. Blastopore

C. Blastocoel

D. Coelom

**Answer: C**



**Watch Video Solution**

**170.** Number of autosomes in human primary spermatocyte is

A. 46

B. 44

C. 23

D. 22

**Answer: B**



**Watch Video Solution**

**171.** Which of the following organs is devoid of glands?

A. Uterus

B. Vagina

C. Vulva

D. Oviduct

**Answer: B**



**Watch Video Solution**

172. Primary spermatocyte differs from spermatogonium in

A. Number of chromosomes

B. size and volume

C. DNA content

D. Size of chromosomes

**Answer: B**



**Watch Video Solution**

**173.** In human, cleavage divisions are:

- A. slow and synchronous
- B. Fast and synchronous
- C. Slow and asynchronous
- D. Fast and asynchronous

**Answer: C**



**Watch Video Solution**



**174.** Vertebrate brain differentiates from

A. endoderm

B. mesoderm

C. ectoderm

D. blastoderm

**Answer: C**



**Watch Video Solution**

**175.** Which one is not a placental hormone

A. HCG

B. HPL

C. Progesterone

D. melatonin

**Answer: D**



**Watch Video Solution**

**176.** In human females, the ovarian cycle begins when the:

A. levels of oestrogen reach their maximum

B. hypothalamus stimulates the anterior pituitary to increase its output of FSH and LH

C. level of progesterone drops precipitously

D. hypothalamus increases its release of  
FSH and LH

**Answer: C**



**Watch Video Solution**

**177.** Sperm of animal species *a* cannot fertilise  
ovum of species *b* because

A. Fertilizin of A and antifertilizin of B are  
not compatible

B. antifertilizin of A and fertilizin of B are not compatible

C. fertilizin of A and B are not compatible.

D. antifertilizin of A and B are not compatible

**Answer: B**



**Watch Video Solution**

**178.** In spermatogenesis, reduction division of chromosomes occurs during conversion of

A. spermatogonia to primary

spermatocytes

B. primary spermatocytes to secondary

spermatocytes

C. secondary spermatocytes to spermatids

D. spermatids to sperms

**Answer: B**



[Watch Video Solution](#)

**179.** Presence of which of the following hormones in the urine confirms pregnancy?

A. Progesterone

B. oestrogen

C. Human chorionic gonadotropin

D. prolactin

**Answer: C**



[Watch Video Solution](#)

**180.** What is the correct sequence of sperm formation?

A. Spermatogonia, spermatocyt, spermatid, spermatozoa,

B. spermatogonia, spermatocyte, spermatid, spermatozoa

C. spermatid, spermatocyte, spermatogonia, Spermatozoa



D. spermatogonia,

spermatocyte,

spermatozoa, spermatid

**Answer: B**



**Watch Video Solution**

**181.** Which one of the following is not the function of placenta?It:

A. facilitates removal of carbon dioxide and waste material from embryo

B. Secretes oxytocin during parturition.

C. Facilitates supply of oxygen and nutrients to embryo.

D. secretes estrogen.

**Answer: B**



**Watch Video Solution**

**182.** Product of sexual reproduction generally generates

A. new genetic combination leading to variation

B. Large biomass

C. Longer viability of seeds

D. Prolonged dormancy

**Answer: A**



**Watch Video Solution**

**183.** Menstrual flow occurs due to lack of:

A. Oxytocin released from maternal  
pituitary

B. Vasopressin

C. Progesterone

D. FSH

**Answer: C**



**Watch Video Solution**

**184.** The shared terminal duct of the reproductive and urinary system in the human male is

A. vasa efferentia

B. urethra

C. ureter

D. vas deferens

**Answer: B**



**Watch Video Solution**

**185.** Select the correct option describing gonadotropin activity in a normal pregnant female.

A. High level of hCG stimulates the thickening of endometrium

B. High level of FSH and LH stimulates the thickening of endometrium

C. High level of FSH and LH facilitate implantation of the embryo

D. High level of hCG stimulates the synthesis of estrogen and progesterone

**Answer: D**



**Watch Video Solution**

**186.** Which of these is not an important component of initiation of parturition in humans?

A. Synthesis of prostaglandins

B. release of oxytocin

C. release of prolactin

D. increase in estrogen and progesterone  
ratio

**Answer: C**



**Watch Video Solution**

**187.** Capacitation refers to changes in the

A. ovum before fertilization



B. ovum after fertilization

C. sperm after fertilization

D. sperm before fertilization

**Answer: D**



**Watch Video Solution**

**188.** Hysterectomy is surgical removal of

A. prostate gland

B. vas-deferens

C. Mammary glands

D. Uterus

**Answer: D**



**Watch Video Solution**

**189.** Which of the following cells during gametogenesis is normally diploid?

A. Spermatid

B. Spermatogonia

C. Secondary polar body

D. Primary polar body

**Answer: B**



**Watch Video Solution**

**190.** A childless couple can be assisted to have a child through a technique called GIFT. The full form of this technique is:-

A. Gamete inseminated fallopian transfer

B. Gamete intra fallopian transfer

C. Gamete internal fertilization and transfer

D. Germ cell internal fallopian transfer

**Answer: B**



**Watch Video Solution**

**191.** Ectopic pregnancies are referred to as

A. pregnancies with genetic abnormality

B. implantation of embryo at site other than uterus

C. implantation of defective embryo in the uterus

D. pregnancies terminated due to hormonal imbalance

**Answer: B**



**Watch Video Solution**

**192.** Which of the following events is not associated with ovulation in human female?

- A. Decrease in estradiol
- B. Full development of graafian follicle
- C. release of secondary oocyte
- D. LH surge

**Answer: A**



**Watch Video Solution**

**193.** Which of the following layers in an antral follicle is acellular?

- A. Granulosa
- B. Theca interna
- C. Stroma
- D. Zona pellucida

**Answer: D**



**Watch Video Solution**

**194.** In human females, meiosis-II is not completed until

A. puberty

B. fertilization

C. uterine implantation

D. birth

**Answer: B**



**Watch Video Solution**



**195.** Which of the following approaches does not give the defined action of contraceptive?

A. Intra uterine devices-Increase phagocytosis of sperms, suppress sperm motility and fertilizing capacity of sperms

B. Hormonal contraceptives- Prevent-retard entry of sperms, prevent ovulation & fertilization

C. Vasectomy-Prevents spermatogenesis

D. Barrier methods-prevent fertilization

**Answer: C**



**Watch Video Solution**

**196.** Fertilization in humans is practically feasible only if

A. the ovum and sperms are transported simultaneously to ampullary-isthmic junction of the fallopian tube

B. the ovum and sperms are transported simultaneously to ampullary-isthmic junction of the cervix

C. the sperms are transported into cervix within 48 hrs of release of ovum in uterus

D. the sperms are transported into vagina just after the release of ovum in fallopian tube.

**Answer: A**



Watch Video Solution

**197.** Identify the correct statement on 'inhibin'

- A. is produced by granulosa cells in ovary
- B. Is produced by granulose cells in ovary  
and inhibits the secretion of LH.
- C. Is produced by nurse cells in testes and  
inhibits the secretion of LH.
- D. Inhibits the secretion of H, FSH and  
prolactin

**Answer: A**



**Watch Video Solution**

**198.** Select the incorrect statement

A. LH triggers ovulation in ovary.

B. LH and FSH decrease gradually during the follicular phase.

C. LH triggers secretion of androgens from the Leydig cells.

D. FSH stimulates the sertoli cells which help in spermiogenesis

**Answer: B**



**Watch Video Solution**

**199.** Changes in HbRH pulse frequency in females is controlled by circulating levels of

A. Estrogen and inhibin

B. Progesterone only

C. progesterone and inhibin

D. Estrogen and progesterone

**Answer: D**



**Watch Video Solution**

**200.** Embryo with more than 16 blastomeres formed due to in vitro fertilization is transferred into

A. Uterus

B. Fallopian tube

C. Fimbriae

D. Cervix

**Answer: A**



**Watch Video Solution**

**201.** Which of the following depicts the correct pathway of transport of sperms?



A. Rete testis → Efferent ductules → Epididymis → vas deferens.

B. Rete testis → Epididymis → efferent ductules → vas deferens

C. Rete testis → vas deferens → efferent ductules → Epididymis

D. Efferent ductules → Rete testis → Vas deferens → Epididymis

**Answer: A**



**Watch Video Solution**

**202.** Match column-I with column-II and select the correct option using the codes given below:-

column - I	Column - II
1. Mons pubi	(i) Embryo formation
2. Antrum	(ii) Sperm
3. Trophoctoderm	(iii) Female external genitalia
4. Nebenkern	(iv) Graafian follicle

A. 1-iii,2-iv,3-ii,4-i

B. 1-iii,2-iv,3-i,4-ii

C. 1-iii,2-i,3-iv,4-ii

D. 1-i,2-iv,3-iii,4-ii

**Answer: B**



**Watch Video Solution**

**203.** Several hormones like hCG, hPL, estrogen, progesterone are produced by

A. Ovary

B. Placenta

C. Fallopian tube

## D. Pituitary

**Answer: B**



**Watch Video Solution**

**204.** Select the correct route for the passage of sperms in male frogs:-

A. Testes → Bidder's canal → kidney →

vasa efferentia → Urinogenital duct

→ cloaca

B. Testes → vasa efferentia → kidney →  
seminal vesicle → urinogenital duct →  
cloaca

C. Testes → vasa efferentia → Bidder's  
canal → ureter → cloaca

D. Testes → vasa efferentia → kidney →  
bidder's canal → urinogenital duct →  
cloaca

**Answer: D**



**Watch Video Solution**

**205.** A temporary endocrine gland in humans is

- A. Pineal gland
- B. corpus cardiacum
- C. Corpus luteum
- D. corpus allatum

**Answer: C**



**Watch Video Solution**

**206.** GnRH, a hypothalamic hormone, needed in reproduction, acts on

A. anterior pituitary gland and stimulates secretion of LH and oxytocin.

B. Anterior pituitary gland and stimulates secretion of LH and FSH

C. posterior pituitary gland and stimulates secretion of oxytocin and FSH

D. posterior pituitary gland and stimulates secretion of LH and relaxin.

**Answer: B**



**Watch Video Solution**

**207.** Capacitation of sperms occurs in

A. Rete testis

B. Epididymis

C. Vas deferens



D. Female reproductive tract

**Answer: D**



**Watch Video Solution**