

BIOLOGY

BOOKS - TRUEMAN BIOLOGY

PLANT KINGDOM

Multiple Choice Question

1. A full mature plant body, undifferentiated into root, stem and leaf is called.

A. thallus

B. coenocytic

- C. hyphae
- D. callus

Answer: A



Watch Video Solution

- 2. Algae are included in
 - A. Thallophytes
 - B. spermatophytes
 - C. Embryophytes
 - D. Tracheophytes

Answer: A



- 3. Algae are characterised by
 - A. thalloid plant body
 - B. ph/ycobilins
 - C. unicellular sex organs
 - D. mechanical tissue.

Answer: A



Watch Video Solution

4. Algae resemble fungi in the presence of similar

B. cell wall
C. similar reserve food
D. similar mode of nutrition
Answer: A
Watch Video Solution
5. Green algae are ancestors of angiosperms/land plants
because
A. both have vascular bundles
B. both have starch as reserve food

A. reproductive structure

- C. both have jacketed sex organs
- D. all of the above

Answer: B



- **6.** Classification of Algae into 11 classes was made by Fritsch on the basis of flagellation, pigmentation and type of reserve food. Out of this, the main criteria used in algal classification grouping of algae is
 - A. chemical composition of cell wall
 - B. type of pigmentation
 - C. nature of food storage

D. shape and colony formation.

Answer: B



Watch Video Solution

7. Study of Algae is called

A. phycology

B. phytology

C. mycology

D. phenology

Answer: A



8. Choose the correct statement.

A. Algae show embryo stage

B. Algae show only haplontic life cycle

C. Algae are plants as they possess cell wall, chl a and starch food.

D. all of the above

Answer: C



- 9. All algae possess
 - A. chl a and b
 - B. chl a, carotenes and phycobilins
 - C. chl b and carotenes
 - D. chl a and carotenoids

Answer: D



Watch Video Solution

10. Largest acellular, green,marine alga ,popularly called umbrella plant is

A. Ulva B. Acetabularia C. Spirogyra D. Volvox **Answer: B Watch Video Solution** 11. Kelps are A. fresh water algae

B. marine green algae

C. large marine parenchymatous brown algae

D. large marine parenchymatous and algae.

Answer: C



Watch Video Solution

12. Red eye spot (stigma) is meant for

A. movement

B. vision

C. photoreception

D. photosynthesis

Answer: C



13. Red snow is caused by

A. stigma

B. chloroplast

C. hypnospore

D. aplanospore

Answer: C



Watch Video Solution

14. The diploid stage is represented by one celled zygote only in

- A. Chlamydomonas
- B. Spirogyra
- C. both 1 and 2
- D. Funaria

Answer: C



- 15. Meiosis in spirogyra/Chlamydomonas/Ulothrix occurs
 - A. as the zygospore germinates
 - B. during conjugation
 - C. during palmella formation

D. during formation of gametes

Answer: A



Watch Video Solution

16. The product of conjugation of Spirogyra is called

A. zoospore

B. zygospore

C. endospore

D. akinete

Answer: B



17. Asexual spores with flagella in algae are called

- A. zoospore
- B. zygospores
- C. aplanospores
- D. hypnospores

Answer: A



Watch Video Solution

18. In spriogyra lateral conjugation takes place in the cells

of

B. two cells of different filaments C. among three filaments D. within one cell of a filament **Answer: B Watch Video Solution 19.** What is the shape of chloroplast in Spirogyra? A. spiral, bank like B. cup like C. stellate like

A. two adjacent cells of same filament

D. U-like

Answer: A



Watch Video Solution

20. Gametes of Spirogyra are

- A. biflagellated
- B. uniflagellated
- C. non-flagellated
- D. diploid

Answer: C



21. Choose the correct statement in Spirogyra.

A. Filaments showing scalariform conjugation are always dioecious

B. Filaments showing lateral conjugation are always monoecious

C. Filaments showing lateral conjugation may be monoecious

D. Asexual reproduction occurs by zoospores.

Answer: B



22. Agar-agar,an important solidyfying agent used in culture medium, is obtained from

- A. red algae
- B. green algae
- C. kelps
- D. yellow algae

Answer: A



23. Red algae show maximum photosynthesis in blue green light. Their main pigment to trap this light is

- A. chlorophyll
- B. phycoerythrin
- C. chl a and xanthophyll
- D. carotenes.

Answer: B



Watch Video Solution

24. Main pigment in phaeophyceae (Brown algae) is

- A. phycocyanin
- B. phycoerythrin
- C. fucoxanthin
- D. chlorophyll b

Answer: C



- **25.** Food reserve in Rhodophyta is :-
 - A. floridean starch
 - B. laminarian starch
 - C. animal starch

D. none of these

Answer: A



Watch Video Solution

26. Reserve food in Phaeophyceae is

- A. laminarin and manitol
- B. glycogen and sorbitol
- C. floridean starch
- D. glucose and starch.

Answer: A



27. Sieved septa/trumpet hyphae occur in

- A. kelps
- B. All green algae
- C. All red algae
- D. Marine algae.

Answer: A



Watch Video Solution

28. Agar,Alginic acid,Carragenin and Funori in sea weeds are

A. proteins
B. phycocolloids
C. acids
D. fats
Answer: B
Watch Video Solution
29. Red algae are similar to blue-green algae in
possession of
A. similar reserve food
A. Sillillar Teserve 1000
B. nucleus

- C. phycobilins
- D. gas-vascuoles

Answer: C



Watch Video Solution

30. Red algae are able to grow deep in sea as

- A. they can trap blue green light of short wave length in deept layer of water
- B. they have clorophyll b to trap blue green light in deept layer of water
- C. Both 1 and 2

D. both wrong.

Answer: A



Watch Video Solution

31. Irish Moss is

- A. Chondrus (a red alga)
- B. Ulva (green alga)
- C. Porphyra (red alga)
- D. Gelidium (red alga)

Answer: A



32. A colourless parasitic red algae is

- A. Harveyella
- B. Batrachospermum
- C. Porphyra (red alga)
- D. Cephaleuros

Answer: A



Watch Video Solution

33. Red rust of tea is caused by

A. Cephaleuros virescence

B. Puccinia graminis

- C. Harveyella
- D. Chlamydomonas gametos

Answer: A



Watch Video Solution

34. A student collected an alga and found that its cells contain both chl a and chl d as well as phycoerythrin but no chl b and flagella. The alga bolongs to

A. Rhodophyceae

- B. Phaeophyceae
- C. Bacillariophyceae
- D. Chlorophyceae

Answer: A



Watch Video Solution

35. Allophycocyanin is found in

- A. Chlorophyceae
- B. Rhodophyceae
- C. Both 1 and 2
- D. All marine algae

Answer: B



Watch Video Solution

- **36.** The algal forms are mostly marine in
 - A. Chlorophyceae
 - B. Phaeophyceae
 - C. Ascomycetes
 - D. All of these

Answer: B



37. Algae and fungi are characterised by the possession of

- A. mitospores
- B. chloroplast
- C. multicelluarl jacketed sex organs
- D. unicellular jacketed sex organs

Answer: A



Watch Video Solution

38. What is common in Thallphytes, Bryophytes and pteridophytes?

B. Presence of conductive system C. Presence of cones D. Absence of vascular tissue. **Answer: A Watch Video Solution 39.** Most of the green algae are A. fresh water B. marine C. terrestrial

A. Dependence on water

D. epiphytic/Epizoic

Answer: A



Watch Video Solution

40. Chlorophyll common between phaeophyceae and bacillariophyceae but absent in rhodophyceae is

- A. chlorophyll a
- B. chlorophyll b
- C. chlorophyll c
- D. chlorophyll e

Answer: C

- **41.** The pigment found only in some algae
 - A. chlorophyll a
 - B. xanthophyll
 - C. carotene
 - D. phycobilins

Answer: D



42. The structure in algae helping in perennation to tide over drought are

- A. zoospores, hypnospores and aplanospores
- B. zoospores, akinetes, zygospores
- C. aplanospores, hypnospores and zygospores.
- D. none of the above

Answer: C



Watch Video Solution

43. A single thylakoid per granum found in the chloroplast of

B. green algae
C. brown algae
D. yellow algae
Answer: A
Watch Video Solution
44. Sexual reproduction where a smaller and motile male
gamete fuses with larger and motile female gamete
known as
A. isogamy

A. red algae

B. anisogamy C. oogamy D. heterogamy **Answer: B Watch Video Solution** 45. Life cycle in Chlamydomonas / Ulothrix/ Spirogya is A. haplontic B. haplobiontic C. diplontic D. diplobiontic

Answer: A



Watch Video Solution

- 46. In Spirogyra, ladder like structure is formed in
 - A. scalariform conjugation
 - B. lateral conjugation
 - C. direct conjugation
 - D. parthenogenesis

Answer: A



- B. jacketed and unicellular
- C. nonjacketed and multicellular
- D. non-jacketed and unicellular

Answer: D



- **49.** Alginic acid is present in the cell wall of
 - A. bacillariophyceae
 - B. Euglena
 - C. Laminaria
 - D. Diatoms

Answer: C



Watch Video Solution

50. In phaeophyceae, thylakoids are found in group of

- A. 3's
- B. 4's
- C. 5's
- D. 6's

Answer: A



51.	Plants	lacking	seed	and	vascular	tissue	but	forming
sp	ores an	d embry	o are					

- A. bryophytes
- B. pteridophytes
- C. angiosperms
- D. gymnosperms

Answer: A



- 52. Bryophytes grow in moist, humid places as
 - A. they have no cuticle

- B. require water for fertilization
- C. they do not have roots
- D. all of the above

Answer: D



- 53. Bryophytes don't attain much height because
 - A. they do not have vascular supply
 - B. they do not have roots and mechanical tissue
 - C. they require water for transport of sperms
 - D. all of the above.

Answer: D



Watch Video Solution

- **54.** Bryophytes can be distinguished from algae/fungi/thallophytes because they have
 - A. thallus, haploid, gametophytic body
 - B. chloroplast
 - C. no conducting tissue
 - D. sterile jacket of cells around multicellular sex organs.

Answer: D

55. Bryophytes show

- A. asexual reproduction & zygotic meiosis
- B. asexual reproduction & sporic meiosis
- C. no asexual reproduction but sporic meiosis
- D. gametophytic dominance and sporic meiosis

Answer: D



- A. rhizoids are always unicellular but roots are multicellular
- B. rhizoids arise in gametophytes but roots in sporophytes
- C. roots are branched but rhizoids are always unbranched.
- D. all of the above.

Answer: B



Watch Video Solution

57. Predominant and largest gaemtophyte is of

A. Selaginella
B. Pinus
C. Moss
D. Rice
Answer: C
Watch Video Solution
58. Which one of the following is true moss
58. Which one of the following is true moss A. Cord moss
A. Cord moss

D. All of the above.

Answer: A



Watch Video Solution

59. The spores produced in capsule in moss, germinate to form haploid structure called

- A. leafy gametophyte
- B. protonema
- C. prothallus
- D. peristome

Answer: B

60. Sex organ of Funaria are

- A. projected and sessile
- B. projected and stalked
- C. embedded and stalked
- D. embedded and sessile

Answer: B



- A. it is rootless
- B. it is without vascular supply
- C. its sporophyte is parasite and attached to gametophyte
- D. it grows near water.

Answer: C



- **62.** Sporophyte of Funaria is
 - A. Parasite on gametophyte as it is non green
 - B. semiparasite on gametophyte as it is green

- C. independent of gametophyte as it is green
- D. none of the above

Answer: B



Watch Video Solution

- **63.** Archegonium of Funaria secretes mucilage rich in
 - A. sucrose
 - B. glucose
 - C. malic acid
 - D. citric acid

Answer: A



64. Polytrichum has

- A. heart shaped prothallus
- B. foot,seta and capsule
- C. vascular bundles
- D. mitospores

Answer: B



Watch Video Solution

65. The sporophyte of Funaria begins development within

B. archegonium C. capsule D. protonema **Answer: B Watch Video Solution 66.** Sexual reproduction in bryophytes is A. oogamous B. isogamous C. anisogamous

A. antheridium

D. hologamy

Answer: A



Watch Video Solution

- 67. Bryophytes called amphibians of plant king-dom as
 - A. they inhabits damp places
 - B. they have tracheids
 - C. their sex organs are multicellular and jacketed
 - D. their reproductive phase requires water.

Answer: D



68. Bryophytes differ from pteridophytes in

A. having mitospores

B. the absence of vascular tissue system

C. lacking embryo stage

D. all of the stage

Answer: B



Watch Video Solution

69. A liver wort is

- A. a parasite causing infection of liver
- B. a kind of virus infecting the liver
- C. a flowering plant for treating liver disorders
- D. a plant without differentiation into roots, stem and leaves.

Answer: D



Watch Video Solution

70. Branched rhizoids and leafy gametophytes are characteristics of

A. all bryophytes

B. some bryophytes like mossesC. algaeD. liverworts.

Answer: B



Watch Video Solution

71. In the life cycle of Funaria, spores are beginning of the generation

- A. gametophyte
- B. sporophyte
- C. peristome

D. both 1 and 2

Answer: A



Watch Video Solution

72. Sphagnum is also called 'Peat Moss' because it

A. it grown in acidic marshes (bogs) and helps in peat

formation

B. it is found in peat

C. it retains water

D. it is fossilized quickly.

Answer: A

73. Moss plant is a

A. gametophyte

B. sporophyte

C. sometimes gametophyte and sometimes

sporophyte

D. predominantly gametophyte with sporophyte

attached to it.

Answer: D



74. Bryophytes are close to pteridophytes in having

A. multicellualr jacketed sex organs

B. flagellated oosphere

C. mitospores in life cycle

D. all of the above

Answer: A



Watch Video Solution

75. To which would you assign a plant that has xylem and phloem, produces meiospores, embryo but lacks seeds and flowers?

B. Pteridophytes C. Tracheophytes D. Spermatophytes. **Answer: B Watch Video Solution** 76. Vascular cryptogams and botanical snakes of plant kingdom are A. pteridophytes B. tracheophytes

A. Bryophytes

- C. angiosperms
- D. spermatophytes

Answer: A



Watch Video Solution

77. Tracheophyte include

- A. bryophytes, pteridophytes and spermatophytes
- B. bryophytes,pteridophytes,gymnoperms and angiosperms
- C. pteridophytes and spermatophytes (gymnosperms and monocotyledons)

D. pteridophytes and spermatophytes (gymnosperms and angiosperms)

Answer: D



Watch Video Solution

78. Gametophytic and sporophytic phases are independent in

- A. bryophyte
- B. pteridophyte
- C. gymnosperms
- D. all of the above.

Answer: B



79. Which one of the following is considered important in the development of seed habit

Or

Seloginella has the character of evolutionary importance.

That character is

- A. seed habit and heterosporous nature
- B. viviparity
- C. rhizophore
- D. spikes.

Answer: A



Watch Video Solution

- **80.** Heterosporous pteridophytes produce
 - A. dioecious gametophytes
 - B. monoecious gametophytes
 - C. homothallic gametophytes
 - D. hermaphrodite gametophytes

Answer: A



A. root
B. leaf
C. organ sui generis
D. stem
Answer: D
Watch Video Solution
82. Adiantum, Pteris/ferns are
A. heterosporous

81. Rhizophore in Selaginella is

- B. homosporous
- C. homogametic
- D. isomerous

Answer: B



- **83.** The main plant body/dominant plant in life cycle of a fern/Pleridophytes is
 - A. gametophyte (x)
 - B. sporophyte (2x)
 - C. apogamous

D. aposporous

Answer: B



Watch Video Solution

84. Stem in ferns is underground, monopodial root stock and called

- A. rhizome
- B. corm
- C. bulb
- D. sucker

Answer: A

85. Leaves of ferns are

- A. fronds with reticulate venation
- B. haploid
- C. pinnately compound and grow by apical meristem
- D. all of the above.

Answer: C



A. fronds with reticulate venation	
B. leaf bases	
C. ramenta	
D. root stock	
Answer: C	
Watch Video Solution	
87. Maiden Hair Fern is	
A. Adiantum	
B. Cheilanthis	
C. Pteris	

D. Dryopteris

Answer: A



Watch Video Solution

88. Sporangia bearing leaves in Pteridophytes are called

A. sporophylls

B. megaphylls

C. fern

D. sorus

Answer: A



89. Fern sperms (antherozoids) are

A. multicillated or multi flagellated and spirally coiled

B. biciliated and coiled

C. uninucleated and uniflagellated

D. multiciliated sickle shaped

Answer: A



Watch Video Solution

90. What represents the gametophytic generation in pteridophytes?

B. Heart shaped prothallus			
C. Indusium			
D. Stomium			
Answer: B			
Allswei. D			
Watch Video Solution			
91. A collection of sporangia attached to placenta and			
covered over by indusium is known as			
A. sorus			
B. sporophyll			

A. Main plant body

- C. ramenta
- D. spike

Answer: A



Watch Video Solution

- **92.** Archegonium of Funaria secretes mucilage rich in
 - A. sucrose
 - B. malic acid
 - C. glucose
 - D. citric acid

Answer: B

93. A thick, multicelluar covering called indusium in ferns over sori is meant for

- A. protection
- B. producing spores
- C. help in dispersal
- D. has no role

Answer: A



94. Antheridium in ferns is

A. multicellular and jacketed

B. stalked, hemispherical and borne ventrally in posterior part between rhizoids

C. its jacket hasd 32-48 cells

D. all of the above.

Answer: A



Watch Video Solution

95. Life cycle in ferns is

A. haplontic B. diplohaplontic C. haplo-haplontic D. diplontic **Answer: B Watch Video Solution** 96. First land plant was represented by A. Ferns **B.** Grasses C. Gymnosperms

D. Bryophytes

Answer: A



Watch Video Solution

97. Spike moss is

- A. Selaginella
- B. Funaria
- C. Lycopodium
- D. Adiantum

Answer: A



98. Ferns are characterised by

- A. cambium
- B. xylem vessles
- C. young leaves with circinate ptyxis and presence of

ramenta

D. all of the above.

Answer: C



99. Ferns resemble with mosses in one respect

- A. both are embryophytes
- B. both are tracheophytes
- C. both are sporophytes
- D. both are aquatic

Answer: A



Watch Video Solution

100. Heteromorphic alternation of generation is found in

A. Spirogyra

- B. Mucor
- C. Selaginella
- D. All of these.

Answer: C



- 101. In ferns, circinate ptyxis (vernation) is defined as
 - A. arrangement of leaf gaps in stems
 - B. coiling of young leaves
 - C. arrangement of sori on leaves
 - D. attachement of ramenta on young parts.

Answer: B



102. If sperms of moss and fern are put together near the archegonium of fern, only the sperms of fern find entry into archeogonium, the reason being that

- A. sperms of moss are killed by larger sperms of fern
- B. archegonia of fern secretes a toxoid to kill moss of fern.
- C. archegonia of fern secrete mucilage rich in malic acid to attract sperms of fern only

D. sperms of moss are less motile and find difficulty in entering archegonium.

Answer: C



Watch Video Solution

103. What may be the possible advantage occurring out of the presence of antheridia and archegonia on the underside of a fern prothallus?

- A. They are protected from direct rays of the sun.
- B. Capillary water accumulates on the underside of prothallus between its lower surface and the soil

surface sex organs projecting in this water can be readily fertilised by the cilliated sperms which are chemotactically attracted by the archegonia

- C. The sex organs remain protected from wind
- D. Nutrients manufactured by the green prothallus can radily seep downwards to the sex organs due to the action of gravity.

Answer: B



Watch Video Solution

104. Walking fern is named so as

- A. it knows walking
- B. it is dispersed through walking of animals in forests
- C. its spores are able to move with wind
- D. it speads and propagates vegetatively by its leaf tips.

Answer: D



Watch Video Solution

105. A water fern capable of fixing atmospheric nitrogen and used as biofertilizer.

A. Azolla
B. Nostoc
C. Adiantum
D. Spirulina
Answer: A Watch Video Solution
106. Which group of gymnosperms is close to
angiosperms?
A. Gnetales
B. Ginkgoales

- C. Cycadales
- D. Confierales

Answer: A



Watch Video Solution

107. Gymnosperms are different from angiosperms in

- A. absence of seeds
- B. absence of ovary
- C. absence of ovule
- D. absence of sieve cells.

Answer: B

108. Which one constitutes the dominant vegetation in colder regions?

- A. Dicots
- B. Gymnosperms
- C. Monocots
- D. Tracheophytes

Answer: B



109. Which one of the following has not changed for the last several thousand years

- A. Pinus excelsa
- B. Ginkgo biloba
- C. Welwitschia
- D. Sequola

Answer: B



Watch Video Solution

110. Largest ovules, largest male and female gametes and tallest trees are found among

- A. angiosperms
- B. tree ferns and some monocots
- C. gymnosperms
- D. dicots

Answer: C



- 111. Siphonogamy in traceophytes
 - A. eliminates dependence on water
 - B. brings pollen grains together
 - C. carries spores

D. protects embryo

Answer: A



Watch Video Solution

112. Cycas is a

A. living fossil

B. fossil

C. endangered species

D. exotic species.

Answer: A



113. The cortex and pith of stem Cycas serves as a source of 'Sago' which is a

A. proteins

B. cellulose

C. starch

D. mixture of starch and protein.

Answer: C



- A. ovule and female cone
 - B. largest male gametophyte
- C. sperm
- D. all of the above.

Answer: C



Watch Video Solution

115. Cycas is

- A. hermaphrodite
- B. dioecious
- C. monoecious

D. none of these

Answer: B



Watch Video Solution

- **116.** Organised female cone is absent in
 - A. Ephedra
 - B. Pinus
 - C. Cycas
 - D. None of these

Answer: C



117. Cycas differs from Pteris in having

A. vessels and tracheids

B. motile sperms

C. pollen tube

D. archegonia

Answer: B



Watch Video Solution

118. Coralloid roots of Cycas are useful in

B. fixation C. absorption and fixation of nitrogen D. respiration from air. **Answer: C Watch Video Solution** 119. Coralloid roots of Cycas possess a symbiotic alga A. Aulosira B. Anabaena C. Oscillatoria

A. absorption of water

D. Chlorella

Answer: B



Watch Video Solution

- 120. Phloem of gymnosperms differ from angiosperms in
 - A. having no companion cells
 - B. having no sieve cells
 - C. having phloem fibre
 - D. having phloem parenchyma

Answer: A



121. Foliage leaves (needles) in Pinus are brone by

A. dwarf shoots

B. long shoots

C. both 1 and 2

D. female strobilus

Answer: A



Watch Video Solution

122. At the time of dehiscence winged pollen grain of

Pinus is

A. 1 celled B. 4 celled C. 3 celled D. 2 celled **Answer: B Watch Video Solution** 123. Simplest and highly reduced archegonium is found in A. Pinus B. Fern C. Bryophytes

D. Liver worts/mosses

Answer: A



Watch Video Solution

124. Microsporangia of Cycas occur over microsporophyll

A. abaxial

B. adaxial

C. lateral

D. margin.

Answer: A



125. Number of generations present in Pinus seed are

A. 2

B. 3

C. 1

D. 4

Answer: B



Watch Video Solution

126. Ploidy in wing of pollen grain and wing of seed of

Pinus is

- A. x in pollen grain and 2x in seed
- B. 2x in pollens and 2x in seed
- C. being an outgrowth, chromosomes are absent
- D. 1/2 x in wing of pollens and polyploidy in seed.

Answer: A



- **127.** The phenomenon of sulphur shower in pine forest is due to
 - A. dispersal of winged pollens of Pinus
 - B. dispersal of winged seeds of Pinus

- C. bursting of thrid year female cone of Pinus
- D. none of the above

Answer: A



Watch Video Solution

128. Which is homologous?

- A. Leaves of Moss and Selaginella
- B. Roots of ferns and moss
- C. Endosperm of Pinus and prothallus of Adianlum
- D. Pinus endosperm & endosperm of Maize

Answer: C

129. In pinus

- A. seeds and ovules are winged
- B. endosperm is triploid
- C. fruits and flowers absent and seeds are naked
- D. all of the above

Answer: C



130. Chilgoza' is a fruit, obtained from a gymnosperm which is

- A. Pinus Roxburghii
- B. Pinus geradiana
- C. Cycas revoluta
- D. Abies balsamiana

Answer: B



Watch Video Solution

131. Ephedrine obtained from the stem of Ephedra is given to cure

- A. gastric disorders
- B. respiratory disorders
- C. arhritis
- D. all of the above.

Answer: B



Watch Video Solution

132. No Gymnosperms is

- A. annual and herbaceous
- B. perennial and herbaceous
- C. xerophyte and woody

D. tree and shrubby

Answer: A



Watch Video Solution

133. Vessels are absent in the xylem of

A. angiosperms

B. monocots

C. gymnosperms excluding gnetales

D. pteridophytes excluding ferns.

Answer: C



134. One of the main evolutionary features of the alternation of generations from algae to flowering plants is.

- A. gradual elaboration of gametophyte
- B. gradual elaboration of sporophyte
- C. elinination of sporophytic tissue
- D. eliminination of gametophyte.

Answer: B



135. In which of the following the angiosperms resemble the Gymnosperms


- A. nature of endosperm
- B. presence of vessels
- C. siphonogamy
- D. double fertilization.

Answer: C



Watch Video Solution

136. In gymnosperms and angiosperms the pollen tube carries the male gamete to the site of fertilization and



- B. vascular cryptogams are all homosporous
- C. Archegoniates are all homosporous
- D. Spermatophytes are all heterosporous.

Answer: D



Watch Video Solution

138. Cyas has 2 cotyledons yet it is classified as Gymnosperm and not as dicot plant because

- A. it resembles data palm/monocot in appearance.
- B. it has compound leaves
- C. its ovules are naked

D. it bears megasporophylls

Answer: C



Watch Video Solution

139. In Cycas pollen grains shed at

- A. 2 celled stage
- B. 3 celled stage
- C. 4 celled stage
- D. 5 celled stage

Answer: B



140. What is true for Cycas?

- A. Cycas is dioecious
- B. Ovules without integument
- C. Largest female cone
- D. All of the above.

Answer: A



Watch Video Solution

141. Which is incorrect about Cycas

- A. Its xylem has vessels B. Its young leaves show circinate vernation C. It does not have organised female flower D. motile sperms **Answer: A Watch Video Solution**
- **142.** Cycas reproduces vegetatively by
 - A. sporophylis
 - B. bulbils
 - C. fragmentation

D. rhizome

Answer: B



Watch Video Solution

- 143. Which one is common between Funaria and Pinus?
 - A. No fruits are produced
 - B. No seeds are produced
 - C. Pollen tube is formed
 - D. Antheridia and archegonia are present

Answer: A



144. Among the following, which does not belong to sporophyte generation in Pinus?

- A. Long shoot
- B. Dwarf shoot
- C. roots are branched but rhizoids are always unbranched.
- D. Endosperm

Answer: D



145. Needle in Pinus represents
A. root
B. scale leaf
C. follage leaf
D. spur
Answer: C



Watch Video Solution

146. The male cone of Pinus is formed of

Or

In pinus male cone bears is large number of

- A. antheridia
- B. megasporophylls/green
- C. microsporophylls
- D. ligules

Answer: C



Watch Video Solution

147. Seeds of Pinus are

- A. adaxial, endospermic and polycotyledonous
- B. abaxial, monocotyledonous and endospermic
- C. hypogeal, endospermic and monocot

D. monocotyledonous, epigeal.

Answer: A



Watch Video Solution

148. Which of the following is not a correct match?

A. Maiden hari fern: Ginkgo

B. Bog Moss : Sphagnum

C. Cord/green moss: Funaria

D. Walking fern : Adiantum

Answer: A



149. Following are are given some trends in the evolution of plants

- A. origin of vascular system
- B. origin of rhizoids
- C. origin of seeds
- D. origin of flowers.

Answer: C



150. Characteristic of Angiosperms which distinguish them from gymnosperms

- A. presence of fruits and flowers
- B. double fertilization and triploid endosperm formed after double fertilization
- C. companion cells in phloem and vessels in xylem
- D. all of the above.

Answer: D



151. Beginning with germination of a moss spore, what is the sequence of structures that develop after germination?

I. embryo II. Gametes III. Sporophyte IV. Potonema V. gametophore

- A. IV,V,III,I,II
- B. V,IV,III,II,I
- C. III,IV,V,II,I
- D. IV,V,II,I,III

Answer: D



152. Angiosperms resemble gymnosperms in

- A. presence of companion cells
- B. type of fertilization
- C. presence of ovules
- D. nature of endosperm

Answer: C



Watch Video Solution

153. An example of a marine angiosperm thriving in shallow seas is

A. Zostera

- B. Wolffia
- C. Pistia
- D. Rhizophora

Answer: A



Watch Video Solution

154. Which one of the following is considered important in the development of seed habit

Or

Seloginella has the character of evolutionary importance.

That character is

A. Haplontic life cycle

- B. Fee-living gametophyte
- C. Dependent sporophyte
- D. Heterospory

Answer: D



Watch Video Solution

155. Which of the following is grouped under phanerogams?

- A. Algae show embryo stage
- B. Bryophytes
- C. Gymnosperms

D. Pteridophytes

Answer: C



Watch Video Solution

156. The stamen in angiosperms is homologous to which part in gymnosperm and pteridophytes?

- A. Microsporangium
- B. Micorosporophyll
- C. Megasporophyll
- D. Male gametophyte

Answer: B

157. The megasporophyll of vascular plants is analogous to which structure in angiosperms

- A. stamen
- B. ovule
- C. carpel
- D. leaf

Answer: C



158. To which group of plants does the Banyan tree belong

- A. angiosperms
- B. Gymnosperms
- C. cryptogams
- D. phaeophyta

Answer: A



Watch Video Solution

159. Mark the national tree

- A. Mangifera indica (Mango tree)
- B. Ficus benghalensis (banyan tree)
- C. Ficus religiosa (pipal tree)
- D. Azadirachta indica (Neem tree)

Answer: B



Watch Video Solution

160. Of the four widely known systems of classification one remains less phylogenetic and more natural, which is of

A. Engler & Prantl

- B. Benthan & Hooker
- C. Rendle
- D. Hutchinson

Answer: B



Watch Video Solution

161. Genera Plantarum was written by

- A. Benthan and Hooker
- B. Hutchinson
- C. Rendle
- D. Engler and Prantl

Answer: A



Watch Video Solution

- **162.** Dryopteris differs from Funaria in having
 - A. an independent gametophyte
 - B. an independent sporophyte
 - C. swimming antherozoids
 - D. archegonia

Answer: B



163. Angiosperms have dominated the land flora primarily by their

- A. nature of self pollination
- B. property of producing large number of seeds
- C. domestication by man
- D. power of adaptability in diverse habitat.

Answer: D



Watch Video Solution

164. In which of the following group, all the organisms are homosporous


- A. Funaria, Salvinia, Dryopteris
- B. Azolla, Selaginella, Anthoceros
- C. Salvinia, Cedrus, Funaria
- D. Sphagnum, Pteris, Adiantum

Answer: D



Watch Video Solution

165. See the given life cycle pattern and choose the option which it correctly represents.



- A. Haplontic
- B. Diplontic
- C. Haplo-diplontic
- D. 1 or 3

Answer: B



166. Read the following statements carefully

(i) Protonema is found in mosses and is absent in liverworts.

(ii) In Marchantia, the sporophyte is divided into the foot, seta and capsule

(iii) Archegonia are partially embedded in pteridophytes
(iv) Gymnosperms do not possess vessels in xylem except
in some gnetophytes.

A. i,ii,iii

B. ii,iv

C. i,iii,iv

D. All are correct.

Answer: C



Watch Video Solution

167. The algal group which most biologists believe to have led to evolution of land plants is

- A. phaeophyta
- B. Rhodophyta
- C. Chlorophyta
- D. Chrysophyta

Answer: C



168. Phycocolloids are found in the cell walls of

A. Diatoms

B. Spirogyra

C. Red and brown algae

D. All algae

Answer: C



Watch Video Solution

169. Which one of the following is not characteristic of all divisions of vascular plants?

- A. An alternation of generations
- B. The development of seeds
- C. Differentiation into roots, stems and leaves
- D. Xylem and phloem for transporting materials between roots and leaves.

Answer: B



- 170. Red algae differs from green and brown algae in
 - A. No chlorophyll a
 - B. No differentiated cells

- C. No phycocyanin within their cells
- D. No flagellated stages in their life cycles

Answer: D



Watch Video Solution

171. Bryophytes resemble resemble algae in the following aspects

- A. Differentiation of the plant body into root,stem and heterotrophic mode of nutrition
- B. Thallus-like plant body, lack of vascular tissue: absence of root: and autotrophic mode of nutrition

- C. Thallus-like plant body, presence of roots, and autotrophic mode of nutrition
- D. Filamentous body: presence of vascular tissue: and autotrophic mode of nutrition

Answer: B



- 172. Funaria differs from Pteridium in the absence of
 - A. Root
 - B. Seed
 - C. Archegonia

D. Embryo

Answer: A



Watch Video Solution

173. Prothallus is

- A. Gametophyte,monoecious,autotroph present in pteridophytes
- B. Gametophyte,monoecious,autotroph found in bryophytes
- C. Sporophyte, dioecious, heterotroph found in bryophytes

D. Gametophyte,dioecious,heterotroph present in pteridophytes

Answer: A



Watch Video Solution

174. In gymnosperms how many male gametes are produced by each pollen grain

A. 4

B. 3

C. 2

D. 1

Answer: C



175. Which arrangements of the organisms represents a rank ordering (based on size/importance) from dominant gametophyte on the left to dominant sporophyte on the right?

- A. Polytrichum, Cycas, Equisetum
- B. Lycopodium, Zostera, Riccia
- C. Sphagnum, Pteridium, Cedrus
- D. Pinus, Selaginella, Marchantia

Answer: C



Watch Video Solution

176. Red algae do not have

- A. zoospores, hypnospores and aplanospores
- B. Cellulose in cell wall
- C. Floridean starch
- D. Sexual reproduction.

Answer: A



177. Select the wrong statement

- A. Gymnosperm do not have antheridia
- B. Cycas and ferns have multiflagellate gametes
- C. Selaginella is a heterosporous non seed plant
- D. Pinus tree is dioecious

Answer: D



Watch Video Solution

178. Ovule of Pinus can be called

A. Female gametophyte

- B. Male gametophyte
- C. Megasporangium
- D. Integumented megasoporangium



- **179.** Conifers differ from grasses in the
 - A. Production of seeds from ovules
 - B. Lack of xylem tracheids
 - C. Absence of pollen tubes
 - D. Formation of endosperm before fertilization.



- 180. "Ordines Anomali" of Benthan and Hooker includes
 - A. Seed plants showing abnormal forms of growth and development
 - B. Plants represented only in fossil state.
 - C. Plants described in the literature but which

 Bentham and Hooker did not see in original
 - D. A few orders which could not be placed satisfactorily in the classification.



- 181. Which of the following is a wrong combination?
 - A. Haploid endosperm, archegonia present but antheridium absent-Gymnosperms
 - B. Triploid endosperm and 7-celled female gametophyte called embryo sac-Angiosperms
 - C. Embryo stage absent, reproduction by mitospores-Algae

D. Gametophyte independent, biflagellate gametes and reproduction by accessory spores-Mosses.

Answer: D



Watch Video Solution

182. Green algae are also included among the items carried by astronauts in a space ship Which of the following are the reasons for this?

1. These being very simple plants, are amenable to a variety of experiments. 2. These may provide nourishment to the persons on the space ship. 3. These act as a source of oxygen.

Select the correct answer using the codes given below.

- A. 1,2 and 3
- B. 1 and 3
- C. 1 and 2
- D. 2 and 3



Watch Video Solution

183. Consider the following pigments 1. Chlorophyll a 2.

Chlorophyll b 3. Chlorophyll c 4. Carotenes 5. Biliproteins

6. Xanthophylls

The characteristic photosynthetic pigments present in Chlorophyceae would include

- A. 1,2,3 and 4
- B. 1,3,4, and 5
- C. 2,3,5 and 6
- D. 1,2,4 and 6



Watch Video Solution

184. Which one of the following pairs is not correctly matched?

A. Red Sea: Trichodesmium erythreum, a

cyanobacterium

- B. Red Tides: Rhodymenia, a red algae
- C. Red Snow: Chalmydomonas nivalis, a green algae
- D. Red Rust of Tea: Cephaleuros virescens, a green algae

Answer: B



Watch Video Solution

185. Consider the following statements

- 1. In pteridophytes, the phloem lacks companion cells.
- 2. The xylem lacks vessels in majority of pteridophytes.
- 3. Heterosporous pteridophytes have monoecious gametophytes.

4. Water is essential for fertilization in pteridophytes.

Which of these statements are correct?

- A. 1,2,3 and 4
- B. 2,3 and 4
- C. 1 and 3
- D. 1,2 and 4

Answer: D



Watch Video Solution

186. Which one of the statements is correct?

A. Leaves of conifers are perennial

- B. All vascular plants have Sieve Tube Members
- C. The red and edible flesh of strawberry is the carpel
- D. Conifers have motile gametes.



- **187.** Which of the following structures in Pinus are haploid
 - A. Megaspore,endosperm and embryo
 - B. Microspore, megaspore and endosperm

- C. Megaspore, integument and root
- D. Microspore, leaf and endosperm

Answer: B



Watch Video Solution

- **188.** More than one answer may be correct : Marchantia polymorpha
- 1. Is dioecious
- 2. Possesses antheridiophores and archegoniophores
- 3. Lacks foot and seta in its sporophyte
- 4. Is heterosporous.

A. 1 and 2

- B. 3 and 4
- C. 1,2 and 3
- D. 2 and 3



Watch Video Solution

189. Consider the following statements

- 1. The thylakoids of blue-green algae are arranged singly in the stroma of chloroplasts.
- 2. In blue-green algae, the thylkoids are not only the sites for photosynthesis but also for respiration

Which of the statements given above is/are correct?

- A. 1 only
- B. 2 only
- C. both 1 and 2
- D. Neither 1 nor 2

Answer: B



Watch Video Solution

190. In gymnosperms the pollen chamber represents

A. A cavity in the ovule in which pollen grains are stored after pollination

- B. An opening in the megagametophyte through which the pollen tube approaches the egg
- C. The microsporangium in which pollen grains develop
- D. A cell in the pollen grain in which the sperms are formed.



- **191.** Bryophytes can be compared with amphibians since both are
- (i) Autotrophic (ii) Dependent on water for reproduction

(iii) Without endoskeleton (iv) Devoid of impervious body surface (v) With separate sexes

A. ii,iv

B. i,ii,iv

C. ii,iii,iv

D. i,iii,iv

Answer: A



Watch Video Solution

192. In which of the following groups all the organisms contain chlorophyll c

- A. Ectocarpus, Fucus, Sargassum, Laminaria
- B. Macrocystis, Batrachospermum, Polysliphonia, Characium
- C. Chlamydomonas, Asterocystis, Ulothrix, Gelidium
- D. Harveyella, Chondrus, Volvox, Chlorella



- **193.** In which of the following group all the orgaisms contain phycobilins?
 - A. Anabaena, Gelidium, Nereocystis, Zoochlorella
 - B. Nostoc, Porphyra, Chondrus, Batrachospermum

- C. Oscillatoria, Sargassum, Volvox, Fucus
- D. None of the above.

Answer: B



Watch Video Solution

194. In which of the following group all the organisms lack motle stages in the life cycle?

- A. Chlamydomonas, Ulothrix, Sargassum
- B. Chondrus, Gelidium, Batrachospermum
- C. Chlorella, Polysiphonia, Volvox
- D. All of the above

Answer: B



Watch Video Solution

195. Go through the following statements

- (i) IN chlorophyceae the flagella re 2 in number, unequal in size and lateral in position
- (ii) In phaeophyceae flagella are 2-8 in number, equal in size and apical in position
- (iii) Chlorella and Spirullina re rich in proteins and are used as food supplements even by space travellers.
- (iv) The leaves in pteridophyta are microphylls as in Selaginella or macrophylls as in ferns.

Which of these are correct

- A. i,ii,iii
- B. ii,iii
- C. i,ii,iv
- D. iii,iv



Watch Video Solution

196. In a moss the sporophyte

A. arises from a spore produced from the gametophyte

- B. manufactures food for itself, as well as for the gametophyte
- C. is partially parasitic on the gametophyte
- D. Produces gametes that give rise to the gametophyte

Answer: C



- 197. Which of these is mismatched
 - A. phaneros-visible
 - B. kryptos-concealed

- C. gymno-naked
- D. thallusu-diploid



Watch Video Solution

- **198.** Which of the following is a vascular cryptogam
 - A. Marchantia
 - B. Cedrus
 - C. Equisetum
 - D. Ginkgo

Answer: C

199. Which one of the following plants is monoecious

- A. Cycas
- B. Papaya
- C. Marchantia
- D. Pinus

Answer: D



A. Ustilago	
B. Wheat	
C. Funaria	
D. Polytrichum	
Answer: A	
Watch Video Solution	
201. Mannitol (sugar alcohol) is the stored food in	
201. Mannitol (sugar alcohol) is the stored food in A. Fucus	
A. Fucus	

D. Porphyra

Answer: A



Watch Video Solution

202. Young leaves of Cycas show

- A. Simple venation
- B. Circinate venation
- C. Alternate arrangement
- D. Opposite arrangement

Answer: B



203. Algae have cell wall made up of

A. Cellulose, hemicellulose and pectins

B. Cellulose, galactans and mannans

C. Hemicellulose, pectins and proteins

D. Pectins, cellulose and proteins

Answer: B



Watch Video Solution

204. Which of the following algal groups has no motile stage

A. Brown
B. Yellow
C. Red
D. Green
Answer: C
Watch Video Solution
205. Sphagnum is an example of
205. Sphagnum is an example of A. Moss
A. Moss

D. Gymnosperm

Answer: A



Watch Video Solution

206. Consider the following four statements whether they are correct or worng.

- A. The sporophyte in liverworts is more elaborate than that in mosses
- B. Salvinia is heterosporous
- C. The life-cycle in all seed-bearing plants is diplontic

D. In Pinus male and female cones are borne on different trees

Answer: B



Watch Video Solution

207. Selaginella and Salvinia are considered to represent a significant step toward evolution of seed habit because


- A. Female gametophyte is free and gets dispersed like seeds
- B. Female gametophyte lacks archegonia

- C. Megaspores possess endosperm and embryo surrounded by seed coat
- D. Embryo develops in female gametophyte which is retained on parent sporophyte



Watch Video Solution

208. Examine the figure given below and select the right option given all the four parts (a,b,c,d) correctly



identified.

A.

Archegoniphore Female thallus Bud Foot

C. $\frac{a}{\text{Seta}} \frac{b}{\text{Sporophyte}} \frac{c}{\text{Protonema}} \frac{d}{\text{Rhizoids}}$

D. $\frac{a}{\text{Anteridiophore}}$ $\frac{b}{\text{Male thallus}}$ $\frac{c}{\text{Globule}}$ Roots



Watch Video Solution

209. Archegoniophore is present in

- A. Marchantia
- B. Chara
- C. Adiantum
- D. Funaria

Answer: A



210. Compared with the gametophytes of the bryophytes, the gametophytes of vascular plants tend to be:-

- A. smaller but to have larger sex organs
- B. larger but to have smaller sex organs
- C. larger and to have larger sex organs
- D. smaller and to have smaller sex organs

Answer: D



Watch Video Solution

211. The gametophyte is not an independent, free-living generation in

B. Adiantum C. Marchantia D. Pinus **Answer: D Watch Video Solution** 212. Asexual reproduction in Liverworts takes place by the formation of specialized structures called A. Gemmae B. zoospores, akinetes, zygospores

A. Polytrichum, Cycas, Equisetum

- C. Sporangia
- D. Microspores



Watch Video Solution

213. Pinus belongs to the class

- A. gnetopsida
- B. cycadopsida
- C. coniferopsida
- D. sphenopsida

Answer: C

- **214.** Read the following five statements (A-E) and answer as asked next to them
- (A) In Equisetum the female gametophyte is retained on the parent sporophyte
- (B) In ginkgo male gametophyte is not independent
- (C) Sexual reproduction in Volvox is isogamous
- (E) The spores of slime moulds lack cell walls

How many of the above statements are correct

- A. Four
- B. One
- C. Two

D. Three

Answer: B



Watch Video Solution

215. Which one of the following pairs is wrongly matched?

- A. viroids-RNA
- B. Mustard-Synergids
- C. Ginkgo-Archegonia
- D. Salvinia-Prothallus

Answer: D

216. How many organism in the list given below are autotrophs Lactobacillus, Nostoc, Chara, Nitrosomonas, Nitrobacter, Streptomuces, Sacharomyces, Trypanosoma, Porphyra Wolfia

- A. Six
- B. Three
- C. Four
- D. Five

Answer: A



217. Cycas and Adiantum resemble each other in having

- A. Motile sperms
- B. Cambium
- C. Vessles
- D. Seeds

Answer: A



Watch Video Solution

218. Which one of the following is common to multicellular fungi, filamentous algae and protonema of

mosses
A. Members of kingdom plantae
B. Mode of Nutrition
C. Multiplication by fragmentation
D. Diplontic life cycle
Answer: C



Watch Video Solution

219. Which one of the following is a correct statement

A. In gymnosperms female gametophyte is free-living

- B. Anteridiophores and archegoniophores are present in pteridophytes
- C. Origin of seed habit can be traced in pteridophytes
- D. Pteridophyte gametophyte has a protonemal and leafy stage.

Answer: C



- 220. Sex-organs of pteridophytes are
 - A. Multicelluar and jacketed
 - B. Unicellular and jacketed

- C. Unicellular and non jacketed
- D. Multicellular and non jacketed

Answer: A



Watch Video Solution

221. The chloroplasts of algae usually lack

- A. Quantasomes
- B. Lamellae
- C. Pigments reticulum
- D. Grana

Answer: D

222. First amphibian plants of the plant kingdom are

- A. Thallophytes
- B. Bryophytes
- C. pteridophytes
- D. Gymnosperms

Answer: B



B. microsporophylls C. sporophylls D. megasporophylls **Answer: A Watch Video Solution** 224. The tallest tree species of the gymnosperms is A. Cycas B. Pinus C. Sequoia

A. macrophylls

D. None of the Above

Answer: C



Watch Video Solution

225. Besides paddy fields, cyanobacteria are also found inside vegetative part of:

- A. Equisetum
- B. Psilotum
- C. Pinus
- D. Cycas

Answer: D

226. Select the wrong statement

- A. In Oomycetes female gamete is smaller and motile, while male gamete is larger and non-motile
- B. Chlamydomonas exhibits both isogamy and anisogamy and Fucus shows oogamy
- C. Isogametes are similar in structure, function and behaviour
- D. Anisogametes differ either in structure, function or behaviour.

Answer: A



Watch Video Solution

227. Which of the following represent maximum number of species among global biodiversity

- A. Fungi
- B. Mosses and Ferns
- C. Algae
- D. Lichens

Answer: A



228. Male gametophyte with least number of cells is present in

- A. Pinus
- B. Pteris
- C. Funaria
- D. Lilium

Answer: D



229. Which one of the following shows isogamy with non-flagellated gametes

- A. Spirogyra
- B. Sargassum
- C. Ectocarpus
- D. Ulothrix

Answer: A



Watch Video Solution

230. An alga which can be employed as food for humna being is

B. Ulothrix C. Chlorella D. Spirogyra **Answer: C Watch Video Solution** 231. Which of the following is responsible for peat formation? A. Sphagnum B. Marchantia

A. polysiphonia

- C. Riccia
- D. Funaria

Answer: A



Watch Video Solution

232. In which of the following, gametophyte is not independent in free living?

- A. Marchantia
- B. Pteris
- C. Pinus
- D. Funaria

Answer: C



- **233.** Read the following five statements (A to E) and select the option with all correct statement
- (A) Mosses and Lichens are the first organisms to colonise a bare rock.
- (B) Seloginella is a homosporous pteridophyte.
- (C) Coralloid roots in Cycas have VAM.
- (D) Main plant body in bryophytes is gametophytic whereas in pteridophytes it is sporophytic.
- (E) In Gymnoperms, male and female gametophytes are present within sporangla located on sporophyte.

- A. B,C and D
- B. A,D and E
- C. B,C and E
- D. A,C and D

Answer: B



Watch Video Solution

234. Male gametes are flagellated in

- A. Anabaena
- B. Ectocarpus
- C. Spirogyra

D. Polysiphonia

Answer: B



Watch Video Solution

235. Which one of the following statements is wrong?

- A. Agar-agar is obtained from Gelidium and Gracilaria
- B. Chlorella and Spirullina are used as space food
- C. Mannitol is stored food in Rhodophyceae
- D. Algin and carrogen are products of algae.

Answer: C



236. Which one is a wrong statement?

A. Archegonia are found in Bryophyta, Pteridophyta and Gymnosperms

B. Mucor has biflagellate zoospores

C. Haploid endosperm is typical feature of gymnosperms

D. Brown algae have chlorophyll a and c,and fucoxanthin

Answer: B



237. Male gametophyte in angiosperms produces:

- A. two sperms and a vegetative cell
- B. single sperm and a vegetative cell
- C. single sperm and two vegetative cells
- D. three sperms

Answer: A



Watch Video Solution

238. Select the correct statement

A. Salvinia, Ginkgo and Pinus all are gymnosperms

- B. Sequoia is one of the tallest trees
- C. The leaves of gymnosperms are not well adapted to extremes of climate
- D. Gymnosperms are both homosporous and heterosporous

Answer: B



239. In bryophytes and and pteridophytes, transport of male gametes requires

A. Insects

- B. Birds
- C. Water
- D. Wind

Answer: C



Watch Video Solution

240. Conifers are adapted to tolerate extreme environmental conditions because of

- A. Broad hardy leaves
- B. Superficial stomata
- C. Thick cuticle

D. Presence of vessels.

Answer: C



Watch Video Solution

241. Which one of the following statements is wrong.

- A. Algae increase the level of dissolved oxygen in the immediate environment.
- B. Algin is obtained from brown algae, and carrageenin from red algae.
- C. Agar-agar is obtained from Gelidium and Gracilaria
- D. Laminaria and Sargassum are used as food.

Answer: B



Watch Video Solution

242. Zygotic meiosis is characteristic of

- A. Marchantia
- B. Fucus
- C. Funaria
- D. Chlamydomonas

Answer: D



A. Chlorella
B. Volvox
C. Ulothrix
D. Spirogyra
Answer: B
Watch Video Solution
244. Select the mismatch
A. Pinus-Dioecious

243. An example of colonial alga is

- **B.** Cycas-Dioecious
- C. Salvinia -Heterosporous
- D. Equisetum-Homosporous

Answer: A



Watch Video Solution

245. Life cycle of Ectocarpus and Fucus respectivley are

- A. haplontic, diplontic
- B. diplontic, haplodiplontic
- C. haplodiplontic, diplontic
- D. haplodiplontic, haplontic

Answer: C



Watch Video Solution

246. Which one is wrongly matched

- A. Unicellular organism -Chlorella
- B. Gemma cups-Marchantia
- C. Biflagellate Zoospores-Brown algae
- D. Uniflagellate gametes-Polysiphonia

Answer: D



247. Which of the following statements is correct?

A. Stems are usually unbranched in both Cycas and Cedrus.

B. Horsetails are gymnosperms.

C. Selaginella is heterosporous, while salvinia is homosporous.

D. Ovules are not enclosed by ovary wall in gymnosperms.

Answer: D



Assertion And Reason

1. [A]: Flagella found in green algae are of tinsel type.

[R]: Flagella in algae are whiplash type.

A. If both A and R are true and R is the correct explanation of A

B. If both A and R are true but R is not the correct

explanation of A

C. If A is true and R is false

D. If both A and R are false

Answer:



2. [A]: The mosses form dense growth.

[R]: They produce a large number of spores.

A. If both A and R are true and R is the correct explanation of A

B. If both A and R are true but R is not the correct explanation of A

C. If A is true and R is false

D. If both A and R are false

Answer:



3. [A] : Pteridophytes and Gymnosperms are included in Archegoniatae .

[R]: They have archegonia as female reproductive organ .

A. If both A and R are true and R is the correct explanation of A

B. If both A and R are true but R is not the correct explanation of A

C. If A is true and R is false

D. If both A and R are false

Answer:



4. [A]: Pinus bears resin canals.

[R]: Resin canals are found in cortex and bounded by a glandular epithelial layer that secretes resin.

A. If both A and R are true and R is the correct explanation of A

B. If both A and R are true but R is not the correct explanation of A

C. If A is true and R is false

D. If both A and R are false

Answer:



5. [A]: In terms, prothallus is club shaped.

[R] : Prothallus is derived from club shaped zoosporangia

A. If both A and R are true and R is the correct

explanation of A

B. If both A and R are true but R is not the correct

explanation of A

C. If A is true and R is false

D. If both A and R are false

Answer:



6. [A] : Either megasporophylls or microsporophylls are found in Gymnosperms .

[R]: Both mega and microsporophylls are common in Gymnosperms . Megasporophylls are smaller than microsporophylls.

A. If both A and R are true and R is the correct explanation of A

B. If both A and R are true but R is not the correct explanation of A

C. If A is true and R is false

D. If both A and R are false

Answer:

7. [A]: Rhizophore is considered to be stem structure.

[R]: Rhizophore develops from angle meristem present between the two branches of stem.

A. If both A and R are true and R is the correct explanation of A

B. If both A and R are true but R is not the correct explanation of A

C. If A is true and R is false

D. If both A and R are false

Answer:

8. [A] : In Selaginella/Pteris reduction division occurs during microspore formation only .

[R] : It was first shown by Zacharis and Strasburger in 1963 .

A. If both A and R are true and R is the correct explanation of A

B. If both A and R are true but R is not the correct explanation of A

C. If A is true and R is false

D. If both A and R are false

Answer:



9. [A] : Neottia and Monotrapa are saprophytic angiosperms .

[R]: The association of a fungus with roots of higher plants is mycorrhiza.

A. If both A and R are true and R is the correct explanation of A

B. If both A and R are true but R is not the correct explanation of A

C. If A is true and R is false

D. If both A and R are false

Answer:



Watch Video Solution

10. [A]: Fossil plants serve as an index for paleoclimates.

[R] : Because plant adaptations to varied environmental conditions are well known .

A. If both A and R are true and R is the correct explanation of A

B. If both A and R are true but R is not the correct explanation of A

- C. If A is true and R is false
- D. If both A and R are false

Answer:



- **11.** [A] : Coralloid root of Cycas is confined to bacterial zone.
- [R]: Nucleus is absent in young sieve cells.
 - A. If both A and R are true and R is the correct explanation of A

B. If both A and R are true but R is not the correct explanation of A

C. If A is true and R is false

D. If both A and R are false

Answer:



Watch Video Solution

12. [A] : Heterospory has led to the reduction and specialization in spores .

[R]: Incipient heterospory has been found to occur in the fern Platyzoma microphylla.

A. If both A and R are true and R is the correct explanation of A

B. If both A and R are true but R is not the correct explanation of A

C. If A is true and R is false

D. If both A and R are false

Answer:



Watch Video Solution

13. [A]: In Pinus only the dwarf roots become mycorrhizal

[R]: Seeds of Pinus are polycotyledonous.

A. If both A and R are true and R is the correct explanation of A

B. If both A and R are true but R is not the correct explanation of A

C. If A is true and R is false

D. If both A and R are false

Answer:



14. [A] : The mosses like liverworts , do not exhibit alternation of generation .

[R] : The adult gametophyte is conspicuous leafy green , photosynthetic plant in ferns .

A. If both A and R are true and R is the correct explanation of A

B. If both A and R are true but R is not the correct explanation of A

C. If A is true and R is false

D. If both A and R are false

Answer:



15. [A]: Mosses do not seem to resist infestations by most fungi and are often seriously damaged by insects.

[R]: They do not possess antibiotic material.

A. If both A and R are true and R is the correct explanation of A

B. If both A and R are true but R is not the correct explanation of A

C. If A is true and R is false

D. If both A and R are false

Answer:



16. [A]: Monotrapa is a parasitic plant.

[R]: It does not show proper mycorrhizal association.

A. If both A and R are true and R is the correct explanation of A

B. If both A and R are true but R is not the correct explanation of A

C. If A is true and R is false

D. If both A and R are false

Answer:



17. [A]: Some of the Sea kelps are used as fertilizers.

[R]: They are important source of minerals.

A. If both A and R are true and R is the correct explanation of A

B. If both A and R are true but R is not the correct explanation of A

C. If A is true and R is false

D. If both A and R are false

Answer:



18. [A]: Angiosperms and gymnosperms both are flowering plants .

[R]: They both form covered seeds.

A. If both A and R are true and R is the correct explanation of A

B. If both A and R are true but R is not the correct explanation of A

C. If A is true and R is false

D. If both A and R are false

Answer:



