

**MATHS****BOOKS - ACCURATE PUBLICATION****SAMPLE QUESTION PAPER - II (UNSOLVED)****Section A Choose The Correct Option In The Following Questions**

1. Range of the function $f(x) = \frac{|x - 2|}{x - 2}$ is

A. $\{ - 1, 1 \}$

B. $\{ - 1, 2 \}$

C. $\{ - 2, 2 \}$

D. None of these

Answer: A



2. $\tan^{-1}(\sqrt{3}) - \cos^{-1}\left(\frac{1}{2}\right)$ is equal to :

A. $\frac{\pi}{3}$

B. $\frac{2\pi}{3}$

C. 0

D. $\frac{\pi}{6}$

Answer: C



Watch Video Solution

3. If $AB = C$ where A is a matrix of order 2×2 and C is a matrix of order 2×5 , then the order of B is :

A. 3×5

B. 2×5

C. 3×3

D. 5×5

Answer: B



Watch Video Solution

4. For what value of 'x', is the matrix:

$$A = \begin{bmatrix} 0 & 1 & -2 \\ -1 & 0 & 3 \\ x & -3 & 0 \end{bmatrix} \text{ a skew-symmetric matrix.}$$

A. -2

B. -3

C. -4

D. -5

Answer: A



Watch Video Solution

5. If area of triangle is 35 sq. units with vertices (2,-6), (5, 4) and (k,4) then k is :

A. 12

B. - 2

C. - 12, - 2

D. 12, - 2

Answer: D



Watch Video Solution

6. If $f(x) = \begin{cases} \frac{x^2 - 25}{x - 5} & , \quad x \neq 5 \\ k & , \quad x = 5 \end{cases}$ is continuous at $x=5$, then k is equal to:

A. 10

B. 5

C. 0

D. 4

Answer: A



Watch Video Solution

7. Derivative of $(\sin^{-1} x + \cos^{-1} x)$ w.r.t 'x' is equal to :

A. -1

B. 0

C. 1

D. 2

Answer: B



Watch Video Solution

8. If $y = (3x^2 + 2)^2$, then $\frac{dy}{dx}$ is

A. $3x^2 + 2$

B. $x(3x^2 + 2)$

C. $10x(3x^2 + 2)$

D. $12x(3x^2 + 2)$

Answer: D



Watch Video Solution

9. $\int \frac{10x^9 + 10^x \log_e 10}{x^{10} + 10^x} dx$ is equal to :

A. $\log|x^{10} + 10^x| + c$

B. $10^x + 10^{10} + c$

C. $10^x - x^{10} + x$

D. $(10^x - x^{10})^{-1} + c$

Answer: A



Watch Video Solution

10. Choose the correct answer: $\int e^x \sec x (1 + \tan x) dx$ equals :

A. $e^x \cos x + C$

B. $e^x \sec x + C$

C. $e^x \sin x + C$

D. $e^x \tan x + C$

Answer: B



Watch Video Solution

11. The number of arbitrary constants in the particular solution of a differential equation of Third order is :

A. 0

B. 2

C. 3

D. 5

Answer: A



Watch Video Solution

12. Solve the following differential equations

$$\frac{dy}{dx} - \frac{y}{x} = 2x^2$$

A. x

B. x^2

C. $\frac{1}{x}$

D. $\frac{2}{x}$

Answer: C



Watch Video Solution

13. The projection of $\vec{a} = 2\hat{i} - \hat{j} + \hat{k}$ on $\vec{b} = \hat{i} - 2\hat{j} + \hat{k}$ is equal to :

A. $\frac{5\sqrt{6}}{3}$

B. $\frac{5}{\sqrt{6}}$

C. $\frac{6}{\sqrt{14}}$

D. $\frac{\sqrt{6}}{5}$

Answer: B



Watch Video Solution

14. If $|\vec{a}| = 5$, $|\vec{b}| = 4$ and $\vec{a} \cdot \vec{b} = 16$, then $|\vec{a} \times \vec{b}|$ is

A. 10

B. 12

C. 14

D. 16

Answer: B



[Watch Video Solution](#)

15. The distance between the planes, $3x + 2y - 6z - 14 = 0$ and $3x + 2y - 6z + 21 = 0$ is,

A. 35

B. 7

C. 1

D. 5

Answer: D



[Watch Video Solution](#)

16. In a single throw of two dice, the probability of getting total of 7 or 9 is :

A. 0

B. $\frac{1}{36}$

C. $\frac{1}{9}$

D. $\frac{1}{6}$

Answer: D



[Watch Video Solution](#)

Section A Fill In The Blanks From The Given Options

1. Consider the set $A = \{1, 2, 3\}$ and R be the smallest equivalence relation on A , then $R = \dots\dots\dots$.



[Watch Video Solution](#)

2.

$$\frac{1}{5}, R = \{(1, 1), (2, 2), (3, 3)\}, 0, \frac{2 \tan^{-1} x}{1 + x^2}, \text{ Parallel } , \left(\vec{b}_1 \times \vec{b}_2 \right) \cdot \left(\vec{a} \right)$$

The value of the det. $\begin{vmatrix} x + y & y + z & z + x \\ z & x & y \\ 1 & 1 & 1 \end{vmatrix}$ is

 [Watch Video Solution](#)

3.

$$0, \frac{\pi}{2}, \frac{x}{(1 - x^2)^{\frac{3}{2}}}, R = \{(3, 8), (6, 6), (9, 4), (12, 2)\}, 2x - y + 1 = 0, \left| \vec{b} \right|$$

Let the relation R be defined in N by aRb if $2a + 3b = 30$, Then R =

 [Watch Video Solution](#)

4. Show that the tangents to the curve $y = 7x^3 + 11$ at the points $x = 2$ and $x = -2$ are parallel.

 [Watch Video Solution](#)

5. $\int_0^{\pi/2} \left(\frac{\sqrt{\cos x}}{\sqrt{\sin x} + \sqrt{\cos x}} \right) dx$ is equal to

 [Watch Video Solution](#)

6. The number of arbitrary constants in the general solution of a differential equation of fourth order are:

 [Watch Video Solution](#)

7. The two lines $\vec{r}_1 = \vec{a}_1 + \lambda \vec{b}$, $\vec{r}_2 = \vec{a}_2 + \mu \vec{b}$ will intersect if $d = \dots$

 [Watch Video Solution](#)

8. If A and B are independent events and if $P(A) = \frac{1}{2}$, $P(B) = \frac{2}{5}$, then $P(A \cap B)$ is equal to :

 [Watch Video Solution](#)

Section A State True Or False For The Following Statements

1. The value of $\sin^{-1}(\sin(2\pi/3))$ is :

 [Watch Video Solution](#)

2. Value of $\begin{vmatrix} 1 & \omega & \omega^2 \\ \omega & \omega^2 & 1 \\ \omega^2 & 1 & \omega \end{vmatrix}$ is zero, where ω, ω^2 are imaginary cube roots of unity.

 [Watch Video Solution](#)

3. Derivative of $\sin^{-1}(\sin 2x)$ w.r.t x is 2

 [Watch Video Solution](#)

4. Solve $\int \frac{2x}{1+x^2} dx$

 [Watch Video Solution](#)

5. If $\vec{a} = \hat{i} + 5\hat{j} + 4\hat{k}$ and $\vec{b} = 5\hat{i} - \hat{j} + 2\hat{k}$, then $\vec{a} \cdot \vec{b}$ is equal to 8

 [Watch Video Solution](#)

6. If a line makes angles α, β, γ respectively with positive directions of the coordinate axes, then the value of $\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma = 1$.

 [Watch Video Solution](#)

7. Let $P(A) > 0$ and $P(B) > 0$. Then A and B can be both mutually exclusive and independent."

 [Watch Video Solution](#)

8. Quadrant represented by the region $x \geq 0, y \leq 0$ is

 [Watch Video Solution](#)

Section B

1. Find the values of a, b, c and d from the equation :

$$\begin{bmatrix} a - b & 2a + c \\ 2a - b & 3c + d \end{bmatrix} = \begin{bmatrix} -1 & 5 \\ 0 & 13 \end{bmatrix} \text{ and write correct answer from the}$$

following:

 [Watch Video Solution](#)

2. If A is a square matrix such that $A^2 = I$, then find the simplified value of $(A - I)^3 + (A + I)^3 - 7A$.

 [Watch Video Solution](#)

3. Show that the function f given by $f(x) = x^3 - 3x^2 + 4x$, $x \in \mathbb{R}$ is strictly increasing on \mathbb{R} .

 [Watch Video Solution](#)

4. Maximum value of the function $\sin x + \cos x$ is .

 [Watch Video Solution](#)

5. Evaluate the following integrals : $\int \left(\frac{x + \cos 6x}{3x^2 + \sin 6x} \right) dx$

 [Watch Video Solution](#)

6. Find the area of the region bounded by $x^2 = 4y$, $y = 2$, $y = 4$ and the y -axis in the first quadrant.

 [Watch Video Solution](#)

7. Show that the vectors $2\hat{i} - \hat{j} + \hat{k}$ and $\hat{i} - 3\hat{j} - 5\hat{k}$ are at right angles.

 [Watch Video Solution](#)

8. Find λ if $(2\hat{i} + 6\hat{j} + 14\hat{k}) \times (\hat{i} - \lambda\hat{j} + 7\hat{k}) = \vec{0}$.

 [Watch Video Solution](#)

Section C

1. Prove that $\cot^{-1} \left(\frac{\sqrt{1+x} - \sqrt{1-x}}{\sqrt{1+x} + \sqrt{1-x}} \right) = \frac{\pi}{4} + \frac{1}{2} \cos^{-1} x$

 [Watch Video Solution](#)

2. Differentiate the following w.r.t. x :

$$x^{\sin x} + (\sin x)^{\cos x}$$

 [Watch Video Solution](#)

3. If $x = a \cos \theta + b \sin \theta$ and $y = a \sin \theta - b \cos \theta$, then prove that

$$y^2 \frac{d^2 y}{dx^2} - \frac{dy}{dx} + y = 0$$

 [Watch Video Solution](#)

4. If $[x]$ stands for integral part of x , then show that $\int_0^1 [5x] dx = 2$.

 [Watch Video Solution](#)

5. Find : $\int \frac{x^3 dx}{x^4 + 3x^2 + 2} dx$.

 [Watch Video Solution](#)

6. Find the particular solution of the following differential equation :

$$\frac{dy}{dx} = 1 + x^2 + y^2 + x^2 y^2, \text{ given that } y = 1 \text{ when } x = 0.$$

 [Watch Video Solution](#)

7. Suppose a girl throws a die. If she gets a 5 or 6, she tosses a coin three times and notes the numbers of heads. If she gets 1,2,3, or 4, she tosses a coin once and notes whether a head or a tail is obtained. If she attained exactly one head what is the probability that she threw 1,2,3, or 4 with the die?

 [Watch Video Solution](#)

8. Two numbers are selected at random (without replacement) from the first six positive integers. Let X denote the larger of the two numbers obtained. Find $E(X)$.

 [Watch Video Solution](#)

1. If $A = \begin{bmatrix} 1 & -2 & 1 \\ 0 & -1 & 1 \\ 2 & 0 & -3 \end{bmatrix}$, find A^{-1} and solve the system of equations

$$x - 2y + z = 0, \quad -y + z = -2, \quad 2x - 3z = 10$$

 [Watch Video Solution](#)

2. Prove the following identities :

$$\begin{vmatrix} a & a^2 & a^3 \\ b & b^2 & b^3 \\ c & c^2 & c^3 \end{vmatrix} = abc(a - b)(b - c)(c - a).$$

 [Watch Video Solution](#)

3. Find the direction ratios of the normal to the plane, which passes through the points $(1, 0, 0)$ and $(0, 1, 0)$ and makes angle $\frac{\pi}{4}$ with the plane $x + y = 3$. Also find the equation of the plane.

 [Watch Video Solution](#)

4. Find the coordinates of the foot of perpendicular and the length of the perpendicular drawn from the point $P(5, 4, 2)$ to the line $\vec{R} = -\hat{i} + 3\hat{j} + \hat{k} + \lambda(2\hat{i} + 3\hat{j} - \hat{k})$. Also find the image of P in this line.

 [Watch Video Solution](#)

5. Solve the following linear programming problems graphically

Minimize $Z = 5x + 7y$ subject to the constraints
 $2x + y \geq 8, x + 2y \geq 10, x, y \geq 0$

 [Watch Video Solution](#)

6. Maximize $z = 9x + 3y$ subject to the constraints

$$2x + 3y \leq 13$$

$$2x + y \leq 5$$

$$x, y \geq 0$$

 [Watch Video Solution](#)

