



MATHS

BOOKS - ACCURATE PUBLICATION

SAMPLE QUESTION PAPER-II (SOLVED)

Section A

1. If $A = \{a, b, c, d\}$ then a relation

$R = \{(a, a), (b, b), (c, c), (d, d)\}$ on A is :

A. Symmetric

B. Transitive

C. Reflexive

D. None of these

Answer: C



Watch Video Solution

2. The value of $\operatorname{cosec}^{-1}(-2)$ is equal to :

A. $\frac{\pi}{3}$

B. $-\frac{\pi}{6}$

C. $-\frac{\pi}{3}$

D. $\frac{\pi}{6}$

Answer: B



Watch Video Solution

3. If $AB = C$ where A is a matrix of order 2×4 and C is a matrix of order 2×5 , then the order of B is :

A. 3×5

B. 4×5

C. 3×3

D. 5×5

Answer: B



[Watch Video Solution](#)

4. The number of all possible matrices of order 3×3 with each entry 0 or 1 is:

A. 27

B. 18

C. 81

D. 512

Answer: D



Watch Video Solution

5. If $A = [[321]]$, then AA' is equal to :

A. $(9 \quad 4 \quad 1)$

B. $\begin{pmatrix} 9 \\ 4 \\ 1 \end{pmatrix}$

C. -14

D. -6

Answer: C



Watch Video Solution

6. If $f(x) = \begin{cases} \frac{\sin 5x}{2x} & x \neq 0 \\ k & x = 0 \end{cases}$ is continuous at $x = 0$ then

value of k is :

A. 5

B. $\frac{3}{5}$

C. $\frac{5}{3}$

D. 0

Answer: B



Watch Video Solution

7. If $y = 3^x$, then $\frac{dy}{dx}$ is :

A. 3^x

B. $3^x \log 3$

C. 3

D. $\frac{3^x}{\log 3}$

Answer: B



Watch Video Solution

8. If $y = \tan x$ then at $x = 0$, y_2 is equal to :

A. -1

B. 1

C. 0

D. $\frac{1}{2}$

Answer: C



Watch Video Solution

9. $\int \frac{\sin x}{\cos^2 x} dx$ equals :

A. $\sec x + c$

B. $\tan x + c$

C. $\operatorname{cosec} x + c$

D. $\sec^2 x + c$

Answer: A



Watch Video Solution

10. $\int_0^1 \frac{1}{\sqrt{1-x^2}} dx$ is equal to :

A. $\frac{\pi}{4}$

B. $\frac{\pi}{3}$

C. $\frac{\pi}{2}$

D. $\frac{\pi}{6}$

Answer: C



Watch Video Solution

11. The number of arbitrary constants in the particular solution of a differential equation of fifth order is :

A. 0

B. 2

C. 3

D. 5

Answer: A



Watch Video Solution

12. The Integrating Factor of the differentiate equation

$$\frac{dy}{dx} - 2y = 3x \text{ is :}$$

A. e^{2x}

B. e^{-2x}

C. e^x

D. $2x$

Answer: B



Watch Video Solution

13. If $\vec{a} = \hat{i} + 2\lambda\hat{j} + \hat{k}$ and $\vec{b} = 2\hat{i} + \hat{j} - 3\hat{k}$ are perpendicular to each other, the value of λ is :

A. 0

B. 1

C. 2

D. $\frac{1}{2}$

Answer: D



Watch Video Solution

14. If θ is the angle between any two vectors \vec{a} and \vec{b} , then $\left| \vec{a} \cdot \vec{b} \right| = \left| \vec{a} \times \vec{b} \right|$ when θ is equal to :

A. 0

B. $\frac{\pi}{4}$

C. $\frac{\pi}{2}$

D. π

Answer: B



[Watch Video Solution](#)

15. The distance of the plane $\vec{r} \cdot (2\hat{i} + 3\hat{j} - 6\hat{k}) = 7$ from origin is :

A. -1

B. 0

C. $\frac{1}{7}$

D. 1

Answer: D



Watch Video Solution

16. In a single throw of two dice, the chances of throwing a sum of 5 is :

A. 0

B. $\frac{1}{36}$

C. $\frac{1}{9}$

D. $\frac{5}{36}$

Answer: C



Watch Video Solution

17. If f is a bijection, then it is.....



Watch Video Solution

18. If $|A| = 3$, where A is a 2×2 matrix, $|\text{Adj } A| = \dots\dots\dots$



Watch Video Solution

19. $\frac{d}{dx} (x^2 + 2x + 5)^2 = \dots\dots\dots$



Watch Video Solution

20. The slope of tangent to the curve $y = 2 - x^2$ at $x = 1$ is



Watch Video Solution

21. The value of $\int_{-\pi}^{\pi} \sin^{2019} x \cos^{2020} x dx$ is equal to.....

 [Watch Video Solution](#)

22. Degree of differential equation $\left(\frac{dy}{dx}\right)^3 + \frac{d^3y}{dx^3} = 5$ is

 [Watch Video Solution](#)

23. The distance between the planes $3x + 2y - 6z - 18 = 0$ and $3x + 2y - 6z + 10 = 0$ is

 [Watch Video Solution](#)

24. If A and B are mutually exclusive, then $P(A \cap B)$ is equal to _____.

 [Watch Video Solution](#)

25. The value of the expressions $(\cos^{-1} x)^2$ is equal to $\sec^2 x$.

 [Watch Video Solution](#)

26. $(A^3)^{-1} = (A^{-1})^3$, where A is a square matrix and $|A| \neq 0$.

 [Watch Video Solution](#)

27. Derivative of $\sin^{-1}(\cos x)$ w.r.t. x is 1

 [Watch Video Solution](#)

28. prove: $\int \frac{\sin^2 x}{1 + \cos x} dx = x + \sin x + c.$



Watch Video Solution

29. If $\vec{a} = \hat{i} + 4\hat{j} + 4\hat{k}$ and $\vec{b} = 4\hat{i} - \hat{j} + 2\hat{k}$, then $\vec{a} \cdot \vec{b}$ is equal to 8.



Watch Video Solution

30. If $\cos \alpha$, $\cos \beta$, $\cos \gamma$ are the direction-cosines of a line, then the value of

$\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma$ is ____ .



Watch Video Solution

31. Quadrant represented by the region $x \geq 0, y \geq 0$ is first.



Watch Video Solution

32. If A and B are two events such that

$P(A) > 0$ and $P(A) + P(B) > 1$, then $P(B | A) \geq 1 - \frac{P(B')}{P(A)}$.



Watch Video Solution

Section B

1. If $A = \begin{bmatrix} -2 & 4 \\ -1 & 3 \end{bmatrix}$, then find AA .



Watch Video Solution

2. If $A = \begin{bmatrix} 1 & -2 \\ 3 & 2 \end{bmatrix}$ and $f(x) = x^2 - 2x + 3$, then find $f(A)$.



[Watch Video Solution](#)

3. Find the equation of the tangent to the curve $y = 2x^2 + 3 \sin x$ at $x = 0$.



[Watch Video Solution](#)

4. Show that the function f given by $f(x) = x^3 - 3x^2 + 4x$, $x \in \mathbb{R}$ is strictly increasing on \mathbb{R} .



[Watch Video Solution](#)

5. Evaluate : $\int \sin 5x \sin 3x dx$.



Watch Video Solution

6. Using integration, find the area of the region bounded by the curve $x^2 + y^2 = 16$ in the first quadrant.



Watch Video Solution

7. Find a vector in direction of vector $4\hat{i} - \hat{j} + 3\hat{k}$ which has magnitude 7 units.



Watch Video Solution

8. If $\vec{a} + \vec{b} + \vec{c} = 0$ and $|\vec{a}| = 3$, $|\vec{b}| = 5$, $|\vec{c}| = 7$, find the angle between \vec{a} and \vec{b} .



Watch Video Solution

Section C

1. Show that $\sin^{-1}\left(\frac{12}{13}\right) + \cos^{-1}\left(\frac{4}{5}\right) + \tan^{-1}\left(\frac{63}{16}\right) = \pi$

 [Watch Video Solution](#)

2. Differentiate : $x^{\sin x} + (\sin x)^x$ w. r. tx :

 [Watch Video Solution](#)

3. If $y = e^{\tan^{-1}x}$, then prove that $(1 + x^2)y_2 + (2x - 1)y_1 = 0$

 [Watch Video Solution](#)

4. Evaluate : $\int \frac{(x - 4)e^x}{(x - 2)^3} dx.$

 [Watch Video Solution](#)

5. Evaluate : $\int \frac{1}{(x - 1)(x + 2)(x - 3)} dx.$

 [Watch Video Solution](#)

6. Solve the differential equation:

$$\left[\frac{e^{-2\sqrt{x}}}{\sqrt{x}} - \frac{y}{\sqrt{x}} \right] \frac{dx}{dy} = 1, (x \neq 0)$$

 [Watch Video Solution](#)

7. An insurance company insured 3000 scooters, 4000 cars and 5000 trucks. The probabilities of an accident involving a scooter, a car and a truck are 0.02, 0.03, 0.04 respectively. One of the insured vehicles meets with an accident. Find the probability that it is a car.



[Watch Video Solution](#)

8. The probability of getting an ace from a well shuffled deck of 52 playing cards.



[Watch Video Solution](#)

1. Using matrix method, solve the following system of equations

$$x + 2y - 3z = 1, 2x - 3z = 2, x + 2y = 3.$$

 [Watch Video Solution](#)

2. If $A = \begin{bmatrix} 3 & 4 & 2 \\ 2 & 3 & 5 \\ 1 & 0 & 1 \end{bmatrix}$ find A^{-1} and hence solve the equations

$$3x + 4y + 2z = -1, 2x + 3y + 5z = 7, x + z = 2.$$

 [Watch Video Solution](#)

3. Find the equation of the plane passing through the line of

intersection of the planes $2x + y - z = 3$ and

$5x - 3y + 4z = 9$ and parallel to the lines

$$\frac{x - 1}{2} = \frac{y - 3}{4} = \frac{z - 5}{5}$$

 [Watch Video Solution](#)

4. Find the shortest distance between the lines given by

$$\vec{r} = 3\hat{i} + 8\hat{j} + 3\hat{k} + \lambda(3\hat{i} - \hat{j} + \hat{k}) \text{ and}$$

$$\vec{r} = -3\hat{i} - 7\hat{j} + 6\hat{k} + \mu(-3\hat{i} + 2\hat{j} + 4\hat{k}).$$



Watch Video Solution

5. Graphically maximize $Z = 5x + 2y$ subject to the constraints :

$$x - 2y \leq 2, 3x + 2y \leq 12, -3x + 2y \leq 3, x \geq 0, y \geq 0$$



Watch Video Solution

6. Solve the following L.P.P graphically :

$$\text{Maximise } Z = 20x + 10y$$

subject to the constraints

$$x + 2y \leq 28$$

$$3x + y \leq 24$$

$$x \geq 2$$

$$x, y \geq 0$$



Watch Video Solution