



MATHS

BOOKS - ACCURATE PUBLICATION

SAMPLE QUESTION PAPER-V

Section A

1. Show that the relation R in the set $\{1,2,3\}$ given by $R = \{(1,2),(2,1)\}$ is symmetric but neither reflexive nor transitive.

- A. Symmetric
- B. Reflective
- C. Transitive
- D. Symmetric and transitive

Answer: A



Watch Video Solution

2. Choose the correct option in the question :

$\sin(\tan^{-1} x), |x| < 1$ is equal to

A. $\frac{x}{\sqrt{1-x^2}}$

B. $\frac{1}{\sqrt{1-x^2}}$

C. $\frac{1}{\sqrt{1+x^2}}$

D. $\frac{x}{\sqrt{1+x^2}}$

Answer: D



Watch Video Solution

3. Choose the correct option in the question :

If $\begin{bmatrix} x + y + z \\ x + z \\ y + z \end{bmatrix} = \begin{bmatrix} 9 \\ 5 \\ 7 \end{bmatrix}$, then $(x + y)$ equals

A. 4

B. 5

C. 6

D. 7

Answer: C



Watch Video Solution

4. Choose the correct option in the question :

If $A = (3 \ 2 \ 1)$ then $A A'$ is equal to

A. $(9 \ 4 \ 1)$

B. $\begin{pmatrix} 9 \\ 4 \\ 1 \end{pmatrix}$

C. -14

D. -6

Answer: C



Watch Video Solution

5. Choose the correct option in the question :

$$\text{If } \begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix} \begin{pmatrix} 3 & 1 \\ 2 & 5 \end{pmatrix} = \begin{pmatrix} 7 & 11 \\ k & 23 \end{pmatrix}, \text{ then } k \text{ equals}$$

A. 15

B. 16

C. 17

D. 18

Answer: C



Watch Video Solution

6. Find k , so that $f(x) = \begin{cases} \frac{\sin kx}{x}, & x \neq 0 \\ 4 + x, & x = 0 \end{cases}$ is continuous at $x = 0$

A. 4

B. 5

C. 6

D. 7

Answer: A



[Watch Video Solution](#)

7. Choose the correct option in the question :

The derivative of $f(x) = |x|$ at $x = 3$ equals

A. 1

B. -1

C. 0

D. 2

Answer: A



[Watch Video Solution](#)

8. Choose the correct option in the question :

Derivative of $\sin^{-1}(e^x)$ w.r.t. x is

A. $\frac{e^x}{\sqrt{1+e^{2x}}}$

B. $\frac{e^x}{\sqrt{1-e^{2x}}}$

C. $\frac{1}{\sqrt{1+e^{2x}}}$

D. $\frac{1}{\sqrt{1-e^{2x}}}$

Answer: B



Watch Video Solution

9. Choose the correct option in the question :

$\int \frac{dx}{e^x + e^{-x}}$ is equal to

A. $\tan^{-1}(e^x) + C$

B. $\tan^{-1}(e^{-x}) + C$

$$C. \log(e^x - e^{-x}) + C$$

$$D. \log(e^x + e^{-x}) + C$$

Answer: A



Watch Video Solution

$$10. \int_0^1 \frac{2x}{5x^2 + 1} dx = \log 6$$

A. $\frac{1}{2} \log 6$

B. $\frac{1}{3} \log 6$

C. $\frac{1}{4} \log 6$

D. $\frac{1}{5} \log 6$

Answer: D



Watch Video Solution

11. A homogeneous differential equation of the form $\frac{dx}{dy} = h\left(\frac{x}{y}\right)$ can be solved by making the substitution.

A. $y = ux$

B. $u = yx$

C. $x = uv$

D. $x = v$

Answer: C



[Watch Video Solution](#)

12. Choose the correct option in the question :

If of $\frac{dy}{dx} + \frac{y}{x} = x^2$ s

A. x

B. $2x$

C. $\log x$

D. $\log 2x$

Answer: A



[Watch Video Solution](#)

13. Choose the correct option in the question :

The projection of $\vec{a} = 2\hat{i} - 2\hat{j} - \hat{k}$ on $\vec{b} = 3\hat{i} - \hat{j} + 2\hat{k}$ is equal to

A. $\frac{5\sqrt{6}}{3}$

B. $\frac{6}{\sqrt{14}}$

C. $\frac{\sqrt{6}}{14}$

D. $\frac{5}{\sqrt{6}}$

Answer: B



[Watch Video Solution](#)

14. If $\vec{a} = 5\hat{i} - \hat{j} - 3\hat{k}$ and $\vec{b} = \hat{i} + 3\hat{j} - 5\hat{k}$, then show that the vectors $(\vec{a} + \vec{b})$ and $(\vec{a} - \vec{b})$ are perpendicular.

A. 30°

B. 45°

C. 60°

D. 90°

Answer: D



[Watch Video Solution](#)

15. Distance between the planes $2x + 3y + 4z = 4$ and $4x + 6y + 8z = 12$ is :

A. 2 units

B. 4 units

C. 8 units

D. $\frac{2}{\sqrt{29}}$ units

Answer: D



Watch Video Solution

16. If $P(A) = \frac{6}{11}$, $P(B) = \frac{5}{11}$ and $P(A \cup B) = \frac{7}{11}$ find: $P(A | B)$

A. $\frac{1}{5}$

B. $\frac{2}{5}$

C. $\frac{3}{5}$

D. $\frac{4}{5}$

Answer: D



Watch Video Solution

17. Fill in the blank from the given option

0.32, _____ symmetric _____ ,
 $\frac{1}{2}\sec^2 \frac{x}{2}, x + y - 3 = 0, x + y - = 2, \frac{1}{2}\log|2x^2 + 6x + 5| + C, \{-1, 1\}$

Range of the function $f(x) = \frac{|x - 3|}{x - 3}$ is

 [Watch Video Solution](#)

18. Fill in the blank from the given option

0.32, _____ symmetric _____ ,
 $\frac{1}{2}\sec^2 \frac{x}{2}, x + y - 3 = 0, x + y - = 2, \frac{1}{2}\log|2x^2 + 6x + 5| + C, \{-1, 1\}$

The matrix $A \begin{bmatrix} 1 & -1 & 5 \\ -1 & 2 & 1 \\ 5 & 1 & 3 \end{bmatrix}$ is a.....matrix.

 [Watch Video Solution](#)

19. Fill in the blank from the given option

0.32, _____ symmetric _____ ,

$$\frac{1}{2}\sec^2 \frac{x}{2}, x + y - 3 = 0, x + y - = 2, \frac{1}{2}\log|2x^2 + 6x + 5| + C, \{ - I, 1 \}$$

Differentiation of: $\sqrt{\frac{1 - \cos x}{1 + \cos x}}$ w.r.t.x is

 [Watch Video Solution](#)

20. Find the equation of the normal to curve $y^2 = 4x$ at the point (1,2)

 [Watch Video Solution](#)

21. Fill in the blank from the given option

Find $\int \frac{2x + 3}{2x^2 + 6x + 5} dx = \dots\dots$

 [Watch Video Solution](#)

22. Fill in the blank from the given option

0.32, symmetric ,

$$\frac{1}{2}\sec^2 \frac{x}{2}, x + y - 3 = 0, x + y - = 2, \frac{1}{2}\log|2x^2 + 6x + 5| + C, \{ - I, 1 \}$$

I.F. of $\frac{dy}{dx} + 2y = \sin x$ is



Watch Video Solution

23. Find the Cartesian equation of the following plane:

$$\vec{r} \cdot (\hat{i} + \hat{j} - \hat{k}) = 2$$



Watch Video Solution

24. Fill in the blank from the given option

0.32,

symmetric

$$\frac{1}{2} \sec^2 \frac{x}{2}, x + y - 3 = 0, x + y - = 2, \frac{1}{2} \log|2x^2 + 6x + 5| + C, \{-I, 1\}$$

If $P(A) = 0.8$, $P(B) = 0.5$, $P(B | A) = 0.4$ then $P(A \cap B)$ is



Watch Video Solution

25. State true or false for the statement

$$\sin^{-1}(2 \sin^{-1} x) = 2x \sqrt{1 - x^2}$$



Watch Video Solution

26. State true or false for the statement

The matrix $\begin{bmatrix} 3 - 2x & x + 1 \\ 2 & 4 \end{bmatrix}$ is singular for $x = 2$.



Watch Video Solution

27. Differentiate the following w.r.t.x.

$\sin^{-1}(3x - 4x^3)$, $-\frac{1}{2} < x < \frac{1}{2}$



Watch Video Solution

28. Solve the following integration

$\int \frac{\sin x + \cos x}{\sqrt{1 + \sin 2x}} \cdot dx$, here $(\sin x + \cos x) > 0$



Watch Video Solution

29. State true or false for the statement

The value of p if $(2\hat{i} + 6\hat{j} + 27\hat{k}) \times (\hat{i} + 3\hat{j} + p\hat{k}) = \vec{0}$ is $\frac{11}{2}$



[Watch Video Solution](#)

30. State true or false for the statement

The equation of a plane parallel to $2x - 3y + z + 9 = 0$ and passing through origin is $2x - 3y + z = 0$.



[Watch Video Solution](#)

31. State true or false for the statement

If a leap year is selected at random, the chance that it will contain 53 Tuesdays is $\frac{2}{7}$.




[Watch Video Solution](#)

32. State true or false for the statement

The feasible solution for LPP is shown in Let. $Z = 3x - 4y$ be objective

function. (Maximum value of Z + Minimum value of Z) is equal - 13



[Watch Video Solution](#)

1. If $3A - B = \begin{bmatrix} 5 & 0 \\ 1 & 1 \end{bmatrix}$ and $B = \begin{bmatrix} 4 & 3 \\ 2 & 5 \end{bmatrix}$, then find the matrix A.

 [Watch Video Solution](#)

2. If $A = \begin{bmatrix} 2 & 3 \\ 5 & -2 \end{bmatrix}$, show that $A^{-1} = \frac{1}{19}A$

 [Watch Video Solution](#)

3. Find the slope of the tangent to the curve $y = 3x^4 - 4x$ at $x = 1$

 [Watch Video Solution](#)

4. Prove that the function $f(x) = x^3 - 6x^2 + 12x - 18$ is increasing on \mathbb{R} .

 [Watch Video Solution](#)

5. Evaluate $\int \frac{1}{x + x \log x} dx$.

 [Watch Video Solution](#)

6. Find the area of the region bounded by the curve $y^2 = 4x$ and the line $x = 3$.

 [Watch Video Solution](#)

7. The two vectors $\hat{j} + \hat{k}$ and $3\hat{i} - \hat{j} + 4\hat{k}$ represent the two sides AB and AC respectively of a $\triangle ABC$. Find the length of the median through A

 [Watch Video Solution](#)

8. If a unit vector \vec{a} makes angle $\frac{\pi}{4}$ with \hat{i} , $\frac{\pi}{3}$ with \hat{j} and acute angle θ with \hat{k} then find the component of \vec{a} along the angle θ .



[Watch Video Solution](#)

Section C

1. Show that : $\sin^{-1}\left(2x\sqrt{1-x^2}\right) = 2\sin^{-1}x$, $-\frac{1}{\sqrt{2}} \leq x \leq \frac{1}{\sqrt{2}}$

 Watch Video Solution

2. Prove that $\cos^{-1}\frac{12}{13} + \sin^{-1}\frac{3}{5} = \sin^{-1}\frac{56}{65}$

 Watch Video Solution

3. If $x^{16} \cdot y^9 = (x^2 + y)^{17}$, prove that $\frac{dy}{dx} = \frac{2y}{x}$.

 Watch Video Solution

4. Evaluate $\int \frac{2}{(1-x)(1+x^2)} dx$

[Watch Video Solution](#)

5. Show that: $\int_0^{\frac{3}{2}} |x \cos \pi x| dx = \frac{5\pi - 2}{2\pi^2}$

[Watch Video Solution](#)

6. Solve the following differential equation $(y^2 - x^2) dy = 3xy dx$.

[Watch Video Solution](#)

7. A and B throw a pair of dice alternately. A wins the game if he gets a total of 7 and B wins if he gets a total of 10. If A starts the game, find the probability that B wins.

[Watch Video Solution](#)

8. A bag contains $(2n + 1)$ coins. It is known that n of these coins have a head on both side whereas the rest of the coins are fair A coin is plaked up at random from the bag and is tossed. If the probability that the toss results in a head is $\frac{31}{42}$, determine the value of n .

 [Watch Video Solution](#)

Section D

1. If x, y, z are different and $\Delta = \begin{vmatrix} x & x^2 & 1 + x^3 \\ y & y^2 & 1 + y^3 \\ z & z^2 & 1 + z^3 \end{vmatrix} = 0$, show that $xyz = -1$

 [Watch Video Solution](#)

2. Using matrix method , solve the equations

$$x - y + z = 4$$

$$2x + y - 3z = 0$$

$$x + y + z = 2$$



Watch Video Solution

3. Find the value of equation of a line passing through the point (2,3,2) and parallel to the line $\vec{r} = (-2\hat{i} + 3\hat{j}) + \lambda(2\hat{i} - 3\hat{j} + 6\hat{k})$. Also, find the distance between these two lines



Watch Video Solution

4. Find the shortest distance between the lines

$$\vec{r} = (\hat{i} + 2\hat{j} + \hat{k}) + \lambda(\hat{i} - \hat{j} + \hat{k}) \quad \text{and}$$

$$\vec{r} = 2\hat{i} - \hat{j} - \hat{k} + \mu(2\hat{i} + \hat{j} + 2\hat{k})$$



Watch Video Solution

5. Maximum and minimise $z = 3x + 2y - 3$ subject to the constraints

$$x + y \geq 4, x + y \leq 12, x \leq 9, y \leq 9, x, y \geq 0$$



Watch Video Solution

6. Graphically minimize and maximize $z = 7x + 5y$ subject to the constraints: $x + y \leq 50$, $x + 2y \leq 100$, $x - y \geq 10$, $y \geq 0$.



[Watch Video Solution](#)

Section A

1. Let R be the relation in the set $\{1, 2, 3, 4\}$ given by $R = \{(1, 2), (2, 2), (1, 1), (4, 4), (1, 3), (3, 3), (3, 2)\}$. Choose the correct answer.

- A. R is reflexive and symmetric but not transitive.
- B. R is reflexive and transitive but not symmetric.
- C. R is symmetric and transitive but not reflexive.
- D. R is an equivalence relation.

Answer: B



[Watch Video Solution](#)

2. Find the principal value of $\cot^{-1}(\sqrt{3})$

A. $\sqrt{2}$

B. $\sqrt{3}$

C. $\sqrt{5}$

D. 1

Answer: B



[Watch Video Solution](#)

3. If the matrix A is both symmetric and skew symmetric, then :

A. A is a diagonal matrix

B. A is zero matrix

C. A is a square matrix

D. None of these

Answer: B



Watch Video Solution

4. If $A = \begin{pmatrix} 1 & 2 \\ 2 & 1 \end{pmatrix}$, then $A^2 - 3A$ is:

A. $2A$

B. A is zero matrix

C. $3A$

D. $4A$

Answer: A



Watch Video Solution

5. The value of the determinant $\begin{vmatrix} 2 & 3 & 4 \\ 5 & 6 & 8 \\ 6x & 9x & 12x \end{vmatrix}$ is

A. 1

B. 2

C. 0

D. 4

Answer: C



Watch Video Solution

6. The value of k so that the function $f(x) = \begin{cases} \frac{\sin 2x}{x} & \text{if } x \neq 0 \\ k & \text{if } x = 0 \end{cases}$ is continuous at $x=0$ is:

A. 0

B. 1

C. 3

D. 2

Answer: D



Watch Video Solution

7. The derivative of $\sqrt{\log(\sin x)}$ w.r.t. x is

A. $\frac{1}{2\sqrt{\cot x}}$

B. $\frac{1}{2}\sec^2 x$

C. $\frac{\sec x}{2 \sin x \sqrt{\log(\tan x)}}$

D. None of these

Answer: C



Watch Video Solution

8. If $x = 4at$, $y = at^4$, then $\frac{dy}{dx}$ is equal to :

A. 4

B. 4a

C. 4at

D. t^3

Answer: D



Watch Video Solution

9. $\int \frac{dx}{x^2 + 2x + 2}$ equals :

A. $x \tan^{-1}(x + 1) + C$

B. $\tan^{-1}(x + 1) + C$

C. $(x + 1)\tan^{-1} + C$

D. $\tan^{-1} x + C$

Answer: B



Watch Video Solution

10. $\int_{-3}^3 (x + \sin x)dx$ is equal to :

A. 0

B. 1

C. 2

D. 3

Answer: A



[Watch Video Solution](#)

11. Find the $\frac{dy}{dx} + y \cot x = 2 \cos x$

A. $\cos x$

B. $\sin x$

C. $-\cos x$

D. $-\sin x$

Answer: B



[Watch Video Solution](#)

12. The Integrating Factor of the differentiate equation $\frac{dy}{dx} - 2y = 3x$ is

:

A. e^{2x}

B. e^{-2x}

C. e^x

D. $2x$

Answer: B



Watch Video Solution

13. Let \vec{a} and \vec{b} be two unit vectors and θ is the angle between them.

Then $\vec{a} + \vec{b}$ is a unit vector if :

A. $\theta = \frac{\pi}{4}$

B. $\theta = \frac{\pi}{3}$

$$C. \theta = \frac{\pi}{2}$$

$$D. \theta = \frac{2\pi}{3}$$

Answer: D



Watch Video Solution

14. Find $\left| \vec{a} \times \vec{b} \right|$, if $\vec{a} = \hat{i} - 7\hat{j} + 7\hat{k}$ and $\vec{b} = 3\hat{i} - 2\hat{j} + 2\hat{k}$

A. $7\sqrt{2}$

B. $10\sqrt{2}$

C. $19\sqrt{2}$

D. $21\sqrt{2}$

Answer: C



Watch Video Solution

15. Distance between plane defined by $3y + 4z + 10 = 0$ and the point $(7, 5, 0)$ is :

- A. 3 units
- B. 4 units
- C. 5 units
- D. 6 units

Answer: D



[Watch Video Solution](#)

16. Probability that A speaks truth is $\frac{4}{5}$. A coin is tossed. A reports that a head appears. The probability that actually there was head is

- A. $\frac{4}{5}$
- B. $\frac{1}{2}$
- C. $\frac{1}{5}$

D. $\frac{2}{5}$

Answer: A

 [Watch Video Solution](#)

Section A Fill In The Blanks

1. Fill in the blanks from the given options:

$$0.15, \frac{x+2}{3} = \frac{y-4}{5} = \frac{z+8}{6}, \frac{1}{4\sqrt{a+\sqrt{a+x}\sqrt{a+x}}}, -\frac{1}{3}, \frac{1}{2}(e-1)$$

, Symmetric, $\frac{19}{2}$ sq. units,

Let L be the set of all lines in a plane and R be the relation in L defined as

$R = \{(L_1, L_2) : L_1 \text{ is perpendicular to } L_2\}$. Show that R is.....

 [Watch Video Solution](#)

2. Fill in the blanks from the given options:

$$0.15, \frac{x+2}{3} = \frac{y-4}{5} = \frac{z+8}{6}, \frac{1}{4\sqrt{a+\sqrt{a+x}\sqrt{a+x}}}, -\frac{1}{3}, \frac{1}{2}(e-1)$$

, Symmetric, $\frac{19}{2}$ sq. units,

The area of the triangle, whose vertices are (3,1), (4, 3) and (-5,4) is

 [Watch Video Solution](#)

3. If $y = \sqrt{a + \sqrt{a + x}}$, then $\frac{dy}{dx}$ is.....

 [Watch Video Solution](#)

4. Fill in the blanks from the given options:

0.15, $\frac{x + 2}{3} = \frac{y - 4}{5} = \frac{z + 8}{6}$, $\frac{1}{4\sqrt{a + \sqrt{a + x}\sqrt{a + x}}}$, $-\frac{1}{3}$, $\frac{1}{2}(e - 1)$

, Symmetric, $\frac{19}{2}$ sq. units,

The slope of the normal to the curves $= 2x^2 + 3 \sin x$ at $x=0$ is.....

 [Watch Video Solution](#)

5. Fill in the blanks from the given options:

0.15, $\frac{x + 2}{3} = \frac{y - 4}{5} = \frac{z + 8}{6}$, $\frac{1}{4\sqrt{a + \sqrt{a + x}\sqrt{a + x}}}$, $-\frac{1}{3}$, $\frac{1}{2}(e - 1)$

, Symmetric, $\frac{19}{2}$ sq. units,

$\int_0^1 x e^{x^2} dx$ equals to.....

 [Watch Video Solution](#)

6. Fill in the blanks from the given options:

0.15, $\frac{x+2}{3} = \frac{y-4}{5} = \frac{z+8}{6}$, $\frac{1}{4\sqrt{a+\sqrt{a+x}}\sqrt{a+x}}$, $-\frac{1}{3}$, $\frac{1}{2}(e-1)$

, Symmetric, $\frac{19}{2}$ sq. units, 1

The degree of the differential equation $5x \left(\frac{dy}{dx} \right)^2 - \frac{d^2y}{dx^2} - 6y = \log x$ is.....

 [Watch Video Solution](#)

7. Find the cartesian equation of the line which passes through the point

$(-2, 4, -5)$ and parallel to the line given by $\frac{x+3}{3} = \frac{y-4}{5} = \frac{z+8}{6}$

 [Watch Video Solution](#)

8. Let X denote the number of hours you study during a randomly selected school day. The probability that X can take the values x , has the following form, where k is some unknown constant.:

$$P(X = x) = \begin{cases} 0.1 & \text{if } x = 0 \\ kx & \text{if } x = 1 \text{ or } 2 \\ k(5 - x) & \text{if } x = 3 \text{ or } 4 \\ 0 & \text{otherwise} \end{cases}$$

Find the value of k .

 [Watch Video Solution](#)

Section A True Or False

1. Write in the simplest form : $\tan^{-1}(\sin x/(1 + \cos x))$

 [Watch Video Solution](#)

2. Prove inverse of a matrix, if it exists, is unique.

 [Watch Video Solution](#)

3. Differentiate the following w.r.t.x.

$$\tan^{-1} \left(\frac{\sqrt{1 - \cos 2x}}{\sqrt{1 + \cos 2x}} \right)$$



Watch Video Solution

4. $\int e^x (\sin x + \cos x) dx$ is equal to :



Watch Video Solution

5. Find λ and μ if: $(2\hat{i} + 6\hat{j} + 27\hat{k}) \times (\hat{i} + \lambda\hat{j} + \mu\hat{k}) = \vec{0}$.



Watch Video Solution

6. Find the direction cosines of the unit vector perpendicular to the plane

$\vec{r} \cdot (6\hat{i} - 3\hat{j} - 2\hat{k}) + 1 = 0$ passing through the origin.




Watch Video Solution

7. If $P(A) = \frac{3}{5}$, $P(B) = \frac{1}{5}$ find $P(A \cap B)$ if A and B are independent events.

[Watch Video Solution](#)

8. The feasible solution for a LPP is shown in Fig. Let $Z = 4x + y$ be the objective function. Maximum value of Z is 130.



[Watch Video Solution](#)

1. Find a matrix A such that $2A - 3B + 5C = O$, where

$$B = \begin{bmatrix} -2 & 2 & 0 \\ 3 & 1 & 4 \end{bmatrix} \text{ and } C = \begin{bmatrix} 2 & 0 & -2 \\ 7 & 1 & 6 \end{bmatrix}$$



[Watch Video Solution](#)

2. For what value of x , the matrix $\begin{bmatrix} 1+x & 7 \\ 3-x & 8 \end{bmatrix}$ is a singular matrix?



[Watch Video Solution](#)

3. Find the equations of the tangent and normal to the curve given by :

$$x = a \sin^3 \theta, y = a \cos^3 \theta \text{ at a point, where } \theta = \frac{\pi}{4}$$



[Watch Video Solution](#)

4. Prove that $f(x) = 4x^3 - 6x^2 + 3x + 12$ is strictly increasing function on \mathbb{R} .

 [Watch Video Solution](#)

5. Evaluate, $\int \sin^3 x \cos^2 x \, dx$

 [Watch Video Solution](#)

6. Using integration, find the area of the circle $x^2 + y^2 = 4$

 [Watch Video Solution](#)

7. X and Y are two points with position vectors $3\vec{a} + \vec{b}$ and $\vec{a} - 3\vec{b}$ respectively. Write the position vector of point Z which divides the line segment XY in the ratio 2:1 externally.

 [Watch Video Solution](#)

8. Find λ if $(2\hat{i} + 6\hat{j} + 14\hat{k}) \times (\hat{i} - \lambda\hat{j} + 7\hat{k}) = \vec{0}$.

 [Watch Video Solution](#)

9. Find x and y , if $2 \begin{bmatrix} 1 & 3 \\ 0 & x \end{bmatrix} + \begin{bmatrix} y & 0 \\ 1 & 2 \end{bmatrix} = \begin{bmatrix} 5 & 6 \\ 1 & 8 \end{bmatrix}$

 [Watch Video Solution](#)

10. If $A = \begin{bmatrix} 1 & 5 \\ 7 & 12 \end{bmatrix}$ and $B = \begin{bmatrix} 9 & 1 \\ 7 & 8 \end{bmatrix}$, find a matrix C such that $3A + 5B + 2C$ is a null matrix.

 [Watch Video Solution](#)

11. Show that the normal at any point θ to the curve $x = a \cos \theta + a\theta \sin \theta$, $y = a \sin \theta - a\theta \cos \theta$ is at a constant distance from the origin.

 [Watch Video Solution](#)

12. Prove that the function f defined by $f(x) = x^2 - x + 1$ is neither increasing nor decreasing in $(-1, 1)$. Hence, find the intervals in which $f(x)$

is:

(i) strictly increasing , (ii) strictly decreasing.

 [Watch Video Solution](#)

13. Evaluate: $\int \frac{\cos(x + a)}{\sin(x + b)} dx$.

 [Watch Video Solution](#)

14. Using definite integrals, find the area of the circle $x^2 + y^2 = 9$

 [Watch Video Solution](#)

15. If $\vec{a} = \hat{i} + 2\hat{j} - 3\hat{k}$, $\vec{b} = 3\hat{i} - \hat{j} + 2\hat{k}$, show that $\vec{a} + \vec{b}$ and $\vec{a} - \vec{b}$ are perpendicular to each other.

 [Watch Video Solution](#)

16. Find a unit vector perpendicular to both \vec{a} and \vec{b} if

$$\vec{a} = 4\hat{i} + 3\hat{j} + 2\hat{k} \text{ and } \vec{b} = 2\hat{i} + 5\hat{j} - 3\hat{k}$$

 [Watch Video Solution](#)

17. If matrix $A = \begin{bmatrix} 1 & -1 \\ -1 & 1 \end{bmatrix}$ and $A^2 = kA$, then write the value of 'k'.

 [Watch Video Solution](#)

18. Find the adjoint of the following matrices :

$$\begin{bmatrix} 2 & -1 \\ 4 & 3 \end{bmatrix}$$

 [Watch Video Solution](#)

19. Find the equation of the tangent line to the curve $y = x^2 - 2x + 7$ which is parallel to the line $2x + y + 9 = 0$.

 [Watch Video Solution](#)

20. Evaluate $\int \frac{dx}{\sqrt{5 - 4x - 2x^2}}$

 [Watch Video Solution](#)

21. Integrate the following functions : $\frac{(x - 3)e^x}{(x - 1)^3}$.

 [Watch Video Solution](#)

22. Using definite integrals, find the area of the circle $x^2 + y^2 = 16$.

 [Watch Video Solution](#)

23. For what values of λ are the vectors $\vec{a} = 2\hat{i} + \lambda\hat{j} - \hat{k}$ and $\vec{b} = 4\hat{i} - 2\hat{j} - 2\hat{k}$ perpendicular to each other?

 [Watch Video Solution](#)

24. Find the angle between two vectors \vec{a} and \vec{b} with magnitudes 1 and 2 respectively and when $|\vec{a} \times \vec{b}| = \sqrt{3}$.

 [Watch Video Solution](#)

Section C

1. Simplify: $\tan^{-1}\left(\frac{\sqrt{1+x^2}-1}{x}\right)$

 [Watch Video Solution](#)

2.

Show

that:

$$\tan^{-1} \left[\frac{\sqrt{1+x^2} + \sqrt{1-x^2}}{\sqrt{1+x^2} - \sqrt{1-x^2}} \right] = \frac{\pi}{4} + \frac{1}{2} \cos^{-1} x^2, \quad -1 < x < 1$$

[Watch Video Solution](#)

3. Find $\frac{dy}{dx}$, if $y^x + x^y + x^x = a^b$.

[Watch Video Solution](#)

4. Find $\int \frac{x^2 dx}{(x \sin x + \cos x)^2}$

[Watch Video Solution](#)

5. Evaluate: $\int_1^4 [|x-1| + |x-2||x-4|] dx$.

[Watch Video Solution](#)

6. Solve $xdy - ydx = \sqrt{x^2 + y^2}dx$, given that $y = 0$ when $x=1$.



Watch Video Solution

7. Solve the differential equation:

$$\left[\frac{e^{-2\sqrt{x}}}{\sqrt{x}} - \frac{y}{\sqrt{x}} \right] \frac{dx}{dy} = 1, (x \neq 0)$$



Watch Video Solution

8. A, B and C throw a pair of dice in that order alternately till one of them gets a total of 9 and wins the game. Find their respective probabilities of winning, if A starts first.



Watch Video Solution

9. In a class, 5% of the boys and 10% of the girls have an IQ of more than 150. In the class 60% of the students are boys and rest girls. If a student

is selected at random and found to have an IQ of more than 150, find the probability that the student is a boy.

 [Watch Video Solution](#)

10. Compute $(AB)^{-1}$, where

$$A \begin{bmatrix} 5 & 0 & 4 \\ 2 & 3 & 2 \\ 1 & 2 & 1 \end{bmatrix}, B^{-1} = \begin{bmatrix} 1 & 3 & 3 \\ 1 & 4 & 3 \\ 1 & 3 & 4 \end{bmatrix}$$

 [Watch Video Solution](#)

11. Use matrix method to solve the following system of equations :

$$x - y + 2z = 7, 3x + 4y - 5z = -5, 2x - y + 3z = 12$$

 [Watch Video Solution](#)

12. Find the direction ratios of the normal to the plane, which passes through the points $(1, 0, 0)$ and $(0, 1, 0)$ and makes angle $\frac{\pi}{4}$ with the

plane $x + y = 3$. Also find the equation of the plane.

 [Watch Video Solution](#)

13. Show that the lines :
 $\frac{x+1}{3} = \frac{y+3}{5} = \frac{z+5}{7}$ and $\frac{x-2}{1} = \frac{y-4}{3} = \frac{z-6}{5}$ intersect each other. Also find their point of intersection.

 [Watch Video Solution](#)

14. Maximise $z = 22x + 44y$ subject to the constraints

$$x + y \geq 3, 3x + 8y \leq 24, x - y \geq 0, x, y \geq 0$$

 [Watch Video Solution](#)

15. Solve the following linear programming problem graphically: Minimise

$$Z = 200x + 500y \quad \text{subject to the constraints:}$$

$$x + 2y \geq 10, 3x + 4y \leq 24, x \geq 0, y \geq 0$$



Watch Video Solution

Section A

1. Let $R = \{(a, a^3) : a \text{ is a prime number less than } 5\}$ be a relation. Then the range of R is

A. $\{8,27\}$

B. $\{9,27\}$

C. $\{5,18\}$

D. $\{6,19\}$

Answer: A



Watch Video Solution

2. The principal value of $\cos^{-1}[\cos(680^\circ)]$ is:

A. 20°

B. 40°

C. 45°

D. None of these

Answer: B

 [Watch Video Solution](#)

3. Matrices A and B will be inverse of each other only if

A. $AB = BA$

B. $AB = BA = 0$

C. $AB=0, BA = I$

D. $AB = BA = I$

Answer: D

 [Watch Video Solution](#)

4. If $\begin{bmatrix} x - y & 2y \\ 2y + z & x + y \end{bmatrix} = \begin{bmatrix} 1 & 4 \\ 9 & 5 \end{bmatrix}$, then the value of $(x + y + z)$ is:

A. 5

B. 8

C. 10

D. 12

Answer: C



[Watch Video Solution](#)

5. For what value of x , is the matrix $A = \begin{bmatrix} 0 & 1 & 4 \\ -1 & 0 & 3 \\ x & -3 & 0 \end{bmatrix}$ a skew-symmetric matrix?

A. 2

B. 4

C. 3

D. 5

Answer: B



[Watch Video Solution](#)

6. The value of the constant k so that the function

$$f(x) = \begin{cases} \frac{1 - \cos 2x}{2x^2} & x \neq 0 \\ k & x = 0 \end{cases} \text{ is continuous at } x=0 \text{ is:}$$

A. 2

B. 3

C. 4

D. 1

Answer: D



[Watch Video Solution](#)

7. If $y = x^a$, $a \in R$ then $\frac{dy}{dx}$ is equal to :

A. $x^a \log x$

B. $x^a \log a$

C. $a^x \log a$

D. ax^{a-1}

Answer: D



[Watch Video Solution](#)

8. When $x = at^2$, $y = 3at$, then $\frac{dy}{dx}$ is:

A. $\frac{1}{t}$

B. $\frac{3}{t}$

C. $\frac{2}{t}$

D. None of these

Answer: A



Watch Video Solution

9. Choose the correct answer: $\int \frac{e^x(1+x)}{\cos^2(xe^x)} dx$ is equal to:

A. $-\cot(xe^x) + C$

B. $\tan(xe^x) + C$

C. $\tan(e^x) + C$

D. $\cot(e^x) + C$

Answer: B



Watch Video Solution

10. $\int_0^{\frac{\pi}{2}} \sqrt{1 - \sin 2x} dx$ is equal to

A. $\sqrt{21}$

B. $\sqrt{2} - 1$

C. $\sqrt{2} + 3$

D. $\sqrt{2} - 3$

Answer: B



Watch Video Solution

11. Which of the following is a homogeneous differential equation?

A. $(4x + 6y + 5)dy - (3y + 2x + 4)dx = 0$

B. $(xy)dx - (x^3 + y^3)dy = 0$

C. $(x^3 + 2y^2)dx + 2xydy = 0$

D. $y^2dx + (x^2 - xy - y^2)dy = 0$

Answer: D



Watch Video Solution

12. Find the particular solution of the differential equation

$$\log\left(\frac{dy}{dx}\right) = 3x + 4y \text{ given that } y = 0 \text{ when } x = 0$$

A. $\frac{1}{3}e^{-4y} = \frac{1}{4}e^{3x} - \frac{7}{12}$

B. $\frac{1}{4}e^{-4y} = \frac{1}{3}e^{3x} - \frac{7}{12}$

C. $\frac{1}{4}e^{-4y} = \frac{1}{5}e^{3x} - \frac{7}{12}$


D. $\frac{1}{5}e^{-4y} = \frac{1}{3}e^{3x} - \frac{7}{12}$

Answer: B



Watch Video Solution

13. In triangle ABC, which of the following is not true :



A. $\vec{AB} + \vec{BC} + \vec{CA} = \vec{0}$

B. $\vec{AB} + \vec{BC} - \vec{AC} = \vec{0}$

C. $\vec{AB} + \vec{BC} - \vec{CA} = \vec{0}$

D. $\vec{AB} - \vec{CB} + \vec{CA} = \vec{0}$

Answer: C



Watch Video Solution

14. If \vec{a} and \vec{b} are unit vectors and $(\sqrt{3}\vec{a} - \vec{b})$ is a unit vector, then the angle between \vec{a} and \vec{b} is:

A. 30°

B. 40°

C. 45°

D. 60°

Answer: A



[Watch Video Solution](#)

15. Distance between plane defined by $3x + 4z + 15 = 0$ and the point $(5, 2, 0)$ is:

A. 3 units

B. 4 units

C. 5 units

D. 6 units

Answer: B



[Watch Video Solution](#)

16. If A and B are two events such that $A \subset B$ and $P(B) \neq 0$, then which of the following is correct?

A. $P(A | B) = \frac{P(B)}{P(A)}$

B. $P(A | B) < P(A)$

C. $P(A | B) \geq P(A)$

D. None of these

Answer: C



[Watch Video Solution](#)

17. Let $A = \{0, 1, 2, 3\}$ and define a relation R on A as follows :

$$R = \{(0, 0), (0, 1), (0, 3), (1, 0), (1, 1), (2, 2), (3, 0), (3, 3)\}$$

Is R reflexive ? symmetric ? transitive ?

- A. Reflexive
- B. Symmetric
- C. Transitive
- D. None of these

Answer: C



[Watch Video Solution](#)

18. Principal value of $\sin^{-1}\left(\frac{1}{2}\right) + \cos^{-1}\left(-\frac{1}{2}\right)$ is

- A. $\frac{\pi}{2}$
- B. $-\frac{\pi}{2}$
- C. $\frac{3\pi}{2}$

D. None of these

Answer: A

 [Watch Video Solution](#)

19. If $A = \begin{bmatrix} \alpha & \beta \\ \gamma & -\alpha \end{bmatrix}$ is such that $A^2 = I$, then

a. $1 + \alpha^2 + \beta\gamma = 0$

b. $1 - \alpha^2 + \beta\gamma = 0$

c. $1 - \alpha^2 - \beta\gamma = 0$

d. $1 + \alpha^2 - \beta\gamma = 0$

A. $1 + \alpha^2 + \beta\gamma = 0$

B. $1 - \alpha^2 + \beta\gamma = 0$

C. $1 - \alpha^2 - \beta\gamma = 0$

D. $1 + \alpha^2 - \beta\gamma = 0$

Answer: C



Watch Video Solution

20. Solve the matrix equation : $(1 \quad x \quad 1) \begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 3 & 2 & 5 \end{pmatrix} \begin{pmatrix} 1 \\ -2 \\ 3 \end{pmatrix} = (0)$

A. $-\frac{9}{8}$

B. $-\frac{5}{8}$

C. $-\frac{4}{9}$

D. $-\frac{5}{7}$

Answer: A



Watch Video Solution

21. The matrix $A = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$ is

A. a unit matrix

B. a diagonal matrix

C. a symmetric matrix

D. a skew-symmetric matrix

Answer: D



[Watch Video Solution](#)

22. The value of k so that the function $f(x) = \begin{cases} kx^2 & , x \geq 1 \\ 4 & , x < 1 \end{cases}$ is

continuous at $x = 1$ is

A. 1

B. 2

C. 3

D. 4

Answer: D



[Watch Video Solution](#)

23. Derivative of $\cos^{-1}(\sin x)$ w.r.t. x equals

A. -1

B. 1

C. $\cos x$

D. $\sin x$

Answer: A



[Watch Video Solution](#)

24. Derivative of $\tan^{-1}\left(\frac{1 - \cos x}{\sin x}\right)$ w.r.t. x is

A. $\frac{1}{2}$

B. $\frac{1}{3}$

C. $\frac{1}{4}$

D. $\frac{1}{5}$

Answer: A



Watch Video Solution

25. $\int \frac{\cos 2x}{(\sin x + \cos x)^2} dx$ is equal to

A. $\frac{-1}{\sin x + \cos x} + C$

B. $\log|\sin x + \cos x| + C$

C. $\log|\sin x - \cos x| + C$

D. $\frac{1}{(\sin x + \cos x)^2}$

Answer: B



Watch Video Solution

26. $\int_1^2 (\log x^2) dx$ equals

A. $4 \log 2 - 2$

B. $4 \log 2 - 3$

C. $3 \log 2 - 2$

D. $3 \log 2 - 4$

Answer: A



Watch Video Solution

27. Solution of $x^5 \frac{dy}{dx} = -y^5$ is

A. $x^{-2} + y^{-2} = c$

B. $x^{-4} + y^{-4} = c$

C. $x^{-3} + y^{-3} = c$

D. $x^{-5} + y^{-5} = c$

Answer: B



Watch Video Solution

28. The general solution of differential equation : $ydy + xdx = 0$ is :

A. $x^2 + y^2 = c$

B. $y^2 = x^2 + c$

C. $y^2 = cx^2$

D. $x^2y^2 = c^2$

Answer: A



Watch Video Solution

29. Vector in the direction of vector $2\hat{i} - 3\hat{j} + 6\hat{k}$ which has magnitude 21

units is

A. $2(2\hat{i} - 3\hat{j} + 6\hat{k})$

B. $4(2\hat{i} - 3\hat{j} + 6\hat{k})$

C. $3(2\hat{i} - 3\hat{j} + 6\hat{k})$

D. $5(2\hat{i} - 3\hat{j} + 6\hat{k})$

Answer: C



Watch Video Solution

30. If $\sqrt{3}|\vec{a} \cdot \vec{b}| = |\vec{a} \times \vec{b}|$, then angle between \vec{a} and \vec{b} is

A. $\frac{\pi}{3}$

B. $\frac{\pi}{6}$

C. $\frac{\pi}{4}$

D. $\frac{\pi}{2}$

Answer: A



Watch Video Solution

31. The distance of the point (3, 4, 5) from the plane

$$\vec{r} \cdot (2\hat{i} - 5\hat{j} + 3\hat{k}) = 13$$

A. $\frac{8}{\sqrt{38}}$

B. $\frac{10}{\sqrt{38}}$

C. $\frac{12}{\sqrt{38}}$

D. $\frac{14}{\sqrt{38}}$

Answer: C

 [Watch Video Solution](#)

32. If $P\left(\frac{A}{B}\right) > P(A)$, then which of the following is correct ::

A. $P(B | A) < P(B)$

B. $P(A \cap B) < P(A) \cdot P(B)$

C. $P(B | A) > P(B)$

D. $P(B | A) = P(B)$

Answer: C

 [Watch Video Solution](#)

33. Range of $f(x) = \begin{cases} 1 & \text{if } x > 0 \\ 0 & \text{if } x = 0 \\ -1 & \text{if } x < 0 \end{cases}$ is.....

 [Watch Video Solution](#)

34. The matrix $A = \begin{bmatrix} 0 & 1 & -1 \\ -1 & 0 & 1 \\ 1 & -1 & 0 \end{bmatrix}$ is a matrix.

 [Watch Video Solution](#)

35. Derivative of $\tan^{-1}\left(\frac{\cos x}{1 + \sin x}\right)$ w.r.t. x is.....

 [Watch Video Solution](#)

36. The interval in which $y = x^2 e^{-x}$ is strictly increasing is

 [Watch Video Solution](#)

37. $\int \frac{\sin^{-1} x}{\sqrt{1-x^2}} dx = \dots\dots\dots$

 [Watch Video Solution](#)

38. I.F. of $\frac{dy}{dx} + \frac{y}{x} = e^x, (x > 0)$ is.....

 [Watch Video Solution](#)

39. Find the equation of the plane with intercept 3 on the y-axis and parallel to ZOx plane.

 [Watch Video Solution](#)

40. If $2P(A) = P(B) = \frac{7}{15}$ and $P(A/B) = \frac{2}{7}$ then find $P(A \cup B)$.

 [Watch Video Solution](#)

41. $\tan^{-1} \sqrt{3} - \sec^{-1}(-2)$ is equal to $-\frac{\pi}{3}$.

 [Watch Video Solution](#)

42. Value of $\begin{vmatrix} 42 & 1 & 6 \\ 28 & 7 & 4 \\ 14 & 3 & 2 \end{vmatrix}$ is 1?

 [Watch Video Solution](#)

43. If $y = \sin^{-1} x$, then prove that

$$\frac{d^2y}{dx^2} = \frac{x}{(1-x^2)^{\frac{3}{2}}}.$$

 [Watch Video Solution](#)

44. $\int [f(x)]^n f'(x) dx = \frac{[f(x)]^{n+1}}{n+1} + c, n \neq -1$

 [Watch Video Solution](#)

45. If $\vec{a} = 2\hat{i} - \hat{j} + \hat{k}$ and $\vec{c} = -\hat{i} + 3\hat{j} + \hat{k}$, then $\vec{a} \cdot (\vec{c} \times \vec{a})$ is 2.

 [Watch Video Solution](#)

46. Show that the D.C.'s of the perpendicular from origin to the plane $\vec{r} \cdot (-2\hat{i} - 3\hat{j} + 6\hat{k}) + 14 = 0$ are $\frac{2}{7}, \frac{3}{7}, -\frac{6}{7}$.

 [Watch Video Solution](#)

47. Let E and F be two events associated with the same random experiment, then E and F are said to be independent if $P(E \cap F) = P(E) \cdot P(F)$.

 [Watch Video Solution](#)

48. Corner points of the feasible region for an LPP are (0, 2), (3, 0), (6, 8) and (0, 5).

Let $F = 4x + 6y$ be the objective function.

The Maximum of F - Minimum of $F = 60$

 [Watch Video Solution](#)

Section A Fill In The Blanks

1. Fill in the blanks from the given options:

$$t^2, 2, -2, 0, \frac{2}{9}, x \sin x, \frac{1}{11} \tan^{11} x + \frac{1}{9} \tan^9 x + C, 1, 2, 4$$

Let $A = \{1, 2, 3\}$. Then number of equivalence relations containing (1,2) is

.....

 [Watch Video Solution](#)

2. Fill in the blanks from the given options:

$$t^2, 2, -2, 0, \frac{2}{9}, x \sin x, \frac{1}{11} \tan^{11} x + \frac{1}{9} \tan^9 x + C, 1, 2, 4$$

Value of $\begin{vmatrix} a-b & b-c & c-a \\ b-c & c-a & a-b \\ c-a & a-b & b-c \end{vmatrix}$ is.....

 [Watch Video Solution](#)

3. If $x = 3at$, $y = at^3$, then $\frac{dy}{dx}$ is.....

 [Watch Video Solution](#)

4. The slope of the tangent to the curve $y = 3 - x^2$ at $x = 1$ is.....

 [Watch Video Solution](#)

5. Fill in the blanks from the given options:

$t^2, 2, -2, 0, \frac{2}{9}, x \sin x, \frac{1}{11} \tan^{11} x + \frac{1}{9} \tan^9 x + C, 1, 2, 4$

$\int \tan^8 x \sec^4 x dx$

 [Watch Video Solution](#)

6. I.F of $x \frac{dy}{dx} + y - xy \cot x = 0 (x \neq 0)$ is.....

 [Watch Video Solution](#)

7. If the equations of a line AB are $\frac{3-x}{1} = \frac{y+2}{-2} = \frac{z-5}{4}$, then the direction ratios of a line parallel to AB are

 [Watch Video Solution](#)

8. If $P(A/B) = \frac{2}{5}$, $2P(A) = P(B) = \frac{5}{9}$, then $P(A \cap B)$ is.....

 [Watch Video Solution](#)

Section A True Or False

1. $2 \tan^{-1} x = \sin^{-1} \frac{2x}{1+x^2}$, $|x| \leq 1$.

 [Watch Video Solution](#)

2. prove for matrices A, B and C, $(ABC)' = C'B'A'$

 [Watch Video Solution](#)

3. Derivative of $\sin^{-1}(\cos x)$ w.r.t. x is 1

 [Watch Video Solution](#)

4. Prove $\int \frac{dx}{\sqrt{a^2 - x^2}} = \sin^{-1}\left(\frac{x}{a}\right) + c, a > 0.$

 [Watch Video Solution](#)

5. The value of p for which $\vec{a} = 3\hat{i} + 2\hat{j} + 9\hat{k}$ and $\vec{b} = \hat{i} + p\hat{j} + 3\hat{k}$ are parallel is .

 [Watch Video Solution](#)

6. The Cartesian equations of a line are $\frac{x-5}{3} = \frac{y+4}{7} = \frac{z-6}{2}$.


Vector equation for the line is

 [Watch Video Solution](#)

7. If $P(\text{not } A) = 0.4$, $P(A \cup B) = 0.75$ and A, B are given to be independent events, then the value of $P(B)$ is $\frac{3}{8}$.

 [Watch Video Solution](#)

8. The feasible solution for a LPP is shown in Fig. Let. $Z = 4x + 7y$ be the objective function. Maximum value of Z is 160:



[▶ Watch Video Solution](#)

Section C

1. Prove that

$$\cos^{-1} \frac{12}{13} + \cos^{-1} \frac{4}{3} = \tan^{-1} \frac{56}{33}$$

[▶ Watch Video Solution](#)

2. If $(x = a \sin(2t)(1 + \cos(2t)))$ and $(y = b \cos(2t)(1 - \cos(2t)))$, then show that $\left(\left(\frac{dy}{dx}\right)_{t=\frac{\pi}{4}} = \frac{b}{a}\right)$.

 [Watch Video Solution](#)

3. If $p^2 = a^2 \cos^2 \theta + b^2 \sin^2 \theta$, prove that $p + \left(\frac{d^2p}{d\theta^2}\right) = \frac{a^2b^2}{p^3}$

 [Watch Video Solution](#)

4. Integrate the function: $\frac{\sin^{-1} \sqrt{x} - \cos^{-1} \sqrt{x}}{\sin^{-1} \sqrt{x} + \cos^{-1} \sqrt{x}}, (x \in [0, 1])$

 [Watch Video Solution](#)

5. Using properties of definite integrals, show that

$$\int_0^{\pi/2} \frac{\sin^2 x}{\sin x + \cos x} dx = \frac{1}{\sqrt{2}} \log(\sqrt{2} + 1)$$

 [Watch Video Solution](#)

6. Solve the following differential equation

$$(x^2 - y^2)dx + 2xydy = 0 \text{ given that } y=1 \text{ when } x=1$$

 [Watch Video Solution](#)

7. An urn contains 2 white and 2 black balls. A ball is drawn at random. If it is white, it is not replaced into urn, otherwise it is replaced along with another ball of the same colour. The process is repeated, find the probability that the third ball drawn is black.

 [Watch Video Solution](#)

8. Three events A, B and C have probabilities $\frac{2}{5}$, $\frac{1}{3}$ and $\frac{1}{2}$, respectively. Given that $P(A \cap C) = \frac{1}{5}$ and $P(B \cap C) = \frac{1}{4}$, find the values of $P(C | B)$ and $P(A' \cap C')$

 [Watch Video Solution](#)

9. Prove that

$$\sin^{-1} \frac{5}{13} + \cos^{-1} \frac{3}{5} = \sin^{-1} \frac{63}{65}$$

 [Watch Video Solution](#)

10. Write the principle value of $\tan^{-1}(1) + \cos^{-1}\left(\frac{1}{2}\right)$.

 [Watch Video Solution](#)

11. Find $\frac{dy}{dx}$ when :

$$y = x^{\sin x - \cos x} + \frac{x^2 - 1}{x^2 + 1}$$

 [Watch Video Solution](#)

12. if $y = \sin^{-1}\left(\frac{5x + 12\sqrt{1-x^2}}{13}\right)$ then find $\frac{dy}{dx}$

 [Watch Video Solution](#)

13. Evaluate: $\int e^{-\frac{x}{2}} \frac{\sqrt{1 - \sin x}}{1 + \cos x} dx$



Watch Video Solution

14. Prove that: $\int_0^1 x(1-x)^n dx = \frac{1}{(n+1)(n+2)}$



Watch Video Solution

15. Solve the following differential equations

$$x dy - (y + 2x^2) dx = 0$$



Watch Video Solution

16. A bag contains 4 red and 4 black balls, another bag contains 2 red and 6 black balls. One of the two bags is selected at random and a ball is drawn from the bag which is found to be red. Find the probability that the ball is drawn from the first bag.

 [Watch Video Solution](#)

Section D

1. Solve the following system of linear equations by matrix method :

$$4x + 3y + z = 10, 3x - y + 2z = 8, x - 2y - 3z = 10.$$

 [Watch Video Solution](#)

2. Prove that

$$\begin{vmatrix} (b+c)^2 & a^2 & bc \\ (c+a)^2 & b^2 & ca \\ (a+b)^2 & c^2 & ab \end{vmatrix} = (a-b)(b-c)(c-a)(a+b+c)(a^2+b^2+c^2).$$

 [Watch Video Solution](#)

3. The plane $ax + by = 0$ is rotated through an angle α about its line of intersection with the plane $z = 0$. Show that the equation to the plane in

new position is $ax + by \pm z\sqrt{a^2 + b^2} \tan \alpha = 0$.



[Watch Video Solution](#)

4. Find the position vector of the foot of perpendicular and the perpendicular distance from the point P with position vector $2\hat{i} + 3\hat{j} + 4\hat{k}$ to the plane $\vec{r} \cdot (2\hat{i} + \hat{j} + 3\hat{k}) - 26 = 0$. Also find the image of P in the plane.



[Watch Video Solution](#)

5. Maximise and minimise $Z = 15x + 3y$ subject to constraints.

$$x + y \leq 8$$

$$2x + y \geq 8$$

$$x - 2y \geq 0$$

$$x, y \geq 0$$



[Watch Video Solution](#)

6. Maximize $z = 4x + 7y$ such that
 $x + 2y \leq 20, x + y \leq 15, x \geq 0, y \geq 0.$

 [Watch Video Solution](#)

7. Use product $\begin{bmatrix} 1 & -1 & 2 \\ 0 & 2 & -3 \\ 3 & -2 & 4 \end{bmatrix} \begin{bmatrix} -2 & 0 & 1 \\ 9 & 2 & -3 \\ 6 & 1 & -2 \end{bmatrix}$ to solve the system of equations $x - y + 2z = 1, 2y - 3z = 1, 3x - 2y + 4z = 2$

 [Watch Video Solution](#)

8. A school wants to award its students for regularity and hard work with a total cash award of Rs. 6,000. If three times the award money for hard work added to that given for regularity amounts to Rs. 11,000, represent the above situation algebraically and find the award money for each value, using matrix method.

 [Watch Video Solution](#)

9. Find the equation of plane passing through the points A(2, -1, 1), B(4, 3, 2) and C(6, 5, -2). Also prove that point $\left(5, -1, -\frac{25}{2}\right)$ lies on the plane given by points A, B and C.

 [View Text Solution](#)

10. Find the shortest distance between the lines :

$$\frac{x+1}{4} = \frac{y-3}{-6} = \frac{z+1}{1} \text{ and } \frac{x+3}{3} = \frac{y-5}{2} = \frac{z-7}{6}.$$

 [Watch Video Solution](#)

11. Maximize $z = 15x + 5y$, subject to the constraints

$$2x + 3y \leq 12$$

$$3x + 2y \leq 12$$

$$x \geq 0, y \geq 0$$

 [Watch Video Solution](#)

12. Solve the following problem graphically :

Minimise and Maximise $Z = 3x + 9y$

subject to the constraints :

$$x + 3y \leq 60$$

$$x + y \geq 10$$

$$x \leq y$$

$$x \geq 0, y \geq 0$$



[Watch Video Solution](#)