

MATHS

BOOKS - ACCURATE PUBLICATION

SAMPLE QUESTION PAPER-VII

Section A

1. Let $R = \{(a, a^3) : a \text{ is a prime number less than } 5\}$ be a relation.

Then the range of R is

A. $\{8,27\}$

B. $\{9,27\}$

C. $\{5,18\}$

D. $\{6,19\}$

Answer: A

 [Watch Video Solution](#)

2. The principal value of $\cos^{-1}[\cos(680^\circ)]$ is:

A. 20°

B. 40°

C. 45°

D. None of these

Answer: B

 [Watch Video Solution](#)

3. Matrices A and B will be inverse of each other only if

A. $AB = BA$

B. $AB = BA = 0$

C. $AB=0, BA = I$

D. $AB = BA = I$

Answer: D



Watch Video Solution

4. If $\begin{bmatrix} x - y & 2y \\ 2y + z & x + y \end{bmatrix} = \begin{bmatrix} 1 & 4 \\ 9 & 5 \end{bmatrix}$, then the value of $(x + y + z)$ is:

A. 5

B. 8

C. 10

D. 12

Answer: C



Watch Video Solution

5. For what value of x , is the matrix $A = \begin{bmatrix} 0 & 1 & 4 \\ -1 & 0 & 3 \\ x & -3 & 0 \end{bmatrix}$ a key-

symmetric matrix?

A. 2

B. 4

C. 3

D. 5

Answer: B



Watch Video Solution

6. The value of the constant k so that the function

$$f(x) = \begin{cases} \frac{1 - \cos 2x}{2x^2} & x \neq 0 \\ k & x = 0 \end{cases} \text{ is continuous at } x=0 \text{ is:}$$

A. 2

B. 3

C. 4

D. 1

Answer: D



Watch Video Solution

7. If $y = x^a$, $a \in R$ then $\frac{dy}{dx}$ is equal to :

A. $x^a \log x$

B. $x^a \log a$

C. $a^x \log a$

D. ax^{a-1}

Answer: D



Watch Video Solution

8. When $x = at^2$, $y = 3at$, then $\frac{dy}{dx}$ is:

A. $\frac{1}{t}$

B. $\frac{3}{t}$

C. $\frac{2}{t}$

D. None of these

Answer: A



Watch Video Solution

9. Choose the correct answer: $\int \frac{e^x(1+x)}{\cos^2(xe^x)} dx$ is equal to:

A. $-\cot(xe^x) + C$

B. $\tan(xe^x) + C$

C. $\tan(e^x) + C$

D. $\cot(e^x) + C$

Answer: B



Watch Video Solution

10. $\int_0^{\frac{\pi}{2}} \sqrt{1 - \sin 2x} dx$ is equal to

A. $\sqrt{2}1$

B. $\sqrt{2} - 1$

C. $\sqrt{2} + 3$

D. $\sqrt{2} - 3$

Answer: B

 [Watch Video Solution](#)

11. Which of the following is a homogeneous differential equation?

A. $(4x + 6y + 5)dy - (3y + 2x + 4)dx = 0$

B. $(xy)dx - (x^3 + y^3)dy = 0$

C. $(x^3 + 2y^2)dx + 2xydy = 0$

D. $y^2dx + (x^2 - xy - y^2)dy = 0$

Answer: D

 [Watch Video Solution](#)

12. Find the particular solution of the differential equation

$$\log\left(\frac{dy}{dx}\right) = 3x + 4y \text{ given that } y = 0 \text{ when } x = 0$$

A. $\frac{1}{3}e^{-4y} = \frac{1}{4}e^{3x} - \frac{7}{12}$

B. $\frac{1}{4}e^{-4y} = \frac{1}{3}e^{3x} - \frac{7}{12}$

C. $\frac{1}{4}e^{-4y} = \frac{1}{5}e^{3x} - \frac{7}{12}$


D. $\frac{1}{5}e^{-4y} = \frac{1}{3}e^{3x} - \frac{7}{12}$

Answer: B



Watch Video Solution

13. In triangle ABC, which of the following is not true :



A. $\vec{AB} + \vec{BC} + \vec{CA} = \vec{0}$

B. $\vec{AB} + \vec{BC} - \vec{AC} = \vec{0}$

C. $\vec{AB} + \vec{BC} - \vec{CA} = \vec{0}$

D. $\vec{AB} - \vec{CB} + \vec{CA} = \vec{0}$

Answer: C

 Watch Video Solution

14. If \vec{a} and \vec{b} are unit vectors and $(\sqrt{3}\vec{a} - \vec{b})$ is a unit vector, then the angle between \vec{a} and \vec{b} is:

A. 30°

B. 40°

C. 45°

D. 60°

Answer: A



Watch Video Solution

15. Distance between plane defined by $3x + 4z + 15 = 0$ and the point $(5, 2, 0)$ is:

A. 3 units

B. 4 units

C. 5 units

D. 6 units

Answer: B



[Watch Video Solution](#)

16. If A and B are two events such that $A \subset B$ and $P(B) \neq 0$, then which of the following is correct?

A. $P(A | B) = \frac{P(B)}{P(A)}$

B. $P(A | B) < P(A)$

C. $P(A | B) \geq P(A)$

D. None of these

Answer: C



[Watch Video Solution](#)

17. Let $A = \{0, 1, 2, 3\}$ and define a relation R on A as follows :

$$R = \{(0, 0), (0, 1), (0, 3), (1, 0), (1, 1), (2, 2), (3, 0), (3, 3)\}$$

Is R reflexive ? symmetric ? transitive ?

A. Reflexive

B. Symmetric

C. Transitive

D. None of these

Answer: C

 [Watch Video Solution](#)

18. Principal value of $\sin^{-1}\left(\frac{1}{2}\right) + \cos^{-1}\left(-\frac{1}{2}\right)$ is

A. $\frac{\pi}{2}$

B. $-\frac{\pi}{2}$

C. $\frac{3\pi}{2}$

D. None of these

Answer: A

 [Watch Video Solution](#)

19. If $A = \begin{bmatrix} \alpha & \beta \\ \gamma & -\alpha \end{bmatrix}$ is such that $A^2 = I$, then

a. $1 + \alpha^2 + \beta\gamma = 0$

b. $1 - \alpha^2 + \beta\gamma = 0$

c. $1 - \alpha^2 - \beta\gamma = 0$

d. $1 + \alpha^2 - \beta\gamma = 0$

A. $1 + \alpha^2 + \beta\gamma = 0$

B. $1 - \alpha^2 + \beta\gamma = 0$

$$C. 1 - \alpha^2 - \beta\gamma = 0$$

$$D. 1 + \alpha^2 - \beta\gamma = 0$$

Answer: C



Watch Video Solution

20. Solve the matrix equation :

$$(1 \ x \ 1) \begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 3 & 2 & 5 \end{pmatrix} \begin{pmatrix} 1 \\ -2 \\ 3 \end{pmatrix} = (0)$$

A. $-\frac{9}{8}$

B. $-\frac{5}{8}$

C. $-\frac{4}{9}$

D. $-\frac{5}{7}$

Answer: A



Watch Video Solution

21. The matrix $A = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$ is

- A. a unit matrix
- B. a diagonal matrix
- C. a symmetric matrix
- D. a skew-symmetric matrix

Answer: D

[Watch Video Solution](#)

22. The value of k so that the function $f(x) = \begin{cases} kx^2 & , x \geq 1 \\ 4 & , x < 1 \end{cases}$ is continuous at $x = 1$ is

- A. 1

B. 2

C. 3

D. 4

Answer: D



Watch Video Solution

23. Derivative of $\cos^{-1}(\sin x)$ w.r.t. x equals

A. -1

B. 1

C. $\cos x$

D. $\sin x$

Answer: A



Watch Video Solution

24. Derivative of $\tan^{-1}\left(\frac{1 - \cos x}{\sin x}\right)$ w.r.t. x is

A. $\frac{1}{2}$

B. $\frac{1}{3}$

C. $\frac{1}{4}$

D. $\frac{1}{5}$

Answer: A



Watch Video Solution

25. $\int \frac{\cos 2x}{(\sin x + \cos x)^2} dx$ is equal to

A. $\frac{-1}{\sin x + \cos x} + C$

B. $\log|\sin x + \cos x| + C$

C. $\log|\sin x - \cos x| + C$

D. $\frac{1}{(\sin x + \cos x)^2}$

Answer: B

 [Watch Video Solution](#)

26. $\int_1^2 (\log x^2) dx$ equals

A. $4 \log 2 - 2$

B. $4 \log 2 - 3$

C. $3 \log 2 - 2$

D. $3 \log 2 - 4$

Answer: A

 [Watch Video Solution](#)

27. Solution of $x^5 \frac{dy}{dx} = -y^5$ is

A. $x^{-2} + y^{-2} = c$

B. $x^{-4} + y^{-4} = c$

C. $x^{-3} + y^{-3} = c$

D. $x^{-5} + y^{-5} = c$

Answer: B



[Watch Video Solution](#)

28. The general solution of differential equation : $ydy + xdx = 0$ is

:

A. $x^2 + y^2 = c$

B. $y^2 = x^2 + c$

C. $y^2 = cx^2$

D. $x^2y^2 = c^2$

Answer: A



Watch Video Solution

29. Vector in the direction of vector $2\hat{i} - 3\hat{j} + 6\hat{k}$ which has magnitude 21 units is

A. $2(2\hat{i} - 3\hat{j} + 6\hat{k})$

B. $4(2\hat{i} - 3\hat{j} + 6\hat{k})$

C. $3(2\hat{i} - 3\hat{j} + 6\hat{k})$

D. $5(2\hat{i} - 3\hat{j} + 6\hat{k})$

Answer: C



Watch Video Solution

30. If $\sqrt{3}|\vec{a} \cdot \vec{b}| = |\vec{a} \times \vec{b}|$, then angle between \vec{a} and \vec{b} is

A. $\frac{\pi}{3}$

B. $\frac{\pi}{6}$

C. $\frac{\pi}{4}$

D. $\frac{\pi}{2}$

Answer: A



Watch Video Solution

31. The distance of the point (3, 4, 5) from the plane

$$\vec{r} \cdot (2\hat{i} - 5\hat{j} + 3\hat{k}) = 13$$

A. $\frac{8}{\sqrt{38}}$

B. $\frac{10}{\sqrt{38}}$

C. $\frac{12}{\sqrt{38}}$

D. $\frac{14}{\sqrt{38}}$

Answer: C



Watch Video Solution

32. If $P\left(\frac{A}{B}\right) > P(A)$, then which of the following is correct : :

A. $P(B | A) < P(B)$

B. $P(A \cap B) < P(A) \cdot P(B)$

C. $P(B | A) > P(B)$

D. $P(B | A) = P(B)$

Answer: C



Watch Video Solution

33. Range of $f(x) = \begin{cases} 1 & \text{if } x > 0 \\ 0 & \text{if } x = 0 \text{ is.....} \\ -1 & \text{if } x < 0 \end{cases}$

 [Watch Video Solution](#)

34. The matrix $A = \begin{bmatrix} 0 & 1 & -1 \\ -1 & 0 & 1 \\ 1 & -1 & 0 \end{bmatrix}$ is a matrix.

 [Watch Video Solution](#)

35. Derivative of $\tan^{-1}\left(\frac{\cos x}{1 + \sin x}\right)$ wr.t. x is.....

 [Watch Video Solution](#)

36. The interval in which $y = x^2 e^{-x}$ is strictly increasing is

 [Watch Video Solution](#)

37. $\int \frac{\sin^{-1} x}{\sqrt{1-x^2}} dx = \dots\dots\dots$

 [Watch Video Solution](#)

38. I.F. of $\frac{dy}{dx} + \frac{y}{x} = e^x$, ($x > 0$) is.....

 [Watch Video Solution](#)

39. Find the equation of the plane with intercept 3 on the y-axis and parallel to ZOY plane.

 [Watch Video Solution](#)

40. If $2P(A) = P(B) = \frac{7}{15}$ and $P(A/B) = \frac{2}{7}$ then find $P(A \cup B)$.

 [Watch Video Solution](#)

41. $\tan^{-1} \sqrt{3} - \sec^{-1}(-2)$ is equal to $-\frac{\pi}{3}$.

 [Watch Video Solution](#)

42. Value of $\begin{vmatrix} 42 & 1 & 6 \\ 28 & 7 & 4 \\ 14 & 3 & 2 \end{vmatrix}$ is 1?

 [Watch Video Solution](#)

43. If $y = \sin^{-1} x$, then prove that

$$\frac{d^2y}{dx^2} = \frac{x}{(1-x^2)^{\frac{3}{2}}}$$

 [Watch Video Solution](#)

$$44. \int [f(x)]^n f'(x) dx = \frac{[f(x)]^{n+1}}{n+1} + c, n \neq -1$$

 [Watch Video Solution](#)

45. If $\vec{a} = 2\hat{i} - \hat{j} + \hat{k}$ and $\vec{c} = -\hat{i} + 3\hat{j} + \hat{k}$, then $\vec{a} \cdot (\vec{c} \times \vec{a})$ is 2.

 [Watch Video Solution](#)

46. Show that the D.C.'s of the perpendicular from origin to the plane $\vec{r} \cdot (-2\hat{i} - 3\hat{j} + 6\hat{k}) + 14 = 0$ are $\frac{2}{7}, \frac{3}{7}, -\frac{6}{7}$.

 [Watch Video Solution](#)

47. Let E and F be two events associated with the same random experiment, then E and F are said to be independent if

$$P(E \cap F) = P(E) \cdot P(F).$$



[Watch Video Solution](#)

48. Corner points of the feasible region for an LPP are $(0, 2)$, $(3, 0)$, $(6, 8)$ and $(0, 5)$.

Let $F = 4x + 6y$ be the objective function.

The Maximum of F - Minimum of $F = 60$



[Watch Video Solution](#)

Section A Fill In The Blanks

1. Fill in the blanks from the given options:

$$t^2, 2, -2, 0, \frac{2}{9}, x \sin x, \frac{1}{11} \tan^{11} x + \frac{1}{9} \tan^9 x + C, 1, 2, 4$$

Let $A = \{1, 2, 3\}$. Then number of equivalence relations containing $(1, 2)$ is

 [Watch Video Solution](#)

2. Fill in the blanks from the given options:

$$t^2, 2, -2, 0, \frac{2}{9}, x \sin x, \frac{1}{11} \tan^{11} x + \frac{1}{9} \tan^9 x + C, 1, 2, 4$$

Value of $\begin{vmatrix} a-b & b-c & c-a \\ b-c & c-a & a-b \\ c-a & a-b & b-c \end{vmatrix}$ is.....

 [Watch Video Solution](#)

3. If $x = 3at$, $y = at^3$, then $\frac{dy}{dx}$ is.....

 [Watch Video Solution](#)

4. The slope of the tangent to the curve $y = 3 - x^2$ at $x = 1$ is.....

 [Watch Video Solution](#)

5. Fill in the blanks from the given options:

$$t^2, 2, -2, 0, \frac{2}{9}, x \sin x, \frac{1}{11} \tan^{11} x + \frac{1}{9} \tan^9 x + C, 1, 2, 4$$

$$\int \tan^8 x \sec^4 x dx \dots\dots\dots$$

 [Watch Video Solution](#)

6. I.F of $x \frac{dy}{dx} + y - xy \cot x = 0 (x \neq 0)$ is.....

 [Watch Video Solution](#)

7. If the equations of a line AB are $\frac{3-x}{1} = \frac{y+2}{-2} = \frac{z-5}{4}$, then the direction ratios of a line parallel to AB are

 [Watch Video Solution](#)

8. If $P(A/B) = \frac{2}{5}$, $2P(A) = P(B) = \frac{5}{9}$, then $P(A \cap B)$ is.....



 Watch Video Solution

Section A True Or False

1. $2 \tan^{-1} x = \sin^{-1} \frac{2x}{1+x^2}, |x| \leq 1.$

 Watch Video Solution

2. prove for matrices A, B and C, $(ABC)' = C'B'A'$

 Watch Video Solution

3. Derivative of $\sin^{-1}(\cos x)$ w.r.t. x is 1

 Watch Video Solution

4. Prove $\int \frac{dx}{\sqrt{a^2 - x^2}} = \sin^{-1}\left(\frac{x}{a}\right) + c, a > 0.$

 [Watch Video Solution](#)

5. The value of p for which $\vec{a} = 3\hat{i} + 2\hat{j} + 9\hat{k}$ and $\vec{b} = \hat{i} + p\hat{j} + 3\hat{k}$ are parallel is .

 [Watch Video Solution](#)

6. The Cartesian equations of a line are $\frac{x - 5}{3} = \frac{y + 4}{7} = \frac{z - 6}{2}.$


Vector equation for the line is

 [Watch Video Solution](#)

7. If $P(\text{not } A) = 0.4$, $P(A \cup B) = 0.75$ and A, B are given to be independent events, then the value of $P(B)$ is $\frac{3}{8}.$

[Watch Video Solution](#)

8. The feasible solution for a LPP is shown in Fig. Let. $Z = 4x + 7y$ be the objective function. Maximum value of Z is 160:



[Watch Video Solution](#)

1. Find x and y , if $2 \begin{bmatrix} 1 & 3 \\ 0 & x \end{bmatrix} + \begin{bmatrix} y & 0 \\ 1 & 2 \end{bmatrix} = \begin{bmatrix} 5 & 6 \\ 1 & 8 \end{bmatrix}$

 [Watch Video Solution](#)

2. If $A = \begin{bmatrix} 1 & 5 \\ 7 & 12 \end{bmatrix}$ and $B = \begin{bmatrix} 9 & 1 \\ 7 & 8 \end{bmatrix}$, find a matrix C such that $3A + 5B + 2C$ is a null matrix.

 [Watch Video Solution](#)

3. Show that the normal at any point θ to the curve $x = a \cos \theta + a\theta \sin \theta$, $y = a \sin \theta - a\theta \cos \theta$ is at a constant distance from the origin.

 [Watch Video Solution](#)

4. Prove that the function f defined by $f(x) = x^2 - x + 1$ is neither increasing nor decreasing in $(-1, 1)$. Hence, find the intervals in which

$f(x)$ is:

(i) strictly increasing , (ii) strictly decreasing.

 [Watch Video Solution](#)

5. Evaluate: $\int \frac{\cos(x + a)}{\sin(x + b)} dx$.

 [Watch Video Solution](#)

6. Using definite integrals, find the area of the circle $x^2 + y^2 = 9$

 [Watch Video Solution](#)

7. If $\vec{a} = \hat{i} + 2\hat{j} - 3\hat{k}$, $\vec{b} = 3\hat{i} - \hat{j} + 2\hat{k}$, show that $\vec{a} + \vec{b}$ and $\vec{a} - \vec{b}$ are perpendicular to each other.

 [Watch Video Solution](#)

8. Find a unit vector perpendicular to both \vec{a} and \vec{b} if

$$\vec{a} = 4\hat{i} + 3\hat{j} + 2\hat{k} \text{ and } \vec{b} = 2\hat{i} + 5\hat{j} - 3\hat{k}$$

 [Watch Video Solution](#)

9. If matrix $A = \begin{bmatrix} 1 & -1 \\ -1 & 1 \end{bmatrix}$ and $A^2 = kA$, then write the value of 'k'.

 [Watch Video Solution](#)

10. Find the adjoint of the following matrices :

$$\begin{bmatrix} 2 & -1 \\ 4 & 3 \end{bmatrix}$$

 [Watch Video Solution](#)

11. Find the equation of the tangent line to the curve $y = x^2 - 2x + 7$ which is parallel to the line $2x + y + 9 = 0$.

 [Watch Video Solution](#)

12. Evaluate $\int \frac{dx}{\sqrt{5 - 4x - 2x^2}}$

 [Watch Video Solution](#)

13. Integrate the following functions : $\frac{(x - 3)e^x}{(x - 1)^3}$.

 [Watch Video Solution](#)

 Watch Video Solution

14. Using definite integrals, find the area of the circle $x^2 + y^2 = 16$.

 Watch Video Solution

15. For what values of λ are the vectors $\vec{a} = 2\hat{i} + \lambda\hat{j} - \hat{k}$ and $\vec{b} = 4\hat{i} - 2\hat{j} - 2\hat{k}$ perpendicular to each other?

 Watch Video Solution

16. Find the angle between two vectors \vec{a} and \vec{b} with magnitudes 1 and 2 respectively and when $\left| \vec{a} \times \vec{b} \right| = \sqrt{3}$.

 Watch Video Solution

1. Prove that

$$\cos^{-1} \frac{12}{13} + \cos^{-1} \frac{4}{3} = \tan^{-1} \frac{56}{33}$$

 [Watch Video Solution](#)

2. If $(x = a \sin(2t)(1 + \cos(2t)))$ and $(y = b \cos(2t)(1 - \cos(2t)))$,

then show that $\left(\left(\frac{dy}{dx} \right)_{t=\frac{\pi}{4}} = \frac{b}{a} \right)$.

 [Watch Video Solution](#)

3. If $p^2 = a^2 \cos^2 \theta + b^2 \sin^2 \theta$, prove that $p + \left(\frac{d^2p}{d\theta^2} \right) = \frac{a^2 b^2}{p^3}$

 [Watch Video Solution](#)

4. Integrate the function: $\frac{\sin^{-1} \sqrt{x} - \cos^{-1} \sqrt{x}}{\sin^{-1} \sqrt{x} + \cos^{-1} \sqrt{x}}, (x \in [0, 1])$

 [Watch Video Solution](#)

5. Using properties of definite integrals, show that

$$\int_0^{\pi/2} \frac{\sin^2 x}{\sin x + \cos x} dx = \frac{1}{\sqrt{2}} \log(\sqrt{2} + 1)$$

 [Watch Video Solution](#)

6. Solve the following differential equation

$$(x^2 - y^2) dx + 2xy dy = 0 \text{ given that } y=1 \text{ when } x=1$$

 [Watch Video Solution](#)

7. An urn contains 2 white and 2 black balls. A ball is drawn at random. If it is white, it is not replaced into urn, otherwise it is replaced along with another ball of the same colour. The process is repeated, find the probability that the third ball drawn is black.

 [Watch Video Solution](#)

8. Three events A, B and C have probabilities $\frac{2}{5}$, $\frac{1}{3}$ and $\frac{1}{2}$, respectively. Given that $P(A \cap C) = \frac{1}{5}$ and $P(B \cap C) = \frac{1}{4}$, find the values of $P(C | B)$ and $P(A' \cap C')$

 [Watch Video Solution](#)

9. Prove that

$$\sin^{-1} \frac{5}{13} + \cos^{-1} \frac{3}{5} = \sin^{-1} \frac{63}{65}$$

 [Watch Video Solution](#)

10. Write the principle value of $\tan^{-1}(1) + \cos^{-1}\left(\frac{1}{2}\right)$.

 [Watch Video Solution](#)

11. Find $\frac{dy}{dx}$ when :

$$y = x^{\sin x - \cos x} + \frac{x^2 - 1}{x^2 + 1}$$

 [Watch Video Solution](#)

12. if $y = \sin^{-1}\left(\frac{5x + 12\sqrt{1-x^2}}{13}\right)$ then find $\frac{dy}{dx}$

 [Watch Video Solution](#)

13. Evaluate: $\int e^{-\frac{x}{2}} \frac{\sqrt{1 - \sin x}}{1 + \cos x} dx$

 [Watch Video Solution](#)

 Watch Video Solution

14. Prove that : $\int_0^1 x(1-x)^n dx = \frac{1}{(n+1)(n+2)}$

 Watch Video Solution

15. Solve the following differential equations

$$x dy - (y + 2x^2) dx = 0$$

 Watch Video Solution

16. A bag contains 4 red and 4 black balls, another bag contains 2 red and 6 black balls. One of the two bags is selected at random and a ball is drawn from the bag which is found to be red. Find the probability that the ball is drawn from the first bag.

 Watch Video Solution

1. Solve the following system of linear equations by matrix method :

$$4x + 3y + z = 10, 3x - y + 2z = 8, x - 2y - 3z = 10.$$

 [Watch Video Solution](#)

2. Prove that

$$\begin{vmatrix} (b+c)^2 & a^2 & bc \\ (c+a)^2 & b^2 & ca \\ (a+b)^2 & c^2 & ab \end{vmatrix} = (a-b)(b-c)(c-a)(a+b+c)(a^2 + b^2 + c^2)$$

 [Watch Video Solution](#)

3. The plane $ax + by = 0$ is rotated through an angle α about its line of intersection with the plane $z = 0$. Show that the equation to the plane in new position is $ax + by \pm z\sqrt{a^2 + b^2} \tan \alpha = 0$.



Watch Video Solution

4. Find the position vector of the foot of perpendicular and the perpendicular distance from the point P with position vector $2\hat{i} + 3\hat{j} + 4\hat{k}$ to the plane $\vec{r} \cdot (2\hat{i} + \hat{j} + 3\hat{k}) - 26 = 0$. Also find the image of P in the plane.



Watch Video Solution

5. Maximise and minimise $Z = 15x + 3y$ subject to constraints.

$$x + y \leq 8$$

$$2x + y \geq 8$$

$$x - 2y \geq 0$$

$$x, y \geq 0$$



Watch Video Solution

6. Maximize $z = 4x + 7y$ such that

$$x + 2y \leq 20, x + y \leq 15, x \geq 0, y \geq 0.$$

 [Watch Video Solution](#)

7. Use product $\begin{bmatrix} 1 & -1 & 2 \\ 0 & 2 & -3 \\ 3 & -2 & 4 \end{bmatrix} \begin{bmatrix} -2 & 0 & 1 \\ 9 & 2 & -3 \\ 6 & 1 & -2 \end{bmatrix}$ to solve the system of equations $x - y + 2z = 1$, $2y - 3z = 1$, $3x - 2y + 4z = 2$

 [Watch Video Solution](#)

8. A school wants to award its students for regularity and hard work with a total cash award of Rs. 6,000. If three times the award money for hard work added to that given for regularity amounts to Rs. 11,000, represent the above situation algebraically and find the award money for each value, using matrix method.

 [Watch Video Solution](#)

9. Find the equation of plane passing through the points A(2, -1, 1), B(4, 3, 2) and C(6, 5, -2). Also prove that point $\left(5, -1, -\frac{25}{2}\right)$ lies on the plane given by points A, B and C.

 [View Text Solution](#)

10. Find the shortest distance between the lines :

$$\frac{x+1}{4} = \frac{y-3}{-6} = \frac{z+1}{1} \text{ and } \frac{x+3}{3} = \frac{y-5}{2} = \frac{z-7}{6}.$$

 [Watch Video Solution](#)

11. Maximize $z = 15x + 5y$, subject to the constraints

$$2x + 3y \leq 12$$

$$3x + 2y \leq 12$$

$$x \geq 0, y \geq 0$$

[Watch Video Solution](#)

12. Solve the following problem graphically :

Minimise and Maximise $Z = 3x + 9y$

subject to the constraints :

$$x + 3y \leq 60$$

$$x + y \geq 10$$

$$x \leq y$$

$$x \geq 0, y \geq 0$$

[Watch Video Solution](#)