



PHYSICS

BOOKS - PSEB

MAGNETISM AND MATTER

Exercise

1. Answer the following questions regarding earth's magnetism: A vector needs three quantities for its specification. Name the three

Independent quantities conventionally used to specify the earth's magnetic field.



[Watch Video Solution](#)

2. Answer the following questions regarding earth's magnetism: The angle of dip at a location in southern India is about 18° . Would you expect a greater or smaller dip angle in Britain?



[Watch Video Solution](#)

3. Answer the following questions regarding earth's magnetism: f you made a map of magnetic field lines at Melbourne In Australia, would the lines seem to go Into the ground or come out of the ground?



[Watch Video Solution](#)

4. Answer the following questions regarding earth's magnetism: In which direction would a compass free point to, if located right on the geographic north or south pole?



[Watch Video Solution](#)

5. Answer the following questions regarding earth's magnetism: The earth's field, it is claimed, roughly approximates the field due to a dipole of magnetic moment $8 \times 10^{22} \text{ J T}^{-1}$ located at its centre. Check the order of magnitude of this number in some way.



[Watch Video Solution](#)

6. Answer the following questions regarding earth's magnetism: Geologists claim that besides the main magnetic N-S poles, there are several local poles on the earth's surface oriented in different directions. How is such a thing possible at all?



[Watch Video Solution](#)

7. Answer the following questions: The earth's magnetic field varies from point to point in

space. Does it also change with time? If so, on what time scale does it change appreciably?



Watch Video Solution

8. Answer the following questions: The earth's core is known to contain Iron. Yet geologists do not regard this as a source of the earth's magnetism. Why?



Watch Video Solution

9. Answer the following questions: The charged currents in the outer conducting regions of the earth's core are thought to be responsible for earth's magnetism. What might be the 'battery' (i.e., the source of energy) to sustain these currents?



Watch Video Solution

10. Answer the following questions: The earth may have even reversed the direction of its

field several times during its history of 4 to 5 billion years. How can geologists know about the earth's field in such distant past?



[Watch Video Solution](#)

11. Answer the following questions: The earth's field departs from its dipole shape substantially at large distances (greater than about 30,000 km). What agencies may be responsible for this distortion?



[Watch Video Solution](#)

12. Answer the following questions: Interstellar space has an extremely weak magnetic field of the order of 10^{-12} T. Can such a weak field be of any significant consequence? Explain.



Watch Video Solution

13. A short bar magnet placed with its axis at 30° with a uniform external magnetic field of 0.25 T experiences a torque of magnitude

equal to $4.5 \times 10^{-2} J$. What is the magnitude of magnetic moment of the magnet?



[Watch Video Solution](#)

14. A short bar magnet of magnetic moment $m = 0.32 JT^{-1}$ is placed in a uniform magnetic field of 0.15 T. If the bar is free to rotate in the plane of the field, which orientation would correspond to its stable equilibrium? What is the potential energy of the magnet in this case?



[Watch Video Solution](#)

15. A short bar magnet of magnetic moment $m = 0.32 \text{ JT}^{-1}$ is placed in a uniform magnetic field of 0.15 T. If the bar is free to rotate in the plane of the field, which orientation would correspond to its unstable equilibrium? What is the potential energy of the magnet in this case?



[Watch Video Solution](#)

16. A closely wound solenoid of 800 turns and area of cross section $2.5 \times 10^{-4} m^2$ carries a current of 3.0 A. Explain the sense in which the solenoid acts like a bar magnet. What is its associated magnetic moment?



Watch Video Solution

17. A closely wound solenoid of 800 turns and area of cross section $2.5 \times 10^{-4} m^2$ carries a current of 3.0 A. If the solenoid is free to turn

about the vertical direction and a uniform horizontal magnetic field of 0.25 T is applied, what is the magnitude of torque on the solenoid when its axis makes an angle of 30° with the direction of applied field?



[Watch Video Solution](#)

18. A bar magnet of magnetic moment $1.5JT^{-1}$ lies aligned with the direction of a uniform magnetic field of 0.22 T. What is the amount of work required by an external

torque to turn the magnet so as to align its magnetic moment: normal to the field direction?



[Watch Video Solution](#)

19. A bar magnet of magnetic moment $1.5JT^{-1}$ lies aligned with the direction of a uniform magnetic field of 0.22 T. What is the amount of work required by an external torque to turn the magnet so as to align its

magnetic moment: opposite to the field direction?



[Watch Video Solution](#)

20. A bar magnet of magnetic moment $1.5JT^{-1}$ lies aligned with the direction of a uniform magnetic field of 0.22 T. Consider orientations (i) normal to the field direction, (ii) opposite to the field direction? What is the torque on the magnet in cases (i) and (ii)?



[Watch Video Solution](#)

21. A closely wound solenoid of 2000 turns and area of cross-section $1.6 \times 10^{-1} \text{m}^2$. carrying a current of 4.0 A, is suspended through its centre allowing it to turn in a horizontal plane. What is the magnetic moment associated with the solenoid?



Watch Video Solution

22. A closely wound solenoid of 2000 turns and area of cross-section $1.6 \times 10^{-1} \text{m}^2$.

carrying a current of 4.0 A, is suspended through its centre allowing it to turn in a horizontal plane. What is the force and torque on the solenoid if a uniform horizontal magnetic field of 7.5×10^2 T is set up at an angle of 30° with the axis of the solenoid?



[Watch Video Solution](#)

23. A circular coil of 16 turns and radius 10 cm carrying a current of 0.75 A rests with its plane normal to an external field of magnitude

5.0×10^{-2} T. The coil is free to turn about an axis in its plane perpendicular to the field direction. When the coil is turned slightly and released, it oscillates about its stable equilibrium with a frequency of 2.0s^{-1} . What is the moment of inertia of the coil about its axis of rotation?



[Watch Video Solution](#)

24. A magnetic needle free to rotate in a vertical plane parallel to the magnetic

meridian has its north tip pointing down at 22° with the horizontal. The horizontal component of the earth's magnetic field at the place is known to be 0.35 G. Determine the magnitude of the earth's magnetic field at the place.



[Watch Video Solution](#)

25. At a certain location in Africa, a compass points 12° west of the geographic north. The north tip of the magnetic needle of a dip circle

placed in the plane of magnetic meridian points 60° above the horizontal. The horizontal component of the earth's field is measured to be 0.16 G. Specify the direction and magnitude of the earth's field at the location.



[Watch Video Solution](#)

26. A short bar magnet has a magnetic moment of $0.48JT^{-1}$. Give the direction and magnitude of the magnetic field produced by

the magnet at a distance of 10 cm from the centre of the magnet on the axis.



[Watch Video Solution](#)

27. A short bar magnet has a magnetic moment of $0.48JT^{-1}$. Give the direction and magnitude of the magnetic field produced by the magnet at a distance of 10 cm from the centre of the magnet on the equatorial lines (normal bisector) of the magnet.



[Watch Video Solution](#)

28. A short bar magnet placed in a horizontal plane has its axis aligned along the magnetic north-south direction. Null points are found on the axis of the magnet at 14 cm from the centre of the magnet. The earth's magnetic field at the place is 0.36 G and the angle of dip is zero. What is the total magnetic field on the normal bisector of the magnet at the same distance as the null-point (i.e., 14 cm) from the centre of the magnet? (At null points, field due to a magnet is equal and opposite to the

horizontal component of earth's magnetic field.)



[Watch Video Solution](#)

29. A short bar magnet placed in a horizontal plane has its axis aligned along the magnetic north-south direction. Null points are found on the axis of the magnet at 14 cm from the center of the magnet. The earth's magnetic field at the place is 0.36 G and the angle of dip is zero. If the bar magnet is turned around by

180° , where will the new null points be located?



[Watch Video Solution](#)

30. A short bar magnet of magnetic moment $5.25 \times 10^{-2} \text{ JT}^{-1}$ is placed with its axis perpendicular to the earth's field direction. At what distance from the centre of the magnet, the resultant field is inclined at 45° with earth's field on its normal bisector. Magnitude of the earth's field at the place is given to be

0.42 G. Ignore the length of the magnet in comparison to the distances involved.



Watch Video Solution

31. A short bar magnet of magnetic moment $5.25 \times 10^{-2} \text{ JT}^1$ is placed with its axis perpendicular to the earth's field direction. At what distance from the centre of the magnet, the resultant field is inclined at 45° with earth's field on its axis. Magnitude of the earth's field at the place is given to be 0.42 G.

Ignore the length of the magnet in comparison to the distances involved.



[Watch Video Solution](#)

32. Answer the following questions: Why does a paramagnetic sample display greater magnetisation (for the same magnetising field) when cooled?



[Watch Video Solution](#)

33. Answer the following questions: Why is diamagnetism, in contrast, almost independent of temperature?



Watch Video Solution

34. Answer the following questions: If a toroid uses bismuth for its core, will the field in the core be (slightly) greater or (slightly) less than when the core is empty?



Watch Video Solution

35. Answer the following questions: Is the permeability of a ferromagnetic material independent of the magnetic field? If not, is it more for lower or higher fields?



Watch Video Solution

36. Answer the following questions: Magnetic field lines are always nearly normal to the surface of a ferromagnet at every point. (This fact is analogous to the static electric field

lines being normal to the surface of a conductor at every point.) Why?



[Watch Video Solution](#)

37. Answer the following questions: Would the maximum possible magnetisation of a paramagnetic sample be of the same order of magnitude as the magnetisation of a ferromagnet?



[Watch Video Solution](#)

38. Answer the following questions: Explain qualitatively on the basis of domain picture the irreversibility in the magnetisation curve of a ferromagnet.



Watch Video Solution

39. Answer the following questions: The hysteresis loop of a soft iron piece has a much smaller area than that of a carbon steel piece. If the material is to go through repeated

cycles of magnetisation, which piece will dissipate greater heat energy?



[Watch Video Solution](#)

40. Answer the following questions: A system displaying a hysteresis loop such as a ferromagnet, is a device for storing memory? Explain the meaning of this statement.



[Watch Video Solution](#)

41. Answer the following questions: What kind of ferromagnetic material is used for coating magnetic tapes in a cassette player, or for building 'memory stores' in a modern computer?



Watch Video Solution

42. Answer the following questions: A certain region of space is to be shielded from magnetic fields. Suggest a method.



43. A long straight horizontal cable carries a current of 2.5 A in the direction 10° south of west to 10° north of east. The magnetic meridian of the place happens to be 10° west of the geographic meridian. The earth's magnetic field at the location is 0.33 G, and the angle of dip is zero. Locate the line of neutral points (ignore the thickness of the cable)? (At neutral points, magnetic field due to a current-carrying cable is equal and

opposite to the horizontal component of earth's magnetic field.)



[Watch Video Solution](#)

44. A telephone cable at a place has four long straight horizontal wires carrying a current of 1.0 A in the same direction east to west. The earth's magnetic field at the place is 0.39 G, and the angle of dip is 35° . The magnetic declination is nearly zero. What are the

resultant magnetic fields at points 4.0 cm below and above the cable?



[Watch Video Solution](#)

45. A compass needle free to turn in a horizontal plane is placed at the centre of circular coil of 30 turns and radius 12 cm. The coil is in a vertical plane making an angle of 45° with the magnetic meridian. When the current in the coil is 0.35 A, the needle points west to east. Determine the horizontal

component of the earth's magnetic field at the location.



[Watch Video Solution](#)

46. A compass needle free to turn in a horizontal plane is placed at the centre of circular coil of 30 turns and radius 12 cm. The coil is in a vertical plane making an angle of 45° with the magnetic meridian. When the current in the coil is 0.35 A, the needle points west to east. Determine the horizontal

component of the earth's magnetic field at the location.



[Watch Video Solution](#)

47. A magnetic dipole is under the influence of two magnetic fields. The angle between the field directions is 60° , and one of the fields has a magnitude of $1.2 \times 10^{-2} T$. If the dipole comes to stable equilibrium at an angle of 15° with this field, what is the magnitude of the other field?



[Watch Video Solution](#)

48. A monoenergetic (18 keV) electron beam initially in the horizontal direction is subjected to a horizontal magnetic field of 0.04 G normal to the initial direction. Estimate the up or down deflection of the beam over a distance of 30 cm ($m_e = 9.11 \times 10^{-31} \text{ kg}$).



[Watch Video Solution](#)

49. A sample of paramagnetic salt contains 2.0×10^{24} atomic dipoles each of dipole moment $1.5 \times 10^{-23} \text{ JT}^{-1}$. The sample is placed under a homogeneous magnetic field of 0.64 T, and cooled to a temperature of 4.2 K. The degree of magnetic saturation achieved is equal to 15%. What is the total dipole moment of the sample for a magnetic field of 0.98 T and a temperature of 2.8 K? (Assume Curie's law)



Watch Video Solution

50. A Rowland ring of mean radius 15 cm has 3500 turns of wire wound on a ferromagnetic core of relative permeability 800. What is the magnetic field B in the core for a magnetising current of 1.2 A?



Watch Video Solution

51. The magnetic moment vectors μ_s and μ_l (associated with the intrinsic spin angular momentum S and orbital angular momentum

1, respectively, of an electron are predicted by quantum theory (and verified experimentally to a high accuracy) to be given by:

$$\mu_s = - (e/m)S, \mu_l = - (e/2m)l$$

Which of these relations is in accordance with the result expected classically? Outline the derivation of the classical result.



Watch Video Solution