



PHYSICS

BOOKS - PSEB

WAVE OPTICS

Exercise

1. Monochromatic light of wavelength 589 nm is incident from air on a water surface. What

are the wavelength, frequency and speed of refracted light. Refractive index of water is 1.33



Watch Video Solution

2. Monochromatic light of wavelength 589 nm is incident from air on a water surface. What are the wavelength, frequency and speed of refracted light. Refractive index of water is 1.33



Watch Video Solution

3. What is the shape of the wavefront in each of the following cases: Light diverging from a point source.



Watch Video Solution

4. What is the shape of the wavefront in each of the following cases: Light emerging out of a convex lens when a point source is placed at its focus.



Watch Video Solution

5. What is the shape of the wavefront in each of the following cases: The portion of the wavefront of light from a distant star intercepted by the Earth.



Watch Video Solution

6. The refractive index of glass is 1.5. What is the speed of light in glass? (Speed of light in vacuum is $3.0 \times 10^8 \text{ ms}^{-1}$)



Watch Video Solution

7. Is the speed of light in glass independent of the colour of light? If not, which of the two colours red and violet travels slower in a glass prism?



Watch Video Solution

8. In a Young 0.28 mm and the screen is placed 1.4 m away. The distance between the central bright fringe and the fourth bright fringe is

measured to be 1.2 cm. Determine the wavelength of light used in the experiment.



Watch Video Solution

9. In Young's experiment with wavelength λ , the intensity of light at a point on the screen where path difference is λ , is K units. What is the intensity of light at a point where path difference is $\frac{\lambda}{3}$?



Watch Video Solution

10. A beam of light consisting of two wavelengths, 650 nm and 520 nm, is used to obtain interference fringes in a Young experiment.- Find the distance of the third bright fringe on the screen from the central maximum for wavelength 650 nm.



Watch Video Solution

11. A beam of light consisting of two wavelengths, 650 nm and 520 nm, is used to

obtain interference fringes in a Young experiment.-What is the least distance from the central maximum where the bright fringes due to both the wavelengths coincide?



Watch Video Solution

12. In a double-slit experiment the angular width of a fringe is found to be 0.2° on a screen placed 1 m away. The wavelength of light used is 600 nm. What will be the angular width of the fringe if the entire apparatus is

immersed in water? Take refractive index of water to be $\frac{4}{3}$.



Watch Video Solution

13. What is the Brewster angle for air to glass transition? (Refractive index of glass = 1.5.)



Watch Video Solution

14. Light of wavelength 5000 \AA falls on a plane reflecting surface. What are the wavelength

and frequency of the reflected light? For what angle of incidence is the reflected ray normal to the incident ray?



Watch Video Solution

15. Estimate the distance for which ray optics is good approximation for an aperture of 4 mm and wavelength 400 nm.



Watch Video Solution

16. Explain how Corpuscular theory predicts the speed of light in a medium, say, water, to be greater than the speed of light in vacuum. Is the prediction confirmed by experimental determination of the speed of light in water ? If not, which alternative picture of light is consistent with experiment?



Watch Video Solution

17. You have learnt in the text how Huygens of reflection and refraction. Use the same principle to deduce directly that a point object placed in front of a plane mirror produces a virtual image whose distance from the mirror is equal to the object distance from the mirror.



Watch Video Solution

18. 4 Let us list some of the factors, which could possibly influence the speed of wave

propagation:(i) nature of the source.(ii) direction of propagation.(iii) motion of the source and/or observer.(iv) wavelength. (v) intensity of the wave. On which of these factors, if any, does,depend? :the speed of light in vacuum



Watch Video Solution

19. 5 Let us list some of the factors, which could possibly influence the speed of wave propagation: (i) nature of the source (ii)

direction of propagation. (iii) motion of the source and/or observer. (iv) wavelength. (v) intensity of the wave. On which of the factors, if any, does the speed of light in a medium (say, glass or water)



Watch Video Solution

20. For sound waves, the Doppler formula for frequency shift differs slightly between the two situations: 1. source at rest, observer moving 2. source moving, observer at rest. The

exact Doppler formulas for the case of light waves in vacuum are, however, strictly identical for these situations. Explain why this should be so. Would you expect the formulas to be strictly identical for the two situations in case of light travelling in a medium?



Watch Video Solution

21. For sound waves, the Doppler formula for frequency shift differs slightly between the two situations: 1. source at rest, observer

moving 2. source moving, observer at rest. The exact Doppler formulas for the case of light waves in vacuum are, however, strictly identical for these situations. Explain why this should be so. Would you expect the formulas to be strictly identical for the two situations in case of light travelling in a medium?



Watch Video Solution

22. In double-slit experiment using light of wavelength 600 nm, the angular width of a

fringe formed on a distant screen is 0.1° .

What is the spacing between the two slits?



Watch Video Solution

23. Answer the following questions: In a single slit diffraction experiment, the width of the slit is made double the original width. How does this affect the size and intensity of the central diffraction band?



Watch Video Solution

24. Answer the following questions: In what way is diffraction from each slit related to the interference pattern in a double-slit experiment?



Watch Video Solution

25. Answer the following questions: When a tiny circular obstacle is placed in the path of light from a distant source, a bright spot is seen at the centre of the shadow of the obstacle. Explain why?



[Watch Video Solution](#)

26. Answer the following questions: Two students are separated by a 7 m partition wall in a room 10 m high. If both light and sound waves can bend around obstacles, how is it that the students are unable to see each other even though they can converse easily.



[Watch Video Solution](#)

27. Answer the following questions: Ray optics is based on the assumption that light travels in a straight line. Diffraction effects (observed when light propagates through small apertures/slits or around small obstacles) disprove this assumption. Yet the ray optics assumption is so commonly used in understanding location and several other properties of images in optical instruments. What is the justification?



Watch Video Solution

28. Two towers on top of two hills are 40 km apart. The line joining them passes 50 m above a hill halfway between the towers. What is the longest wavelength of radio waves, which can be sent between the towers without appreciable diffraction effects?



Watch Video Solution

29. 9 A parallel beam of light of wavelength 500 nm falls on a narrow slit and the resulting

diffraction pattern is observed on a screen 1 m away. It is observed that the first minimum is at a distance of 2.5 mm from the centre of the screen. Find the width of the slit.



Watch Video Solution

30. Answer the following questions: When a low flying aircraft passes overhead, we sometimes notice a slight shaking of the picture on our TV screen. Suggest a possible explanation.



[Watch Video Solution](#)

31. Answer the following questions: As you have learnt in the text, the principle of linear superposition of wave displacement is basic to understanding intensity distributions in diffraction and interference patterns. What is the justification of this principle?



[Watch Video Solution](#)

32. In deriving the single slit diffraction pattern, it was stated that the intensity is zero at angles of $n\frac{\lambda}{a}$. Justify this by suitably dividing the slit to bring out the cancellation.



Watch Video Solution