



MATHS

BOOKS - ARIHANT MATHS (HINGLISH)

CONTINUITY AND DIFFERENTIABILITY

EXAMPLE

1. Determine the value of the constant k , so that the function $f(x) = \begin{cases} kx, & \text{if } x < 0 \\ 3, & \text{if } x \geq 0 \end{cases}$ is continuous at $x = 0$

 [Watch Video Solution](#)

2. Prove that every rational function is continuous.

 [Watch Video Solution](#)

3. Discuss the continuity of the function f given by $f(x) = |x|$ at $x = 0$.

 [Watch Video Solution](#)

4. Prove that the greatest integer function $[x]$ is continuous at all points except at integer points.

 [Watch Video Solution](#)

5. Find all points of discontinuity of the function 'f' defined by :

$$f(x) = \begin{cases} x + 2, & x \leq 1 \\ x - 2, & 1 < x < 2 \\ 0, & x \geq 2. \end{cases}$$

 [Watch Video Solution](#)

6. Find the values of k so that the function f is continuous at the indicated point in $f(x) = \begin{cases} kx + 1, & \text{if } x \leq 5 \\ 3x - 5, & \text{if } x > 5 \end{cases}$ at $x = 5$



Watch Video Solution

7. Find the value of a for which the function f defined by

$$f(x) = \begin{cases} a \frac{\sin \pi}{2} (x + 1), & x \leq 0 \\ \frac{\tan x - \sin x}{x^3}, & x > 0 \end{cases}$$

is continuous at $x = 0$.



Watch Video Solution

8. Find the values of a and b such that the function defined by

$$f(x) = \begin{cases} 5, & x < 2 \\ 2a + b, & x = 2 \\ 2ax + b, & x > 2 \end{cases}$$

is continuous at $x = 2$.



Watch Video Solution

9. Find the relationship between a and b so that the function f defined by

$$f(x) = \begin{cases} ax + 1, & \text{if } x \leq 3 \\ bx + 3, & \text{if } x > 3 \end{cases}$$

is continuous at $x = 3$.



Watch Video Solution

10. Show that the function $f(x)$ defined by :

$$f(x) = \begin{cases} \frac{\sin x}{x} + \cos x, & x > 0 \\ 2, & x = 0 \\ \frac{4(1 - \sqrt{1-x})}{x}, & x < 0 \end{cases}$$

is continuous at $x = 0$.



[Watch Video Solution](#)

11. Given that $f(x) = \begin{cases} \frac{\sqrt{1+kx} - \sqrt{1-kx}}{x} & \text{if } -1 \leq x < 0 \\ \frac{2x+1}{x-1} & \text{if } 0 \leq x \leq 1 \end{cases}$ is

continuous at $x = 0$. Find k



[Watch Video Solution](#)

12. If $f(x) = \begin{cases} \frac{\sin(a+1)x + 2\sin x}{x}, & x < 0 \\ 2, & x = 0 \\ \frac{\sqrt{1+bx} - 1}{x}, & x > 0 \end{cases}$

is continuous at $x = 0$, then find the values of 'a' and 'b'.



[Watch Video Solution](#)

13. Show that the function f defined by $f(x) = |1 - x| + |x|$, where x is any real number, is a continuous function.

 [Watch Video Solution](#)

14. Is $f(x) = |x - 1| + |x - 2|$ differentiable at $x = 2$?

 [Watch Video Solution](#)

15. Find 'a' and 'b', if the function given by:

$$f(x) = \begin{cases} ax^2 + b, & \text{if } x < 1 \\ 2x + 1, & \text{if } x \geq 1 \end{cases}$$

is differentiable at $x = 1$.

 [Watch Video Solution](#)

16. If f is derivable at $x = a$, then $\lim_{x \rightarrow a} \left(\frac{xf(a) - af(x)}{x - a} \right)$

 [Watch Video Solution](#)

17. If $f(x) = \begin{cases} x^2 \sin\left(\frac{1}{x}\right) & x \neq 0 \\ 0 & x = 0 \end{cases}$, then

 [Watch Video Solution](#)

18. If $f(x) = x + 1$, find $\frac{d}{dx}(\text{fof})(x)$.

 [Watch Video Solution](#)

19. Find $\frac{dy}{dx}$ when $\sin^2 x + \cos^2 y = 1$.

 [Watch Video Solution](#)

20. Use Chain rule of find $\frac{dy}{dx}$, if $y = \left(\frac{2x - 1}{2x + 1}\right)^2$.

 [Watch Video Solution](#)

21. Differentiate $s \in (\cos(x^2))$ with respect to x .

 [Watch Video Solution](#)

22. If $y = \sqrt{\frac{1-x}{1+x}}$, prove that $(1-x^2)\frac{dy}{dx} + y = 0$

 [Watch Video Solution](#)

23. If $y = \{x + \sqrt{x^2 + a^2}\}^n$, prove that $\frac{dy}{dx} = \frac{ny}{\sqrt{x^2 + a^2}} + a$

 [Watch Video Solution](#)

24. Find $\frac{dy}{dx}$, if $y + \sin y = \cos x$.

 [Watch Video Solution](#)

25. If $(x^2 + y^2)^2 = xy$, find $\frac{dy}{dx}$

 [Watch Video Solution](#)

26. If $x \sin(a + y) + \sin a \cdot \cos(a + y) = 0$, then prove that

$$\frac{dy}{dx} = \frac{\sin^2(a + y)}{\sin a}$$

 [Watch Video Solution](#)

27. If $x^2 + y^2 = t$ and $x^4 + y^4 = t^2 + \frac{1}{t^2}$, then prove that

$$\frac{dy}{dx} = \frac{1}{x^3 y}$$

 [Watch Video Solution](#)

28. Differentiate $\sin^{-1}(\sqrt{\cos x})$ w.r.t. x , using chain rule.

 [Watch Video Solution](#)

29. If $y = \sqrt{\cot^{-1} \sqrt{x}}$, find $\frac{dy}{dx}$.



Watch Video Solution

30. If $y = \sin(2 \sin^{-1} x)$, show that:

$$\frac{dy}{dx} = 2\sqrt{\frac{1-y^2}{1-x^2}}$$



Watch Video Solution

31. If $y = \sec^{-1}\left(\frac{1}{2x^2 - 1}\right)$ then $\frac{dy}{dx} = ?$



Watch Video Solution

32. If $y = \sin^{-1}\left[x(1-x) - \sqrt{x}\sqrt{1-x^2}\right]$, find $\frac{dy}{dx}$,



Watch Video Solution

33. Differentiate $\tan^{-1}\left(\frac{1 + \cos x}{\sin x}\right)$ with respect to x .



Watch Video Solution

34. If $y = \sin^{-1}\left(6x\sqrt{1-9x^2}\right)$, $-\frac{1}{3\sqrt{2}} < x < \frac{1}{3\sqrt{2}}$, then find $\frac{dy}{dx}$.

 Watch Video Solution

35. Find $\frac{dy}{dx}$ if $y = \sin^{-1}\left[\frac{6x - 4\sqrt{1-4x^2}}{5}\right]$

 Watch Video Solution

36. सिद्ध करें कि (Prove that) :

$$\frac{d}{dx} \left[\frac{x}{2} \sqrt{a^2 - x^2} + \frac{a^2}{2} \sin^{-1} \frac{x}{a} \right] = \sqrt{a^2 - x^2}$$

 Watch Video Solution

37. If $\log(x^2 + y^2) = 2 \tan^{-1}\left(\frac{y}{x}\right)$, show that $\frac{dy}{dx} = \frac{x+y}{x-y}$.

 Watch Video Solution

38. If $\sqrt{1-x^2} + \sqrt{1-y^2} = a(x-y)$, then prove that

$$\frac{dy}{dx} = \sqrt{\frac{1-y^2}{1-x^2}}$$

 [Watch Video Solution](#)

39. Is it true that $x = e^{\log x}$ for all real

 [Watch Video Solution](#)

40. Differentiate the following w.r.t. x : 3^{x+2} .

 [Watch Video Solution](#)

41. If $xy = e^{x-y}$, prove that $\frac{dy}{dx} = \frac{y(x-1)}{x(y+1)}$.

 [Watch Video Solution](#)

42. Differentiate $\sin^{-1}\left(\frac{2^{x+1}}{1+4^x}\right)$ w.r.t. x .

 [Watch Video Solution](#)

43. Differentiate the following with respect to x : $\sin^{-1}\left(\frac{2^{x+1} \cdot 3^x}{1+(36)^x}\right)$

 [Watch Video Solution](#)

44. If $y = \log \tan\left(\frac{\pi}{4} + \frac{x}{2}\right)$, show that :

$$\frac{dy}{dx} - \sec x = 0.$$

 [Watch Video Solution](#)

45. If $y = \log_{10} x + \log_x 10 + \log_x x + \log_{10} 10$, find $\frac{dy}{dx}$.

 [Watch Video Solution](#)

46. If $e^x + e^y = e^{x+y}$, prove that :

$$\frac{dy}{dx} = -\frac{e^x(e^y - 1)}{e^y(e^x - 1)}.$$



Watch Video Solution

47. Find the derivative of $\log(\sin(\log x))$



Watch Video Solution

48. Find $\frac{dy}{dx}$, when $x = \frac{1-t^2}{1+t^2}$ and $y = \frac{2t}{1+t^2}$



Watch Video Solution

49. Find $\frac{dy}{dx}$ when :

$$x = a(1 + \cos \theta), y = a(\theta + \sin \theta).$$



Watch Video Solution

50. Find $\frac{dy}{dx}$, if $x^{\frac{2}{3}} + y^{\frac{2}{3}} = a^{\frac{2}{3}}$.

 [Watch Video Solution](#)

51. Find $\frac{dy}{dx}$, when $x = e^{\theta} \left(\theta + \frac{1}{\theta} \right)$ and $y = e^{-\theta} \left(\theta - \frac{1}{\theta} \right)$

 [Watch Video Solution](#)

52. If $x = a(2\theta - \sin 2\theta)$ and $y = a(1 - \cos 2\theta)$ find $\frac{dy}{dx}$ when $\theta = \frac{\pi}{3}$

 [Watch Video Solution](#)

53. Find the value of $\frac{dy}{dx}$ at $\theta = \frac{\pi}{4}$, if :

$x = ae^{\theta}(\sin \theta - \cos \theta)$ and $y = ae^{\theta}(\sin \theta + \cos \theta)$.

 [Watch Video Solution](#)

54. If $x = a \sin 2t(1 + \cos 2t)$ and $y = b \cos 2t(1 - \cos 2t)$, find the values of $\frac{dy}{dx}$ at $t = \frac{\pi}{4}$ and $t = \frac{\pi}{3}$.

 [Watch Video Solution](#)

55. Differentiate $\sin^2(x^2)$ w.r.t. x^2 .

 [Watch Video Solution](#)

56. Differentiate $\sin^{-1}\left(\frac{x}{\sqrt{1+x^2}}\right)$ w.r.t. $\tan^{-1} x$.

 [Watch Video Solution](#)

57. Differentiate $\sin^{-1}\left(2x\sqrt{1-x^2}\right)$ w.r.t. $\cos^{-1}\left(\frac{1-x^2}{1+x^2}\right)$.

 [Watch Video Solution](#)

58.

if

$$x \in \left(\frac{1}{\sqrt{2}}, 1 \right).$$

Differentiate

$$\tan^{-1} \left(\frac{\sqrt{1-x^2}}{x} \right) \text{ w. r. t. } \cos^{-1} \left(2x\sqrt{1-x^2} \right).$$

[Watch Video Solution](#)

59. If $y = x^x$, find $\frac{dy}{dx}$.

[Watch Video Solution](#)

60. Differentiate x^{x^x} w.r.t. x .

[Watch Video Solution](#)

61. Find $f'(x)$, if $f(x) = (\sin x)^{\sin x}$ for all $0 < x < \pi$.

[Watch Video Solution](#)

62. Differentiate $x^{\tan x} + (\sin x)^{\cos x}$ w.r.t. x .



Watch Video Solution

63. If $x^y y^x = 1$, prove that $\frac{dy}{dx} = -\frac{y(y + x \log y)}{x(y \log x + x)}$



Watch Video Solution

64. If $x^y = e^{x-y}$ then prove that $\frac{dy}{dx} = \frac{\log x}{(1 + \log x)^2}$



Watch Video Solution

65. If $x^y = e^{x-y}$, show that $\frac{dy}{dx} = \frac{\log x}{\{\log(xe)\}^2}$



Watch Video Solution

66. If $x^y + y^x = a^b$, then find $\frac{dy}{dx}$.

 [Watch Video Solution](#)

67. Find $\frac{dy}{dx}$, if $y^x + x^y + x^x = a^b$.

 [Watch Video Solution](#)

68. If $f(x) = \left(\frac{3+x}{1+x}\right)^{2+3x}$, find $f'(0)$

 [Watch Video Solution](#)

69. If $y = \sqrt{2^x + \sqrt{2^x + \sqrt{2^x + \dots \text{to } \infty}}}$, then prove that :

$$(2y - 1) \frac{dy}{dx} = 2^x \log 2.$$

 [Watch Video Solution](#)

70. $\sqrt{\log x}$



Watch Video Solution

71. If $y = e^x + e^x + e\left(\left(x + \rightarrow \infty\right)\right)$, show that $\frac{dy}{dx} = \frac{y}{1 - y}$.



Watch Video Solution

72. If $y = \sqrt{x}^{\sqrt{x}^{\sqrt{x} \dots \infty}}$ then prove that $\frac{dy}{dx} = \frac{y^2}{x(2 - y \log x)}$.



Watch Video Solution

73. Find the second derivative of $\sin^{-1} x$.



Watch Video Solution

74. If $y = 500 e^{7x} + 600 e^{-7x}$, show that $\frac{d^2y}{dx^2} = 49 y$.



Watch Video Solution

 Watch Video Solution

75. If $x = a(\cos t + t \sin t)$ and $y = a(\sin t - t \cos t)$, find $\frac{d^2y}{dx^2}$.

 Watch Video Solution

76. If $y = \sin(\sin x)$, prove that $\frac{d^2y}{dx^2} + \tan x \frac{dy}{dx} + y \cos^2 x = 0$.

 Watch Video Solution

77. If $x = \tan\left(\frac{1}{a} \log y\right)$, show that $(1 + x^2) \frac{d^2y}{dx^2} + (2x - a) \frac{dy}{dx} = 0$.

 Watch Video Solution

78. If $x = a \cos \theta + b \sin \theta$, $y = a \sin \theta - b \cos \theta$, show that $y^2 \frac{d^2y}{dx^2} - x \frac{dy}{dx} + y = 0$.

 Watch Video Solution

79. If $x = \sin t$ and $y = \sin pt$, prove that

$$(1 - x^2) \frac{d^2y}{dx^2} - x \frac{dy}{dx} + p^2y = 0.$$

 [Watch Video Solution](#)

80. If $y = a \sin(\log x) + b \cos(\log x)$, prove that :

$$x^2 \frac{d^2y}{dx^2} + x \frac{dy}{dx} + y = 0.$$

 [Watch Video Solution](#)

81. If $x \cos(a + y) = \cos y$, then prove that $\frac{dy}{dx} = \frac{\cos^2(a + y)}{\sin a}$.

Hence, show that $\sin a \frac{d^2y}{dx^2} + \sin 2(a + y) \frac{dy}{dx} = 0$.

 [Watch Video Solution](#)

82. If $y = \left[\log \left(x + \sqrt{1 + x^2} \right) \right]^2$, show that :

$$(1 + x^2) \frac{d^2y}{dx^2} + x \frac{dy}{dx} - 2 = 0.$$

 [Watch Video Solution](#)

83. If $y = \log(1 + 2t^2 + t^4)$, $x = \tan^{-1} t$, find $\frac{d^2y}{dx^2}$.

 [Watch Video Solution](#)

84. If $y = x^x$, prove that $\frac{d^2y}{dx^2} - \frac{1}{y} \left(\frac{dy}{dx} \right)^2 - \frac{y}{x} = 0$

 [Watch Video Solution](#)

85. If $y = \operatorname{cosec}^{-1} x$, $x > 1$, then show that

$$x(x^2 - 1) \frac{d^2y}{dx^2} + (2x^2 - 1) \frac{dy}{dx} = 0.$$

 [Watch Video Solution](#)

86. If $y = e^{2 \tan^{-1} x}$, then show that :

$$(1 + x^2)^2 \frac{d^2 y}{dx^2} + 2x(1 + x^2) \frac{dy}{dx} = 4y.$$

 [Watch Video Solution](#)

87. At what points on the curve $y = (\cos x - 1)$ in $[0, 2\pi]$, is the tangent parallel to the x-axis?

 [Watch Video Solution](#)

88. It is given that for the function f given by $f(x) = x^3 + bx^2 + ax$, $x \in [1, 3]$. Rolle's theorem holds with $c = 2 + \frac{1}{\sqrt{3}}$. Find the values of a and b .

 [Watch Video Solution](#)

89. Discuss the applicability of Lagrange's Mean Value Theorem to:

$$f(x) = x(x - 1)(x - 2) \text{ in } \left[0, \frac{1}{2}\right].$$

 [Watch Video Solution](#)

90. Check whether Lagrange's Mean Value Theorem is applicable on :

$$f(x) = \sin x + \cos x \text{ in interval } \left[0, \frac{\pi}{2}\right].$$

 [Watch Video Solution](#)

91. Find a point on the parabola $y = (x - 3)^2$, where the tangent is parallel to the chord joining $(3, 0)$ and $(4, 1)$.

 [Watch Video Solution](#)

92. Using Lagrange's mean value theorem, find a point on the curve $y = \sqrt{x - 2}$ defined on the interval $[2, 3]$, where the tangent is parallel to

the chord joining the end points of the curve.

 [Watch Video Solution](#)

93. $f(x) = \begin{cases} \frac{x^3 + x^2 - 16x + 20}{(x - 2)^2}, & x \neq 2 \\ k, & x = 2 \end{cases}$. Find the value of k , so that the following function is continuous at $x=2$.

 [Watch Video Solution](#)

94. Let $f(x) = x|x|$, for all $x \in \mathbb{R}$. Discuss the derivability of $f(x)$ at $x = 0$.

 [Watch Video Solution](#)

95. If $y = (\tan x + \sec x)$, prove that $\frac{d^2y}{dx^2} = \frac{\cos x}{(1 - \sin x)^2}$

 [Watch Video Solution](#)

96. If $f(x) = |\cos x|$, then $f' \left(\frac{3\pi}{4} \right)$ equal to -

 [Watch Video Solution](#)

97. If $f(x) = \frac{\sqrt{2} \cos x - 1}{\cot x - 1}$, $x \neq \frac{\pi}{4}$. Find the value of $f\left(\frac{\pi}{4}\right)$ so that $f(x)$ becomes continuous at $x = \frac{\pi}{4}$.

 [Watch Video Solution](#)

98. Given that , $f(x) = \begin{cases} \frac{1 - \cos 4x}{x^2} & \text{if } x < 0 \\ a & \text{if } x = 0 \\ \frac{\sqrt{x}}{\sqrt{16 + \sqrt{x}} - 4} & \text{if } x > 0 \end{cases}$

If $f(x)$ is continuous at $x = 0$, find the value of a

 [Watch Video Solution](#)

99. Examine the differentiability of the function 'f' defined by :

$$f(x) = \begin{cases} 2x + 3, & \text{if } -3 \leq x < -2 \\ x + 1, & \text{if } -2 \leq x < 0 \\ x + 2, & \text{if } 0 \leq x \leq 1 \end{cases} .$$

 [Watch Video Solution](#)

100. Differentiate $\tan^{-1}\left(\frac{\sqrt{1-x^2}}{x}\right)$ with respect to $\cos^{-1}(2x\sqrt{1-x^2})$, when $x \neq 0$.

 [Watch Video Solution](#)

EXERCISE 5(a) (SHORT ANSWER TYPE QUESTIONS)

1. If $f(x) = \begin{cases} kx^2 & \text{if } x \leq 2 \\ 3 & \text{if } x > 2 \end{cases}$ is continuous at $x = 2$, then the value of k is

 [Watch Video Solution](#)

2. Check the continuity of the function f given by $f(x) = 2x + 3atx = 1$.

 [Watch Video Solution](#)

3. Check the continuity of the following functions :

$$f(x) = x^2 \text{ at } x = 0.$$

 [Watch Video Solution](#)

4. Examine the continuity of the function $f(x) = 2x^2 - 1$ at $x = 3$.

 [Watch Video Solution](#)

5. Examine the function $f(x) = 2x^2 - 5$ for its continuity at the point $x = 3$.

 [Watch Video Solution](#)

6. is $f(x) = |x|$ a continuous function?

 [Watch Video Solution](#)

7. Examine the following functions for continuity :

$$f(x) = x - 5$$

 [Watch Video Solution](#)

8. Examine the following functions for continuity :

$$f(x) = x^3 + x^2 - 1$$

 [Watch Video Solution](#)

9. Examine the following functions for continuity :

$$f(x) = \frac{1}{x - 5}, x \neq 5$$

 [Watch Video Solution](#)

10. Examine the following functions for continuity :

$$f(x) = \frac{x^2 - 25}{x + 5}, x \neq -5$$



Watch Video Solution

11. Examine the following functions for continuity:

(d) $f(x) = |x - 5|$.



Watch Video Solution

12. Prove that the following functions are continuous at all points of their domains :

$$f(x) = \cos x$$



Watch Video Solution

13. Prove that the following functions are continuous at all points of their domains :

$$f(x) = e^x + e^{-x}$$



[Watch Video Solution](#)

14. Prove that the following functions are continuous at all points of their domains :

$$f(x) = \tan x$$



[Watch Video Solution](#)

15. Prove that the following functions are continuous at all points of their domains :

$$f(x) = \cos(x^2).$$



[Watch Video Solution](#)

16. Discuss the continuity of the following functions a)

$$f(x) = \sin x + \cos x$$



Watch Video Solution

17. Discuss the continuity of $f(x) = \sin x - \cos x$



Watch Video Solution

18. Discuss the continuity of $f(x) = \sin x \cos x$



Watch Video Solution

19. Discuss the continuity of the following functions :

$$f(x) = \frac{\sin x}{\cos x}.$$



Watch Video Solution

20. Prove that $f(x) = |\sin x|$ is continuous at all points of its domain.



Watch Video Solution

21. Examine if $\sin|x|$ is a continuous function.



Watch Video Solution

22. Is the function defined by $f(x) = x^2 - \sin x + 5$ continuous at $x = \pi$?



Watch Video Solution

23. Show that $f(x) = x - |x|$, $x \in \mathbb{R}$ is continuous at $x = 0$.



Watch Video Solution

24. Show that the function defined by $g(x) = x - [x]$ is discontinuous at all integral points which $[x]$ denotes the greatest integer function.

 [Watch Video Solution](#)

25. Find all points of discontinuity of f , where f is defined by

$$f(x) = \begin{cases} x + 1 & \text{if } x \geq 1 \\ x^2 + 1 & \text{if } x < 1 \end{cases}$$

 [Watch Video Solution](#)

26. Find all points of discontinuity of f , where f is defined by

$$f(x) = \begin{cases} x^{10} - 1 & \text{if } x \leq 1 \\ x^2 & \text{if } x > 1 \end{cases}$$

 [Watch Video Solution](#)

27. Find all points of discontinuity of f , where f is defined by

$$f(x) = \begin{cases} 2x + 3 & \text{if } x \leq 2 \\ 2x - 3 & \text{if } x > 2 \end{cases}$$



Watch Video Solution

28. Find all points of discontinuity of f , where f is defined by

$$f(x) = \begin{cases} \frac{|x|}{x}, & \text{if } x \neq 0 \\ 0, & \text{if } x = 0 \end{cases}$$



Watch Video Solution

29. Find all points of discontinuity of f , where f is defined by

$$f(x) = \begin{cases} \frac{x}{|x|}, & \text{if } x < 0 \\ 1, & \text{if } x \geq 0 \end{cases}$$



Watch Video Solution

30. Is the function defined by

$$f(x) = \begin{cases} x + 5, & \text{if } x \leq 1 \\ x - 5, & \text{if } x > 1 \end{cases}$$
 a continuous function?



Watch Video Solution

31. Is the function f defined by

$$f(x) = \begin{cases} x & \text{if } x \leq 1 \\ 5 & \text{if } x > 1 \end{cases}$$

continuous at $x = 0$? At $x = 2$?



Watch Video Solution

32. Show that the function f given by

$$f(x) = \begin{cases} x^3 + 3 & \text{if } x \neq 0 \\ 1 & \text{if } x = 0 \end{cases}$$
 is not continuous at $x = 0$.



Watch Video Solution

33. Discuss the continuity of the function f defined by $f(x) = \frac{1}{x}$, $x \neq 0$.



Watch Video Solution

34. Discuss the continuity of the function :

$$f(x) = \begin{cases} x, & \text{if } x \geq 0 \\ x^2, & \text{if } x < 0 \end{cases}$$

 [Watch Video Solution](#)

35. Discuss the continuity of the function defined by

$$f(x) = \begin{cases} x + 2, & \text{if } x < 0 \\ -x + 2, & \text{if } x > 0 \end{cases}$$

 [Watch Video Solution](#)

36. Examine the continuity of the function :

$$f(x) = \begin{cases} x + 1, & x \leq 2 \\ 2x - 1, & x > 2 \end{cases} \text{ at } x = 2.$$

 [Watch Video Solution](#)

37. $f(x) = \begin{cases} \frac{x^2 - 25}{x - 5}, & \text{when } x \neq 5 \\ 10, & \text{when } x = 5 \end{cases}$ is continuous at $x = 5$

 [Watch Video Solution](#)

38. Discuss the continuity of the function :

$$f(x) = \begin{cases} \frac{|x-2|}{x-2}, & x \neq 2 \\ 1 & ,x = 2 \end{cases} \text{ at } x = 2$$



Watch Video Solution

39. Discuss the continuity of the function :

$$f(x) = \begin{cases} \frac{|x-2|}{2-x}, & x \neq 2 \\ -1 & ,x = 2 \end{cases} \text{ at } x = 2$$



Watch Video Solution

40. Discuss the continuity of the function :

$$f(x) = \begin{cases} \frac{|x-a|}{x-a}, \text{when } x \neq a \\ 1 & , \text{when } x = a \end{cases} \text{ at } x = a.$$



Watch Video Solution

41. Discuss the continuity of the function f , where f is defined by

$$f(x) \begin{cases} 3, & \text{if } 0 \leq x \leq 1 \\ 4, & \text{if } 1 < x < 3 \\ 5, & \text{if } 3 \leq x \leq 10 \end{cases}$$



Watch Video Solution

42. Discuss the continuity of the function f , where f is defined by $f(x) = \begin{cases} -2, & \text{if } x < -1 \\ 2x, & \text{if } -1 \leq x \leq 1 \end{cases}$



Watch Video Solution

43. Discuss the continuity of the function :

$$f(x) = \begin{cases} x, & 0 \leq x < \frac{1}{2} \\ \frac{1}{2}, & x = \frac{1}{2} \\ 1 - x, & \frac{1}{2} < x \leq 1 \end{cases} \quad \text{at } x = \frac{1}{2}$$



Watch Video Solution

44. Discuss the continuity of the function :

$$f(x) = \begin{cases} \frac{1 - \cos x}{x^2}, & x \neq 0 \\ 1, & x = 0 \end{cases} \text{ at } x = 0.$$

 [Watch Video Solution](#)

45. Show that the function $f(x)$ defined as $f(x) = x \cos\left(\frac{1}{x}\right)$, $x \neq 0$; 0 , $x = 0$ is continuous at $x = 0$ but not differentiable at $x = 0$.

 [Watch Video Solution](#)

46. Show that the following functions are continuous at $x = 0$:

$$f(x) = \begin{cases} x \cos \frac{1}{x} & \text{when } x \neq 0 \\ 0 & \text{when } x = 0 \end{cases}$$

 [Watch Video Solution](#)

47. Test the continuity of the function $f(x)$:

$$f(x) = \begin{cases} x^2 \sin. \frac{1}{x} & \text{when } x \neq 0 \\ 0 & \text{when } x = 0 \end{cases}$$

 [Watch Video Solution](#)

48. $f(x) = \{(\cos x, \text{ when } x \geq 0,), (-\cos x, \text{ when } x < 0)\}$ is discontinuous at $x=0$

 [Watch Video Solution](#)

49. Examine the continuity of the function $f(x)$ at $x = 0$.

$$f(x) = \begin{cases} \frac{\sin x}{x} & \text{when } x \neq 0 \\ 2 & \text{when } x = 0 \end{cases}$$

 [Watch Video Solution](#)

50. Examine the continuity of the function

$$f(x) = \begin{cases} \frac{|\sin x|}{x}, & x \neq 0 \\ 1, & x = 0 \end{cases}$$

 [Watch Video Solution](#)

51. Examine the continuity of the function $f(x)$ at $x = 0$.

$$f(x) = \begin{cases} \frac{\tan 2x}{3x} & \text{when } x \neq 0 \\ \frac{3}{2} & \text{when } x = 0 \end{cases}$$

 [Watch Video Solution](#)

52. Discuss the continuity of the cosine, cosecant, secant and cotangent functions.

 [Watch Video Solution](#)

EXERCISE 5(a) (LONG ANSWER TYPE QUESTIONS (I))

1. Test for continuity of the following function at $x = a$: $f(x) = \begin{cases} (x-a)^2, & x \neq a \\ 0, & x = a \end{cases}$

$$\sin\left(\frac{1}{x-a}\right), \text{ where } x \neq a \quad 0, \text{ when } x = a$$



[Watch Video Solution](#)

2. Examine the function for continuity at $x = 0$:

$$f(x) = \begin{cases} \frac{\sin x}{x} & \text{when } x < 0 \\ x + 1 & \text{when } x \geq 0 \end{cases}.$$



[Watch Video Solution](#)

3. Discuss the continuity of $f(x)$ at $x = 0$ if:

$$f(x) = \begin{cases} \frac{\sqrt{1+x} - \sqrt{1-x}}{\sin x}, & \text{if } x \neq 0 \\ 1 & , \text{if } x = 0 \end{cases}.$$



[Watch Video Solution](#)

4. $f(x) = \begin{cases} kx^2, & \text{if } x \leq 2 \\ 3, & \text{if } x > 2 \end{cases}$ at $x = 2$



[Watch Video Solution](#)

5. The function is defined by $f(x) = \begin{cases} kx + 1 & \text{if } x \leq \pi \\ \cos x & \text{if } x > \pi \end{cases}$ at $x = \pi$.

 [Watch Video Solution](#)

6. If the function $f(x) = \begin{cases} kx + 5, & \text{when } x \leq 2 \\ x - 1, & \text{when } x > 2 \end{cases}$ is continuous at $x = 2$

then $k = ?$

 [Watch Video Solution](#)

7. $f(x) = \begin{cases} 3k - 2x, & \text{when } x < 1 \\ 2k + 1, & \text{when } x \geq 1 \end{cases}$ at $x = 1$

 [Watch Video Solution](#)

8. $f(x) = \begin{cases} 3x - 8 & \text{if } x \leq 5 \\ 2k & \text{if } x > 5 \end{cases}$ at $x = 5$

 [Watch Video Solution](#)

$$9. f(x) = \begin{cases} \frac{x-1}{x+1}, & x \neq 1 \\ \lambda - 1, & x = 1 \end{cases} \text{ at } x = 1.$$

 [Watch Video Solution](#)

$$10. f(x) = \begin{cases} \frac{1 - \cos Ax}{x \sin x} & \text{if } x \neq 0 \\ \frac{1}{2} & \text{if } x = 0 \end{cases} \text{ at } x = 0$$

 [Watch Video Solution](#)

11. If the function $f(x) = \begin{cases} \frac{1 - \cos(ax)}{x^2} & \text{when } x \neq 0 \\ 1 & \text{when } x = 0 \end{cases}$ be continuous at $x=0$, then find the value of a .

 [Watch Video Solution](#)

$$12. f(x) = \begin{cases} \frac{\sin 2x}{5x}, & \text{when } x \neq 0 \\ m, & \text{when } x = 0 \end{cases} \text{ at } x = 0$$

 [Watch Video Solution](#)

13. Let $f(x) = \begin{cases} \frac{k \cos x}{\pi - 2x}, & x \neq \frac{\pi}{2} \\ 3, & x = \frac{\pi}{2}. \end{cases}$

If $\lim_{x \rightarrow \frac{\pi}{2}} f(x) = f\left(\frac{\pi}{2}\right)$, find the value of k.

 [Watch Video Solution](#)

14. $f(x) = \begin{cases} \frac{k \cos x}{(\pi - 2x)}, & \text{if } x \neq \frac{\pi}{2} \\ 3, & \text{if } x = \frac{\pi}{2} \end{cases}$

at $x = \frac{\pi}{2}$.

 [Watch Video Solution](#)

15. $f(x) = \begin{cases} \frac{x^2 - 9}{x - 3}, & \text{when } x \neq 3 \\ k, & \text{when } x = 3 \end{cases}$ at $x = 3$

 [Watch Video Solution](#)

16. $f(x) = \begin{cases} \frac{(x+3)^2 - 36}{x-3}, & x \neq 3 \\ k, & x = 3 \end{cases}$ at $x = 3$.



17. For what value of 'k' is the function defined by :

$$f(x) = \begin{cases} k(x^2 + 2), & \text{if } x \leq 0 \\ 3x + 1 & , \text{if } x > 0 \end{cases}$$

continuous at $x = 0$? Also, write whether the function is continuous at $x =$

1.

[View Text Solution](#)

18. If the function defined by :

$$f(x) = \begin{cases} 2x - 1, & x < 2 \\ a, & x = 2 \\ x + 1, & x > 2 \end{cases}$$

is continuous at $x = 2$, find the value of 'a'.

Also discuss the continuity of $f(x)$ at $x = 3$.

[Watch Video Solution](#)

19. Given that , $f(x) = \begin{cases} \frac{1 - \cos 4x}{x^2} & \text{if } x < 0 \\ a & \text{if } x = 0 \\ \frac{\sqrt{x}}{\sqrt{16 + \sqrt{x}} - 4} & \text{if } x > 0 \end{cases}$

If $f(x)$ is continuous at $x = 0$, find the value of a

 [Watch Video Solution](#)

20. Find the values of a and b such that the function defined by

$$f(x) = \begin{cases} 5 & \text{if } x \leq 2 \\ ax + b & \text{if } 2 < x < 10 \\ 21 & \text{if } x \geq 10 \end{cases} \text{ is continuous function.}$$

 [Watch Video Solution](#)

21. Determine the constants ' a ' and ' b ' so that the function ' f ' defined

below is continuous everywhere :

$$f(x) = \begin{cases} x + 2, & x \leq 2 \\ ax + b, & 2 < x < 5. \\ 3x - 2, & x \geq 5 \end{cases}$$

 [Watch Video Solution](#)

22. Determine the constants 'a' and 'b' so that the function 'f' defined below is continuous everywhere :

$$f(x) = \begin{cases} 5a, & x \leq 0 \\ a \sin x + \cos x, & 0 < x < \frac{\pi}{2} \\ b - \frac{\pi}{2}, & x \geq \frac{\pi}{2} \end{cases}$$

 [Watch Video Solution](#)

23. If the function $f(x) = \begin{cases} 3ax + b & \text{for } x > 1 \\ 11 & \text{when } x = 1 \\ 5ax - 2b & \text{for } x < 1 \end{cases}$

is continuous at $x = 1$, find the values of a and b .

 [Watch Video Solution](#)

24. Find the values of 'a' and 'b' so that the following function is continuous at $x = 3$ and $x = 5$:

$$f(x) = \begin{cases} 1, & \text{if } x \leq 3 \\ ax + b, & \text{if } 3 < x < 5 \\ 7, & \text{if } 5 \leq x \end{cases}$$

 [Watch Video Solution](#)

25. Find 'a' and 'b' if the function :

$$f(x) = \begin{cases} \frac{\sin x}{x}, & -2 \leq x < 0 \\ a \cdot 2^x, & 0 \leq x \leq 1 \\ b + x, & 1 < x \leq 2 \end{cases}$$

is a continuous function on $[-2, 2]$.



[Watch Video Solution](#)

26. Determine the values of 'a' and 'b' such that the following function is continuous at $x = 0$:

$$f(x) = \begin{cases} \frac{x + \sin x}{\sin(a+1)x}, & \text{if } -\pi < x < 0 \\ 2, & \text{if } x = 0 \\ 2 \frac{e^{\sin bx} - 1}{bx}, & \text{if } x > 0 \end{cases} .$$



[Watch Video Solution](#)

$$27. \text{ Let } f(x) = \begin{cases} \frac{1 - \sin^3 x}{3 \cos^2 x} & \text{if } x < \frac{\pi}{2} \\ a & \text{if } x = \frac{\pi}{2} \\ \frac{b(1 - \sin x)}{(\pi - 2x)^2} & \text{if } x > \frac{\pi}{2} \end{cases},$$

if $f(x)$ is continuous at $x = \frac{\pi}{2}$, find a and b .



Watch Video Solution

28. If the following function $f(x)$ is continuous at $x=0$, find the values of a , b and c , where

$$f(x) = \begin{cases} \frac{\sin(a+1)x + \sin x}{x} & \text{if } x < 0 \\ c & \text{if } x = 0 \\ \frac{\sqrt{x+bx^2} - \sqrt{x}}{bx^{\frac{3}{2}}} & \text{if } x > 0 \end{cases}$$



Watch Video Solution

29. If $f(x) = \begin{cases} x^2 + ax + b, & \text{if } 0 \leq x < 2 \\ 3x + 2, & \text{if } 2 \leq x \leq 4 \\ 2ax + 5b, & \text{if } 4 < x \leq 8 \end{cases}$ is continuous on $[0, 8]$ then

$$a - b = \dots\dots$$



Watch Video Solution

30. Find all the points of discontinuity of the function 'f' defined by :

$$f(x) = \begin{cases} x + 2, & \text{if } x < 1 \\ 0, & \text{if } x = 1 \\ x - 2, & \text{if } x > 1 \end{cases}$$



[View Text Solution](#)

31. Find all points of discontinuity of f, where f is defined by $f(x) = \begin{cases} |x| + 3, & \text{if } x \geq -3 \\ -3 - 2x, & \text{if } x < -3 \end{cases}$



[Watch Video Solution](#)

32. Find all points of discontinuity of f, where f is defined by

$$f(x) = \begin{cases} 2x + 3, & \text{if } x \leq 2 \\ 2x - 3, & \text{if } x > 2 \end{cases}$$



[Watch Video Solution](#)

33. Show that the function defined by $f(x) = s \in (x^2)$ is a continuous function.

 [Watch Video Solution](#)

34. Show that the function defined by $f(x) = \cos(x^2)$ is a continuous function.

 [Watch Video Solution](#)

35. Show that the function defined by $f(x) = |\cos x|$ is a continuous function.

 [Watch Video Solution](#)

36. Show that the function 'f' given by :

$$f(x) = |x| + |x - 1|, x \in R$$

is continuous both at $x = 0$ and $x = 1$



[Watch Video Solution](#)

37. Discuss the continuity of the function of given by

$$f(x) = + |x - 1| + |x - 2| \text{ at } x = 1 \text{ and } x = 2$$



[Watch Video Solution](#)

38. Find the points of discontinuity, if any, of the following function:

$$f(x) = \begin{cases} \frac{x^4 - 16}{x - 2} & , \quad \text{if } x \neq 2 \\ 2 & , \quad \text{if } x = 2 \end{cases}$$



[Watch Video Solution](#)

39. Discuss continuity of $|x|$.



[Watch Video Solution](#)

EXERCISE 5(b) (LONG ANSWER TYPE QUESTIONS (I))

1. Examine the derivability of the following functions at the specified points :

$$|x| \text{ at } x = 0$$

 [Watch Video Solution](#)

2. Examine the derivability of the following functions at the specified points :

$$[x] \text{ at } x = 1$$

 [Watch Video Solution](#)

3. Examine the derivability of the following functions at the specified points :

$$|x|^2 \text{ at } x = 0$$

 [Watch Video Solution](#)

4. Examine the derivability of the following functions at the specified points :

$$x^3 \text{ at } x = 2$$

 [Watch Video Solution](#)

5. If $f(x)$ is differentiable at $x = a$, find $\lim_{x \rightarrow a} \frac{x^2 f(a) - a^2 f(x)}{x - a}$.

 [Watch Video Solution](#)

6. If $F(x) = f(ax)$ and $f(ax)$ is differentiable, then prove that $F'(x) = af'(ax)$, $a \neq 0$.

 [Watch Video Solution](#)

7. Show that $f(x) = \begin{cases} x \sin \frac{1}{x}, & \text{when } x \neq 0 \\ 0, & \text{when } x = 0 \end{cases}$ is continuous but not differentiable at $x = 0$

 [Watch Video Solution](#)

8. Show that the function defined by $f(x) = (3 - 2x), x < 2$ and $f(x) = 3x - 7, x \geq 2$ is not derivable at $x = 2$

 [Watch Video Solution](#)

9. Discuss continuity $f(x) = \begin{cases} \frac{e^{1/x}}{1+e^{1/x}} & \text{if } x \neq 0 \\ 0 & \text{if } x = 0 \end{cases}$ at $x = 0$

 [Watch Video Solution](#)

10. Consider the following in respect of the function

$$f(x) = \begin{cases} 2 + x, & x \geq 0 \\ 2 - x, & x < 0 \end{cases}$$

1. $\lim_{x \rightarrow 1} f(x)$ does not exist.

2. $f(x)$ is differentiable at $x=0$

3. $f(x)$ is continuous at $x=0$

Which of the above statements is /are correct?



[Watch Video Solution](#)

11. Show that the function 'f' defined as follows, is continuous at $x = 2$, but not differentiable there at :

$$f(x) = \begin{cases} 3x - 2, & 0 < x \leq 1 \\ 2x^2 - x, & 1 < x \leq 2. \\ 5x - 4, & x > 2 \end{cases}$$



[Watch Video Solution](#)

12. The function 'f' defined as :

$$f(x) = \begin{cases} x^2 + 3x + a, & \text{if } x \leq 1 \\ bx + c, & \text{if } x > 1 \end{cases}$$

is derivable for every x . Find the value of 'a' and 'b'.



[Watch Video Solution](#)

13. For what choice of a and b is the function

$$f(x) = \begin{cases} x^2, & x \leq a \\ ax + b, & x > a \end{cases} \Leftrightarrow \text{differentiable at } x = a.$$

 [Watch Video Solution](#)

14. Let $f: \mathbb{R} \rightarrow \mathbb{R}$ (\mathbb{R} is the set of real numbers) be defined as follows :

$$f(x) = \begin{cases} 2 - x, & \text{for } 1 \leq x \leq 2 \\ x - \frac{1}{2}x^2, & \text{for } x > 2 \end{cases}.$$

Examine the continuity and differentiability of $f(x)$ at $x = 2$.

 [Watch Video Solution](#)

15. Show that the function $f(x) = |x - 3|$, $x \in \mathbb{R}$, is continuous but not differentiable at $x = 3$.

 [Watch Video Solution](#)

16. Show that $f(x) = |x - 5|$ is continuous but not differentiable at $x = 5$.

 [Watch Video Solution](#)

17. Write an example of a function which is everywhere continuous but fails to be differentiable exactly at five points.

 [Watch Video Solution](#)

EXERCISE 5(c) (SHORT ANSWER TYPE QUESTIONS)

1. If
 $f(x) = x + 7$, and $g(x) = x - 7$, $x \in R$ them find $\frac{d}{dx}(f \circ g)(x)$.

 [Watch Video Solution](#)

2. $f(x) = \cos \sqrt{x}$. find $\frac{dy}{dx}$

 [Watch Video Solution](#)

3. $f(x) = (3x^2 + 2)^3(5x - 1)^2$ find $\frac{dy}{dx}$

 [Watch Video Solution](#)

4. $f(x) = (2x^2 + 3)^{\frac{5}{3}}(x + 5)^{\frac{-1}{3}}$

 [Watch Video Solution](#)

5. $f(x) = \frac{3}{2 - x}$, $x \neq 2$ find $\frac{dy}{dx}$

 [Watch Video Solution](#)

6. $h(x) = (x + 1)(x + 2)(x + 3)$. find $\frac{dy}{dx}$



Watch Video Solution

7. Use Chain Rule to find $\frac{dy}{dx}$, if $y = \left(\frac{3x - 1}{3x + 1}\right)^2$.



Watch Video Solution

8. $y = (4x^3 - 5x^2 + 1)^4$, Find $\frac{dy}{dx}$.



Watch Video Solution

9. Find $\frac{dy}{dx}$ if :

$$y = (x - 1)^{-3}, x \neq 1$$



Watch Video Solution

10. Find $\frac{dy}{dx}$ if :

$$y = 1 - 2\left(\frac{5x}{3x + 2}\right)^2 + \left(\frac{5x}{3x + 2}\right)^3$$



Watch Video Solution

11. Find $\frac{dy}{dx}$ if :

$$ax + by^2 = \cos y$$



Watch Video Solution

12. Differentiate the following w.r.t.x :

$$\sin(x^2)$$



Watch Video Solution

13. Differentiate the following w.r.t.x :

$$\sin(x^2 + 5)$$



Watch Video Solution

14. Differentiate the following w.r.t.x :

$$\tan(2x + 3)$$

 [Watch Video Solution](#)

15. Differentiate the following w.r.t.x : $\sin(x^2)$

 [Watch Video Solution](#)

16. Differentiate the following w.r.t.x :

$$\sin^4(ax + b)^2$$

 [Watch Video Solution](#)

17. Differentiate the following w.r.t.x :

$$\sin(\cot x)$$

 [Watch Video Solution](#)

18. Differentiate the following w.r.t.x :

$$\cos ec(\cot \sqrt{x})$$

 [Watch Video Solution](#)

19. Differentiate the following w.r.t.x :

$$\sin^2(x^5)$$

 [Watch Video Solution](#)

20. Differentiate the following w.r.t.x :

$$\cos^2(x^3)$$

 [Watch Video Solution](#)

21. Differentiate the following w.r.t.x :

$$\cos x^3 \cdot \sin^2(x^5)$$



Watch Video Solution

22. Differentiate the following w.r.t.x :

$$2\sqrt{\cot(x^2)}$$



Watch Video Solution

23. Differentiate the following w.r.t x. (i) $\cos^{-1}(\sin x)$ (ii)

$$\tan^{-1}\left(\frac{\sin x}{1 + \cos x}\right) \text{ (iii) } \sin^{-1}\left(\frac{2^{x+1}}{1 + 4^x}\right)$$



Watch Video Solution

24. Differentiate the following w.r.t.x :

$$\sqrt{15x^2 - x + 1}$$



Watch Video Solution

25. Differentiate the following w.r.t. x :

$$\frac{\sin(ax + b)}{\cos(cx + d)}$$



Watch Video Solution

26. Differentiate w.r.t. x the function

$\cos(a \cos x + b \sin x)$, for some constant a and b .



Watch Video Solution

27. Find $\frac{dy}{dx}$ if :

$$y = 9u^2, u = 1 - \frac{3}{2}x^2$$



Watch Video Solution

28. Find $\frac{dy}{dx}$ if :

$$y = \frac{3 - v}{2 + v}, v = \frac{4x}{1 - x^2}$$

 [Watch Video Solution](#)

29. Find $\frac{dy}{dx}$ if :

$$y = at^2, t = \frac{x}{2a}$$

 [Watch Video Solution](#)

30. Find $\frac{dy}{dx}$ at $x=1, y=\pi/4$ if $\sin^2 y + \cos xy = 0$

 [Watch Video Solution](#)

31. If $x^{16}y^9 = (x^2 + y)^{17}$, prove that $\frac{dy}{dx} = \frac{2y}{x}$.

 [Watch Video Solution](#)

32. Differentiate the following w.r.t. x :

$$|2x - 1|$$

 [Watch Video Solution](#)

33. Differentiate the following w.r.t. x :

$$|2x^2 - 3|$$

 [Watch Video Solution](#)

34. If $y + \sin y = \cos x$, then find the values of 'y' for which $\frac{dy}{dx}$ is valid.

 [Watch Video Solution](#)

EXERCISE 5(d) (SHORT ANSWER TYPE QUESTIONS)

1. Find the derivative of y w.r.t. x in each of the following :

$$xy + y^2 = \tan x + y$$

 [Watch Video Solution](#)

2. Find the derivative of y w.r.t. x in each of the following :

$$x^2 + y^2 - 3xy + 1$$

 [Watch Video Solution](#)

3. Find the derivative of y w.r.t. x in each of the following :

$$\sin^2 y + \cos xy = \pi.$$

 [Watch Video Solution](#)

4. Find the derivative of y w.r.t. x in each of the following :

$$\sqrt{x} + \sqrt{y} = 20$$





[Watch Video Solution](#)

5. Find the derivative of y w.r.t. x in each of the following :

$$y(y + 1) = x(x + 1)(x + 2)$$



[Watch Video Solution](#)

6. Find the derivative of y w.r.t. x in each of the following :

$$x^2 = \frac{x + 3y}{x - 3y}$$



[Watch Video Solution](#)

7. Find the derivative of y w.r.t. x in each of the following :

$$\frac{y}{x + y} = 3 + x^3.$$



[Watch Video Solution](#)

8. Find the derivatives of $f(x)$ w.r.t. x in the following :

$$f(x) = \sqrt[3]{2x^4 + x^2 - x}$$



[Watch Video Solution](#)

9. Find the derivatives of $f(x)$ w.r.t. x in the following :

$$f(x) = \sqrt[3]{ax + b}$$



[Watch Video Solution](#)

10. Find the derivatives of $f(x)$ w.r.t. x in the following :

$$f(x) = (x^2 + x + 5)^{1/3} (x^3 + 1)^{2/3}$$



[Watch Video Solution](#)

11. Find the derivatives of $f(x)$ w.r.t. x in the following :

$$g(x) = \sqrt[3]{5x - 9} \sqrt[3]{3x - 4}$$





Watch Video Solution

12. Obtain $\frac{dy}{dx}$ when :

$$x^2 + y^2 + 2axy = 0$$



Watch Video Solution

13. Find $\frac{dy}{dx}$ if $x^3 + y^3 - 3axy = 0$.



Watch Video Solution

14. If $x^2 + y^2 + 2gx + 2fy + c = 0$ then $\frac{dy}{dx} =$



Watch Video Solution

15. Obtain $\frac{dy}{dx}$ when :

$$x^4 + y^4 + 4xy - 100 = 0$$





Watch Video Solution

16. If $ax^2 + 2hxy + by^2 = 0$ then $\frac{dy}{dx} =$



Watch Video Solution

17. If $\sqrt{x} + \sqrt{y} = 5$, find $\frac{dy}{dx}$ at $(4, 9)$.



Watch Video Solution

18. If $x\sqrt{1+y} + y\sqrt{1+x} = 0$, prove that $\frac{dy}{dx} = -\frac{1}{(x+1)^2}$.



Watch Video Solution

19. If $y = \sqrt{x} + \frac{1}{\sqrt{x}}$, then the value of $\left(2x \frac{dy}{dx} + y\right)$ is-



Watch Video Solution

20. Find $\frac{dy}{dx}$ for each of the following

$$y = (x^2 + 3x + 5)(x^2 - 2)^2$$

 [Watch Video Solution](#)

21. Find $\frac{dy}{dx}$ for each of the following

$$y = \left(\sqrt{x} + \frac{1}{\sqrt{x}} \right) (1 + x + x^2)$$

 [Watch Video Solution](#)

22. Find $\frac{dy}{dx}$ for each of the following

$$y = \left(\frac{x - \sqrt{x}}{1 - 2x} \right)^2$$

 [Watch Video Solution](#)

23. Find $\frac{dy}{dx}$ for each of the following

$$y = \left(\frac{1}{1+x} \right) \left(x^{-2} + \frac{2}{x} - 1 \right) + \sqrt[3]{x} - \frac{1}{\sqrt[3]{x}}$$

 [Watch Video Solution](#)

24. Find $\frac{dy}{dx}$ for each of the following

$$y = \sqrt[3]{x^2(x^2 + 3)}.$$

 [Watch Video Solution](#)

25. If $\cos y = x \cos(a + y)$, with $\cos a \neq \pm 1$, prove that

$$\frac{dy}{dx} = \frac{\cos^2(a + y)}{\sin a}.$$

 [Watch Video Solution](#)

26. If $\sin y = x \sin(a + y)$, prove that, $\frac{dy}{dx} = \frac{\sin a}{1 - 2x \cos a + x^2}$.

 [Watch Video Solution](#)

EXERCISE 5(e) (SHORT ANSWER TYPE QUESTIONS)

1. Differentiate the following w.r.t. x :

$$\cos^{-1}\left(\frac{x}{x+1}\right)$$



[Watch Video Solution](#)

2. $\sin^{-1}(x\sqrt{x}), 0 \leq x \leq 1$



[Watch Video Solution](#)

3. Differentiate the following w.r.t. x :

$$(\tan^{-1} x)^2$$



[Watch Video Solution](#)

4. Differentiate the following w.r.t. x :

$$x \sec^{-1} x.$$



[Watch Video Solution](#)

5. Differentiate the following w.r.t. x :

$$\cos^{-1}(\sin x)$$



[Watch Video Solution](#)

6. Differentiate the following w.r.t. x :

$$(i) \sin^{-1} 2x \quad (ii) \tan^{-1} \sqrt{x} \quad (iii) \cos^{-1}(\cot x)$$



[Watch Video Solution](#)

7. Differentiate the following w.r.t. x :

$$(\cot^{-1} x)^2$$



[Watch Video Solution](#)

8. Differentiate the following w.r.t. x :

$$\sin(2 \sin^{-1} x)$$

[Watch Video Solution](#)

9. Find the derivatives w.r.t. x :

$$\tan^{-1} \left(\frac{\sin x}{1 + \cos x} \right)$$

[Watch Video Solution](#)

10. $\tan^{-1} \frac{\cos x}{1 + \sin x} =$

[Watch Video Solution](#)

11. $\cot^{-1} \left(\frac{1 + \cos x}{\sin x} \right)$

[Watch Video Solution](#)

12. Differentiate the following w.r.t. x :

$$\sin^{-1}(1 - 2x^2)$$

[Watch Video Solution](#)

13. $\sin^{-1}(3x - 4x^3)$

[Watch Video Solution](#)

14. Differentiate the following w.r.t. x :

$$\cos^{-1}(4x^3 - 3x), \quad -1 < x < 1$$

[Watch Video Solution](#)

15. Differentiate the following w.r.t. x :

$$\sin^{-1}\left(\frac{2x}{1+x^2}\right), \quad -1 < x < 1$$

 [Watch Video Solution](#)

16. $\operatorname{cosec}^{-1}\left(\frac{1+x^2}{2x}\right)$

 [Watch Video Solution](#)

17. Differentiate the following w.r.t. x :

$$\sin^{-1}\left(\frac{1-x^2}{1+x^2}\right), \quad 0 < x < 1$$

 [Watch Video Solution](#)

18. Differentiate the following w.r.t. x :

$$\cos^{-1}\left(\frac{1-x^2}{1+x^2}\right), \quad 0 < x < 1$$

 [Watch Video Solution](#)

19. Differentiate the following w.r.t. x :

$$\cos^{-1}\left(\frac{2x}{1+x^2}\right), -1 < x < 1$$



Watch Video Solution

20. Differentiate the following w.r.t. x :

$$\tan^{-1}\left(\frac{2x}{1-x^2}\right), 0 < x < 1$$



Watch Video Solution

21. $\tan^{-1}\left(\frac{3x-x^3}{1-3x^2}\right)$



Watch Video Solution

22. निम्न फलन का x के सापेक्ष अवकल गुणांक ज्ञात कीजिए - $\tan^{-1}\left[\frac{x^{1/3}+a^{1/3}}{1-x^{1/3}a^{1/3}}\right]$



 [Watch Video Solution](#)

23. $\tan^{-1} \frac{x}{\sqrt{a^2 - x^2}}$

 [Watch Video Solution](#)

EXERCISE 5(e) (LONG ANSWER TYPE QUESTIONS (II))

1. Differentiate $\tan^{-1} \left\{ \frac{\sqrt{1+x^2} - 1}{x} \right\}$ w.r.t. x .

 [Watch Video Solution](#)

2. $\tan^{-1} \left(\frac{\sqrt{1+x^2} + 1}{x} \right)$

 [Watch Video Solution](#)

3. Differentiate the following w.r.t. x :

$$\tan^{-1} \left(\frac{\sqrt{1 + a^2 x^2} - 1}{ax} \right)$$

 [Watch Video Solution](#)

4. Differentiate the following w.r.t. x :

$$\cot^{-1} \left(\frac{\sqrt{1 + x^2} - 1}{x} \right), x \neq 0$$

 [Watch Video Solution](#)

5. Differentiate the following w.r.t. x :

$$\cot^{-1} \left(\frac{1 + x}{1 - x} \right)$$

 [Watch Video Solution](#)

6. Differentiate the following w.r.t. x :

$$\cot^{-1}\left(\sqrt{1+x^2} - x\right).$$



Watch Video Solution

7. Differentiate the following w.r.t. x :

$$\tan^{-1}(\sec x + \tan x).$$



Watch Video Solution

8. Differentiate the following w.r.t. x :

$$\tan^{-1} \sqrt{\frac{1 - \cos x}{1 + \cos x}}$$



Watch Video Solution

9. Differentiate w.r.t. x :

$$(i) \tan^{-1} \left\{ \sqrt{\frac{1 + \cos x}{1 - \cos x}} \right\}$$

$$(ii) \tan^{-1} \left\{ \sqrt{\frac{1 + \sin x}{1 - \sin x}} \right\}$$

 [Watch Video Solution](#)

10. Differentiate the following w.r.t. x :

$$\tan^{-1} \sqrt{\frac{1 + \sin x}{1 - \sin x}}$$

 [Watch Video Solution](#)

11. Differentiate the following w.r.t. x :

$$\sin^{-1} \left(\sqrt{\frac{1 + x^2}{2}} \right).$$

 [Watch Video Solution](#)

12. If $y = \tan^{-1} \left(\frac{2y}{1 - x^2} \right) + \sec^{-1} \left(\frac{1 + x^2}{1 - x^2} \right)$, $x > 0$, prove that

$$\frac{dy}{dx} = \frac{4}{1 + x^2}.$$

 [Watch Video Solution](#)

13. Find $\frac{dy}{dx}$ of $y = \cot^{-1} \left[\frac{\sqrt{1 + \sin x} + \sqrt{1 - \sin x}}{\sqrt{1 + \sin x} - \sqrt{1 - \sin x}} \right]$

 [Watch Video Solution](#)

14. Find $\frac{dy}{dx}$ of $y = \cot^{-1} \left[\frac{\sqrt{1 + \sin x} + \sqrt{1 - \sin x}}{\sqrt{1 + \sin x} - \sqrt{1 - \sin x}} \right]$

 [Watch Video Solution](#)

15. If $y = \tan^{-1} \frac{(\sqrt{1 + \sin x} + \sqrt{1 - \sin x})}{(\sqrt{1 + \sin x} - \sqrt{1 - \sin x})}$, find $\frac{dy}{dx}$.

 [Watch Video Solution](#)

16. $\frac{d}{dx} \left[\cos^{-1} \sqrt{\frac{(1+x)}{2}} \right]$

 [Watch Video Solution](#)

17. If $y = \tan^{-1} \left\{ \frac{\sqrt{1+x^2} + \sqrt{1-x^2}}{\sqrt{1+x^2} - \sqrt{1-x^2}} \right\}$, then

 [Watch Video Solution](#)

18. If $y = \tan^{-1} \frac{\sqrt{1+x^2} - \sqrt{1-x^2}}{\sqrt{1+x^2} + \sqrt{1-x^2}}$, then the value of $\frac{dy}{dx}$ is -

 [Watch Video Solution](#)

19. If $y = \sin \left[2 \tan^{-1} \left\{ \sqrt{\frac{1-x}{1+x}} \right\} \right]$, find $\frac{dy}{dx}$

 [Watch Video Solution](#)

20. If $y = \tan^{-1} \left(\frac{5ax}{a^2 - 6x^2} \right)$, then $\frac{dy}{dx} =$

 [Watch Video Solution](#)

21. Find $\frac{dy}{dx}$ if $y = \frac{\tan^{-1}(4x)}{1+5x^2} + \frac{\tan^{-1}(2+3x)}{3-2x}$



Watch Video Solution

22. If $y = \cos^{-1}\left(\frac{3x + 4\sqrt{1-x^2}}{5}\right)$, find $\frac{dy}{dx}$



Watch Video Solution

EXERCISE 5(e) (LONG ANSWER TYPE QUESTIONS (II))

1. If $f(x) = \tan^{-1} x$, $g(x) = \tan^{-1}\left(\frac{1+x}{1-x}\right)$ for $|x| < 1$, show that $f'(x) = g'(x)$ and $g(x) - f(x) = \frac{\pi}{4}$.



Watch Video Solution

2. If $y = \frac{2}{\sqrt{a^2 - b^2}} \tan^{-1}\left[\sqrt{\frac{a-b}{a+b}} \tan \frac{x}{2}\right]$, prove that $\frac{dy}{dx} = \frac{1}{a + b \cos x}$, $a > b > 0$.



Watch Video Solution

3. If $\sqrt{1-x^6} + \sqrt{1-y^6} = a(x^3 - y^3)$, then prove that

$$\frac{dy}{dx} = \frac{x^2}{y^2} \sqrt{\frac{1-y^6}{1-x^6}}$$

 [Watch Video Solution](#)

4. $\frac{d}{dx} [\tan^{-1}(\cot x) + \cot^{-1}(\tan x)] =$

 [Watch Video Solution](#)

5. If $y = \sin^{-1}(\cos x) + \cos^{-1}(\sin x)$, then prove that :

$$\frac{dy}{dx} + 2 = 0.$$

 [Watch Video Solution](#)

EXERCISE 5(f) (SHORT ANSWER TYPE QUESTIONS)

1. Differentiate the following w.r.t. x :

$$e^{-x}$$



Watch Video Solution

2. Differentiate the following w.r.t. x :

$$e^{\sin^{-1} x}$$



Watch Video Solution

3. Differentiate the following w.r.t. x :

$$e^{\cot^{-1} x^2}$$



Watch Video Solution

4. Differentiate the following w.r.t. x :

$$e^{m \log x}$$





[Watch Video Solution](#)

5. Differentiate the following w.r.t. x :

$$\sqrt{e^{\sqrt{x}}}, x > 0$$



[Watch Video Solution](#)

6. Differentiate the following w.r.t. x :

$$\log(\sin x)$$



[Watch Video Solution](#)

7. Differentiate the following w.r.t. x :

$$\log(\cos e^x)$$



[Watch Video Solution](#)

8. Differentiate the following w.r.t. x : $\sin(\tan^{-1} e^{-x})$



Watch Video Solution

9. Differentiate the following w.r.t. x :

$$\frac{\cos x}{\tan x}, x > 0$$



Watch Video Solution

10. Differentiate the following w.r.t. x :

$$\sqrt{\tan x} a^x.$$



Watch Video Solution

11. Differentiate the following w.r.t. x :

$$e^x \sin x$$



Watch Video Solution

12. Differentiate the following w.r.t. x :

$$\sqrt{x} \log x^2$$



[Watch Video Solution](#)

13. Differentiate the following w.r.t. x :

$$x^{-1/3} e^x$$



[Watch Video Solution](#)

14. Differentiate the following w.r.t. x

$$x \cdot \sin x \cdot e^x$$



[Watch Video Solution](#)

15. Differentiate the following w.r.t. x :

$$e^{\sin^{-1}(x+1)}$$



Watch Video Solution

16. Differentiate the following w.r.t. x :

$$\tan\{\log(\sin x)\}$$



Watch Video Solution

17. Differentiate the following w.r.t. x :

$$e^{\sin \sqrt{x}}$$



Watch Video Solution

18. Differentiate the following w.r.t. x :

$$e^{\cos^{-1}(x+1)}$$





[Watch Video Solution](#)

19. Differentiate the following w.r.t. x :

$$e^{\cos^{-1} x^2}$$



[Watch Video Solution](#)

20. Differentiate the following w.r.t. x :

$$\sqrt{1-x^2} \cdot e^{5x}$$



[Watch Video Solution](#)

21. Differentiate the following w.r.t. x :

$$e^{\sqrt{1-x^2}} \cdot \tan x$$



[Watch Video Solution](#)

22. Differentiate the following w.r.t. x :

$$e^{\sin^2} \left(2 \tan^{-1} \sqrt{\frac{1-x}{1+x}} \right)$$

 [Watch Video Solution](#)

23. Differentiate the following w.r.t. x :

$$\frac{\log x}{x}$$

 [Watch Video Solution](#)

24. Differentiate the following w.r.t. x :

$$\frac{e^x}{x}$$

 [Watch Video Solution](#)

25. Differentiate the following w.r.t. x :

$$\frac{\log x}{e^x}$$



[Watch Video Solution](#)

26. Differentiate the following w.r.t. x :

$$\log(\cos 5x)$$



[Watch Video Solution](#)

27. Differentiate the following w.r.t. x :

$$\frac{1}{\log \cos x}$$



[Watch Video Solution](#)

EXERCISE 5(f) (LONG ANSWER TYPE QUESTIONS (I))

1. Differentiate the following w.r.t. x :

$$(x^2 + 7x + 2)(e^x - \sin x)$$



[Watch Video Solution](#)

2. Differentiate the following w.r.t. x :

$$e^{-3x} \sin^2 3x$$



[Watch Video Solution](#)

3. Differentiate the following w.r.t. x :

$$e^{-x^2} \sin(\log x)$$



[Watch Video Solution](#)

4. Differentiate the following w.r.t. x :

$$\tan^{-1} \left(\frac{e^{2x} + 1}{e^{2x} - 1} \right)$$



[Watch Video Solution](#)

5. Differentiate the following w.r.t. x :

$$\log \left((x + 3) + \sqrt{x^2 + 6x + 3} \right)$$



[View Text Solution](#)

6. Differentiate the following w.r.t.x. $\log(x + \sqrt{a^2 + x^2})$



[Watch Video Solution](#)

7. Differentiate the following w.r.t. x :

$$x\sqrt{x^2 + 1} + \log(x + \sqrt{x^2 + 1})$$



[View Text Solution](#)

8. Differentiate the following w.r.t. x :

$$\sqrt{x^2 + 1} - \log\left(\frac{1}{x} + \sqrt{1 + \frac{1}{x^2}}\right)$$



[Watch Video Solution](#)

9. Differentiate the following w.r.t. x :

$$\frac{e^x(x - 1)}{(x^2 + 1)}$$

 [Watch Video Solution](#)

10. Differentiate the following w.r.t. x :

$$\frac{e^{ax}}{\sin(bx + c)}$$

 [Watch Video Solution](#)

11. Differentiate the following w.r.t. x :

$$\frac{1}{3}e^x - 5e$$

 [Watch Video Solution](#)

12. Differentiate the following w.r.t. x :

$$e^x + 2 \cos x$$



[Watch Video Solution](#)

13. Differentiate the following w.r.t. x :

$$x^2 e^x \sin x$$



[Watch Video Solution](#)

14. Differentiate the following w.r.t. x :

$$e^{\sec^2 x} + 3 \cos^{-1} x.$$



[Watch Video Solution](#)

15. Differentiate the following w.r.t. x :

$$\log\left(\sin \sqrt{1+x^2}\right)$$



[Watch Video Solution](#)

16. Differentiate the following w.r.t. x :

$$\sin(\log x), x > 0$$

 [Watch Video Solution](#)

17. Differentiate the following w.r.t. x :

$$\log(\cos 5x)$$

 [Watch Video Solution](#)

18. Differentiate the following w.r.t. x :

$$\cot(\log x + e^{\sqrt{x}})$$

 [View Text Solution](#)

19. Differentiate the following w.r.t. x :

$$2l_n\left(\frac{x-1}{x+1}\right)$$



Watch Video Solution

20. Differentiate the following w.r.t. x :

$$x^2 \ln \left(\sqrt{\frac{x^2 + 9}{x^2 + 4}} \right)$$



Watch Video Solution

21. Differentiate the following w.r.t. x :

$$\ln(\sec x + \tan x)$$



Watch Video Solution

22. Differentiate the following w.r.t. x :

$$\ln \left(\sqrt{\frac{1 - \cos x}{1 + \cos x}} \right)$$



Watch Video Solution

23. Differentiate the following w.r.t. x :

$$\log\left(\frac{1+x}{1-x}\right)$$

 [Watch Video Solution](#)

24. Differentiate the following w.r.t. x :

$$\log \tan\left(\frac{\pi}{4} + \frac{x}{2}\right)$$

 [View Text Solution](#)

25. Differentiate the following w.r.t. x :

$$\log\left(\frac{x + \sqrt{x^2 - a^2}}{x - \sqrt{x^2 - a^2}}\right)$$

 [Watch Video Solution](#)

26. Differentiate the following w.r.t. x :

$$\log \sin^{-1}\left(2x\sqrt{1-x^2}\right)$$



[View Text Solution](#)

27. Differentiate the following w.r.t. x :

$$\sqrt{\log\left(\sin\left(\frac{x^2}{3} - 1\right)\right)}$$



[Watch Video Solution](#)

28. Find $\frac{dy}{dx}$ when :

$$\sin y + \log y = x^2 + 18x + 3$$



[View Text Solution](#)

29. Find $\frac{dy}{dx}$ when :

$$xy + xe^{-y} + ye^x = x^2.$$



[View Text Solution](#)

30. if $e^{x+y} = xy$, show that $\frac{dy}{dx} = \frac{y(1-x)}{x(y-1)}$

 [Watch Video Solution](#)

31. if $y = \frac{\sin^{-1} x}{\sqrt{1-x^2}}$, prove that $(1-x^2)\frac{dy}{dx} = xy + 1$

 [Watch Video Solution](#)

32. if $x = \tan\left(\frac{1}{a}\log y\right)$, show that $(1+x^2)\frac{dy}{dx} = ay$.

 [Watch Video Solution](#)

33. Differentiate $\tan^{-1}\left(\frac{2^{x+1}}{1-4^x}\right)$ with respect to x .

 [View Text Solution](#)

1. Prove that :

$$\frac{d}{dx} \left[\log \left(\frac{x^2 + x + 1}{x^2 - x + 1} \right) + \frac{2}{\sqrt{3}} \tan^{-1} \left(\frac{x\sqrt{3}}{1 - x^2} \right) \right] = \frac{4}{1 + x^2 + x^4}$$

 [Watch Video Solution](#)

2. Prove that :

$$\frac{d}{dx} \left[\frac{x \sin^{-1} x}{\sqrt{1 - x^2}} + \log \sqrt{1 - x^2} \right] = \frac{\sin^{-1} x}{(1 - x^2)^{3/2}}.$$

 [Watch Video Solution](#)

3. If $y = \log_{10} \sin x$, prove that :

$$\frac{dy}{dx} = (\log_{10} e) \cot x.$$

 [Watch Video Solution](#)

4. If $y = \sin[\sin(\log 3x)]$, find $\frac{dy}{dx}$.



[View Text Solution](#)

5. Find, from first principle, the derivative of the following w.r.t. x :

$$e^{2x}$$



[Watch Video Solution](#)

6. Find, from first principle, the derivative of the following w.r.t. x :

$$e^{\sqrt{x}}$$



[Watch Video Solution](#)

7. Find, from first principle, the derivative of the following w.r.t. x :

$$xe^x$$



[Watch Video Solution](#)

8. Find, from first principle, the derivative of the following w.r.t. x :

$$e^{\sin x}$$



Watch Video Solution

9. Find, from first principle, the derivative of the following w.r.t. x :

$$e^{\sqrt{\tan x}}$$



Watch Video Solution

10. Find, from first principle, the derivative of the following w.r.t. x :

$$\log(\sin x)$$



Watch Video Solution

11. Find, from first principle, the derivative of the following w.r.t. x :

$$\log(\cos x)$$





Watch Video Solution

12. Find, from first principle, the derivative of the following w.r.t. x :

$$\log x^2$$



Watch Video Solution

13. Find, from first principle, the derivative of the following w.r.t. x :

$$\cos(\log x), \text{ where } x > 0.$$



Watch Video Solution

EXERCISE 5(g) (SHORT ANSWER TYPE QUESTIONS)

1. If x and y are connected parametrically by the equations given, without

eliminating the parameter, Find $\frac{dy}{dx} \cdot x = 2at^2, y = at^4$



Watch Video Solution

2. Find $\frac{dy}{dx}$, if x and y are connected parametrically by the equations, given below without eliminating the parameter:

$$x = \frac{2t}{1+t^2}, y = \frac{1-t^2}{1+t^2}$$

 [Watch Video Solution](#)

3. Find $\frac{dy}{dx}$, if x and y are connected parametrically by the equations, given below without eliminating the parameter:

$$x = \frac{1-t^2}{1+t^2}, y = \frac{2t}{1+t^2}$$

 [Watch Video Solution](#)

4. Find $\frac{dy}{dx}$, if x and y are connected parametrically by the equations, given below without eliminating the parameter:

$$x = \frac{a(1+t^2)}{1-t^2}, y = \frac{2t}{1-t^2}$$

 [Watch Video Solution](#)

5. Find $\frac{dy}{dx}$, if x and y are connected parametrically by the equations, given below without eliminating the parameter:

$$x = a\sqrt{\frac{t^2 - 1}{t^2 + 1}}, y = at\sqrt{\frac{t^2 - 1}{t^2 + 1}}.$$

 [Watch Video Solution](#)

6. If x and y are connected parametrically by the equations given, without eliminating the parameter, Find $\frac{dy}{dx}$. $x = \sin t, y = \cos 2t$

 [Watch Video Solution](#)

7. Find $\frac{dy}{dx}$, if x and y are connected parametrically by the equations, given below without eliminating the parameter:

$$x = \log t, y = \sin t$$

 [Watch Video Solution](#)

8. Find $\frac{dy}{dx}$, if x and y are connected parametrically by the equations, given below without eliminating the parameter:

$$x = e^t \cos t, y = e^t \sin t \quad \text{at} \quad t = \frac{\pi}{2}.$$



[Watch Video Solution](#)

9. Find $\frac{dy}{dx}$, if $x = a \cos \theta$, $y = a \sin \theta$



[Watch Video Solution](#)

10. Find $\frac{dy}{dx}$, if x and y are connected parametrically by the equations, given below without eliminating the parameter:

$$x = a \cos \theta, y = b \sin \theta$$



[Watch Video Solution](#)

11. If x and y are connected parametrically by the equations given, without eliminating the parameter, Find $\frac{dy}{dx}$. $x = a \cos \theta$, $y = b \cos \theta$

 [Watch Video Solution](#)

12. Find $\frac{dy}{dx}$, if x and y are connected parametrically by the equations, given below without eliminating the parameter:

$$x = a \cos^2 \theta, y = b \sin^2 \theta$$

 [Watch Video Solution](#)

13. Find $\frac{dy}{dx}$, if x and y are connected parametrically by the equations, given below without eliminating the parameter:

$$x = 2 \cos^2 \theta, y = 2 \sin^2 \theta$$

 [Watch Video Solution](#)

14. Find $\frac{dy}{dx}$, if x and y are connected parametrically by the equations, given below without eliminating the parameter:

$$x = 2 \cos^3 \theta, y = 2 \sin^3 \theta$$

 [Watch Video Solution](#)

15. Find $\frac{dy}{dx}$, if x and y are connected parametrically by the equations, given below without eliminating the parameter:

$$x = 3 \cos^3 \theta, y = 3 \sin^3 \theta$$

 [Watch Video Solution](#)

16. Find $\frac{dy}{dx}$, if x and y are connected parametrically by the equations, given below without eliminating the parameter:

$$x = c \tan \theta, y = c \cot \theta$$

 [Watch Video Solution](#)

17. Find $\frac{dy}{dx}$, if x and y are connected parametrically by the equations, given below without eliminating the parameter:

$$x = a \tan^2 \theta, y = b \sec^2 \theta$$

 [Watch Video Solution](#)

18. If x and y are connected parametrically by the equations given, without eliminating the parameter, Find $\frac{dy}{dx}$. $x = a \sec \theta, y = b \tan \theta$

 [Watch Video Solution](#)

19. If x and y are connected parametrically by the equations given, without eliminating the parameter, Find $\frac{dy}{dx}$. $x = \frac{\sin^3 t}{\sqrt{\cos 2t}}, y = \frac{\cos^3 t}{\sqrt{\cos 2t}}$

 [Watch Video Solution](#)

20. If $y = a(\theta + \sin \theta), x = a(1 - \cos \theta)$, then $\frac{dy}{dx} =$

 [Watch Video Solution](#)

 Watch Video Solution

21. Find $\frac{dy}{dx}$ if $x = a(\theta - \sin \theta)$ and $y = a(1 - \cos \theta)$.

 Watch Video Solution

22. If x and y are connected parametrically by the equations given, without eliminating the parameter, Find $\frac{dy}{dx}$.

$$x = a(\theta - \sin \theta), y = a(1 + \cos \theta)$$

 Watch Video Solution

23. Find $\frac{dy}{dx}$, if x and y are connected parametrically by the equations, given below without eliminating the parameter:

$$y = a(\theta + \sin \theta), x = a(1 + \cos \theta)$$

 Watch Video Solution

24. Find $\frac{dy}{dx}$, if x and y are connected parametrically by the equations, given below without eliminating the parameter:

$$x = a(\theta - \sin \theta), y = b(1 + \cos \theta)$$

 [Watch Video Solution](#)

25. If x and y are connected parametrically by the equations given, without eliminating the parameter, Find $\frac{dy}{dx}$.

$$x = \cos \theta - \cos 2\theta, y = \sin \theta - \sin 2\theta$$

 [Watch Video Solution](#)

26. Find $\frac{dy}{dx}$, if x and y are connected parametrically by the equations, given below without eliminating the parameter:

$$x = a(\cos \theta - \cos 2\theta), y = a(\sin \theta - \sin 2\theta)$$

 [Watch Video Solution](#)

27. If x and y are connected parametrically by the equations given, without eliminating the parameter, Find $\frac{dy}{dx}$.

$$x = a(\cos \theta + \theta \sin \theta), y = a(\sin \theta - \theta \cos \theta)$$

 [Watch Video Solution](#)

28. Find $\frac{dy}{dx}$, if x and y are connected parametrically by the equations, given below without eliminating the parameter:

$$x = \cos 2\theta + 2 \cos \theta, y = \sin 2\theta - 2 \sin \theta.$$

 [Watch Video Solution](#)

EXERCISE 5(g) (LONG ANSWER TYPE QUESTIONS (I))

1. Find $\frac{dy}{dx}$, if x and y are connected parametrically by the equations, given below without eliminating the parameter:

$$x = a \sin \theta, y = a \left(\cos \theta + \log \tan \frac{\theta}{2} \right)$$

 [Watch Video Solution](#)

2. Find $\frac{dy}{dx}$, if x and y are connected parametrically by the equations, given below without eliminating the parameter:

$$y = 12(1 - \cos t), x = 10(t - \sin t), -\frac{\pi}{2} < t < \frac{\pi}{2}.$$

 [Watch Video Solution](#)

3. If $x = 2 \cos \theta - \cos 2\theta$, $y = 2 \sin \theta - \sin 2\theta$, find $\frac{dy}{dx}$ at $\theta = \frac{\pi}{2}$.

 [Watch Video Solution](#)

4. If $x = a \left(\cos \theta + \log \tan \frac{\theta}{2} \right)$, $y = a \sin \theta$, find $\frac{dy}{dx}$ at $\theta = \frac{\pi}{3}$.

 [Watch Video Solution](#)

5. Find $\frac{dy}{dx}$, where $x = t^3 + \frac{1}{t}$ and $y = (t + t^2)^3$.

 [Watch Video Solution](#)

6. For a positive constant a find $\frac{dy}{dx}$, where $y = a^{t+\frac{1}{t}}$ and $x = \left(t + \frac{1}{t}\right)^a$.

 [Watch Video Solution](#)

7. If $x = a \sin 2t(1 + \cos 2t)$ and $y = b \cos 2t(1 - \cos 2t)$, show that at $\frac{\pi}{4}$, $\frac{dy}{dx} = \frac{b}{a}$.

 [Watch Video Solution](#)

8. If $x = \frac{1 + \log t}{t^2}$, $y = \frac{3 + 2 \log t}{t}$, $t > 0$, prove that :

$$\frac{dy}{dx} = t$$

 [Watch Video Solution](#)

9. If $x = \frac{1 + \log t}{t^2}$, $y = \frac{3 + 2 \log t}{t}$, $t > 0$, prove that :

$$y \frac{dy}{dx} - 2x \left(\frac{dy}{dx} \right)^2 = 1.$$



Watch Video Solution

10. Find $\frac{dy}{dx}$, when :

$$x = e^t(\sin t + \cos t) \text{ and } y = e^t(\sin t - \cos t).$$



Watch Video Solution

EXERCISE 5(h) (SHORT ANSWER TYPE QUESTIONS)

1. Differentiate the following w.r.t. as indicated :

$$x^2 \text{ w.r.t. } x^3.$$



Watch Video Solution

2. Differentiate the following w.r.t. as indicated :

$$\frac{ax + b}{cx + d} \text{ w.r.t. } \frac{a'x + b'}{c'x + d'}$$

 [Watch Video Solution](#)

3. Differentiate the following w.r.t. as indicated :

$$e^{2x} \text{ w.r.t. } e^x$$

 [Watch Video Solution](#)

4. Differentiate the following w.r.t. as indicated :

$$\frac{x^2}{1 + x^2} \text{ w.r.t. } x^2$$

 [Watch Video Solution](#)

5. Differentiate the following w.r.t. as indicated :

$$(x^2 - 1)^{4/5} \text{ w.r.t. } |x|$$





[Watch Video Solution](#)

6. Differentiate the following w.r.t. as indicated :

$$\cos^{-1} \theta \text{ w.r.t. } \log(1 + \theta)$$



[Watch Video Solution](#)

7. Differentiate the following w.r.t. as indicated :

$$\sin^{-1} \theta \text{ w.r.t. } \log(1 + \theta)$$



[Watch Video Solution](#)

8. Differentiate the following w.r.t. as indicated :

$$\tan x \text{ w.r.t. } \cos x$$



[Watch Video Solution](#)

9. Differentiate the following w.r.t. as indicated :

$\sec x$ w.r.t. $\cos ec x$



Watch Video Solution

10. Differentiate the following w.r.t. as indicated :

$\sin^2 x$ w.r.t. $e^{\cos x}$



Watch Video Solution

11. Differentiate the following w.r.t. as indicated :

$\tan^2 x$ w.r.t. $\sec^2(x^2)$



Watch Video Solution

EXERCISE 5(h) (LONG ANSWER TYPE QUESTIONS (I))

1. Differentiate w.r.t. as indicated :

$$\cos^{-1}\left(\frac{1}{\sqrt{1+x^2}}\right) \text{ w.r.t. } \tan^{-1}x$$



[Watch Video Solution](#)

2. Differentiate w.r.t. as indicated :

$$\sin^{-1}\left(\frac{2x}{1+x^2}\right) \text{ w.r.t. } \tan^{-1}x$$



[Watch Video Solution](#)

3. Differentiate w.r.t. as indicated :

$$\sin^{-1}\left(\frac{2x}{1+x^2}\right) \text{ w.r.t. } \tan^{-1}\left(\frac{2x}{1-x^2}\right)$$



[Watch Video Solution](#)

4. Differentiate w.r.t. as indicated :

$$\tan^{-1}\frac{3x-x^3}{1-3x^2} \text{ w.r.t. } \tan^{-1}\frac{2x}{1-x^2}$$



[Watch Video Solution](#)

5. Differentiate $\cos^{-1}\left(\frac{1-x^2}{1+x^2}\right)$ with respect to $\tan^{-1}\left(\frac{3x-x^3}{1-3x^2}\right)$.



[Watch Video Solution](#)

6. Differentiate w.r.t. as indicated :

$\tan^{-1}\left(\frac{3x-x^3}{1-3x^2}\right)$ w.r.t. $\tan^{-1}\left(\frac{x}{\sqrt{1-x^2}}\right)$



[Watch Video Solution](#)

7. Differentiate w.r.t. as indicated :

$\cos^{-1}\left(\frac{1-x^2}{1+x^2}\right)$ w.r.t. $\sin^{-1}\left(\frac{2x}{1+x^2}\right)$



[Watch Video Solution](#)

8. Differentiate w.r.t. as indicated :

$$\tan^{-1} \left(\frac{\sqrt{1 + a^2 x^2} - 1}{ax} \right) \text{ w.r.t. } \tan^{-1} ax$$

 [Watch Video Solution](#)

9. Differentiate w.r.t. as indicated :

$$\tan^{-1} \left(\frac{\sqrt{1 + x^2} - 1}{x} \right) \text{ w.r.t. } \sin^{-1} \left(\frac{2x}{1 + x^2} \right)$$

 [Watch Video Solution](#)

10. Differentiate w.r.t. as indicated :

$$\tan^{-1} \left(\frac{x}{\sqrt{1 - x^2}} \right) \text{ w.r.t. } \sin^{-1} (2x\sqrt{1 - x^2}).$$

 [Watch Video Solution](#)

11. Differentiate w.r.t. as indicated :

$$\tan^{-1}\left(\frac{x}{1 + \sqrt{1 - x^2}}\right) \text{ w.r.t. } \sin\left(2 \cot^{-1} \sqrt{\frac{1+x}{1-x}}\right)$$

 [Watch Video Solution](#)

12. Write the derivative of e^x wrt. \sqrt{x}

 [Watch Video Solution](#)

13. Differentiate w.r.t. as indicated :

$$\log_{10} x \text{ w.r.t. } x^2$$

 [Watch Video Solution](#)

14. Differentiate w.r.t. as indicated :

$$\sin x^2 \text{ w.r.t. } x^3$$

 [Watch Video Solution](#)

15. Differentiate w.r.t. as indicated :

$$\sqrt{1+x^2} \text{ w.r.t. } \tan^{-1} x$$

 [Watch Video Solution](#)

16. Prove that derivative of $\tan^{-1}\left(\frac{x}{1+\sqrt{1-x^2}}\right)$ w.r.t. $\sin^{-1} x$ is independent of x .

 [Watch Video Solution](#)

17. Prove that the derivative of $\tan^{-1}\left(\frac{\sqrt{1+x^2}-1}{x}\right)$ w.r.t. $\tan^{-1} x$ is independent of x .

 [Watch Video Solution](#)

18. Differentiate $\tan^{-1}\left(\frac{\sqrt{1+x^2}-\sqrt{1-x^2}}{\sqrt{1+x^2}+\sqrt{1-x^2}}\right)$ w. r. t. $\sin^{-1}\left(\frac{2x}{1+x^2}\right)$

 [Watch Video Solution](#)

19. Differentiate $\tan^{-1}\left\{\frac{\sqrt{1+x^2}-\sqrt{1-x^2}}{\sqrt{1+x^2}+\sqrt{1-x^2}}\right\}$ with respect to $\cos^{-1}x^2$

 [Watch Video Solution](#)

20. Differentiate $\tan^{-1}\left(\frac{\sqrt{1+x^2}+1}{x}\right)$ w.r.t. $\tan^{-1}\left(\frac{2x\sqrt{1-x^2}}{1-2x^2}\right)$

at $x = 0$.

 [Watch Video Solution](#)

EXERCISE 5(i) (SHORT ANSWER TYPE QUESTIONS)

1. Differentiate the following w.r.t. x :

$$(\sqrt{x})^{\sqrt{x}}$$



Watch Video Solution

2. Differentiate the following w.r.t. x :

$$\left(\frac{1}{2}\right)^x$$



Watch Video Solution

3. Differentiate the following w.r.t. x :

$$(x^x)^x$$



Watch Video Solution

4. Differentiate the following w.r.t. x :

$$x^{x^2}$$

 [Watch Video Solution](#)

5. Differentiate the following w.r.t. x :

$$(5x)^{3 \cos 2x}$$

 [Watch Video Solution](#)

6. Differentiate the following w.r.t. x :

$$x^{\sin x}, x > 0$$

 [Watch Video Solution](#)

7. Differentiate the following w.r.t. x :

$$(\sin x)^x$$

 [Watch Video Solution](#)

8. Differentiate the following w.r.t. x :

$$x^{\sin^{-1} x}$$



Watch Video Solution

9. Differentiate the following w.r.t. x :

$$x^x \sin^{-1} \sqrt{x}$$



Watch Video Solution

10. Differentiate the following w.r.t. x :

$$(\sin x)^{\log x}, \sin x > 0$$



Watch Video Solution

11. Differentiate the following w.r.t. x :

$$(\sin x)^{\tan x}$$





[Watch Video Solution](#)

12. Differentiate the following w.r.t. x :

$$(\sin x)^{\sin x}$$



[Watch Video Solution](#)

13. Differentiate the following w.r.t. x :

$$(\sec^2 x)^{1/x}$$



[Watch Video Solution](#)

14. Differentiate the following w.r.t. x :

$$(x \cos x)^x$$



[Watch Video Solution](#)

15. Differentiate the following w.r.t. x :

$$(x)^{\log x}$$



Watch Video Solution

16. Differentiate the following w.r.t. x :

$$(\log x)^{\log x}, x > 1$$



Watch Video Solution

17. Differentiate the following w.r.t. x :

$$x^{\sin 2x + \cos 2x}$$



Watch Video Solution

18. Differentiate the following w.r.t. x :

$$x^{\sin x + \cos x}$$





[Watch Video Solution](#)

19. Differentiate the following w.r.t. x :

$$(\log x)^x$$



[Watch Video Solution](#)

20. Differentiate the following w.r.t. x :

$$(\sin^{-1} x)^x$$



[Watch Video Solution](#)

21. Differentiate the following w.r.t. x :

$$(\tan^{-1} x)^x$$



[Watch Video Solution](#)

22. Differentiate the following w.r.t. x :

$$x^{\cos^{-1} x}$$

 [Watch Video Solution](#)

23. Differentiate the following w.r.t. x :

$$(\sin x)^{\cos^{-1} x}$$

 [Watch Video Solution](#)

24. Differentiate the following w.r.t. x :

$$(\sin x - \cos x)^{\sin x - \cos x}, \frac{\pi}{4} < x < \frac{3\pi}{4}$$

 [Watch Video Solution](#)

25. Differentiate the following w.r.t. x :

$$\cos(x^x)$$





Watch Video Solution

26. Differentiate the following w.r.t. x :

$$(1 + x)^{\log x}$$



Watch Video Solution

27. Differentiate the following w.r.t. x :

$$(\log x)^{\cos x}$$



Watch Video Solution

EXERCISE 5(i) (LONG ANSWER TYPE QUESTIONS (I))

1. Differentiate the following w.r.t. x :

$$x^{\sin x} + (\sin x)^x$$



Watch Video Solution

2. Differentiate the following w.r.t. x :

$$(x)^{\log x} + (\log x)^x$$



Watch Video Solution

3. Differentiate the following w.r.t. x :

$$(x)^{\tan x} + (\tan x)^x$$



Watch Video Solution

4. Differentiate the following w.r.t. x :

$$x^{\cot x} + (\cos x)^{\sin x}$$



Watch Video Solution

5. Differentiate the following w.r.t. x :

$$x^{\cos x} + (\cos x)^x$$



[Watch Video Solution](#)

6. Differentiate the following w.r.t. x :

$$x^{\cos x} + (\cos x)^{\sin x}$$

[Watch Video Solution](#)

7. Differentiate the following w.r.t. x :

$$(\sin x)^{\cos x} + (\cos x)^{\sin x}$$

[Watch Video Solution](#)

8. $y = (\sin x)^{\tan x} + (\cos x)^{\sec x}$

[Watch Video Solution](#)

9. Differentiate the following w.r.t. x :

$$(\log x)^x + (x)^{\cos x}$$

 [Watch Video Solution](#)

10. Differentiate the following w.r.t. x :

$$x^{\sin x} + (\log x)^x$$

 [Watch Video Solution](#)

11. Differentiate the following w.r.t. x :

$$(x)^{\sin x} + \sin(x^x)$$

 [Watch Video Solution](#)

12. Differentiate the following w.r.t. x :

$$(x)^x + (\sin x)^x$$



[Watch Video Solution](#)

13. Differentiate the following w.r.t. x :

$$(x)^{\sin x} + (\cos x)^x$$

[Watch Video Solution](#)

14. Differentiate the following w.r.t. x :

$$x^{\sin x} + (\sin x)^{\cos x}$$

[Watch Video Solution](#)

15. Differentiate the following w.r.t. x :

$$(\sin x)^{\sec x} + (\tan x)^{\cos x}$$

[Watch Video Solution](#)

16. Differentiate the following w.r.t. x :

$$(\tan x)^{\cot x} + x^{\tan x}, 0 < x < \frac{\pi}{4}$$

 [Watch Video Solution](#)

17. Differentiate the following w.r.t. x :

$$x^{\sin x} + (\sin x)^{\cos x}$$

 [Watch Video Solution](#)

18. Differentiate the following w.r.t. x :

$$(\sin x)^x + \sin^{-1} \sqrt{x}$$

 [Watch Video Solution](#)

19. Differentiate the following w.r.t. x : $(\log x)^x + x^{\log x}$

 [Watch Video Solution](#)

20. Differentiate the following w.r.t. x :

$$(\log x)^{\cos x} + \frac{x^2 + 1}{x^2 - 1}$$

 [Watch Video Solution](#)

21. Differentiate the following w.r.t. x : $(x \cos x)^x + (x \sin x)^{\frac{1}{x}}$

 [Watch Video Solution](#)

22. Differentiate the following w.r.t. x :

$$(\cos x)^x + (\sin x)^{1/x}$$

 [Watch Video Solution](#)

23. $y = e^{\sin x} + (\tan x)^x$

 [Watch Video Solution](#)

24. Differentiate the following w.r.t. x :

$$x^x - 2^{\sin x}$$



Watch Video Solution

25. Differentiate the functions given w.r.t. x : $\left(x + \frac{1}{x}\right)^x + x^{\left(1 + \frac{1}{x}\right)}$



Watch Video Solution

26. Differentiate the following w.r.t. x :

$$x^{x^2-3} + (x-3)^{x^2}, \text{ for } x > 3$$



Watch Video Solution

27. Differentiate the following w.r.t. x :

$$x^x + (\sin x)^x$$





Watch Video Solution

28. Differentiate the following w.r.t. x :

$$\frac{(ax + b)(cx + d)}{(ax - b)(cx - d)}, x \neq \frac{b}{a}, \frac{d}{c}$$



Watch Video Solution

29. Differentiate the following w.r.t. x :

$$\sqrt{\frac{(x - 3)(x^2 + 4)}{3x^2 + 4x + 5}}$$



Watch Video Solution

30. Differentiate the following w.r.t. x :

$$x^2 e^x \sin x$$



Watch Video Solution

31. Differentiate the following w.r.t. x :

$$e^x \cos^3 x \sin^2 x$$

 [Watch Video Solution](#)

32. Differentiate the following w.r.t. x :

$$(x + 3)^2(x + 4)^3(x + 5)^4$$

 [Watch Video Solution](#)

33. Differentiate the following w.r.t. x :

$$\sqrt{(x - 1)(x - 2)(x - 3)(x - 4)}$$

 [Watch Video Solution](#)

34. If $xy = e^{x-y}$, find $\frac{dy}{dx}$.

 [Watch Video Solution](#)

35. If $(\sin x)^y = (\sin y)^x$, find $\frac{dy}{dx}$.

 [Watch Video Solution](#)

36. Find $\frac{dy}{dx}$ if $(\sin x)^{\cos y} = (\cos y)^{\sin x}$.

 [Watch Video Solution](#)

37. Differentiate $\log(x^x + \cos ec^2 x)$ w.r.t. x .

 [Watch Video Solution](#)

38. If $x^p y^q = (x + y)^{p+q}$, show that $\frac{dy}{dx} = \frac{y}{x}$.

 [Watch Video Solution](#)

39. If $y = x^y$, show that $\frac{dy}{dx} = \frac{y^2}{x(1 - y \log x)}$

 [Watch Video Solution](#)

40. If $y^x = e^{y-x}$, then prove that $\frac{dy}{dx} = \frac{(1 + \log y)^2}{\log y}$

 [Watch Video Solution](#)

41. If $x^x + y^x = 1$, prove that $\frac{dy}{dx} = - \left\{ \frac{x^x(1 + \log x) + y^x \log y}{xy^{(x-1)}} \right\}$

 [Watch Video Solution](#)

42. If $x^y + y^x = 1$, find $\frac{dy}{dx}$

 [Watch Video Solution](#)

43. If $x^y + y^x = a^b$, then find $\frac{dy}{dx}$.

 [Watch Video Solution](#)

44. If $x^y + y^x = 4$, find $\frac{dy}{dx}$

 [Watch Video Solution](#)

45. If $x^y + y^x = \log a$, find $\frac{dy}{dx}$.

 [Watch Video Solution](#)

46. Show that if $x^y + y^x = m^n$, then :

$$\frac{dy}{dx} = - \frac{y^x \log y + yx^{y-1}}{x^y \log x + xy^{x-1}}$$

 [Watch Video Solution](#)

47. Find the derivative of the function given by :

$f(x) = (1 + x)(1 + x^2)(1 + x^4)\dots (1 + x^{2^n})$ and hence, find $f'(0)$.

A. 0

B. -1

C. 1

D. 2

Answer: c



[Watch Video Solution](#)

48. Differentiate $(x^2 - 5x + 8)(x^3 + 7x + 9)$ in three ways mentioned below:(i) by using product rule(ii) by expanding the product to obtain a single polynomial.(iii) by logarithmic differentiation.Do they all give the same answer?



[Watch Video Solution](#)

49. If u , v and w are functions of x , then show that

$$\frac{d}{dx}(uvw) = \frac{du}{dx}vw + u\frac{dv}{dx}w + uv\frac{dw}{dx}$$

in two ways - first by repeated application of product rule, second by logarithmic differentiation.

 [Watch Video Solution](#)

EXERCISE 5(j) (LONG ANSWER TYPE QUESTIONS (I))

1. If $y = \sqrt{x + \sqrt{x + \sqrt{x + \dots \rightarrow \infty}}}$, prove that $\frac{dy}{dx} = \frac{1}{2y - 1}$

 [Watch Video Solution](#)

2. If

$$y = \sqrt{3^x + \sqrt{3^x + \sqrt{3^x + \dots \rightarrow \infty}}}$$

prove that $(2y - 1)\frac{dy}{dx} = 3^x \log 3$

 [Watch Video Solution](#)

3. If $y = \sqrt{\sin x + \sqrt{\sin x + \sqrt{\sin x + \dots}}}$, prove that $\frac{dy}{dx} = \frac{\cos x}{2y - 1}$

 Watch Video Solution

4. If $y = (\sin x)^{(\sin x)^{(\sin x) \dots \infty}}$, prove that $\frac{dy}{dx} = \frac{y^2 \cot x}{(1 - y \log \sin x)}$.

 Watch Video Solution

5. If

$y = (\cos x)^{\cos x \wedge (\cos x) \wedge (((\infty)))}$, prove that $\frac{dy}{dx} = -\frac{y^2 \tan x}{(1 - y \log \cos x)}$

 Watch Video Solution

6. If $y = (\tan x)^{(\tan x)^{(\tan x) \dots \infty}}$, then prove that $\frac{dy}{dx} = 2$ at $x = \frac{\pi}{4}$.

 Watch Video Solution

7. If $y = x^{x^{x^{\dots \infty}}}$ then prove that $x \frac{dy}{dx} = \frac{y^2}{1 - y \log x}$

 Watch Video Solution

8. If $y = a^{x^{a^{x^{\dots \infty}}}}$, prove that $\frac{dy}{dx} = \frac{y^2(\log y)}{x[1 - y(\log x)(\log y)]}$.

 Watch Video Solution

9. If $y = x + \frac{1}{x + \frac{1}{x + \frac{1}{x + \dots \infty}}}$, prove that $\frac{dy}{dx} = \frac{y}{(2y - x)}$.

 Watch Video Solution

10. If $y = \frac{\sin x}{1 + \frac{\cos x}{1 + \frac{\sin x}{1 + \frac{\cos x}{1 + \dots \infty}}}}$, prove that $\frac{dy}{dx} = \frac{(1 + y)\cos x + y \sin x}{1 + 2y + \cos x - \sin x}$

 Watch Video Solution

1. Find (a) $\frac{dy}{dx}$ and (b) $\frac{d^2y}{dx^2}$ when y is given by :

$$1 + 2x.$$

 [Watch Video Solution](#)

2. Find (a) $\frac{dy}{dx}$ and (b) $\frac{d^2y}{dx^2}$ when y is given by :

$$ax^3 + bx^2 + cx + d$$

 [Watch Video Solution](#)

3. Find (a) $\frac{dy}{dx}$ and (b) $\frac{d^2y}{dx^2}$ when y is given by :

$$\frac{1}{2x + 3}, x \neq -\frac{3}{2}$$

 [Watch Video Solution](#)

4. Find (a) $\frac{dy}{dx}$ and (b) $\frac{d^2y}{dx^2}$ when y is given by :

$$\log x - x$$

 [Watch Video Solution](#)

5. Find (a) $\frac{dy}{dx}$ and (b) $\frac{d^2y}{dx^2}$ when y is given by :

$$e^x + x^4.$$

 [Watch Video Solution](#)

6. Find the second derivatives of the following functions :

$$x^{20}$$

 [Watch Video Solution](#)

7. Find the second derivatives of the following functions :

$$x^2 + 3x + 2$$



[Watch Video Solution](#)

8. Find the second derivatives of the following functions :

$$x \cos x$$



[Watch Video Solution](#)

9. Find the second derivatives of the following functions :

$$x^3 + \tan x$$



[Watch Video Solution](#)

10. Find the second derivatives of the following functions :

$$\log x$$



[Watch Video Solution](#)

11. Find the second derivatives of the following functions :

$$x^3 \log x$$



Watch Video Solution

12. Find the second derivatives of the following functions :

$$\log(\log x)$$



Watch Video Solution

13. Find the second derivatives of the following functions :

$$\sin(\log x)$$



Watch Video Solution

14. Find the second derivatives of the following functions :

$$e^x \sin 5x$$





Watch Video Solution

15. Find the second derivatives of the following functions :

$$e^{6x} \cos 3x$$



Watch Video Solution

16. Find the second derivatives of the following functions :

$$e^{-x} \cos x$$



Watch Video Solution

17. Find the second derivatives of the following functions :

$$\tan x + \sec x$$



Watch Video Solution

18. Find the second derivatives of the following functions :

$$\frac{\log x}{x}$$

 [Watch Video Solution](#)

19. Find the second derivatives of the following functions :

$$x^x$$

 [Watch Video Solution](#)

EXERCISE 5(k) (LONG ANSWER TYPE QUESTIONS (I))

1. If $y = \cos^{-1} x$, then show that :

$$(1 - x^2) \frac{d^2y}{dx^2} - x \frac{dy}{dx} = 0$$

 [Watch Video Solution](#)

2. If $y = a \sin^{-1} x$, then show that :

$$(1 - x^2) \frac{d^2 y}{dx^2} - x \frac{dy}{dx} = 0$$

 [Watch Video Solution](#)

3. If $y = ae^{2x} + be^{-x}$, show that, $\frac{d^2 y}{dx^2} - \frac{dy}{dx} - 2y = 0$.

 [Watch Video Solution](#)

4. If $y = 3e^{2x} + 2e^{3x}$, prove that $\frac{d^2 y}{dx^2} - 5\frac{dy}{dx} + 6y = 0$.

 [Watch Video Solution](#)

5. If $y = Ae^{mx} + Be^{nx}$, show that $\frac{d^2 y}{dx^2} - \frac{(m+n)dy}{dx} + mny = 0$.

 [Watch Video Solution](#)

6. If $y = Pe^{ax} + Qe^{bx}$, show that :

$$\frac{d^2y}{dx^2} - (a + b)\frac{dy}{dx} + aby = 0.$$



Watch Video Solution

7. If $y = 2 \sin x + 3 \cos x$, prove that : $y + \frac{d^2y}{dx^2} = 0$



Watch Video Solution

8. If $y = 5 \sin x - 3 \sin x$, prove that : $\frac{d^2y}{dx^2} + y = 0$.



Watch Video Solution

9. If $e^y(x + 1) = 1$, show that $\frac{d^2y}{dx^2} = \left(\frac{dy}{dx}\right)^2$.



Watch Video Solution

10. If $y = A \sin x + B \cos x$, prove that $\frac{d^2y}{dx^2} + y = 0$.



Watch Video Solution

11. Find $\frac{d^2y}{dx^2}$ in the following

$$x = at^2, y = 2at$$



Watch Video Solution

12. Find $\frac{d^2y}{dx^2}$ in the following

$$x = \frac{2at^2}{1+t}, y = \frac{3at}{1+t}$$



Watch Video Solution

13. Find $\frac{d^2y}{dx^2}$ in the following

$$x = a \cos \theta, y = b \sin \theta$$



Watch Video Solution

14. Find $\frac{d^2y}{dx^2}$ in the following

$$x = a \cos^3 \theta, y = a \sin^3 \theta$$



Watch Video Solution

15. Find $\frac{d^2y}{dx^2}$ in the following

$$x = a \cos^3 \theta, y = b \sin^3 \theta$$



Watch Video Solution

16. Find $\frac{d^2y}{dx^2}$ in the following

$$\text{If } x^{\frac{2}{3}} + y^{\frac{2}{3}} = a^{\frac{2}{3}}, \text{ find } \frac{d^2y}{dx^2}$$



Watch Video Solution

17. Find $\frac{d^2y}{dx^2}$ in the following

If $x = a \cos^3 \theta$ and $y = a \sin^3 \theta$, then find the value of $\frac{d^2y}{dx^2}$ at $\theta = \frac{\pi}{6}$.



Watch Video Solution

18. Find $\frac{d^2y}{dx^2}$ in the following

$x = a(\cos t + t \sin t)$, $y = a(\sin t - t \cos t)$



Watch Video Solution

19. Find $\frac{d^2y}{dx^2}$ in the following

$x = a(\theta + \sin \theta)$, $y = a(1 + \cos \theta)$



Watch Video Solution

20. Find $\frac{d^2y}{dx^2}$ at $\theta = \frac{\pi}{2}$ when :

$x = a(\theta + \sin \theta)$, $y = a(1 - \cos \theta)$



Watch Video Solution

21. Find $\frac{d^2y}{dx^2}$ at $\theta = \frac{\pi}{2}$ when :

$$x = a(\theta - \sin \theta), y = a(1 + \cos \theta)$$



Watch Video Solution

22. Find $\frac{d^2y}{dx^2}$ at $\theta = \frac{\pi}{2}$ when :

$$x = a(1 - \cos \theta), y = a(\theta + \sin \theta)$$



Watch Video Solution

23. Find $\frac{d^2y}{dx^2}$ at $\theta = \frac{\pi}{2}$ when :

$$x = a(\theta - \sin \theta), y = a(1 - \cos \theta)$$



Watch Video Solution

24. Find $\frac{d^2y}{dx^2}$ at $\theta = \frac{\pi}{4}$ when :

$$x = a(\cos \theta + \log \tan \theta / 2), y = a \sin \theta$$

 [Watch Video Solution](#)

25. If $x = \cos t + \frac{\log \tan t}{2}$, $y = \sin t$, then find the value of $\frac{d^2y}{dt^2}$ and $\frac{d^2y}{dx^2}$ at $t = \frac{\pi}{4}$.

 [Watch Video Solution](#)

26. Find $\frac{d^2y}{dx^2}$ when :

$$x = 2 \cos \theta - \cos 2\theta \text{ and } y = 2 \sin \theta - \sin 2\theta$$

 [Watch Video Solution](#)

27. If $x = a(\cos 2\theta + 2\theta \sin 2\theta)$ and $y = a(\sin 2\theta - 2\theta \cos 2\theta)$, find $\frac{d^2y}{dx^2}$ at $\theta = \frac{\pi}{8}$.



Watch Video Solution

28. If $x = a \sin t$ and $y = a \left(\cos t + \frac{\log \tan t}{2} \right)$, find $\frac{d^2 y}{dx^2}$



Watch Video Solution

29. If $x + y = \tan^{-1} y$ and $\frac{d^2 y}{dx^2} = f(y) \frac{dy}{dx}$, then $f(y) =$

A. $-\frac{2}{y^3}$

B. $\frac{2}{y^3}$

C. $\frac{1}{y}$

D. $-\frac{1}{y}$

Answer: B



Watch Video Solution

30. If $y = (\sin^{-1} x)^2$ then prove that $(1 - x^2) \frac{d^2y}{dx^2} - x \frac{dy}{dx} - 2 = 0$.

 [Watch Video Solution](#)

31. If $y = (\cos^{-1} x)^2$, then prove that :

$$(1 - x^2)y_2 - xy_1 - 2 = 0.$$

 [Watch Video Solution](#)

32. If $y = (\tan^{-1} x)^2$, show that $(x^2 + 1)^2 y_2 + 2x(x^2 + 1)y_1 = 2$

 [Watch Video Solution](#)

33. If $y = (\cot^{-1} x)^2$, then show that

$$(x^2 + 1)^2 \frac{d^2y}{dx^2} + 2x(x^2 + 1) \frac{dy}{dx} = 2.$$

 [Watch Video Solution](#)

34. If $y = \frac{s \in^{-1} x}{\sqrt{1-x^2}}$, show that $(1-x^2) \frac{d^2 y}{dx^2} - 3x \frac{dy}{dx} - y = 0$

 [Watch Video Solution](#)

35. If $y = \log [x + \sqrt{x^2 + 1}]$, prove that $(x^2 + 1) \frac{d^2 y}{dx^2} + x \frac{dy}{dx} = 0$

 [Watch Video Solution](#)

36. If $y = \log \{x + \sqrt{x^2 + a^2}\}$, prove that: $(x^2 + a^2) \frac{d^2 y}{dx^2} + x \frac{dy}{dx} = 0$.

 [Watch Video Solution](#)

37. If $y = \{x + \sqrt{x^2 + 1}\}^m$, then show that $(x^2 + 1) \frac{d^2 y}{dx^2} + x \frac{dy}{dx} - m^2 y = 0$.

 [Watch Video Solution](#)

38. If $y = \cos(m \cos^{-1} x)$, then prove that :

$$(1 - x^2) \frac{d^2 y}{dx^2} - x \frac{dy}{dx} + m^2 y = 0.$$

 [Watch Video Solution](#)

39. If $y = \sin(m \tan^{-1} x)$, prove that :

$$(1 + x^2)^2 y_2 + 2x(1 + x^2) y_1 + m^2 y = 0.$$

 [Watch Video Solution](#)

40. If $y = e^{m \sin^{-1} x}$, $-1 \leq x \leq 1$, show that :

$$(1 - x^2) y_2 - x y_1 - m^2 y = 0.$$

 [Watch Video Solution](#)

41. If $y = e^{a \cos^{-1} x}$, $-1 \leq x \leq 1$, show that : $\frac{dy}{dx} = \frac{-ae^{a \cos^{-1} x}}{\sqrt{1 - x^2}}$

 [Watch Video Solution](#)

42. If $y = e^a \cos^{(-1)x}$, $-1 \leq x \leq 1$, show that

$$(1 - x^2) \frac{d^2y}{dx^2} - x \frac{dy}{dx} - a^2y = 0.$$

 [Watch Video Solution](#)

43. If $y = e^{m \tan^{-1} x}$, prove that :

$$(1 + x^2) \frac{d^2y}{dx^2} + (2x - m) \frac{dy}{dx} = 0.$$

 [Watch Video Solution](#)

44. If $y = e^{ax} \cos bx$, then prove that :

$$\frac{d^2y}{dx^2} - 2a \frac{dy}{dx} + (a^2 + b^2)y = 0.$$

 [Watch Video Solution](#)

45. If $y = \sin^{-1} x$, show that $(1 - x^2) \frac{d^2y}{dx^2} - x \frac{dy}{dx} = 0$.

 [Watch Video Solution](#)

46. If $y = \tan x$, then show that : $\frac{d^2y}{dx^2} = 2y \frac{dy}{dx}$

 [Watch Video Solution](#)

47. If $y = \tan^{-1} x$, show that $(1 + x^2) \frac{d^2y}{dx^2} + 2x \frac{dy}{dx} = 0$.

 [Watch Video Solution](#)

48. If $x = f(t)$, $y = g(t)$ possess second order derivatives for all t in (a, b) and $f'(t)$ is invertible [$f'(t) \neq 0$], then prove that :

$$\frac{d^2y}{dx^2} = \frac{f'(t)g''(t) - g'(t)f''(t)}{[f'(t)]^3}.$$

 [Watch Video Solution](#)

49. If $y^m x^n = (x + y)^{m+n}$, then find $\frac{dy}{dx}$

 [Watch Video Solution](#)

 Watch Video Solution

50. If $x^m y^n = (x + y)^{m+n}$, prove that :

$$\frac{d^2 y}{dx^2} = 0.$$

 Watch Video Solution

51. If $p^2 = a^2 \cos^2 \theta + b^2 \sin^2 \theta$, then show that :

$$p + \frac{d^2 p}{d\theta^2} = \frac{a^2 b^2}{p^3}.$$

 Watch Video Solution

EXERCISE 5(I) (SHORT ANSWER TYPE QUESTIONS)

1. Verify the truth of Rolle's Theorem for the following functions

$$f(x) = x^2 + 2, a = -2, \text{ and } b = 2$$

 Watch Video Solution

2. Verify the truth of Rolle's Theorem for the following functions

$$f(x) = \frac{x^3}{3} - \frac{5}{3}x^2 + 2x, x \in [0, 3]$$

 [Watch Video Solution](#)

3. Verify the truth of Rolle's Theorem for the following functions

$$f(x) = \frac{x(x-2)}{x-1} \text{ on } [0, 2]$$

 [Watch Video Solution](#)

4. Verify the truth of Rolle's Theorem for the following functions

$$f(x) = x^2 + 2x - 8 \text{ defined in the interval } [-4, 2].$$

 [Watch Video Solution](#)

5. Verify the truth of Rolle's Theorem for the following functions

$$f(x) = x^{3/5} \text{ in } [-1, 1]$$



[Watch Video Solution](#)

6. Verify the truth of Rolle's Theorem for the following functions

$$f(x) = x^{1/3} \text{ in } [-1, 1]$$



[Watch Video Solution](#)

7. Verify the truth of Rolle's Theorem for the following functions

$$f(x) = |x| \text{ in } [-1, 1]$$



[Watch Video Solution](#)

8. Verify the truth of Rolle's Theorem for the following functions

$$f(x) = |x - 1| \text{ in } [1, 2]$$



[Watch Video Solution](#)

9. Verify the truth of Rolle's Theorem for the following functions

$$f(x) = \sqrt{x - 2} \text{ in } [1, 2]$$

 [Watch Video Solution](#)

10. Verify the truth of Rolle's Theorem for the following functions

$$f(x) = [x] \text{ in } [-1, 1]$$

 [Watch Video Solution](#)

11. Verify the conditions of Rolle's Theorem in the following problems. In

each case, find a point in the interval, where the derivative vanishes :

$$x^2 - 1 \text{ on } [-1, 1]$$

 [Watch Video Solution](#)

12. Verify the conditions of Rolle's Theorem in the following problems. In each case, find a point in the interval, where the derivative vanishes :

$$(x^2 - 1)(x - 2) \text{ on } [-1, 2]$$



[Watch Video Solution](#)

13. Verify the conditions of Rolle's Theorem in the following problems. In each case, find a point in the interval, where the derivative vanishes :

$$\sin x - \sin 2x \text{ on } [0, \pi]$$



[Watch Video Solution](#)

14. Verify the conditions of Rolle's Theorem in the following problems. In each case, find a point in the interval, where the derivative vanishes :

$$\log(x^2 + 2) - \log 3 \text{ on } [-1, 1]$$



[Watch Video Solution](#)

15. Verify the conditions of Rolle's Theorem in the following problems. In

each case, find a point in the interval, where the derivative vanishes :

$$e^{1-x^2} \text{ on } [-1, 1]$$



[Watch Video Solution](#)

EXERCISE 5(I) (LONG ANSWER TYPE QUESTIONS (I))

1. Verify the truth of Rolle's Theorem for the following functions :

$$f(x) = x^2 - 5x + 4 \text{ on } [1, 4]$$



[Watch Video Solution](#)

2. Verify the truth of Rolle's Theorem for the following functions :

$$f(x) = x^2 - 4x + 3 \text{ on } [1, 3]$$



[Watch Video Solution](#)

3. Verify the truth of Rolle's Theorem for the following functions :

$$f(x) = x^3 - 2x^2 - 3x \text{ in the interval } [-1, 3].$$

 [Watch Video Solution](#)

4. Verify the truth of Rolle's Theorem for the following functions :

$$f(x) = x^2 + 2x - 8 \text{ in the interval } [-4, 2]$$

 [Watch Video Solution](#)

5. Verify the truth of Rolle's Theorem for the following functions :

$$f(x) = x^2 - x - 12 \text{ in the interval } [-3, 4]$$

 [Watch Video Solution](#)

6. Verify the truth of Rolle's Theorem for the following functions :

$$f(x) = (x - 2)(x - 3)(x - 4) \text{ in the interval } 2 \leq x \leq 4.$$





[Watch Video Solution](#)

7. Verify the truth of Rolle's Theorem for the following functions :

$$f(x) = x^3 - 4x \text{ in the interval } -2 \leq x \leq 2$$



[Watch Video Solution](#)

8. Verify the truth of Rolle's Theorem for the following functions :

$$f(x) = (x - 1)(x - 2)^2 \text{ in the interval } [1, 2]$$



[Watch Video Solution](#)

9. Verify the truth of Rolle's Theorem for the following functions :

$$f(x) = (x - 2)(x - 4)^2 \text{ in the interval } [2, 4].$$



[Watch Video Solution](#)

10. Verify Rolle's Theorem in the interval $[a, b]$ for the function :

$$f(x) = (x - a)^2(x - b)^2. \text{ Find the value of 'c'.$$

 [Watch Video Solution](#)

11. Examine the applicability of Rolle's Theorem for the function :

$$f(x) = 2 + (x - 1)^{2/3} \text{ in the interval } 0 \leq x \leq 2.$$

 [Watch Video Solution](#)

12. Verify Rolle's Theorem for the functions :

$$f(x) = \sin^2 x, \text{ defined in the interval } [0, \pi]$$

 [Watch Video Solution](#)

13. Verify Rolle's Theorem for the functions :

$$f(x) = \cos x, \text{ defined in the interval } \left[-\frac{\pi}{2}, \frac{\pi}{2} \right]$$





[Watch Video Solution](#)

14. Verify Rolle's Theorem for the functions :

$$f(x) = \tan x, \text{ defined in the interval } [0, \pi].$$



[Watch Video Solution](#)

15. Verify Rolle's Theorem for the functions :

$$f(x) = \sin x + \cos x \text{ in the interval } [0, 2\pi]$$



[Watch Video Solution](#)

16. Verify Rolle's Theorem for the functions :

$$f(x) = \sin x + \cos x + 5 \text{ in the interval } [0, 2\pi]$$



[Watch Video Solution](#)

17. Verify Rolle's Theorem for the functions :

$$f(x) = \sin x \cos x \text{ in } \left[0, \frac{\pi}{2}\right]$$

 [Watch Video Solution](#)

18. Verify Rolle's Theorem for the functions :

$$f(x) = \sin^3 x + \cos^3 x \text{ in the interval } \left[0, \frac{\pi}{2}\right].$$

 [Watch Video Solution](#)

19. Verify Rolle's Theorem for the function :

$$f(x) = \begin{cases} -4x + 5, & 0 \leq x \leq 1 \\ 2x - 3, & 1 < x \leq 2 \end{cases}$$

 [Watch Video Solution](#)

20. At what points on the following curve, is the tangent parallel to x-axis

?

$$y = x^2 \text{ on } [-2, 2]$$



Watch Video Solution

21. At what points on the following curve, is the tangent parallel to x-axis ?

$$y = \cos x - 1 \text{ on } [0, 2\pi]$$



Watch Video Solution

22. For the function $f(x) = x^3 - 6x^2 + ax + b$, it is given that $f(1) = f(3) = 0$. Find the values of 'a' and 'b', and hence, verify Rolle's Theorem on $[1, 3]$.



Watch Video Solution

23. Let $f(x) = (x - 1)(x - 2)(x - 3)$ on the interval $[1, 3]$. Prove that there is more than one c in $(1, 3)$ such that $f'(c) = 0$.

 [Watch Video Solution](#)

EXERCISE 5(m) (SHORT ANSWER TYPE QUESTIONS)

1. Discuss the applicability of Lagrange's Mean Value Theorem to the following :

$$f(x) = x^2 - 1 \text{ on } [1, 2].$$

 [Watch Video Solution](#)

2. Discuss the applicability of Lagrange's Mean Value Theorem to the following :

$$f(x) = x^2 \text{ on } [2, 4]$$

 [Watch Video Solution](#)

3. Discuss the applicability of Lagrange's Mean Value Theorem to the following :

$$f(x) = x^2 - 2x + 3 \text{ in } [0, 4]$$

 [Watch Video Solution](#)

4. Verify Lagrange's mean value theorem for the following function:

$$f(x) = x^2 + 2x + 3, \text{ for } [4, 6].$$

 [Watch Video Solution](#)

5. Discuss the applicability of Lagrange's Mean Value Theorem to the following :

$$f(x) = 2x^2 - 10x + 29 \text{ in } [2, 7]$$

 [Watch Video Solution](#)

6. Discuss the applicability of Lagrange's Mean Value Theorem to the following :

$$f(x) = x^2 - 4x - 3 \text{ in } [1, 4]$$



[Watch Video Solution](#)

7. Discuss the applicability of Lagrange's Mean Value Theorem to the following :

$$f(x) = 2x - x^2 \text{ in } [0, 1]$$



[Watch Video Solution](#)

8. Discuss the applicability of Lagrange's Mean Value Theorem to the following :

$$f(x) = x(2 - x) \text{ in } [0, 1]$$



[Watch Video Solution](#)

9. $f(x) = x^3 - 2x^2 - x + 3$ in $[0, 1]$



[Watch Video Solution](#)

10. Discuss the applicability of Lagrange's Mean Value Theorem to the following :

$$f(x) = x^3 - 5x^2 - 3x \text{ in } [1, 3]$$

 [Watch Video Solution](#)

11. Discuss the applicability of Lagrange's Mean Value Theorem to the following :

$$f(x) = (x - 1)(x - 2)(x - 3) \text{ in } [0, 4]$$

 [Watch Video Solution](#)

EXERCISE 5(m) (LONG ANSWER TYPE QUESTIONS (I))

1. Verify the conditions of Mean Value Theorem in the following. In each case, find a point in the interval as stated by the Mean Value Theorem :

$$f(x) = x \text{ on } [a, b]$$

 [Watch Video Solution](#)

2. Verify the conditions of Mean Value Theorem in the following. In each case, find a point in the interval as stated by the Mean Value Theorem :

$$f(x) = x + \frac{1}{x} \text{ on } [1, 3]$$

 [Watch Video Solution](#)

3. Verify the conditions of Mean Value Theorem in the following. In each case, find a point in the interval as stated by the Mean Value Theorem :

$$f(x) = ax^2 + bx + c \text{ on } [0, 1]$$

 [Watch Video Solution](#)

4. Verify the conditions of Mean Value Theorem in the following. In each case, find a point in the interval as stated by the Mean Value Theorem :

$$f(x) = ax^2 + ex + e \text{ on } [0, 1]$$

 [Watch Video Solution](#)

5. Verify the conditions of Mean Value Theorem in the following. In each case, find a point in the interval as stated by the Mean Value Theorem :

$$f(x) = \sin x - \sin 2x \text{ on } [0, 2\pi]$$



[Watch Video Solution](#)

6. Verify the conditions of Mean Value Theorem in the following. In each case, find a point in the interval as stated by the Mean Value Theorem :

$$f(x) = \sin x - \sin 2x \text{ on } [0, \pi]$$



[Watch Video Solution](#)

7. Verify Lagrange's Mean Value Theorem for the functions :

$$f(x) = x^{1/3} \text{ in the interval } [-1, 1]$$



[Watch Video Solution](#)

8. Verify Lagrange's Mean Value Theorem for the functions :

$$f(x) = (x - 1)^{2/3} \text{ in the interval } [1, 2].$$

 [Watch Video Solution](#)

9. Verify Lagrange's Mean Value Theorem for the functions :

$$f(x) = \frac{1}{x} \text{ in the interval } [-1, 2]$$

 [Watch Video Solution](#)

10. $f(x) = \frac{1}{4x - 1}$ in $[1, 4]$

 [Watch Video Solution](#)

11. Verify Lagrange's Mean Value Theorem for the functions :

$$f(x) = |x| \text{ in the interval } [-1, 1].$$

 [Watch Video Solution](#)

12. Verify Lagrange's Mean Value Theorem for the functions :

$$f(x) = \sqrt{x^2 - 4} \text{ in the interval } [2, 4].$$

 [Watch Video Solution](#)

13. Verify Lagrange's Mean Value Theorem for the functions :

$$f(x) = \sqrt{25 - x^2} \text{ in the interval } [-3, 4].$$

 [Watch Video Solution](#)

14. Verify Lagrange's Mean Value Theorem for the functions :

$$f(x) = \log_e x \text{ in the interval } [1, 2]$$

 [Watch Video Solution](#)

15. Verify Lagrange's Mean Value Theorem for the functions :

$$f(x) = x \text{ on } [a, b]$$

 [Watch Video Solution](#)

16. Find 'c' of Lagrange's Mean Value Theorem for the functions :

$$f(x) = 2x^2 - 1 \text{ in the interval } [1, 2]$$

 [Watch Video Solution](#)

17. Find 'c' of Lagrange's Mean Value Theorem for the functions :

$$f(x) = \log x \text{ in the interval } [1, e]$$

 [Watch Video Solution](#)

18. Find 'c' of Lagrange's Mean Value Theorem for the functions :

$$f(x) = e^x \text{ in the interval } [0, 1]$$





Watch Video Solution

19. Verify Mean Value Theorem, if $f(x) = x^3 - 5x^2 - 3x$ in the interval $[a, b]$, where $a = 1$ and $b = 3$. Find all $c \in (1, 3)$ for which $f'(c) = 0$.



Watch Video Solution

20. If mean value theorem holds for the function $f(x) = (x - 1)(x - 2)(x - 3)$, $x \in [0, 4]$, then $c =$



Watch Video Solution

21. Verify Lagrange's Mean Value Theorem for the function :

$$f(x) = \begin{cases} 2 + x^3 & \text{if } x \leq 1 \\ 3x & \text{if } x > 1 \end{cases} \text{ on } [-1, 2].$$



Watch Video Solution

22. Find a point on the parabola $y = (x - 2)^2$, where the tangent is parallel to the chord joining $(2, 0)$ and $(4, 4)$.

 [Watch Video Solution](#)

23. Find a point on the graph of $y = x^3$, where the tangent is parallel to the chord joining $(1, 1)$ and $(3, 27)$.

 [Watch Video Solution](#)

24. Find a point on the curve $y = x^3 - 3x$ where the tangent is parallel to the chord joining $(1, -2)$ and $(2, 2)$.

 [Watch Video Solution](#)

25. Find the co-ordinates of the point at which the tangent to the curve given by $f(x) = x^2 - 6x + 1$ is parallel to the chord joining the points

(1, -4) and (3, -8).



Watch Video Solution

26. Use Lagrange's Mean value Theorem to determine a point P on the curve $y = \sqrt{x - 2}$, where the tangent is parallel to the chord joining (2, 0) and (3, 1).



Watch Video Solution

OBJECTIVE TYPE QUESTIONS (MULTIPLE CHOICE QUESTIONS)

1. The function $f(x) = \begin{cases} \frac{\sin x}{x} + \cos x, & \text{if } x \neq 0 \\ k, & \text{if } x = 0 \end{cases}$ is continuous at $x = 0$,

then the value of 'k' is :

A. 3

B. 2

C. 1

D. 1.5

Answer: B



[Watch Video Solution](#)

2. The function $f(x) = [x]$, where $[x]$ denotes the greatest integer function, is continuous at

A. 4

B. -2

C. 1

D. 1.5

Answer: D



[Watch Video Solution](#)

3. If the function $f(x)$ defined by

$$f(x) = \begin{cases} x \sin \frac{1}{x}, & \text{for } x \neq 0 \\ k, & \text{for } x = 0 \end{cases}$$

is continuous at $x = 0$, then k is equal to

A. 8

B. 8

C. -1

D. None of these

Answer: D



Watch Video Solution

4. Differential coefficient of $\sec(\tan^{-1} x)$ w.r.t x is

A. $\frac{x}{\sqrt{1+x^2}}$

B. $\frac{x}{1+x^2}$

C. $x\sqrt{1+x^2}$

D. $\frac{1}{\sqrt{1+x^2}}$

Answer: A



Watch Video Solution

5. If $y = \log\left(\frac{1-x^2}{1+x^2}\right)$, then $\frac{dy}{dx}$ is equal to

A. $\frac{4x^3}{1-x^4}$

B. $\frac{-4x}{1-x^4}$

C. $\frac{1}{4-x^4}$

D. $\frac{-4x^3}{1-x^4}$

Answer: B



Watch Video Solution

6. If $y = \sqrt{\sin x + y}$, then $\frac{dy}{dx}$ is equal to

A. $\frac{\cos x}{2y - 1}$

B. $\frac{\cos x}{1 - 2y}$

C. $\frac{\sin x}{1 - 2y}$

D. $\frac{\sin x}{2y - 1}$

Answer: A



Watch Video Solution

7. If $u = \sin^{-1}\left(\frac{2x}{1+x^2}\right)$ and $v = \tan^{-1}\left(\frac{2x}{1-x^2}\right)$, where -1

A. $\frac{1}{2}$

B. x

C. $\frac{1-x^2}{1+x^2}$

D. 1

Answer: D



Watch Video Solution

8. If $x = t^2$ and $y = t^3$, then $\frac{d^2y}{dx^2} = \frac{3}{2}$ (b) $\frac{3}{(4t)}$ (c) $\frac{3}{2(t)}$ (d) $\frac{3t}{2}$

A. $\frac{3}{2}$

B. $\frac{3}{4t}$

C. $\frac{3}{2t}$

D. $\frac{3t}{2}$

Answer: B



Watch Video Solution

9. The value of c in Rolle's theorem for the function $f(x) = x^3 - 3x$ in the interval $[0, \sqrt{3}]$ is

A. 1

B. -1

C. $\frac{3}{2}$

D. $\frac{1}{3}$

Answer: A



Watch Video Solution

10. The value of c in Lagrange's mean value theorem for the function

$f(x) = x(x - 2)$ when $x \in [1, 2]$ is

A. $\frac{3}{2}$

B. $\frac{2}{3}$

C. $\frac{1}{2}$

D. $\frac{3}{4}$

Answer: A

 [Watch Video Solution](#)

11. If $f(x) = \begin{cases} \frac{x^2-9}{x-3}, & \text{if } x \neq 3 \\ m, & \text{if } x = 3 \end{cases}$ is continuous at $x = 3$, then the value of

'm' is :

A. 3

B. 6

C. 2

D. 1

Answer: B

 [Watch Video Solution](#)

12. If $y = \log(\tan x)$, then $\frac{dy}{dx}$ is :

A. $\frac{1}{\tan x}$

B. $\frac{\sec^2 x}{\tan x}$

C. $\sec^2 x$

D. 0

Answer: B



Watch Video Solution

13. Derivative of $\cos \sqrt{x}$ is :

A. $-\frac{\sin \sqrt{x}}{2\sqrt{x}}$

B. $\frac{\sin \sqrt{x}}{2\sqrt{x}}$

C. $-\frac{\sin \sqrt{x}}{\sqrt{x}}$

D. $-\frac{\sin \sqrt{x}}{2}$

Answer: A



Watch Video Solution

14. If $x - y = \pi$, then $\frac{dy}{dx}$ is :

A. 1

B. 0

C. π

D. -1

Answer: A



Watch Video Solution

15. $\frac{d}{dx}(\tan^{-1} x)$ is :

A. $\frac{1}{1+x^2}$

B. $\frac{1}{1-x^2}$

C. $\frac{-1}{1+x^2}$

D. $\frac{-1}{1-x^2}$

Answer: A



Watch Video Solution

16. If $y = a^x x^a$, then $\frac{dy}{dx}$ is equal to :

A. $a^x x^{a-1}(a - x \log a)$

B. $a^x x^{a-1}(a + x \log a)$

C. $a^x x^a(a + x \log a)$

D. $a^x x^{a-1}(x + a \log a)$

Answer: B



Watch Video Solution

17. $f(x) = \begin{cases} \frac{\sin x}{x}, & x \neq 0 \\ k - 1, & x = 0 \end{cases}$ is continuous at $x = 0$, then 'k' is :

A. 2

B. 0

C. -1

D. 1

Answer: A



[Watch Video Solution](#)

18. The derivative of a^x is :

A. a^x

B. $\frac{a^x}{\log a}$

C. $a^x \log a$

D. None of these

Answer: C



[Watch Video Solution](#)

19. Derivative of $\sin x^3$ w.r.t. x is :

A. $\cos x^3$

B. $3x^2 \cos x^3$

C. $3x^2 \cos x$

D. None of these

Answer: B



[Watch Video Solution](#)

20. At $x = 2$, $f(x) = [x]$ is

A. continuous but not differentiable

B. differentiable but not continuous

C. continuous as well as differentiable

D. None of these

Answer: B



Watch Video Solution

21. If $y = e^{\sin \sqrt{x}}$, then $\frac{dy}{dx}$ is :

A. $e^{\sin \sqrt{x}} \cos \sqrt{x}$

B. $\frac{e^{\sin \sqrt{x}} \cos \sqrt{x}}{2\sqrt{x}}$

C. $\frac{e^{\sin \sqrt{x}}}{2\sqrt{x}}$

D. None of these

Answer: B



Watch Video Solution

22. $\frac{d(a^x)}{dx} =$

A. a^x

B. $\log(a^x)$

C. $a^x \log a$

D. xa^{x-1}

Answer: C



Watch Video Solution

23. Which one of the following is true ?

For the real function :

$$f(x) = \begin{cases} x + 2 & \text{if } x \leq 1 \\ x - 2 & \text{if } x > 1 \end{cases}$$

A. f is continuous at all real numbers $x > 1$ and $x < 1$

B. f is continuous at all real numbers $x \geq 1$

C. f is continuous at all real numbers $x \leq 1$

D. f is continuous at $x = 1$.

Answer: A



Watch Video Solution

24. If $f(x) = \begin{cases} mx + 1 & x \leq 5 \\ 3x - 5 & x > 5 \end{cases}$ is continuous, then the value of m is :

A. $\frac{9}{5}$

B. $\frac{5}{9}$

C. $\frac{5}{3}$

D. $\frac{3}{5}$

Answer: A



Watch Video Solution

25. If $y = \sin x$, then at $x = \frac{\pi}{2}$, y_2 is equal to :

A. -1

B. 1

C. 0

D. $\frac{1}{2}$

Answer: A



Watch Video Solution

26. The derivative of $\tan 45^\circ$ is :

A. 1

B. $\cot 45^\circ$

C. $-\cot 45^\circ$

D. 0

Answer: D



Watch Video Solution

27. If the function $f(x) = \begin{cases} kx + 1, & \text{if } x \leq \pi \\ \cos x, & \text{if } x > \pi \end{cases}$ is continuous at $x = \pi$, then the value of 'k' is :

A. -1

B. $-\frac{2}{\pi}$

C. -2

D. None of these

Answer: B



[Watch Video Solution](#)

28. If $y = \tan^{-1} \frac{x}{2} - \cot^{-1} \frac{x}{2}$, then $\frac{dy}{dx}$ is :

A. $\frac{4}{4 + x^2}$

B. $\frac{2}{4 + x^2}$

C. $\frac{1}{4 + x^2}$

D. $\frac{2}{1+x^2}$

Answer: A

 [Watch Video Solution](#)

29. let $y = \sqrt{x + \sqrt{x + \sqrt{x \dots \infty}}}$ then dy/dx equals

A. $\frac{1}{2y-1}$

B. $\frac{1}{y^2-1}$

C. $\frac{2y}{y^2-1}$

D. None of these

Answer: A

 [Watch Video Solution](#)

30. The derivative of $e^{\sin^{-1} x}$ is :

A. $\frac{e^{\sin^{-1} x}}{\sqrt{1-x^2}}$

B. $\frac{e^{\sin^{-1} x}}{\sqrt{1+x^2}}$

C. $e^{\sin^{-1} x} (\sqrt{1+x^2})$

D. None of these

Answer: A [Watch Video Solution](#)**OBJECTIVE TYPE QUESTIONS (FILL IN THE BLANKS)**

1. If $f(x) = \begin{cases} kx^2 & \text{if } x \leq 2 \\ 3 & \text{if } x > 2 \end{cases}$ is continuous at $x = 2$, then the value of k is

 [Watch Video Solution](#)

2.

If

$f(x) = x + 7$, and $g(x) = x - 7$, $x \in R$ then find $\frac{d}{dx}(f \circ g)(x)$.

 [Watch Video Solution](#)

 Watch Video Solution

3. If $2x + 3y = \sin x$, then $\frac{dy}{dx} = \text{-----}$.

 Watch Video Solution

4. $\frac{dy}{dx} (\cos ec^{-1} x) = \text{-----}$.

 Watch Video Solution

5. $\frac{d}{dx} \left(\sqrt{e^{\sqrt{x}}} \right) = \text{-----}$.

 Watch Video Solution

6. If $x = at^2$, $y = 2at$, then $\frac{dy}{dx} =$

 Watch Video Solution

7. The derivative of x^x w.r.t. x is _____.



Watch Video Solution

8. If $x^2 + 3x + 2$, then $\frac{d^2y}{dx^2} =$ _____.



Watch Video Solution

9. Find the value of c in Rolle's theorem for the function $f(x) = x^3 - 3x$ in $[-\sqrt{3}, 0]$.



Watch Video Solution

10. Value of ' c ' in LMV Theorem for $f(x) = x^2$ on $[2, 4]$ is _____.



Watch Video Solution

1. Is the function defined by $f(x) = x^2 - \sin x + 5$ continuous at $x = \pi$?

 [Watch Video Solution](#)

2. If $x = 2at^2$, $y = at^4$, then $\frac{dy}{dx} = t^2$.

 [Watch Video Solution](#)

3. Differential of $\sin^2(x^2)$ w.r.t. x^2 is $\sin 2x^2$.

 [Watch Video Solution](#)

OBJECTIVE TYPE QUESTIONS (VERY SHORT ANSWER TYPE QUESTIONS)

1. Find the points at which the function $f(x) = [x]$ is not continuous in $(-1, 4)$, where $[x]$ is the largest integer function.



 [Watch Video Solution](#)

2. Find all points of discontinuity of f , where f is defined by

$$f(x) = \begin{cases} x^3 - 3, & \text{if } x \leq 2 \\ x^2 + 1, & \text{if } x < 2 \end{cases}$$

 [Watch Video Solution](#)

3. Discuss the continuity of the function:

$$f(x) = \begin{cases} 1 + x^2, & 0 \leq x \leq 1 \\ 2 - x, & x > 1 \end{cases} \text{ at } x = 1.$$

 [Watch Video Solution](#)

4. Differentiate $\cos\{\sin(x)^2\}$ w.r.t. x .

 [Watch Video Solution](#)

5. Find $\frac{dy}{dx}$ if $\sin^2 x + \cos^2 y = 1$.





Watch Video Solution

6. If $f(x) = x + 1$, then write the value of $\frac{d}{dx}(f \circ f)(x)$.



Watch Video Solution

7.

If

$f(x) = x + 7$, and $g(x) = x - 7$, $x \in R$ then find $\frac{d}{dx}(f \circ g)(x)$.



Watch Video Solution

8. If $y = \cos ec(\cot \sqrt{x})$, then find $\frac{dy}{dx}$.



Watch Video Solution

9. Differentiate $\sin(x^2 + 5)$ w.r.t. x .



Watch Video Solution

10. Differentiate $\sqrt{\sin(e^x)}$ w.r.t. x .

 [Watch Video Solution](#)

11. If $y = \log(\cos e^x)$, then find $\frac{dy}{dx}$.

 [Watch Video Solution](#)

12. Differentiate $(x^2 + 3x + 4)^n$, $n \in I$ w.r.t. x .

 [Watch Video Solution](#)

13. Find $\frac{dy}{dx}$ when $2x + 3y = \sin x$.

 [Watch Video Solution](#)

14. Find the derivative of the following w.r.t. x :

$$x - y = \pi$$



Watch Video Solution

15. Find the derivative of the following w.r.t. x :

$$x^3 + x^2y + xy^2 + y^3 = 81$$



Watch Video Solution

16. Find the derivative of the following w.r.t. x :

$$2x + 3y = \sin y$$



Watch Video Solution

17. Find the derivative of the following w.r.t. x :

$$ax + by^2 = \cos y$$



[Watch Video Solution](#)

18. Find the derivative of the following w.r.t. x :

$$\frac{1}{x} - \frac{1}{y} - 10 = 0$$

[Watch Video Solution](#)

19. Find the derivative of the following w.r.t. x :

$$y = \frac{4}{3}x^{3/4}$$

[Watch Video Solution](#)

20. Find the derivative of the following w.r.t. x :

$$\tan^{-1} \sqrt{x}$$

[Watch Video Solution](#)

21. Find the derivative of the following w.r.t. x :

$$\tan(\sin^{-1} x)$$

 [Watch Video Solution](#)

22. Find the derivative of the following w.r.t. x :

$$x \tan^{-1} x$$

 [Watch Video Solution](#)

23. Find the derivative of the following w.r.t. x :

$$\cos^{-1}(e^x)$$

 [Watch Video Solution](#)

24. Find the derivative of the following w.r.t. x :

$$\log(\log x), x > 1$$



[Watch Video Solution](#)

25. Find the derivative of the following w.r.t. x :

$$e^{\cos x}$$

[Watch Video Solution](#)

26. Find $\frac{dy}{dx}$, when $x = at^2$ and $y = 2at$

[Watch Video Solution](#)

27. Find $\frac{dy}{dx}$ when

$$x = 4t, y = \frac{4}{t}.$$

[Watch Video Solution](#)

28. Find $\frac{d^2y}{dx^2}$ when

$$y = e^x + \sin x$$

 [Watch Video Solution](#)

29. Find $\frac{d^2y}{dx^2}$ when

$$y = \tan^{-1} x$$

 [Watch Video Solution](#)

30. Find $\frac{d^2y}{dx^2}$ when

$$y = \frac{\log x}{x}.$$

 [Watch Video Solution](#)

31. If $\frac{dy}{dx} = \frac{y}{x}$, prove that $\frac{d^2y}{dx^2} = 0$.

 [Watch Video Solution](#)

32. If $2^x = 3^y$, then find $\frac{dy}{dx}$.

 [Watch Video Solution](#)

33. Find the second derivative of $\sin^{-1} x$.

 [Watch Video Solution](#)

34. Is Rolle's Theorem applicable to the function:

$f(x) = |x|$ in the interval $[-1, 1]$?

 [Watch Video Solution](#)

35. Is LMV Theorem applicable to the function:

$f(x) = \sin x \sin 2x$ in the interval $[0, \pi]$?

 [Watch Video Solution](#)

EXERCISE 5.1

1. Prove that the function $f(x) = 5x - 3$ is continuous at $x = 0$, at $x = -3$ and at $x = 5$.

 [Watch Video Solution](#)

2. Examine the continuity of the function $f(x) = 2x^2 - 1$ at $x = 3$.

 [Watch Video Solution](#)

3. Examine the following functions for continuity. (a) $f(x) = x - 5$ (b)

$f(x) = \frac{1}{x - 5}$ (c) $f(x) = \frac{x^2 - 25}{x + 5}$ (d) $f(x) = |x - 5|$

 [Watch Video Solution](#)

4. Examine the following functions for continuity :

$$f(x) = \frac{x^2 - 25}{x + 5}, x \neq -5$$

 [Watch Video Solution](#)

5. Examine the following functions for continuity:

(d) $f(x) = |x - 5|$.

 [Watch Video Solution](#)

6. Prove that the function $f(x) = x^n$ is continuous at $x = n$, where n is a positive integer.

 [Watch Video Solution](#)

7. Is the function f defined by

$$f(x) = \begin{cases} x & \text{if } x \leq 1 \\ 5 & \text{if } x > 1 \end{cases}$$

continuous at $x = 0$? At $x = 2$?

 [Watch Video Solution](#)

8. Find all points of discontinuity of f , where f is defined by

$$f(x) = \begin{cases} 2x + 3, & \text{if } x \leq 2 \\ 2x - 3, & \text{if } x > 2 \end{cases}$$

 [Watch Video Solution](#)

9. Find all points of discontinuity of f , where f is defined by $f(x) = \begin{cases} |x| + 3, & \text{if } x \geq -3 \\ -3 - 2x, & \text{if } x < -3 \end{cases}$

 [Watch Video Solution](#)

10. Find all points of discontinuity of f , where f is defined by

$$f(x) = \begin{cases} x, & \text{if } x < 0 \\ \frac{x}{|x|}, & \text{if } x \geq 0 \end{cases}$$

 [Watch Video Solution](#)

11. Find all points of discontinuity of f , where f is defined by

$$f(x) = \begin{cases} x & \text{if } x < 0 \\ \frac{1}{|x|} & \text{if } x \geq 0 \end{cases}$$

 [Watch Video Solution](#)

12. Find all points of discontinuity of f , where f is defined by

$$f(x) = \begin{cases} x + 1 & \text{if } x \geq 1 \\ x^2 + 1 & \text{if } x < 1 \end{cases}$$

 [Watch Video Solution](#)

13. Find all points of discontinuity of f , where f is defined by

$$f(x) = \begin{cases} x^3 - 3 & \text{if } x \leq 2 \\ x^2 + 1 & \text{if } x < 2 \end{cases}$$

 [Watch Video Solution](#)

14. Find all points of discontinuity of f , where f is defined by

$$f(x) = \begin{cases} x^{10} - 1 & \text{if } x \leq 1 \\ x^2 & \text{if } x > 1 \end{cases}$$

 [Watch Video Solution](#)

15. Is the function defined by

$$f(x) = \begin{cases} x + 5, & \text{if } x \leq 1 \\ x - 5, & \text{if } x > 1 \end{cases} \text{ a continuous function?}$$

 [Watch Video Solution](#)

16. Discuss the continuity of the function f , where f is defined by

$$f(x) = \begin{cases} 3, & \text{if } 0 \leq x \leq 1 \\ 4, & \text{if } 1 < x < 3 \\ 5, & \text{if } 3 \leq x \leq 10 \end{cases}$$

 [Watch Video Solution](#)

17. Discuss the continuity of the function f , where f is defined by :

$$f(x) = \begin{cases} 2x, & \text{if } x < 0 \\ 0, & \text{if } 0 \leq x \leq 1 \\ 4x, & \text{if } x > 1 \end{cases}$$

 [Watch Video Solution](#)

18. Discuss the continuity of the function f , where f is defined by $f(x) = \begin{cases} -2, & \text{if } x < -1 \\ 2x, & \text{if } -1 \leq x < 1 \\ -1, & \text{if } x = 1 \end{cases}$

 [Watch Video Solution](#)

19. Find the relationship between a and b so that the function f defined by: $f(x) = \begin{cases} ax + 1 & \text{if } x \leq 3 \\ bx + 3 & \text{if } x > 3 \end{cases}$

 [Watch Video Solution](#)

20. For what value of λ is the function defined by $f(x) = \begin{cases} \lambda(x^2 - 2x), & \text{if } x \leq 0 \\ 4x + 1, & \text{if } x > 0 \end{cases}$ continuous at $x = 0$? What about continuity at $x = 1$?

 [Watch Video Solution](#)

21. Show that the function defined by $g(x) = x - [x]$ is discontinuous at all integral points which $[x]$ denotes the greatest integer function.

 [Watch Video Solution](#)

22. Is the function defined by $f(x) = x^2 - \sin x + 5$ continuous at $x = \pi$?

 [Watch Video Solution](#)

23. Discuss the continuity of the following functions a)

$$f(x) = \sin x + \cos x$$

 [Watch Video Solution](#)

24. Discuss the continuity of $f(x) = \sin x - \cos x$

 [Watch Video Solution](#)

25. Discuss the continuity of the following functions :

$$f(x) = \sin x \cdot \cos x$$

 [Watch Video Solution](#)

26. Discuss the continuity of the cosine, cosecant, secant and cotangent functions.

 [Watch Video Solution](#)

27. Find all points of discontinuity of f , where

$$f(x) = \begin{cases} \frac{\sin x}{x}, & \text{if } x < 0 \\ x + 1, & \text{if } x \geq 0 \end{cases}$$

 [Watch Video Solution](#)

28. Determine if f defined by :

$$f(x) = \begin{cases} x^2 \sin \frac{1}{x}, & \text{if } x \neq 0 \\ 0, & \text{if } x = 0 \end{cases} \text{ is a continuous function ?}$$



[Watch Video Solution](#)

29. Examine the continuity of f , where f is defined by :

$$f(x) = \begin{cases} \sin x + \cos x, & \text{if } x \neq 0 \\ 1, & \text{if } x = 0 \end{cases} .$$



[Watch Video Solution](#)

30. Find the value of k so that the function f defined by

$$f(x) = \begin{cases} \frac{k \cos x}{\pi - 2x}, & \text{if } x \neq \frac{\pi}{2} \\ 3, & \text{if } x = \frac{\pi}{2} \end{cases} \text{ is continuous at } x = \frac{\pi}{2}$$



[Watch Video Solution](#)

31. $f(x) = \begin{cases} kx^2, & \text{if } x \leq 2 \\ 3, & \text{if } x > 2 \end{cases}$ at $x = 2$



Watch Video Solution

32. Find the values of k so that the function f is continuous at the indicated point in $f(x) = \begin{cases} kx + 1, & \text{if } x \leq \pi \\ \cos x, & \text{if } x > \pi \end{cases}$ at $x = \pi$



Watch Video Solution

33. Find the values of k so that the function f is continuous at the indicated point in $f(x) = \begin{cases} kx + 1, & \text{if } x \leq 5 \\ 3x - 5, & \text{if } x > 5 \end{cases}$ at $x = 5$



Watch Video Solution

34. Find the value of 'a' and 'b' such that the function defined by :

$$f(x) = \begin{cases} 5, & \text{if } x \leq 2 \\ ax + b, & \text{if } 2 < x < 5 \\ 20, & \text{if } x \geq 5 \end{cases} \text{ is a continuous function.}$$



Watch Video Solution

35. If $f(x) = \left(\frac{3x + \tan^2 x}{x} \right)$ is continuous at $x = 0$, then $f(0)$ is equal to.

A. 3

B. 2

C. 4

D. 0



Watch Video Solution

36. Show that the function defined by $f(x) = |\cos x|$ is a continuous function.



Watch Video Solution

37. Examine that $\sin|x|$ is a continuous function.



Watch Video Solution

38. Find all the points of discontinuity of f defined by

$$f(x) = \frac{|x|}{|x + 1|}.$$



Watch Video Solution

EXERCISE 5.2

1. Differentiate the functions with respect to x in :

$$\sin(x^2 + 5)$$



Watch Video Solution

2. Differentiate the functions with respect to x in :

$$\cos(\sin x)$$

 [Watch Video Solution](#)

3. Differentiate the functions with respect to x in :

$$\sin(ax + b)$$

 [Watch Video Solution](#)

4. Differentiate the functions with respect to x in :

$$\sec(\tan \sqrt{x})$$

 [Watch Video Solution](#)

5. Differentiate the functions with respect to x in :

$$\frac{\sin(ax + b)}{\cos(cx + d)}$$



[Watch Video Solution](#)

6. Differentiate the functions with respect to x in :

$$\cos x^3 \cdot \sin^2(x^5)$$



[Watch Video Solution](#)

7. Differentiate the functions with respect to x in :

$$2\sqrt{\tan(x^2)}$$



[Watch Video Solution](#)

8. Differentiate the functions with respect to x in :

$$\cos(\sqrt{x})$$



[Watch Video Solution](#)

9. Prove that the function f given by $f(x) = |x - 1|$, $x \in \mathbb{R}$ is not differentiable at $x = 1$

 [Watch Video Solution](#)

10. Prove that the greatest integer function defined by $f(x) = [x]$, $0 < x < 3$ is not differentiable at $x = 1$ and $x = 2$.

 [Watch Video Solution](#)

EXERCISE 5.3

1. Find $\frac{dy}{dx}$ in the following :

$$2y + 3x = \sin y$$

 [Watch Video Solution](#)

2. Find $\frac{dy}{dx}$ in the following: $2x + 3y = s \in y$



Watch Video Solution

3. Find $\frac{dy}{dx}$ in the following :

$$ax + by^2 = \cos y$$



Watch Video Solution

4. Find $\frac{dy}{dx}$ in the following: $xy + y^2 = \tan x + y$



Watch Video Solution

5. Find $\frac{dy}{dx}$ in the following :

$$x^2 + y^2 = 5xy$$



Watch Video Solution

6. $x^3 + x^2y + xy^2 + Y^3 = 81$

 [Watch Video Solution](#)

7. Find $\frac{dy}{dx}$ in the following :

$$\cos xy = x^2y$$

 [Watch Video Solution](#)

8. Find $\frac{dy}{dx}$ in the following :

$$\sin^3 x + \cos^3 y = 1$$

 [Watch Video Solution](#)

9. Find $\frac{dy}{dx}$ in the following: $y = \sin^{-1}\left(\frac{2x}{1+x^2}\right)$

 [Watch Video Solution](#)

10. Find $\frac{dy}{dx}$ in the following :

$$y = \tan^{-1}\left(\frac{3x - x^3}{1 - 3x^2}\right), \quad -\frac{1}{\sqrt{3}} < x < \frac{1}{\sqrt{3}}$$

 [Watch Video Solution](#)

11. Find $\frac{dy}{dx}$ in the following: $y = \cos^{-1}\left(\frac{1-x^2}{1+x^2}\right), 0$

 [Watch Video Solution](#)

12. Find $\frac{dy}{dx}$ in the following: $y = \sin^{-1}\left(\frac{1-x^2}{1+x^2}\right), 0$

 [Watch Video Solution](#)

13. Find $\frac{dy}{dx}$ in the following: $y = \cos^{-1}\left(\frac{2x}{1+x^2}\right), -1$

 [Watch Video Solution](#)

14. Find $\frac{dy}{dx}$ in the following :

$$y = \sin^{-1}\left(2x\sqrt{1-x^2}\right), \quad -\frac{1}{\sqrt{2}} < x < \frac{1}{\sqrt{2}}.$$

 [Watch Video Solution](#)

15. Find $\frac{dy}{dx}$ in the following: $y = \sec^{-1}\left(\frac{1}{2x^2-1}\right), 0$

 [Watch Video Solution](#)

EXERCISE 5.4

1. Differentiate the following w.r.t. x :

$$\frac{e^x}{\sin x}$$

 [Watch Video Solution](#)

2. Differentiate the following w.r.t. x :

$$e^{\sin^{-1} x}$$



Watch Video Solution

3. Differentiate the following w.r.t. x :

$$e^{x^3}$$



Watch Video Solution

4. Differentiate the following w.r.t. x :

$$\cos(\tan^{-1} e^{-x})$$



Watch Video Solution

5. Differentiate the following w.r.t. x :

$$\log(\sin e^x)$$





Watch Video Solution

6. Differentiate the following w.r.t. x :

$$e^x + e^{x^2} + \dots + e^{x^7}$$



Watch Video Solution

7. Differentiate the following w.r.t. x :

$$\sqrt{e^{\sqrt{x}}}, x > 0$$



Watch Video Solution

8. Differentiate the following w.r.t. x :

$$\log(\log x), x > 1$$



Watch Video Solution

9. Differentiate the following w.r.t. x :

$$\frac{\cos x}{\log x}, x > 0$$

 [Watch Video Solution](#)

10. Differentiate the following w.r.t. x :

$$\cos(\log x + e^x), x > 0$$

 [Watch Video Solution](#)

EXERCISE 5.5

1. Differentiate the functions given in w.r.t. x :

$$\cos 2x + \sin 3x.$$

 [Watch Video Solution](#)

2. Differentiate the functions given w.r.t. x:

$$\sqrt{\left((x-1) \frac{x-2}{(x-3)(x-4)(x-5)}\right)}$$



Watch Video Solution

3. Differentiate the functions given in w.r.t. x :

$$(\log x)^{\cos x}$$



Watch Video Solution

4. Differentiate the functions given in w.r.t. x :

$$x^x - 2^{\sin x}$$



Watch Video Solution

5. Differentiate the functions given in w.r.t. x :

$$(x+7)^2 \cdot (x+4)^3.$$



Watch Video Solution

6. Differentiate the functions given w.r.t. x : $\left(x + \frac{1}{x}\right)^x + x^{\left(1 + \frac{1}{x}\right)}$



Watch Video Solution

7. Differentiate the functions given in w.r.t. x :

$$(\log x)^x + x^{\log x}$$



Watch Video Solution

8. Differentiate the functions given in w.r.t. x :

$$(\sin x)^x + \sin^{-1} \sqrt{x}$$



Watch Video Solution

9. Differentiate the functions given in w.r.t. x :

$$x^{\sin x} + (\sin x)^{\cos x}$$



Watch Video Solution

10. Differentiate the functions given in w.r.t. x :

$$x^{x \cos x} + \frac{x^2 + 1}{x^2 - 1}$$



Watch Video Solution

11. Differentiate the functions given in w.r.t. x :

$$(x \cos x)^x + (x \sin x)^{1/x}.$$



Watch Video Solution

12. Find $\frac{dy}{dx}$ of the function given in :

$$x^y + y^x = b$$

 [Watch Video Solution](#)

13. Find $\frac{dy}{dx}$ of the function given in :

$$y^x = x^y$$

 [Watch Video Solution](#)

14. Find $\frac{dy}{dx}$, when:

$$(\cos x)^y = (\cos y)^x$$

 [Watch Video Solution](#)

15. Find $\frac{dy}{dx}$ of the functions given $xy = e^{(x-y)}$

 [Watch Video Solution](#)

16. Find the derivative of the function given by $f(x) = (1 + x)(1 + x^2)(1 + x^4)(1 + x^8)$ and hence find $f'(1)$.

 [Watch Video Solution](#)

17. Differentiate $(x^2 - 5x + 8)(x^3 + 7x + 9)$ in three ways mentioned below: (i) by using product rule (ii) by expanding the product to obtain a single polynomial. (iii) by logarithmic differentiation. Do they all give the same answer?

 [Watch Video Solution](#)

18. If u , v and w are functions of x , then show that $\frac{d}{dx}(uvw) = \frac{du}{dx}vw + u\frac{dv}{dx}w + uv\frac{dw}{dx}$ in two ways - first by repeated application of product rule, second by logarithmic differentiation.

 [Watch Video Solution](#)

EXERCISE 5.6

1. If x and y are connected parametrically by the equations given, without eliminating the parameter, Find $\frac{dy}{dx} \cdot x = 2at^2, y = at^4$

 [Watch Video Solution](#)

2. If x and y are connected parametrically by the equations given in, without eliminating the parameter, find $\frac{dy}{dx}$.

$$x = a \cos \theta, y = b \sin \theta.$$

 [Watch Video Solution](#)

3. If x and y are connected parametrically by the equations given in, without eliminating the parameter, find $\frac{dy}{dx}$.

$$x = a(\cos t + t \sin t), y = a(\sin t - t \cos t)$$

 [Watch Video Solution](#)

4. If x and y are connected parametrically by the equations given in, without eliminating the parameter, find $\frac{dy}{dx}$.

$$x = t^2 - 2t, y = t^4 - 4t$$



[Watch Video Solution](#)

5. If x and y are connected parametrically by the equations given in, without eliminating the parameter, find $\frac{dy}{dx}$.

$$x = \cos \phi - \cos 2\phi, y = \sin \phi - \sin 2\phi$$



[Watch Video Solution](#)

6. If x and y are connected parametrically by the equations given, without eliminating the parameter, Find $\frac{dy}{dx}$. $x = a(\theta - \sin \theta), y = a(1 + \cos \theta)$



[Watch Video Solution](#)

7. If x and y are connected parametrically by the equations given, without

eliminating the parameter, Find $\frac{dy}{dx} \cdot x = \frac{\sin^3 t}{\sqrt{\cos 2t}}, y = \frac{\cos^3 t}{\sqrt{\cos 2t}}$



[Watch Video Solution](#)

8. If x and y are connected parametrically by the equations given in,

without eliminating the parameter, find $\frac{dy}{dx}$.

$$x = a \left(\cos \theta + \log \tan \frac{\theta}{2} \right), y = a \sin \theta.$$



[Watch Video Solution](#)

9. If x and y are connected parametrically by the equations given in,

without eliminating the parameter, find $\frac{dy}{dx}$.

$$x = a \sec t, y = b \tan t.$$



[Watch Video Solution](#)

10. If x and y are connected parametrically by the equations given, without eliminating the parameter, Find $\frac{dy}{dx}$.

$$x = a(\cos \theta + \theta \sin \theta), y = a(\sin \theta - \theta \cos \theta)$$

 [Watch Video Solution](#)

11. If $x = \sqrt{a^{\sin^{-1}((-1)t)}}$, $y = \sqrt{a^{\cos^{-1}((-1)t)}}$, show that $\frac{dy}{dx} = -\frac{y}{x}$

 [Watch Video Solution](#)

EXERCISE 5.7

1. Find the first and second order derivatives of the functions given in :

$$x^2 + 3x + 2.$$

 [Watch Video Solution](#)

2. Find the second order derivatives of the functions given. x^{20}



Watch Video Solution

3. Find the second order derivatives of the functions given in :

$x \cdot \cos x$.



Watch Video Solution

4. Find the second order derivatives of the functions given in :

$x \log x$.



Watch Video Solution

5. Find the second order derivatives of the functions given in :

$x^3 \log x$.



Watch Video Solution

6. Find the second order derivatives of the functions given in :

$$e^{2x} \sin 5x.$$

 [Watch Video Solution](#)

7. Find the second order derivatives of the functions given in :

$$e^{5x} \cos 3x.$$

 [Watch Video Solution](#)

8. Find the second order derivatives of the functions given in :

$$\tan^{-1} x.$$

 [Watch Video Solution](#)

9. Find the second order derivatives of the functions given in :

$$\log(\log x)$$



[Watch Video Solution](#)

10. Find the second order derivatives of the functions given in :

$\sin(\log x)$.

[Watch Video Solution](#)

11. If $y = 5 \cos x - 3 \sin x$, prove that $\frac{d^2y}{dx^2} + y = 0$.

[Watch Video Solution](#)

12. If $y = \cos^{-1} x$, find $\frac{d^2y}{dx^2}$ in terms of y alone.

[Watch Video Solution](#)

13. If $y = 3 \cos(\log x) + 4 \sin(\log x)$, show that $x^2 y_2 + x y_1 + y = 0$

[Watch Video Solution](#)

14. If $y = Ae^{mx} + Be^{nx}$, show that $\frac{d^2y}{dx^2} - (m + n)\frac{dy}{dx} + mny = 0$

 [Watch Video Solution](#)

15. If $y = 500e^{7x} + 600e^{-7x}$, show that $\frac{d^2y}{dx^2} = 49y$.

 [Watch Video Solution](#)

16. If $e^y(x + 1) = 1$, show that $\frac{d^2y}{dx^2} = \left(\frac{dy}{dx}\right)^2$.

 [Watch Video Solution](#)

17. If $y = (\tan^{-1} x)^2$, show that $(x^2 + 1)^2 y_2 + 2x(x^2 + 1)y_1 = 2$

 [Watch Video Solution](#)

EXERCISE 5.8

1. Verify Rolles theorem for the function $f(x) = x^2 + 2x - 8$, $x \in [-4, 2]$.

 [Watch Video Solution](#)

2. Examine if Rolles theorem is applicable to any of the following functions. Can you say something about the converse of Rolles theorem from these example?(i) $f(x) = [x]$ for $x \in [5, 9]$ (ii) $f(x) = [x]$ for $x \in [-2, 2]$ (iii) $f(x) = x^2$

 [Watch Video Solution](#)

3. If $f: [-5, 5] \rightarrow \mathbb{R}$ is differentiable function and if $f'(x)$ does not vanish anywhere, then prove that $f(-5) \neq f(5)$.

 [Watch Video Solution](#)

4. Verify Mean Value Theorem, if $f(x) = x^2 - 4x - 3$ in the interval $[a, b]$, where $a = 1$ and $b = 4$.

 [Watch Video Solution](#)

5. Verify Mean Value theorem, if $f(x) = x^3 - 5x^2 - 3x$ in the interval $[a, b]$, where $a = 1$ and $b = 3$. Find all $c \in (1, 3)$ for which $f'(c) = 0$.

 [Watch Video Solution](#)

6. Differentiate $\frac{x^m}{x^n}$

 [Watch Video Solution](#)

MISCELLANEOUS EXERCISE

1. Differentiate w.r.t. x the function in $(3x^2 + 9x + 7)^9$.

 [Watch Video Solution](#)

2. Differentiate w.r.t. x the function in

$$\sin^3 x + \cos^6 x.$$

 [Watch Video Solution](#)

3. Differentiate w.r.t. x the function $(5x)^{3 \cos 2x}$.

 [Watch Video Solution](#)

4. $\sin^{-1}(x\sqrt{x}), 0 \leq x \leq 1$

 [Watch Video Solution](#)

5. Differentiate w.r.t. x the function $\frac{\cos^{-1}x}{\sqrt{2x+7}}, -2$

 [Watch Video Solution](#)

6. Differentiate w.r.t. x the function in

$$\cot^{-1} \left[\frac{\sqrt{1 + \sin x} + \sqrt{1 - \sin x}}{\sqrt{1 + \sin x} - \sqrt{1 - \sin x}} \right], 0 < x < \frac{\pi}{2}$$

 [Watch Video Solution](#)

7. Differentiate w.r.t. x the function in

$$(\log x)^{\log x}, x > 1.$$

 [Watch Video Solution](#)

8. Differentiate w.r.t. x the function

$$\cos (a \cos x + b \sin x), \text{ for some constant } a \text{ and } b.$$

 [Watch Video Solution](#)

9. Differentiate w.r.t. x the function in

$$(\sin x - \cos x)^{(\sin x - \cos x)}, \frac{\pi}{4} < x < \frac{3\pi}{4}.$$

 [Watch Video Solution](#)

10. Differentiate w.r.t. x the function $x^x + x^a + a^x + a^a$, for some fixed

$$a > 0 \text{ and } x > 0.$$

 [Watch Video Solution](#)

11. Differentiate w.r.t. x the function $x^{x^2 - 3} + (x - 3)^{x^2 - 2}$ for $x > 3$.

 [Watch Video Solution](#)

12. Find $\frac{dy}{dx}$, if $y = 12(1 - \cos t)$, $x = 10(t - \sin t)$, $t = \pi/2$

 [Watch Video Solution](#)

13. Find $\frac{dy}{dx}$, if $y = \sin^{-1} x + \sin^{-1} \sqrt{1 - x^2}$, $-1 \leq x \leq 1$

 [Watch Video Solution](#)

14. If $x\sqrt{1+y} + y\sqrt{1+x} = 0$, prove that $\frac{dy}{dx} = -\frac{1}{(x+1)^2}$.

 [Watch Video Solution](#)

15. If $(x-a)^2 + (y-b)^2 = c^2$, for some $c > 0$, prove that

$\frac{\left[1 + \left(\frac{dy}{dx}\right)^2\right]^{\frac{3}{2}}}{\frac{d^2y}{dx^2}}$ is a constant independent of a and b .

 [Watch Video Solution](#)

16. If $\cos y = x \cos(a+y)$, with $\cos a \neq \pm 1$, prove that

$\frac{dy}{dx} = \frac{\cos^2(a+y)}{\sin a}$.



Watch Video Solution

17. If $x = a(\cos t + t \sin t)$ and $y = a(\sin t - t \cos t)$, find $\frac{d^2y}{dx^2}$.



Watch Video Solution

18. If $f(x) = |x|^3$, show that $f(x)$ exists for all real x and find it.



Watch Video Solution

19. Using mathematical induction prove that $\frac{d}{dx}(x^n) = nx^{n-1}$ for all positive integers n .



Watch Video Solution

20. Using the fact that $s \in (A + B) = s \in A \cos B + \cos A s \in B$ and the differentiation, obtain the sum formula for cosines.

 [Watch Video Solution](#)

21. Does there exist a function which is continuous everywhere but not differentiable at exactly two points? Justify your answer.

 [Watch Video Solution](#)

22. If $y = |f(x)g(x)h(x)lmnabc|$, prove that

$$\frac{dy}{dx} = |f'(x)g'(x)h'(x)lmnabc|$$

 [Watch Video Solution](#)

23. If $y = e^a \cos^{(-1)x}$, $-1 \leq x < 1$, show that

$$(1 - x^2) \frac{d^2y}{dx^2} - x \frac{dy}{dx} - a^2y = 0$$

 [Watch Video Solution](#)

1. Examine the continuity of the function :

$$f(x) = \begin{cases} \frac{|x-4|}{(x-4)}, & \text{if } x \neq 4 \\ 0, & \text{if } x = 4 \end{cases} \text{ at } x = 4.$$



Watch Video Solution

2. For what value of k , $f(x) = \begin{cases} \frac{2^{x+2}-16}{4^x-16}, & x \neq 2 \\ k, & x = 2 \end{cases}$ is continuous at $x = 2$?



Watch Video Solution

3. Find the values of a and b such that the function f defined by

$$f(x) = \begin{cases} \frac{x-4}{|x-4|} + a & \text{if } x < 4 \\ a + b & \text{if } x = 4 \\ \frac{x-4}{|x-4|} + b & \text{if } x > 4 \end{cases}$$

is a continuous function at $x = 4$.



Watch Video Solution

4. Given $f(x) = \frac{1}{x-1}$. Find the points of discontinuity of the composite function $f(f(x))$.

 [Watch Video Solution](#)

5. Find $f'(x)$ when

$$f(x) = 2^{\cos^2 x}$$

 [Watch Video Solution](#)

6. Find $f'(x)$ when

$$f(x) = \sin^{-1} \left(\frac{1}{\sqrt{x+1}} \right).$$

 [Watch Video Solution](#)

7. If $\sin x = \frac{2t}{1+t^2}$, $\tan y = \frac{2t}{1-t^2}$, find $\frac{dy}{dx}$

 [Watch Video Solution](#)

8. If $y = \sec^{-1}\left(\frac{1}{4x^3 - 3x}\right)$, find $\frac{dy}{dx}$.

 [Watch Video Solution](#)

9. Find $\frac{dy}{dx}$ when $\tan^{-1}(x^2 + y^2) = 0$.

 [Watch Video Solution](#)

10. Examine the differentiability of f , where f is defined by

$$f(x) = \begin{cases} x[x] & \text{if } 0 \leq x < 2 \\ (x-1)x & \text{if } 2 \leq x < 3 \end{cases} \text{ at } x = 2$$

 [Watch Video Solution](#)

11. Show that $f(x) = |x - 5|$ is continuous but not differentiable at $x = 5$.

 [Watch Video Solution](#)

12. A function $f: R \rightarrow R$ satisfies the equation $f(x + y) = f(x) \cdot f(y)$ for all $x, y \in R$, $f(x) \neq 0$. Suppose that the function is differentiable at $x = 0$ and $f'(0) = 2$, then prove that $f' = 2f(x)$.

 [Watch Video Solution](#)

13. If $x = e^{x/y}$, then prove that $\frac{dy}{dx} = \frac{x - y}{x \log x}$.

 [Watch Video Solution](#)

14. $f(x) = \frac{1}{4x - 1}$ in $[1, 4]$

 [Watch Video Solution](#)

15. Discuss the Lagrange's mean value theorem for the function $f(x) = \sin x - \sin 2x$, in $[0, \pi]$.



Watch Video Solution

16. $f(x) = \sqrt{25 - x^2}$ in $[1, 5]$



Watch Video Solution

REVISION EXERCISE

1. Show that $f(x) = \begin{cases} |2x - 3|[x], & x \geq 1 \\ \sin\left(\frac{\pi x}{2}\right), & x < 1 \end{cases}$ is continuous but not

differentiable at $x = 1$.



Watch Video Solution

2. Does there exist a function which is continuous everywhere but not differentiable at exactly two points? Justify your answer.



Watch Video Solution

3. Is $|\sin x|$ differentiable? What about $\cos|x|$?

 [Watch Video Solution](#)

4. Find, from first principle, the derivative of $\frac{\sin^{-1} x}{x}$ w.r.t. x .

 [Watch Video Solution](#)

5. Differentiate the following w.r.t. x :

$$\sqrt{3x+2} + \frac{1}{\sqrt{2x^2+4}}$$

 [Watch Video Solution](#)

6. Differentiate the following w.r.t. x :

$$e^{\sec^2 x} + 3 \cos^{-1} x.$$

 [Watch Video Solution](#)

7. Differentiate the following w.r.t. x :

$$f(x) = \log \left[(2 - x)^{1/2} (x^2 - 1)^{-1/4} \right]$$



[Watch Video Solution](#)

8. Differentiate the following function w.r.t x :

$$\tan^{-1} \left(\frac{\sqrt{1+x} - \sqrt{1-x}}{(\sqrt{1+x} + \sqrt{1-x})} \right)$$



[Watch Video Solution](#)

9. Differentiate the following w.r.t. x :

$$(3x^2 - 9x + 5)^9$$



[Watch Video Solution](#)

10. Differentiate the following w.r.t. x :

$$\sin^3 x + \cos^6 x$$



Watch Video Solution

11. Differentiate the following w.r.t. x :

$$e^{\log(x + \sqrt{x^2 + a^2})}$$



Watch Video Solution

12. Differentiate the following w.r.t. x :

$$e^{2 \log x + 3x}$$



Watch Video Solution

13. Prove that $\left(\cot^{-1} x + \cot^{-1} \frac{1}{x} \right)$ is a constant.



Watch Video Solution

14. If $y = f\left(\frac{2x - 1}{x^2 + 1}\right)$ and $f'(x) = \sin x^2$, find $\frac{dy}{dx}$.



Watch Video Solution

15. Find the derivative of the following w.e.t. x :

$$\frac{3}{\sqrt[3]{x}} - \frac{5}{\cos x} + \log_3 x + \frac{6}{\sin x} - \frac{2 \tan x}{\sec x} + 7$$



Watch Video Solution

16. Find the derivative of the following w.e.t. x :

$$\log\left(\frac{1}{\sqrt{x}}\right) + 5x^a - 3a^x + \sqrt[3]{x^2} + 6\sqrt[4]{x^{-3}}.$$



Watch Video Solution

17. If $y = \tan^{-1}\left(\frac{e^{2x} + 1}{e^{2x} - 1}\right)$, prove that :

$$\frac{dy}{dx} = -\frac{2e^{2x}}{1 + e^{4x}}.$$



Watch Video Solution

18. If the derivative of $\tan^{-1}(a + bx)$ takes the value 1 at $x = 0$, prove that $1 + a^2 = b$.

 [Watch Video Solution](#)

19. Using the fact that $s \in (A + B) = s \in A \cos B + \cos A s \in B$ and the differentiation, obtain the sum formula for cosines.

 [Watch Video Solution](#)

20. If $\sqrt{y+x} + \sqrt{y-x} = c$ show that $\frac{dy}{dx} = \frac{y}{x} - \sqrt{\left(\frac{y^2}{x^2}\right) - 1}$

 [Watch Video Solution](#)

21. if $\sin x = y \sin(x + b)$ show that $\frac{dy}{dx} = \frac{\sin b}{\sin^2(x + b)}$

 [Watch Video Solution](#)

22. If $x \sin(a + y) + \sin a \cos(a + y) = 0$, provethat

$$\frac{dy}{dx} = \frac{\sin^2(a + y)}{\sin a}$$

 [Watch Video Solution](#)

23. If $y = x \sin(a + y)$, provethat $\frac{dy}{dx} = \frac{\sin^2(a + y)}{\sin(a + y) - y \cos(a + y)}$.

 [Watch Video Solution](#)

24. Differentiate $\log[\log(\log x)]$ w.r.t. x .

 [Watch Video Solution](#)

25. if $y = e^{(x)^{e^x}} + x^{e^{e^x}} + e^{x^{x^e}}$, then dy/dx

$$= e^{(x)^{e^x}} x^{e^x} \left[e^x \log x + \frac{e^x}{x} \right] + x^{e^{e^x}} e^{e^x} \left[\frac{1}{x} + e^x \log x \right] + e^{x^{x^e}} x^{x^e} x^{e-1} [1 + e$$

 [Watch Video Solution](#)

26. $y = \sqrt{x + \sqrt{x + \sqrt{x + \dots + \infty}}}$. Find $\frac{dy}{dx}$

 [Watch Video Solution](#)

27. Find $\frac{dy}{dx}$ when :

$$y = x^{\sin x - \cos x} + \frac{x^2 - 1}{x^2 + 1}$$

 [Watch Video Solution](#)

28. If $y = x^{\cot x} + \frac{2x^2 - 3}{x^2 + x + 2}$, find $\frac{dy}{dx}$.

 [Watch Video Solution](#)

29. If $y = x^{(x^x)}$, prove that :

$$\frac{dy}{dx} = x^{x+x^x} \left[\frac{1}{x} + (1 + \log x) \log x \right].$$

 [Watch Video Solution](#)

30. If $y = (\tan x)^{(\tan x)^{(\tan x) \dots \infty}}$, then prove that $\frac{dy}{dx} = 2$ at $x = \frac{\pi}{4}$.

 [Watch Video Solution](#)

31. Differentiate $\tan x \tan 2x \tan 3x \tan 4x$ in two ways :

(i) by taking logarithms and

(ii) by repeatedly applying product rule.

 [Watch Video Solution](#)

32. If $x = \sec \theta - \cos \theta$ and $y = \sec^n \theta - \cos^n \theta$ then show that

$$(x^2 + 4) \left(\frac{dy}{dx} \right)^2 = n^2 (y^2 + 4)$$

 [Watch Video Solution](#)

33. For what choices of a, b, c, if any, does the function

$$f(x) = \begin{cases} ax^2 + bx + x & , 0 \leq x \leq 1 \\ bx - c & , 1 \leq x \leq 2 \\ x & , x > 2 \end{cases}$$

become differentiable at $x = 1$ and $x = 2$?



Watch Video Solution

34. If $y = Ae^{-kt} \cos(pt + c)$, then prove that $\frac{d^2y}{dt^2} + 2k\frac{dy}{dt} + n^2y = 0$,

where $n^2 = p^2 + k^2$



Watch Video Solution

35. Using mathematical induction prove that :

$$\frac{d}{dx}(x^n) = nx^{n-1} \text{ for all } n \in \mathbb{N}.$$



Watch Video Solution

36. If $(1 + x)^n = C_0 + C_1x + C_2x^2 + \dots + C_nx^n$, then prove that :

(i) $C_1 + 2C_2 + \dots + nC_n = n \cdot 2^{n-1}$

(ii) $C_1 - 2C_2 + \dots + (-1)^{n-1}nC_n = 0$



Watch Video Solution

37. If $y = \frac{ax^2}{(x-a)(x-b)(x-c)} + \frac{bx}{(x-b)(x-c)} + \frac{c}{x-c} + 1$ then

$$\frac{y'}{y} =$$



Watch Video Solution

CHECK YOUR UNDERSTANDING

1. Are the following functions continuous at each point of their domains?

$$e^x$$



Watch Video Solution

2. Are the following functions continuous at each point of their domains?

$\sin x$

 [Watch Video Solution](#)

3. Are the following functions continuous at each point of their domains?

$\cos x$

 [Watch Video Solution](#)

4. Is greatest integer function $[x]$ is continuous everywhere ?

 [Watch Video Solution](#)

5. Find $\frac{dy}{dx}$ when $2x + 3y = \cos x$.

 [Watch Video Solution](#)

6. Find the derivative of $\cos^{-1} x$ assuming that it exists.

 [Watch Video Solution](#)

7. Differentiate $e^{\sin^{-1} x}$, w.r.t. x .

 [Watch Video Solution](#)

8. If $y = \log(\cos e^x)$, then $\frac{dy}{dx} = \text{-----}$.

 [Watch Video Solution](#)

9. Find $\frac{d^2y}{dx^2}$ when $y = \log x + x$.

 [Watch Video Solution](#)

10. Find $\frac{d^2y}{dx^2}$ when $y = e^x + \cos x$



Watch Video Solution

11. Verify Rolle's theorem for $f(x) = x, x \in [1, 2]$.



Watch Video Solution

12. Verify L.M.V. theorem for $f(x) = x(x - 2)$ on $[1, 3]$



Watch Video Solution

COMPETITION FILE

1. Let y be an implicit function of x defined by $x^{2x} - 2x^x \cot y - 1 = 0$.

Then $y'(1)$ equals: 1 b. $\log 2$ c. $-\log 2$ d. -1

A. -1

B. 1

C. $\log 2$

D. $-\log 2$

Answer: A



[Watch Video Solution](#)

2. Let $f : (-1,1) \rightarrow \mathbb{R}$ be a differentiable function with $f(0) = -1$ and $f'(0) = 1$

Let $g(x) = [f(f(2x) + 1)]^2$. Then $g'(0) =$

A. 4

B. -4

C. $\log 2$

D. $-\log 2$

Answer: B



[Watch Video Solution](#)

3. $\frac{d^2x}{dy^2}$ equals

A. $\left(\frac{d^2y}{dx^2}\right)^{-1}$

B. $-\left(\frac{d^2y}{dx^2}\right)^{-1} \left(\frac{dy}{dx}\right)^{-3}$

C. $\left(\frac{d^2y}{dx^2}\right)^{-1} \left(\frac{dy}{dx}\right)^{-2}$

D. $-\left(\frac{d^2y}{dx^2}\right) \left(\frac{dy}{dx}\right)^{-3}$

Answer: D



Watch Video Solution

4. If $f(x)$ is differentiable at $x = a$, find $\lim_{x \rightarrow a} \frac{x^2 f(a) - a^2 f(x)}{x - a}$.

A. $a^2 f(a)$

B. $a f(a) - a^2 f'(a)$

C. $2a f(a) - a^2 f'(a)$

D. $2a f(a) + a^2 f'(a)$

Answer: C



Watch Video Solution

5. If $f: \mathbb{R} \rightarrow \mathbb{R}$ is a function defined by $f(x) = [x] \cos\left(\frac{2x-1}{2}\right)\pi$, where $[x]$ denotes the greatest integer function, then f is-

- A. continuous for every real x
- B. discontinuous only at $x=0$
- C. discontinuous only at non-zero integral values of x
- D. continuous only at $x=0$

Answer: A



Watch Video Solution

6. If $y = \sec(\tan^{-1} x)$, then $\frac{dy}{dx}$ at $x = 1$ is equal to

A. $\frac{1}{2}$

B. 1

C. $\sqrt{2}$

D. $\frac{1}{\sqrt{2}}$

Answer: D



Watch Video Solution

7. If g is the inverse of a function f and $f'(x) = \frac{1}{1+x^5}$, then $g'(x)$ is equal to

A. $5x^4$

B. $\frac{1}{1 + \{g(x)\}^5}$

C. $1 + \{g(x)\}^5$

D. $1 + x^5$

Answer: B



Watch Video Solution

8. if the function $g(x) = \begin{cases} k\sqrt{x+1}, 0 \leq x \leq 3 \\ mx+2, 3 < x \leq 5 \end{cases}$ is differentiable, then

the value of $k+m$ is :

A. 2

B. $\frac{16}{5}$

C. $\frac{10}{3}$

D. 4

Answer: A



Watch Video Solution

9. For $x \in \mathbb{R}$, $f(x) = |\log 2 - \sin x|$ and $g(x) = f(f(x))$, then

A. g is not differentiable at $x=0$

B. $g'(0) = \cos \log 2$

C. $g'(0) = -\cos(\log 2)$

D. g is differentiable at $x = 0$ and $g'(0) = -\sin(\log 2)$

Answer: B

 [Watch Video Solution](#)

10. If for $x \left(0, \frac{1}{4}\right)$, the derivative of $\tan^{-1}\left(\frac{6x\sqrt{x}}{1-9x^3}\right)$ is $\sqrt{x}g(x)$, then $g(x)$ equals: (1) $\frac{3x}{1-9x^3}$ (2) $\frac{3}{1+9x^3}$ (3) $\frac{9}{1+9x^3}$ (4) $\frac{3x\sqrt{x}}{1-9x^3}$

A. $\frac{3x}{1-9x^3}$

B. $\frac{3}{1+9x^3}$

C. $\frac{9}{1+9x^3}$

D. $\frac{3x\sqrt{x}}{1-9x^3}$

Answer: C

 [Watch Video Solution](#)

11. Let $S = \{t \in R: f(x) = |x - \pi|(e^{|x|} - 1)\sin|x|$ is not differentiable at $t\}$ Then the set S is equal to: (1) ϕ (2) $\{0\}$ (3) $\{\pi\}$ (4) $\{0, \pi\}$

A. ϕ (empty set)

B. $\{1\}$

C. $\{\pi\}$

D. $\{0, \pi\}$

Answer: A



Watch Video Solution

12. Let $y = y(x)$ be the solution of the differential equation, $x \left(\frac{dy}{dx} \right) + y = x \log_e x$, ($x > 1$) if $2y(2) = \log_e 4 - 1$, then $y(e)$ is equal to: (a) $-\left(\frac{e}{2}\right)$ (b) $-\left(\frac{e^2}{2}\right)$ (c) $\frac{e}{4}$ (d) $\frac{e^2}{4}$

A. $\frac{e^2}{4}$

B. $-\frac{e}{2}$

C. $-\frac{e^2}{2}$

D. $\frac{e}{4}$

Answer: D



Watch Video Solution

13. for $x > 1$ if $(2x)^{2y} = 4e^{2x-2y}$ then $(1 + \log_e 2x)^2 \frac{dy}{dx}$

A. $\log_e 2x$

B. $x \log_e 2x$

C. $\frac{x \log_e 2x + \log_e 2}{x}$

D. $\frac{x \log_e 2x - \log_e 2}{x}$

Answer: D



Watch Video Solution

14. let S be the set of all points in $(-\pi, \pi)$ at which the function $f(x) = \min \{\sin x, \cos x\}$ is not differentiable Then S is a subset of which of the following ?

A. $\left\{ -\frac{3\pi}{4}, -\frac{\pi}{4}, \frac{3\pi}{4}, \frac{\pi}{4} \right\}$

B. 1

C. 2

D. 3

Answer: A



Watch Video Solution

15. let $f(x) = \begin{cases} \frac{\sqrt{2} \cos x - 1}{\cot x - 1} & x \neq \frac{\pi}{4} \\ k & x = \frac{\pi}{4} \end{cases}$

Find k for which f(x) is continuous

A. $-\frac{1}{2}$

B. $\frac{1}{3}$

C. $\frac{1}{2}$

D. 1

Answer: C



Watch Video Solution

16. Let $f(x) = 15 - |x - 10|$ and $g(x) = f(f(x))$ then $g(x)$ is non differentiable at

A. $\{5,10,15\}$

B. $\{5,10,15,20\}$

C. $\{10\}$

D. $\{5,15\}$

Answer: A



Watch Video Solution

1. A function $f: R \rightarrow R$ is defined as follows : $f(x) = \begin{cases} x, & \text{if } x \leq 1 \\ 5, & \text{if } x > 1 \end{cases}$

Which one of the following is true ?

- A. f is continuous at 0 and 1
- B. f is continuous at 1 and 2
- C. f is continuous at 0 and 2
- D. f is continuous at 0,1 and 2

Answer: C

 [Watch Video Solution](#)

2. if $\sqrt{x} + \sqrt{y} = 4$, then $\frac{dy}{dx}$ is :

- A. $-\sqrt{\frac{x}{y}}$
- B. $-\sqrt{\frac{y}{x}}$

C. $\sqrt{\frac{x}{y}}$

D. $\sqrt{\frac{y}{x}}$

Answer: B



Watch Video Solution

3. If $f(x)=x+7$, and $g(x)=x-7$, $x \in \mathbb{R}$, then $\frac{d}{dx} (g \circ f)(x) = \underline{\hspace{2cm}}$



Watch Video Solution

4. Find $\frac{dy}{dx}$ if $\sin^2 x + \cos^2 y = 1$



Watch Video Solution

5. Find $\frac{dy}{dx}$ when $x=4t$, $y=\frac{4}{t}$



Watch Video Solution

6. The function $f(x)$ is defined as follows :

$$f(x) = \begin{cases} x^2 + ax + b & , 0 \leq x < 2 \\ 3x + 2 & , 2 \leq x \leq 4 . \\ 2ax + 5b & , 4 < x \leq 8 \end{cases}$$

If $f(x)$ is continuous on $[0,8]$, find the values of 'a' and 'b'.



[Watch Video Solution](#)

7. If $y = \tan^{-1} \left(\frac{\sqrt{1+x^2} + \sqrt{1-x^2}}{\sqrt{1+x^2} - \sqrt{1-x^2}} \right)$, $x^2 \leq 1$, then find $\frac{dy}{dx}$



[Watch Video Solution](#)

8. If $x^y = e^{x-y}$, then show that $\frac{dy}{dx} = \frac{\log x}{(1 + \log x)^2}$



[Watch Video Solution](#)

9. If $y = e^{ax} \cos bx$, Show that $\frac{d^2y}{dx^2} - 2a \frac{dy}{dx} + (a^2 + b^2)y = 0$



[Watch Video Solution](#)

10. Verify Rolle's Theorem for the function : $f(x) = \sin x + \cos x$ in the interval $[0, 2\pi]$

 [Watch Video Solution](#)

11. Differentiate $\tan^{-1}\left(\frac{\sqrt{1+x^2} - \sqrt{1-x^2}}{\sqrt{1+x^2} + \sqrt{1-x^2}}\right)$ w. r. t. $\sin^{-1}\left(\frac{2x}{1+x^2}\right)$

 [Watch Video Solution](#)

12. Find $\frac{dy}{dx}$, if $y^x + x^y + x^x = a^b$.

 [Watch Video Solution](#)