



India's Number 1 Education App

MATHS

BOOKS - ARIHANT MATHS (HINGLISH)

DETERMINANTS

Examples

1. If A and B are square matrices of the same order 3, such that $|A| = 2$ and $AB = 2I$, write the value of $|B|$.



Watch Video Solution

2. Evaluate
$$\begin{vmatrix} \cos 15^\circ, \sin 15^\circ \\ \sin 75^\circ, \cos 75^\circ \end{vmatrix}$$

A. 0

B. 1

C. 2

D. 4

Answer: A



Watch Video Solution

3. If $\begin{vmatrix} 3x & 7 \\ -2 & 4 \end{vmatrix} = \begin{vmatrix} 8 & 7 \\ 6 & 4 \end{vmatrix}$, find the value of x.



Watch Video Solution

4. Expand $\begin{vmatrix} 3 & -5 & 4 \\ 7 & 6 & 1 \\ 1 & 2 & 3 \end{vmatrix}$

A. 180

B. 190

C. 18

Answer: A



Watch Video Solution

5. Evaluate :

$$\Delta = \begin{vmatrix} 0 & \sin \alpha & -\cos \alpha \\ -\sin \alpha & 0 & \sin \beta \\ \cos \alpha & -\sin \beta & 0 \end{vmatrix}.$$



Watch Video Solution

6. Find the product of the determinants :

$$\begin{vmatrix} 1 & 2 & 1 \\ -1 & 0 & 0 \\ 1 & 0 & 1 \end{vmatrix} \times \begin{vmatrix} 1 & -1 & 0 \\ 0 & 1 & 2 \\ 2 & -1 & 1 \end{vmatrix}$$



Watch Video Solution

7. If α , β and γ are real numbers without expanding at any stage prove that

$$\begin{vmatrix} 1 & \cos(\beta - \alpha) & \cos(\gamma - \alpha) \\ \cos(\alpha - \beta) & 1 & \cos(\gamma - \beta) \\ \cos(\alpha - \gamma) & \cos(\beta - \gamma) & 1 \end{vmatrix} = 0.$$



Watch Video Solution

8. Find the area of the triangle with vertices :

(3, 8), (-4, 2), (5, 1).



Watch Video Solution

9. By using determinants , find the value of 'y' for which the points

(1, 3), (2, 5) and (3, y) are collinear.



Watch Video Solution

10. A triangle has its three sides equal to a , b and c . If the co-ordinates of its vertices are $A(x_1, y_1)$, $B(x_2, y_2)$ and $C(x_3, y_3)$, show that :

$$\begin{vmatrix} x_1 & y_1 & 2 \\ x_2 & y_2 & 2 \\ x_3 & y_3 & 2 \end{vmatrix}^2 = (a + b + c)(b + c - a)(c + a - b)(a + b - c).$$



Watch Video Solution

11. If $\Delta = \begin{vmatrix} 5 & 3 & 8 \\ 2 & 0 & 1 \\ 1 & 2 & 3 \end{vmatrix}$, write :

(i) the minor of the element a_{23}

(ii) the co-factor of the element a_{32} .



Watch Video Solution

12. Find minors and cofactors of all the elements of the determinant
 $|1 - 243|$



Watch Video Solution

13. If $A = [a_{ij}]$ is a matrix of order 2×2 , such that $|A| = -15$ and c_{ij} represents co-factor of a_{ij} , then find $a_{21}c_{21} + a_{22}c_{22}$.

 Watch Video Solution

14. If A_{ij} is the cofactor of the element a_{ij} of the determinant $[2 - 3 - 7604157]$, then write the value of $a_{32} \cdot A_{32}$.

 Watch Video Solution

15. Find the minors and cofactors of the elements of the determinant

$$\Delta = \begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix}$$

 Watch Video Solution

16. For what values of k , does the system of linear equations $x + y + z = 2$, $2x + y - z = 3$, $3x + 2y + kz = 4$ have a unique

solution ?



Watch Video Solution

17. Using Cramer's Rule , solve the following system of equations :

$$3x - 2y = 5, x - 3y = -3$$



Watch Video Solution

18. Examine the consistency of the system of equations

$$3x + 5y + 2z = 2 \\ 2x + 2y + 3z = 1 \\ 3x - 5y = 3$$



Watch Video Solution

19. Solve

$$x + y + z = 1, 2x + 2y + 2z = 2, 3x + 3y + 3z = 4$$



Watch Video Solution

20. Using Cramer's Rule , solve the following system of equations :

$$3x - 4y + 5z = -6$$

$$x + y - 2z = -1$$

$$2x + 3y + z = 5$$



Watch Video Solution

21. Solve :

$$2x - y + z = 4, x + 3y + 2z = 12, 3x + 2y + 3z = 16$$



Watch Video Solution

22. Solve following system of homogeneous linear equations:

$$x + y - 2z = 0, \quad 2x + y - 3z = 0, \quad 5x + 4y - 9z = 0$$



Watch Video Solution

Frequently Asked Questions

1. The maximum value of $\begin{vmatrix} 1 & 1 & 1 \\ 1 & 1 + \sin \theta & 1 \\ 1 & 1 & 1 + \cos \theta \end{vmatrix}$ is $\frac{1}{2}$



Watch Video Solution

2. Using properties of determinants prove that $((1, 1, 1+3x), (1+3y, 1, 1), (1, 1+3z, 1)) = 9(3xyz + xy + yz + zx)$



Watch Video Solution

3. If $\Delta = \begin{vmatrix} 1 & a & a^2 \\ a & a^2 & 1 \\ a^2 & 1 & a \end{vmatrix} = -4$, then find the value of :

$$\begin{vmatrix} a^3 - 1 & 0 & a - a^4 \\ 0 & a - a^4 & a^3 - 1 \\ a - a^4 & a^3 - 1 & 0 \end{vmatrix}$$


Watch Video Solution

4. Prove that :

$$\begin{vmatrix} a+b & b+c & c+a \\ b+c & c+a & a+b \\ c+a & a+b & b+c \end{vmatrix} = 2 \begin{vmatrix} a & b & c \\ b & c & a \\ c & a & b \end{vmatrix}.$$



Watch Video Solution

5. $\begin{vmatrix} b^2c^2 & bc & b+c \\ c^2a^2 & ca & c+a \\ a^2b^2 & ab & a+b \end{vmatrix} = 0$



Watch Video Solution

6. Without expanding, prove that the following determinants vanish :

$$\begin{vmatrix} a & b & a+b \\ b & c & b+c \\ c & a & c+a \end{vmatrix}$$



Watch Video Solution

7. If $f(x) = |a - 10axa - 1ax^2axa|$, using properties of determinants, find the value of $f(2x) - f(x)$.



Watch Video Solution

8. Using properties of determinants , find the value of k if

$$\begin{vmatrix} x & y & x + y \\ y & x + y & x \\ x + y & x & y \end{vmatrix} = k(x^3 + y^3).$$



Watch Video Solution

9. Prove that :

$$\begin{vmatrix} a + b + 2c & a & b \\ c & b + c + 2a & b \\ c & a & c + a + 2b \end{vmatrix} = 2(a + b + c)^3$$



Watch Video Solution

10. If x, y, z are different and $\Delta = \begin{vmatrix} x & x^2 & 1+x^3 \\ y & y^2 & 1+y^3 \\ z & z^2 & 1+z^3 \end{vmatrix} = 0$, show that :

(i) $1+xyz = 0$

(ii) $xyz = -1$



Watch Video Solution

11. Prove that :

$$\begin{vmatrix} 1+a & 1 & 1 \\ 1 & 1+b & 1 \\ 1 & 1 & 1+c \end{vmatrix} = abc \left(1 + \frac{1}{a} + \frac{1}{b} + \frac{1}{c} \right) = abc + ab + bc + ca$$



Watch Video Solution

12. Show that $\begin{vmatrix} 1 & 1 & 1 \\ a^2 & b^2 & c^2 \\ a^3 & b^3 & c^3 \end{vmatrix} = (b-c)(c-a)(a-b)(bc+ca+ab)$



Watch Video Solution

13. Prove that

$$\begin{vmatrix} a^2 + 1 & ab & ac \\ ab & b^2 + 1 & bc \\ ac & bc & c^2 + 1 \end{vmatrix} = 1 + a^2 + b^2 + c^2.$$



Watch Video Solution

14.

Prove

that

$$|yz - x^2zx - y^2xy - z^2zx - y^2xy - z^2yz - x^2xy - z^2yz - x^2zx - y^2|$$

is divisible by $(x + y + z)$, and hence find the quotient.



Watch Video Solution

15. Prove that

$$\begin{vmatrix} -a^2 & ab & ac \\ ba & -b^2 & bc \\ ca & cb & -c^2 \end{vmatrix} = 4a^2b^2c^2$$



Watch Video Solution

$$16. \begin{vmatrix} x+2 & x+6 & x-1 \\ x+6 & x-1 & x+2 \\ x-1 & x+2 & x+6 \end{vmatrix} =$$



Watch Video Solution

17. Using properties of determinants, prove that :

$$\begin{vmatrix} (x+y)^2 & zx & xy \\ zx & (z+y)^2 & xy \\ zy & xy & (z+x)^2 \end{vmatrix} = 2xyz(x+y+z)^3.$$



Watch Video Solution

Examples Frequency Asked Questions

1. If A is square matrix of order 3 , with $|A| = 9$, then write the value of $|2adj. A|$



Watch Video Solution

2. For what value of x , the matrix $\begin{bmatrix} 5 & -x & +124 \end{bmatrix}$ is singular?



Watch Video Solution

3. If A and B are invertible matrices of order 3, $|A| = 2$ and $\left|(AB)^{-1}\right| = -\frac{1}{6}$, find $|B|$



Watch Video Solution

4. If $A = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 2 \\ 3 & 3 & 4 \end{bmatrix}$, find $\text{adj. } A$



Watch Video Solution

5. Given $A = \begin{pmatrix} 2 & -3 \\ -4 & 7 \end{pmatrix}$ compute A^{-1} and show that $2A^{-1} = 9I - A$



Watch Video Solution

6. For the matrix $A = [1\ 1\ 1\ 2 - 3\ 2 - 1\ 3]$. Show that $A^3 - 6A^2 + 5A + 11 I_3 = O$. Hence, find A^{-1} .

 Watch Video Solution

7. Find the inverse of the matrix :

$$A = \begin{bmatrix} 0 & 1 & 2 \\ 0 & 1 & 1 \\ 1 & 0 & 2 \end{bmatrix}$$

 Watch Video Solution

8. If $A = \begin{bmatrix} 1 & -2 & 3 \\ 0 & -1 & 4 \\ -2 & 2 & 1 \end{bmatrix}$, find $(A')^{-1}$

 Watch Video Solution

9. If $A = [2\ 3\ 1 - 4]$ and $B = [1 - 2 - 1\ 3]$, then verify that $(AB)^{-1} = B^{-1}A^{-1}$

 Watch Video Solution

10. If $A = \begin{bmatrix} 1 & 3 & 3 \\ 1 & 4 & 3 \\ 1 & 3 & 4 \end{bmatrix}$, verify $A \cdot (\text{adj. } A) = |A|I$ and find A^{-1} .

 Watch Video Solution

11. Find the inverse of each of the matrices given below :

Computer

$$(AB)^{-1} \text{ when } A = \begin{bmatrix} 1 & 1 & 2 \\ 0 & 2 & -3 \\ 3 & -2 & 4 \end{bmatrix} \text{ and } B^{-1} = \begin{bmatrix} 1 & 2 & 0 \\ 0 & 3 & -1 \\ 1 & 0 & 2 \end{bmatrix}.$$

Find A^{-1} . Then, $(AB)^{(-1)} = B^{(-1)}A^{(-1)}$.

 Watch Video Solution

12. Find A^{-1} if $A = \begin{vmatrix} 0 & 1 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 0 \end{vmatrix}$ and show that $A^{-1} = \frac{A^2 - 3I}{2}$

 Watch Video Solution

13. $x+2y=9$

$2x+4y=7$



Watch Video Solution

14. Solve the following system of equations by matrix method :

$3x - 4y = 5$ and $4x + 2y = 3$



Watch Video Solution

15. Using matrices, solve the following system of equations:

$4x + 3y + 3z = 60$, $x + 2y + 3z = 45$ and $6x + 2y + 3z = 70$



Watch Video Solution

16. Solve the following system of equations by matrix method :

$x + y - 2 = 3$ $2x + 3y + z = 10$ $3x - y - 7z = 1$ $x + y + z = 3$

$$2x - y + z = -1$$

$$2x + y - 3z = -9$$

$$6x - 12y + 25z = 4$$

$$4x + 15y - 20z = 3$$

$$2x + 18y + 15z = 10$$

$$3x + 4y + 7z = 14$$

$$2x - y + 3z = 4$$

$$x + 2y - 3z = 0$$

$$\frac{2}{x} - \frac{3}{y} + \frac{3}{z} = 10$$

$$\frac{1}{x} + \frac{1}{y} + \frac{1}{z} = 10$$

$$\frac{3}{x} - \frac{1}{y} + \frac{2}{z} = 13$$

$$5x + 3y + z = 16$$

$$2x + y + 3z = 19 \quad x + 2y + 4z = 25 \quad 3x + 4y + 2z = 8 \quad 2y - 3z = 3$$

$$x - 2y + 6z = -2 \quad 2x + y + z = 2 \quad x + 3y - z = 5 \quad 3x + y - 2z = 6$$

$$2x + 6y = 2 \quad 3x - z = -8 \quad 2x - y + z = -3 \quad 2y - z = 1$$

$$x - y + z = 2 \quad 2x - y = 0 \quad 8x + 4y + 3z = 18 \quad 2x + y + z = 5$$

$$x + 2y + z = 5 \quad x + y + z = 6 \quad x + 2z = 7 \quad 3x + y + z = 12$$

$$\frac{2}{x} + \frac{3}{y} + \frac{10}{z} = 4, \quad \frac{4}{x} - \frac{6}{y} + \frac{5}{z} = 1 \quad \frac{6}{x} + \frac{9}{y} - \frac{20}{z} = 2; \quad x, y, z \neq 0$$

$$x - y + 2z = 7 \quad 3x + 4y - 5z = -5 \quad 2x - y + 3z = 12$$



Watch Video Solution

17. Using elementary transformations, find the inverse of the matrix

$A = (843211122)$ and use it to solve the following system of linear

equations : $8x + 4y + 3z = 19$ $2x + y + z = 5$ $x + 2y + 2z = 7$



Watch Video Solution

18. If $A = \begin{bmatrix} 2 & 3 & 4 \\ 1 & -1 & 0 \\ 0 & 1 & 2 \end{bmatrix}$, find A^{-1} . Hence, solve the system of equations

$$x-y=3,$$

$$2x+3y+4z=17,$$

$$y+2z=7$$



Watch Video Solution

19. If $A = \begin{bmatrix} 2 & 3 & 10 \\ 4 & -6 & 5 \\ 6 & 9 & -20 \end{bmatrix}$, find A^{-1} . Using A^{-1} , solve the system of equations : $\frac{2}{x} + \frac{3}{y} + \frac{10}{z} = 2$

$$\frac{4}{x} - \frac{6}{y} + \frac{5}{z} = 5$$

$$\frac{6}{x} + \frac{9}{y} - \frac{20}{z} = -4$$



Watch Video Solution

20. Use product $\begin{bmatrix} 1 & -1 & 2 \\ 0 & 2 & -3 \\ 3 & -2 & 4 \end{bmatrix} \begin{bmatrix} -2 & 0 & 1 \\ 9 & 2 & -3 \\ 6 & 1 & -2 \end{bmatrix}$ to solve the equations :

$$x - 3z = 9, -x + 2y - 2z = 4, 2x - 3y + 4z = -3$$



Watch Video Solution

21. Solve, using matrices :

$$2x - y + 3z = 5, 3x + 2y - z = 7 \text{ and } 4x + 5y - 5z = 9$$



Watch Video Solution

22. Solve the following systems of linear homogenous equations :

$$2x + y - 3z = 0, x + 3y + z = 0 \text{ and } 3x - 2y + z = 0$$



Watch Video Solution

23. Solve the following systems of linear homogenous equations :

$$3x + 2y + 7z = 0, 4x - 3y - 2z = 0 \text{ and } 5x + 9y + 23z = 0$$



Watch Video Solution

24. The sum of three numbers is -1 . If we multiply second number by 2, third number by 3 and add them, we get 5. If we subtract the third number from the sum of first and second numbers, we get -1 . Represent it by a system of equation. Find the numbers, using inverse of a matrix.



Watch Video Solution

25. The monthly incomes of Aryan and Babban are in the ratio $3 : 4$ and their monthly expenditures are in the ratio $5 : 7$. If each saves Rs. 15000 per month, find their monthly incomes using matrix method. This problem reflects which value?



Watch Video Solution

26. A typing charges rs 145 for typing 10 English and 1 Hindi page for typing 3 English and 1 Hindi pages are Rs 180. Using matrices, find the charges of typing 1 English and 1 Hindi pages separately. However, typing charged only Rs 2 per page from a poor students Shyam for 5 Hindi

pages. How much less was charged from this poor boy? Which value are reflected in the problem ?



Watch Video Solution

27. Two schools A and B decided to award prizes to their students for three values honesty (x), punctuality (y) and obedience (z). School A decided to award a total of ₹11000 for the three values to 5, 4 and 3 students respectively while school B decided to award ₹10700 for the three values to 4, 3 and 5 students respectively. If all the three prizes together amount to ₹2700, then :

- (i) Represent the above situation by a matrix equation and form linear equations, using matrix multiplication.
- (ii) Is it possible to solve the system of equations so obtained, using matrix multiplication?



Watch Video Solution

28. The management committee of a residential colony decided to award some of its members (say x) for honesty, (say y) for helping others and some others (say z) for supervising the workers to keep the colony neat and clean. The sum of all the awardees is 12. Three times the sum of the awardees for cooperation and supervision added to two times the number of awardees for honesty is 33. If the sum of the number of awardees for honesty and supervision is twice the number of awardees for helping others. Using matrix method, find the number of awardees of each category.



[Watch Video Solution](#)

Examples Questions From Ncert Exemplar

1. If $\Delta = \begin{vmatrix} 1 & x & x^2 \\ 1 & y & y^2 \\ 1 & z & z^2 \end{vmatrix}$, $\Delta_1 = \begin{vmatrix} 1 & 1 & 1 \\ yz & zx & xy \\ x & y & z \end{vmatrix}$, then prove that $\Delta + \Delta_1 = 0$



[Watch Video Solution](#)

2. Without expanding, find the value of

$$\begin{vmatrix} \cos ec^2\theta & \cot^2\theta & 1 \\ \cot^2\theta & \cos ec^2\theta & -1 \\ 42 & 40 & 2 \end{vmatrix}$$



[Watch Video Solution](#)

3. If A,B and C are the angles of a triangle and

$$\begin{vmatrix} 1 & 1 & 1 \\ 1 + \sin A & 1 + \sin B & 1 + \sin C \\ \sin A + \sin^2 A & \sin B + \sin^2 B & \sin C + \sin^2 C \end{vmatrix} = 0$$

then prove that $\triangle ABC$ must be isoceles.



[Watch Video Solution](#)

4. Show that if the determinant :

$$\Delta = \begin{vmatrix} 3 & -2 & \sin 3\theta \\ -7 & 8 & \cos 2\theta \\ -11 & 14 & 2 \end{vmatrix} = \theta, \text{ then } \sin \theta = 0 \text{ or } \frac{1}{2}.$$



[Watch Video Solution](#)

1. Evaluate the determinants in questions 1 and 2 :

Find the values of x, if

$$(i) \begin{vmatrix} 2 & 4 \\ 5 & 1 \end{vmatrix} = \begin{vmatrix} 2x & 4 \\ 6 & x \end{vmatrix}$$

$$(ii) \begin{vmatrix} 2 & 3 \\ 4 & 5 \end{vmatrix} = \begin{vmatrix} x & 3 \\ 2x & 5 \end{vmatrix}$$



Watch Video Solution

2. Evaluate the determinants in questions 1 and 2 :

Find the values of x, if

$$(i) \begin{vmatrix} 2 & 4 \\ 5 & 1 \end{vmatrix} = \begin{vmatrix} 2x & 4 \\ 6 & x \end{vmatrix}$$

$$(ii) \begin{vmatrix} 2 & 3 \\ 4 & 5 \end{vmatrix} = \begin{vmatrix} x & 3 \\ 2x & 5 \end{vmatrix}$$



Watch Video Solution

3. Solve the equation: $\begin{vmatrix} 3 & x \\ x & 1 \end{vmatrix} = \begin{vmatrix} 3 & 2 \\ 4 & 1 \end{vmatrix}$



Watch Video Solution

4. Find the value of 'x' if :

$$\begin{vmatrix} 3 & x \\ x & 1 \end{vmatrix} = \begin{vmatrix} 5 & 2 \\ 4 & -1 \end{vmatrix}$$



Watch Video Solution

5. Find the value of 'x' if :

$$\begin{vmatrix} 2x & 3 \\ 5 & 2 \end{vmatrix} = \begin{vmatrix} 16 & 3 \\ 5 & 2 \end{vmatrix}, x > 0$$



Watch Video Solution

6. if $\begin{vmatrix} x+1 & x-1 \\ x-3 & x+2 \end{vmatrix} = \begin{vmatrix} 4 & -1 \\ 1 & 3 \end{vmatrix}$. find the value of x.



Watch Video Solution

7. Find the value of 'x' if :

$$\begin{vmatrix} -1 & 2 \\ 4 & 8 \end{vmatrix} = \begin{vmatrix} 2 & x \\ x & -4 \end{vmatrix}$$



Watch Video Solution

8. If $| \times 1x | = |3412|$, write the positive value of x .



Watch Video Solution

9. If $A = \begin{bmatrix} 1 & 3 \\ 4 & 1 \end{bmatrix}$, then find $|3A'|$.



Watch Video Solution

$$10. \begin{vmatrix} 3 & -1 & -2 \\ 0 & 0 & -1 \\ 3 & -5 & 0 \end{vmatrix}$$



Watch Video Solution

11. Write the minor and cofactor of each element of second column in the

following determinants and evaluate them:

$$\begin{vmatrix} 4 & 9 & 7 \\ 3 & 5 & 7 \\ 5 & 4 & 5 \end{vmatrix}$$



Watch Video Solution

12. Evaluate the determinants :

$$(i) \begin{vmatrix} 3 & -1 & -2 \\ 0 & 0 & 1 \\ 3 & -5 & 0 \end{vmatrix}$$

$$(ii) \begin{vmatrix} 3 & -4 & 5 \\ 1 & 1 & -2 \\ 2 & 3 & 1 \end{vmatrix}$$

$$(iii) \begin{vmatrix} 0 & 1 & 2 \\ -1 & 0 & -3 \\ -2 & 3 & 0 \end{vmatrix}$$

$$(iv) \begin{vmatrix} 2 & -1 & -2 \\ 0 & 2 & -1 \\ 3 & -5 & 0 \end{vmatrix}$$



Watch Video Solution

13. Evaluate the determinants :

$$(i) \begin{vmatrix} 3 & -1 & -2 \\ 0 & 0 & 1 \\ 3 & -5 & 0 \end{vmatrix}$$

$$(ii) \begin{vmatrix} 3 & -4 & 5 \\ 1 & 1 & -2 \\ 2 & 3 & 1 \end{vmatrix}$$

$$\begin{array}{l} \text{(iii)} \left| \begin{array}{ccc} 0 & 1 & 2 \\ -1 & 0 & -3 \\ -2 & 3 & 0 \end{array} \right| \\ \text{(iv)} \left| \begin{array}{ccc} 2 & -1 & -2 \\ 0 & 2 & -1 \\ 3 & -5 & 0 \end{array} \right| \end{array}$$



Watch Video Solution

14. Evaluate the following determinants :

$$\left| \begin{array}{ccc} 3 & -4 & 5 \\ 1 & 1 & -2 \\ 2 & 3 & 1 \end{array} \right|$$



Watch Video Solution

15. Evaluate the following determinants by two method :

$$\left| \begin{array}{ccc} 1 & 2 & 4 \\ -1 & 3 & 0 \\ 4 & 1 & 0 \end{array} \right|$$



Watch Video Solution

16. Evaluate the following determinants by two method :

$$\begin{vmatrix} 0 & 2 & 0 \\ 2 & 3 & 4 \\ 4 & 5 & 6 \end{vmatrix}$$



Watch Video Solution

17. Evaluate the determinants in questions 1 and 2 :

If $A = \begin{vmatrix} 1 & 1 & -2 \\ 2 & 1 & -3 \\ 5 & 4 & -9 \end{vmatrix}$, find $|A|$.



Watch Video Solution

18. If $A = \begin{bmatrix} 2 & 1 & 1 \\ 1 & 2 & 1 \\ 1 & 1 & 2 \end{bmatrix}$, then show that : $|4A| = 64|A|$.



Watch Video Solution

19. $\begin{vmatrix} 1! & 2! & 3! \\ 2! & 3! & 4! \\ 3! & 4! & 5! \end{vmatrix} = ?$



Watch Video Solution

20. If $\begin{vmatrix} x+1 & 1 & 1 \\ 1 & x+1 & 1 \\ -1 & 1 & x+1 \end{vmatrix} = 0$, find the value of 'x'.



Watch Video Solution

Exercise 4 B Short Answer Type Questions

1. Using the property of determinants and without expanding, prove that:

$$|a - - - ab - - aa - bc - aa - - c| = 0$$



Watch Video Solution

2.
$$\begin{vmatrix} 2 & a & abc \\ 2 & b & bca \\ 2 & c & cab \end{vmatrix}$$



[Watch Video Solution](#)

3. Use properties of determinants to evaluate :

$$\begin{vmatrix} x+y & y+z & z+x \\ z & x & y \\ 1 & 1 & 1 \end{vmatrix}$$



[Watch Video Solution](#)

4. Use properties of determinants to evaluate :

$$\begin{vmatrix} 102 & 18 & 36 \\ 1 & 3 & 4 \\ 17 & 3 & 6 \end{vmatrix}$$



[Watch Video Solution](#)

5. Use properties of determinants to evaluate :

$$\begin{vmatrix} 2 & 3 & 1 \\ 4 & 6 & 2 \\ 1 & 3 & 2 \end{vmatrix}$$



Watch Video Solution

6. Use properties of determinants to evaluate :

$$\begin{vmatrix} 2 & 3 & 5 \\ 261 & 127 & 388 \\ 20 & 30 & 50 \end{vmatrix}$$



Watch Video Solution

7. Show that $\begin{vmatrix} ma_1 & b_1 & nc_1 \\ ma_2 & b_2 & nc_2 \\ ma_3 & b_3 & nc_3 \end{vmatrix} = -mn \begin{vmatrix} c_1 & b_1 & a_1 \\ c_2 & b_2 & a_2 \\ c_3 & b_3 & a_3 \end{vmatrix}$



Watch Video Solution

8. Show that $x = 1$ is a root of the equation :
$$\begin{vmatrix} x + 1 & 2x & 11 \\ 2x & x + 1 & -4 \\ -3 & 4x - 7 & 6 \end{vmatrix} =$$



Watch Video Solution

9. If p, q, r are in A.P. Write the value of :

$$\begin{vmatrix} x + 1 & x + 2 & x + p \\ x + 2 & x + 3 & x + q \\ x + 3 & x + 4 & x + r \end{vmatrix}$$



Watch Video Solution

10. Show without expanding at any stage that:

$$[(a, b, c), [a + x, b + 2y, c + 2z], [x, y, z]] = 0$$



Watch Video Solution

11. Without expanding the determinant, show that the determinant

$$\begin{vmatrix} a^2 + 10 & ab & ac \\ ab & b^2 + 10 & bc \\ ac & bc & c^2 + 10 \end{vmatrix}$$
 is divisible by 100



[Watch Video Solution](#)

Exercise 4 B Long Answer Type Questions I

1. Using the property of determinants and without expanding, prove that

$$\begin{vmatrix} 2 & 7 & 65 \\ 3 & 8 & 75 \\ 5 & 9 & 86 \end{vmatrix} = 0$$



[Watch Video Solution](#)

2. Without actual expansion, prove that the following determinants vanish :

$$\begin{vmatrix} a & b & c \\ a - b & b - c & c - a \\ b & c & a \end{vmatrix}$$



[Watch Video Solution](#)



Watch Video Solution

3. Using the property of determinants and without expanding in questions 1 to 7 prove that ,

$$\begin{vmatrix} x & a & x+a \\ y & b & y+b \\ z & c & z+c \end{vmatrix} = 0$$



Watch Video Solution

4. Prove that $\begin{vmatrix} 1 & a & a^2 - bc \\ 1 & b & b^2 - ca \\ 1 & c & c^2 - ab \end{vmatrix} = 0$



Watch Video Solution

5. without expanding at any stage Prove that

$$\begin{vmatrix} 0 & a & -b \\ -a & 0 & -c \\ b & c & 0 \end{vmatrix} = 0$$



Watch Video Solution

6. Without actual expansion, prove that the following determinants vanish :

$$\begin{vmatrix} x+y & y+z & z+x \\ z & x & y \\ 1 & 1 & 1 \end{vmatrix}$$



[Watch Video Solution](#)

7. The value of the determinant $\begin{vmatrix} 1 & bc & a(b+c) \\ 1 & ca & b(a+c) \\ 1 & ab & c(a+b) \end{vmatrix}$ doesn't depend on



[Watch Video Solution](#)

8. Without expanding the determinants show that :

$$\left(\frac{1}{a} + \frac{1}{b} + \frac{1}{c} + 1 \right) \text{ is a factor of } \begin{vmatrix} 1+a & 1 & 1 \\ 1 & 1+b & 1 \\ 1 & 1 & 1+c \end{vmatrix}$$



[Watch Video Solution](#)

9. Prove that :
$$\begin{vmatrix} x - y - z & 2x & 2x \\ 2y & y - z - x & 2y \\ 2z & 2z & z - x - y \end{vmatrix}$$



[Watch Video Solution](#)

10. Without expanding the determinant, prove that

$$|aa^2bc \ ^\wedge 2ca \ ^\wedge 2ab| = |1a^2a^31b^2b^31c^2c^3|$$



[Watch Video Solution](#)

11. Show without expanding at any stage that:

$$[0, \sin \alpha - \cos \alpha], [-\sin \alpha, 0, \sin \beta], [\cos \alpha s - \sin \beta, 0] \mid = 0$$



[Watch Video Solution](#)

12. If
$$\begin{vmatrix} x^2 + x & x + 1 & x - 2 \\ 2x^2 + 3x - 1 & 3x & 3x - 3 \\ x^2 + 2x + 3 & 2x - 1 & 2x - 1 \end{vmatrix} = xA + B$$
 then



[Watch Video Solution](#)

13. Evaluate the following :

$$\begin{vmatrix} b+c & a & a \\ b & c+a & b \\ c & c & a+b \end{vmatrix}$$



Watch Video Solution

14. Evaluate the following :

$$\begin{vmatrix} x+\lambda & x & x \\ x & x+\lambda & x \\ x & x & x+\lambda \end{vmatrix}$$



Watch Video Solution

15. Evaluate the following:

$$\begin{vmatrix} 1 & a & b+c \\ 1 & b & c+a \\ 1 & c & a+b \end{vmatrix}$$



Watch Video Solution

16. Evaluate:
$$\begin{bmatrix} x & y & x+y \\ y & x+y & x \\ x+y & x & y \end{bmatrix}$$



Watch Video Solution

17. Evaluate the following:
$$\begin{vmatrix} 1 & x & y \\ 1 & x+y & y \\ 1 & x & x+y \end{vmatrix}$$



Watch Video Solution

18. Prove that:
$$\begin{vmatrix} b+c & c = a & a+b \\ q+r & r+p & p+q \\ y+z & z+x & x+y \end{vmatrix} = 2 \begin{vmatrix} a & b & c \\ p & q & r \\ x & y & z \end{vmatrix}$$



Watch Video Solution

19. Use the properties of determinant and without expanding prove that

$$\begin{vmatrix} b+c & q+r & y+z \\ c+a & r+p & z+x \\ a+b & p+q & x+y \end{vmatrix}.$$



Watch Video Solution



Watch Video Solution

Exercise 4 B Long Answer Type Questions II

1. By using properties of determinants, prove the following:

$$|x + 4 \begin{vmatrix} x & 2 \\ 2 & x \end{vmatrix} + 4 \begin{vmatrix} 2 & x \\ x & 2 \end{vmatrix} + 4| = (5x + 4)(4 - x)^2$$



Watch Video Solution

2. Prove: $|x + 4 \begin{vmatrix} x & x \\ x & x \end{vmatrix} + 4 \begin{vmatrix} x & x \\ x & x \end{vmatrix} + 4| = 16(3x + 4)$



Watch Video Solution

3. Prove, using properties of determinants:

$$|y + k \begin{vmatrix} y & y \\ y & y \end{vmatrix} + k \begin{vmatrix} y & y \\ y & y \end{vmatrix} + k| = k^2(3y + k)$$



Watch Video Solution

4. for $x, y, z > 0$ Prove that $\begin{vmatrix} 1 & \log_x y & \log_x z \\ \log_y x & 1 & \log_y z \\ \log_z x & \log_z y & 1 \end{vmatrix} = 0$



Watch Video Solution

5. Prove that $\begin{vmatrix} 1 & a & a^2 \\ 1 & b & b^2 \\ 1 & c & c^2 \end{vmatrix} = (a - b)(b - c)(c - a)$



Watch Video Solution

6. Prove that

$$\begin{vmatrix} a & b - c & c + b \\ a + c & b & c - a \\ a - b & b + a & c \end{vmatrix} = (a + b + c)(a^2 + b^2 + c^2).$$



Watch Video Solution

7. Prove the following :

$$\begin{vmatrix} 1 & a & a \\ a & 1 & a \\ a & a & 1 \end{vmatrix} = (2a + 1)(1 - a)^2$$



8. Prove that :
$$\begin{vmatrix} 1 & x & x^3 \\ 1 & y & y^3 \\ 1 & z & z^3 \end{vmatrix}$$



9. Prove the following :

$$\begin{vmatrix} 1 & 1 & 1 \\ a & b & c \\ bc & ca & ab \end{vmatrix} = (a - b)(b - c)(c - a)$$



10.
$$\begin{vmatrix} 1 & a & bc \\ 1 & b & ca \\ 1 & c & ab \end{vmatrix} = (a - b)(b - c)(c - a)$$



11. Prove the following :

$$\begin{vmatrix} bc & a & 1 \\ ca & b & 1 \\ ab & c & 1 \end{vmatrix} = (a - b)(b - c)(a - c)$$



Watch Video Solution

12. Prove the following :

$$\begin{vmatrix} a & b & c \\ a^2 & b^2 & c^2 \\ bc & ca & ab \end{vmatrix} = \begin{vmatrix} a & a^2 & bc \\ b & b^2 & ca \\ c & c^2 & ab \end{vmatrix} = (ab + bc + ca)(a - b)(b - c)(c - a).$$



Watch Video Solution

13. Prove the following :

$$\begin{vmatrix} a^2 & a & b + c \\ b^2 & b & c + a \\ c^2 & c & a + b \end{vmatrix} = -(a + b + c)(a - b)(b - c)(c - a)$$



Watch Video Solution

14.

Prove

that

$$\begin{vmatrix} a & b & c \\ a^2 & b^2 & c^2 \\ b+c & c+a & a+b \end{vmatrix} = (a-b)(b-c)(c-a)(a+b+c)$$



Watch Video Solution

15. Prove the following :

$$\begin{vmatrix} x & x^2 & y+z \\ y & y^2 & z+x \\ z & z^2 & x+y \end{vmatrix} = (y-z)(z-x)(x-y)(x+y+z)$$



Watch Video Solution

16. Prove the following :

$$\begin{vmatrix} \alpha & \alpha^2 & \beta+\gamma \\ \beta & \beta^2 & \gamma+\alpha \\ \gamma & \gamma^2 & \alpha+\beta \end{vmatrix} = \begin{vmatrix} \alpha & \beta & \gamma \\ \alpha^2 & \beta^2 & \gamma^2 \\ \beta+\gamma & \gamma+\alpha & \alpha+\beta \end{vmatrix} = (\beta-\gamma)(\gamma-\alpha)(\alpha-\beta)(\alpha+\beta)$$



Watch Video Solution

$$17. \begin{vmatrix} a & b & c \\ a-b & b-c & c-a \\ b+c & c+a & a+b \end{vmatrix} = a^3 + b^3 + c^3 - 3abc$$



Watch Video Solution

$$18. \text{Evaluate the following: } \begin{vmatrix} a^2 + 2a & 2a + 1 & 1 \\ 2a + 1 & a + 2 & 1 \\ 3 & 3 & 1 \end{vmatrix}$$



Watch Video Solution

$$19. \text{Given : } a^2 + b^2 + c^2 = 0$$

Prove the following :

$$\begin{vmatrix} b^2 + c^2 & ab & ca \\ ab & c^2 + a^2 & bc \\ ca & bc & a^2 + b^2 \end{vmatrix} = 4a^2b^2c^2$$



Watch Video Solution

20.
$$\begin{vmatrix} 1 + a^2 - b^2 & 2ab & -2b \\ 2ab & 1 - a^2 + b^2 & 2a \\ 2b & -2a & 1 - a^2 - b^2 \end{vmatrix} = (1 + a^2 + b^2)^3$$



[Watch Video Solution](#)

21. Prove the following :

$$\begin{vmatrix} x & y & z \\ x^2 & y^2 & z^2 \\ x^3 & y^3 & z^3 \end{vmatrix} = \begin{vmatrix} x & x^2 & x^3 \\ y & y^2 & y^3 \\ z & z^2 & z^3 \end{vmatrix} = xyz(x - y)(y - z)(z - x)$$



[Watch Video Solution](#)

22.
$$\begin{bmatrix} x & x^2 & yz \\ y & y^2 & zx \\ z & z^2 & xy \end{bmatrix} = (x - y)(y - z)(z - x)(xy + yz + zx)$$



[Watch Video Solution](#)

23. Prove that :
$$\begin{vmatrix} x + y + 2z & x & y \\ z & y + z + 2x & y \\ z & x & x + a + 2y \end{vmatrix} = 2(x + y +)^3$$



Watch Video Solution

24. Prove that

$$\begin{vmatrix} b+c & c+a & a+b \\ c+a & a+b & b+c \\ a+b & b+c & c+a \end{vmatrix} = 2(a+b+c)(ab+bc+ca - a^2 - b^2 - c^2).$$



Watch Video Solution

25. Using properties of determinants, prove the following:

$$|\begin{array}{ccc} x & +yx+2y & x+2y \\ x+2y & x & +yx+2yx \end{array}| = 9y^2(x+y)$$



Watch Video Solution

$$26. \begin{vmatrix} b+c & a & a \\ b & c+a & b \\ c & c & a+b \end{vmatrix} = 4abc$$



Watch Video Solution

27. Prove the following :

$$\begin{vmatrix} 2ab & a^2 & b^2 \\ a^2 & b^2 & 2ab \\ b^2 & 2ab & a^2 \end{vmatrix} = - (a^3 + b^3)^2.$$



Watch Video Solution

28. Prove the following : $\begin{vmatrix} 1 & x & x^2 - yz \\ 1 & y & y^2 - zx \\ 1 & z & z^2 - xy \end{vmatrix} = 0.$



Watch Video Solution

29. For any scalar p prove that

$$= |x^2(1 + px^3) - y^2(1 + py^3) - z^2(1 + pz^3)| = (1 + pxyz)(x - y)(y - z)(z - x)$$

.



Watch Video Solution

$$30. \begin{vmatrix} x+y+z & -z & -y \\ -z & x+y+z & -x \\ -y & -x & x+y+z \end{vmatrix} = 2(x+y)(y+z)(z+x)$$



Watch Video Solution

$$31. \text{Prove: } |2yy - z - x2y2z2zz - x - yx - y - z2x2x| = (x + y + z)^3$$



Watch Video Solution

$$32. \begin{vmatrix} a-b-c & 2a & 2a \\ 2b & b-c-a & 2b \\ 2c & 2c & c-a-b \end{vmatrix}$$



Watch Video Solution

33. Show that:

$$|3a - a + b - a + c - b + a3b - b + c - c + a - c + b3c| = 3(a + b + c)($$



Watch Video Solution

34. Using properties of determinants. Find the value of 'x'

$$\begin{vmatrix} 4-x & 4+x & 4+x \\ 4+x & 4-x & 4+x \\ 4-x & 4+x & 4+x \end{vmatrix} = 0$$



Watch Video Solution

35. Solve: $|a + xa - xa - xa - xa + xa - xa - xa - xa + x| = 0$



Watch Video Solution

36. Prove that $\begin{vmatrix} x & \sin \theta & \cos \theta \\ -\sin \theta & -x & 1 \\ \cos \theta & 1 & x \end{vmatrix}$ is independent of theta



Watch Video Solution

37. Prove that $\begin{vmatrix} 1+a & 1 & 1 \\ 1 & 1+b & 1 \\ 1 & 1 & 1+c \end{vmatrix}$
=abc $\left(1 + \frac{1}{a} + \frac{1}{b} + \frac{1}{c}\right)$ hence find the value of the determinant, if a,b,

and c are the roots of the equation $px^3 = qx^2 + rx - s = 0$.



Watch Video Solution

38. Prove that :
$$\begin{vmatrix} (y+z)^2 & x^2 & x^2 \\ y^2 & (x+z)^2 & y^2 \\ z^2 & z^2 & (x+y)^2 \end{vmatrix} = 2xyz(x+y+z)^3$$



Watch Video Solution

39. Prove that
$$\begin{vmatrix} (b+c)^2 & a^2 & a^2 \\ b^2 & (c+a)^2 & b^2 \\ c^2 & c^2 & (a+b)^2 \end{vmatrix} = 2abc(a+b+c)^3$$



Watch Video Solution

40.
$$\begin{vmatrix} (b+c)^2 & ab & ca \\ ab & (a+c)^2 & bc \\ ac & bc & (a+b)^2 \end{vmatrix} = 2abc(a+b+c)^3$$



Watch Video Solution

41.

Show

that

$$\Delta = \left| (y+z)^2xyz \times y(x+z)^2yzxzyz(x+y)^2 \right| = 2xyz(x+y+z)^3.$$



Watch Video Solution

42. If a, b, c are positive and unequal, show that value of the determinant

$$\Delta = \begin{vmatrix} a & b & c \\ b & c & a \\ c & a & b \end{vmatrix}$$
 is negative



Watch Video Solution

43. Using properties of determinants, prove that

$$|aa + ba + b + c2a3a + 2b4a + 3b + 2c3a6a + 3b10a + 6b + 3c| = a^3$$



Watch Video Solution

44. Using properties of determinants. Prove that

$$|11 + p1 + p + q23 + 2p4 + 3p + 2q36 + 3p10 + 6p + 3q| = 1$$



Watch Video Solution

45. Q.
$$\begin{vmatrix} x+y & x & x \\ 15x+4y & 4x & 2x \\ 10x+8y & 8x & 3x \end{vmatrix} = x^3$$



Watch Video Solution

46. Prove
$$\begin{vmatrix} -bc & b^2 + bc & c^2 + bc \\ a^2 + ac & -ac & c^2 + ac \\ a^2 + ab & b^2 + ab & -ab \end{vmatrix} = (ab + bc + ca)^2$$



Watch Video Solution

47. Prove that

$$\begin{vmatrix} a^2 & a^2 - (b - c)^2 & bc \\ b^2 & b^2 - (c - a)^2 & ca \\ c^2 & c^2 - (a - b)^2 & ab \end{vmatrix} = (a^2 + b^2 + c^2)(a - b)(b - c)(c - a)(a + b + c)$$



Watch Video Solution

48. If p, q, r are not in G.P. and

$$\begin{vmatrix} 1 & \frac{q}{p} & \alpha + \frac{q}{p} \\ 1 & \frac{r}{p} & \alpha + \frac{r}{q} \\ p\alpha + q & q\alpha + r & 0 \end{vmatrix} = 0 \text{ Show that}$$

$$p\alpha^2 + 2q\alpha + r = 0$$



Watch Video Solution

Exercise 4 C Short Answer Type Questions

1. The value of

$$\begin{vmatrix} a_1x_1 + b_1y_1 & a_1x_2 + b_1y_2 & a_1x_3 + b_1y_3 \\ a_2x_1 + b_2y_1 & a_2x_2 + b_2y_2 & a_2x_3 + b_2y_3 \\ a_3x_1 + b_3y_1 & a_3x_2 + b_3y_2 & a_3x_3 + b_3y_3 \end{vmatrix}, \text{ is}$$



Watch Video Solution

2. The value of

$$\begin{vmatrix} 2y_1z_1 & y_1z_2 + y_2z_1 & y_1z_3 + y_3z_1 \\ y_1z_2y_2z_1 & 2y_2z_2 & y_2z_3 + y_3z_2 \\ y_1z_3 + y_3z_1 & y_2z_3 + y_3z_2 & 2y_3z_3 \end{vmatrix}, \text{ is}$$



Watch Video Solution

3. If $\Delta = \begin{vmatrix} \cos(\alpha_1 - \beta_1) & \cos(\alpha_1 - \beta_2) & \cos(\alpha_1 - \beta_3) \\ \cos(\alpha_2 - \beta_1) & \cos(\alpha_2 - \beta_2) & \cos(\alpha_2 - \beta_3) \\ \cos(\alpha_3 - \beta_1) & \cos(\alpha_3 - \beta_2) & \cos(\alpha_3 - \beta_3) \end{vmatrix}$ then Δ equals

 Watch Video Solution

4. If $l_1, m_1, n_1, l_2, m_2, n_2$ and l_3, m_3, n_3 are direction cosines of three

mutually perpendicular lines then, the value of $\begin{vmatrix} l_1 & m_1 & n_1 \\ l_2 & m_2 & n_2 \\ l_3 & m_3 & n_3 \end{vmatrix}$ is

 Watch Video Solution

Exercise 4 D Short Answer Type Questions

1. Find area of the triangle with vertices at the point given in each of the following :

(i) (1,0), (6,0), (4,3)

(ii) (2,7), (1,1), (10,8)

(iii) (-2,-3), (3,2), (-1,-8)



Watch Video Solution

2. Using determinant, find the area of the triangle with vertices :

(1, -1), (2, 4), (-3, 5)



Watch Video Solution

3. Find the area of the triangle with vertices at the points given in each of the following . Are the following points collinear ?

(0, 0), (6, 0), (4, 3)



Watch Video Solution

4. Find area the triangle with at the point given in each of the following

(2,7),(1,1) (10,8)



Watch Video Solution

Exercise 4 D Long Answer Type Questions

1. Find the area of the triangle with vertices at the points given in each of the following . Are the following points collinear ?

(- 2, - 3), (3, 2), (- 1, - 8)



[Watch Video Solution](#)

2. Using determinants , show that the following points are collinear :

(11, 7), (5, 5) and (- 1, 3).



[Watch Video Solution](#)

3. Use determinants to show that the following points are collinear.

(i) A(2, 3), B(-1, -2) and C(5, 8)

(ii) A(3, 8), B(-4, 2) and C(10, 14)

(iii) P(-2, 5), Q(-6, -7) and R(-5, -4)



[Watch Video Solution](#)

4. If the points $(3, -2)$, $(x, 2)$ and $(8, 8)$ are collinear, find x using determinant.



Watch Video Solution

5. If the area of triangle is 35sq.units with vertices $(2, -6)$, $(5, 4)$ and $(k, 4)$, find the values of 'k'



Watch Video Solution

6. If area of ΔABC is 12 square units and vertices are $A(x, 2)$, $B(4, -1)$ and $C(-3, 7)$, then find the value of x .



Watch Video Solution

7. Find the values of 'k' if area of the triangle is 4 square units and vertices are $(2, 0)$, $(0, 4)$, $(0, k)$.



Watch Video Solution

8. Find the values of k if area of triangle is 4 sq. units and vertices are: $(k, 0)$, $(0, 2)$, $(4, 0)$



Watch Video Solution

9. Determine the value of 'x' for which the area of the triangle formed by joining the points $(x, 4)$, $(0, 8)$ and $(-1, 3)$ will be $4\frac{1}{2}$ square units.



Watch Video Solution

10. Find the equation of the line joining the following points, using determinants :

(1, 2) and (3, 6)



Watch Video Solution

11. (i) Find the equation of line joining (1,2) and (3,4) using determinants,
(ii) Find the equation of the line joining (3,1) and (9,3) using determinants.



Watch Video Solution

12. Find the equation of the line joining A(-1,3) and B (0,0) using determinants and find k if D(k, 0) is a point such that area of triangle ABD is 3sq units.



Watch Video Solution

13. Prove that the area of the triangle whose vertices are :

$$(at_1^2, 2at_1), (at_2^2, 2at_2), (at_3^2, 2at_3) \text{ is } a^2(t_1 - t_2)(t_2 - t_3)(t_3 - t_1).$$



Watch Video Solution

14. Show that points $A(a, b + c)$, $B(b, c + a)$, $C(c, a + b)$ are collinear.



Watch Video Solution

15. If the points (a_1, b_1) , $m(a_2, b_2)$ and $(a_1 - a_2, b_2 - b_1)$ are collinear, then prove that $\frac{a_1}{a_2} = \frac{b_1}{b_2}$



Watch Video Solution

16. If $P(x, y)$ is any point on the line joining the point $A(a, 0)$ and $B(0, b)$, then show that $\frac{x}{a} + \frac{y}{b} = 1$.



Watch Video Solution

17. An equilateral triangle has each side to a. If the coordinates of its vertices are (x_1, y_1) , (x_2, y_2) and (x_3, y_2) and (x_3, y_3) then the square

of the determinant $\begin{vmatrix} x_1 & y_1 & 1 \\ x_2 & y_2 & 1 \\ x_3 & y_3 & 1 \end{vmatrix}$ equals



Watch Video Solution

Exercise 4 E Short Answer Type Questions

1. Find all the co-factors of $\begin{vmatrix} 1 & 3 \\ -2 & 4 \end{vmatrix}$



Watch Video Solution

2. Write Minors and Cofactors of the elements of following determinants:

- (i) $|2 - 403|$ (ii) $|acbd|$



Watch Video Solution

3. Write Minors and Cofactors of the elements of following determinants:

- (i) $|2 - 403|$ (ii) $|acbd|$



Watch Video Solution

4. In the determinant $\begin{vmatrix} 5 & -1 & 4 \\ 2 & 3 & -5 \\ -1 & -2 & 6 \end{vmatrix}$, find the co-factors of the elements
– 5, 3 and 6



Watch Video Solution

5. Write the minor and cofactor of each element of the following determinants and also evaluate the determinant in each case:
- $$\begin{vmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{vmatrix}$$



Watch Video Solution

6. Find the minor and cor-factor of each element of the first column of the following determinants :

$$\begin{vmatrix} 0 & 2 & 6 \\ 1 & 5 & 0 \\ 3 & 7 & 1 \end{vmatrix}$$

Hence or otherwise evaluate them.



Watch Video Solution

7. Find the minor and co-factor of each element of the first column of the following determinants :

$$\begin{vmatrix} 1 & a & bc \\ 1 & b & ca \\ 1 & c & ab \end{vmatrix}$$

Hence or otherwise evaluate them.



Watch Video Solution

8. Write the minor and cofactor of each element of the following determinants and also evaluate the determinant in each case:

$$\begin{vmatrix} 5 & -10 \\ 0 & 3 \end{vmatrix}$$



Watch Video Solution

9. Write the minor and cofactor of each element of the following determinants and also evaluate the determinant in each case:

$$\begin{vmatrix} 1 & 3 & -2 \\ 4 & -5 & 6 \\ 3 & 5 & 2 \end{vmatrix}$$



Watch Video Solution

10. Write minors and cofactors of the elements of the following determinants:

$$\begin{vmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{vmatrix}$$

(ii) $\begin{vmatrix} 1 & 0 & 4 \\ 3 & 5 & -1 \\ 0 & 1 & 2 \end{vmatrix}$



Watch Video Solution

11. Using Cofactors of elements of third column, evaluate
 $\Delta = |1xyz\ 1yzx\ 1zxy|$



Watch Video Solution

12. Write down the co-factors of the elements of the first row of the following determinant and hence evaluate the determinant

$$\begin{vmatrix} 1 & 3 & -3 \\ 2 & -1 & 0 \\ 4 & -2 & 5 \end{vmatrix}$$



[Watch Video Solution](#)

13. Find minors and co-factors of the elements of the determinant :

$$\begin{vmatrix} 2 & -3 & 5 \\ 6 & 0 & 4 \\ 1 & 5 & -7 \end{vmatrix}$$

and verify that $a_{11}A_{31} + a_{12}A_{32} + a_{13}A_{33} = 0$



[Watch Video Solution](#)

Exercise 4 F Short Answer Type Questions

1. Solve the following system of linear equations by Cramers rule:

$$3x + ay = 4, \quad 2x + ay = 2, \quad a \neq 0$$



Watch Video Solution

2. With the help of determinants , solve the following systems of equations :

$$2x + 3y = 9, 3x - 2y = 7$$



Watch Video Solution

3. With the help of determinants , solve the following systems of equations :

$$2x - y = - 2, 3x + 4y + 8 = 0$$



Watch Video Solution

4. Solve the following system of equations by matrix method:

$$3x + y = 19, \quad 3x - y = 23$$



Watch Video Solution

5. Classify the following system of equations as consistent or inconsistent

. If consistent, solve them :

$$x + y = -1, 2x - 3y = 8$$



Watch Video Solution

6. Classify the following system of equations as consistent or inconsistent

. If consistent, solve them :

$$3x - 2y = 4, 6x - 4y = 10$$



Watch Video Solution

7. Classify the following system of equations as consistent or inconsistent

. If consistent, solve them :

$$x + 2y = 5, 3x + 6y = 15$$



Watch Video Solution

8. Use Cramer's rule to show that the following system of equations has infinite number of solutions

$$2x - 3y - z = 0, x + 3y - 2z = 0, x - 3y = 0$$



Watch Video Solution

9. Solve the following system of equations, using Cramer's rule :

$$x + y + 3z = 6, x - 3y - 3z = -4, 5x - 3y + 3z = 8$$



Watch Video Solution

10. Show that the following systems of linear equations are inconsistent:

$$3x + y = 5, \quad -6x - 2y = 9$$



Watch Video Solution

11. Show that the following systems of linear equations are inconsistent:

$$2x - y = 5, \quad 4x - 2y = 7$$



Watch Video Solution

12. Find a quadratic function defined by the equation :

$$f(x) = ax^2 + bx + c$$

if (i) $f(1) = 0$, $f(2) = -2$ and $f(3) = -6$

(ii) $f(0) = f(-1) = 0$ and $f(1) = 2$



Watch Video Solution

13. The system of equations

$\lambda x + y + z = 0$, $-x + \lambda y + z = 0$, $-x - y + \lambda z = 0$ will have a non-zero solution if real values of λ are given by



Watch Video Solution

Exercise 4 F Long Answer Type Questions

1. Solve the following system of equations by Cramer's Rule :

$$x + 3y = 4, y + 3z = 7, 4x + z = 6$$



Watch Video Solution

2. Solve the following system of equations by Cramer's Rule :

$$x + y + z = -1, x + 2y + 3z = -4, x + 3y + 4z = -6$$



Watch Video Solution

3. Solve the following system of equations by Cramer's Rule :

$$3x + y + z = 10, x + y + z = 0, 5x - 9y = 1.$$



Watch Video Solution

4. Solve the following system of equations by Cramer's Rule :

$$(b + c)(y + z) - zx = b - c$$

$$(c + a)(z + x) - by = c - a$$

$$(a + b)(x + y) - cz = a - b$$



[View Text Solution](#)

5. Classify the following system of equations as consistent or inconsistent. If consistent, then solve them :

$$x - y + 3z = 6, x + 3y - 3z = -4, 5x + 3y + 3z = 14$$



[Watch Video Solution](#)

6. Classify the following system of equations as consistent or inconsistent. If consistent, then solve them :

$$x + y + 4z = 6, 3x + 2y - 2z = 9, 5x + y + 2z = 13$$



[Watch Video Solution](#)

7. Classify the following system of equations as consistent or inconsistent. If consistent, then solve them :

$$2x + 5y - z = 9, 3x - 3y + 2z = 7, 2x - 4y + 3z = 1$$



[Watch Video Solution](#)



8. Solve the following system of homogenous linear equations :

$$2x + 3y + 4z = 0, x + y + z = 0, 2x - y + 3z = 0$$



9. Solve the following system of homogenous linear equations :

$$x + y - z = 0, x - 2y + z = 0, 3x + 6y - 5z = 0$$



10. The system of equations:

$$x + ky + 3z = 0, 3x + ky - 2z = 0, 2x + 3y - 4z = 0$$
 has non- trivial

solution. when $k =$



11. Find the cost of sugar and wheat per kg, if the cost of 7kg of sugar and 3kg of wheat is ₹240 and cost of 7kg of wheat and 3kg of sugar is ₹160.
(Use determinants)



Watch Video Solution

12. Represent the following problem by a system of equations :

"₹240 is the cost of 5kg sugar, 5kg of wheat and 2kg of rice. The cost of 4kg rice 2kg sugar and 5kg wheat is ₹190. The cost of 3kg wheat 2kg rice and 4kg sugar is ₹190". Use determinants to find the cost of each per kg.



Watch Video Solution

Exercise 4 G Short Answer Type Questions

1. Verify $A(\text{adj. } A) = (\text{adj. } A)A = |A|I$:

$$\begin{bmatrix} 2 & 3 \\ -4 & -6 \end{bmatrix}$$



[Watch Video Solution](#)



Watch Video Solution

2. Verify $A(\text{adj. } A) = (\text{adj. } A)A = |A|I$:

$$\begin{bmatrix} 1 & -1 & 2 \\ 3 & 0 & -2 \\ 1 & 0 & 3 \end{bmatrix}$$



Watch Video Solution

3. Verify that $A(\text{adj}A) = I$ when :

$$A = \begin{bmatrix} \cos \theta & -\sin \theta & 0 \\ \sin \theta & \cos \theta & 0 \\ 0 & 0 & 1 \end{bmatrix}$$



Watch Video Solution

4. Find the inverse of each of the following matrix :

$$\begin{bmatrix} -1 & 5 \\ -3 & 2 \end{bmatrix}$$



Watch Video Solution

5. Find the inverse of each of the following matrice :

$$\begin{bmatrix} 2 & -2 \\ 4 & 3 \end{bmatrix}$$



Watch Video Solution

6. If $A = \begin{bmatrix} 2 & -1 \\ -1 & 2 \end{bmatrix}$, verify $A^2 - 4A + 3I = 0$, where $I = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ and $O = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$. Hence find A^{-1} .



Watch Video Solution

7. If $A = \begin{bmatrix} 3 & 1 \\ -1 & 2 \end{bmatrix}$, show that $A^2 - 5A + 7I = O$. Hence, find A^{-1} .



Watch Video Solution

8. Consider the matrix $A = \begin{bmatrix} 2 & 3 \\ 4 & 5 \end{bmatrix}$.

Show that $A^2 - 7A - 2I = O$



Watch Video Solution

9. Consider the matrix $A = \begin{bmatrix} 2 & 3 \\ 4 & 5 \end{bmatrix}$.

Hence , find A^{-1} .



[Watch Video Solution](#)

10. If $A = \begin{bmatrix} 2 & 3 \\ 5 & -2 \end{bmatrix}$, write A^{-1} in terms of A.



[Watch Video Solution](#)

11. Verify $(AB)^{-1} = B^{-1}A^{-1}$ for the matrices A and B where

$$A = \begin{bmatrix} 2 & 3 \\ 1 & -4 \end{bmatrix} \text{ and } B = \begin{bmatrix} 1 & -2 \\ -1 & 3 \end{bmatrix}$$



[Watch Video Solution](#)

12. Verify $(AB)^{-1} = B^{-1}A^{-1}$ for the matrices A and B where

$$A = \begin{bmatrix} 3 & 7 \\ 2 & 5 \end{bmatrix} \text{ and } B = \begin{bmatrix} 6 & 8 \\ 7 & 9 \end{bmatrix}$$



Watch Video Solution

13. Verify $(AB)^{-1} = B^{-1}A^{-1}$ for the matrices A and B where

$$A = \begin{bmatrix} 3 & 2 \\ 7 & 5 \end{bmatrix} \text{ and } B = \begin{bmatrix} 6 & 7 \\ 8 & 9 \end{bmatrix}$$



Watch Video Solution

14. Verify $(AB)^{-1} = B^{-1}A^{-1}$ for the matrices A and B where

$$A = \begin{bmatrix} 4 & 1 \\ 6 & 5 \end{bmatrix} \text{ and } B = \begin{bmatrix} 2 & 5 \\ 1 & 6 \end{bmatrix}$$



Watch Video Solution

15. Show that the matrix $A = \begin{bmatrix} 2 & 3 \\ 1 & 2 \end{bmatrix}$ satisfies the equation $A^2 - 4A + I = O$ and hence, find A^{-1} .



Watch Video Solution

16. If $A = \begin{bmatrix} 2 & -1 \\ 1 & 3 \end{bmatrix}$, then show that $A^2 - 5A + 7I_2 = O$, hence find A^{-1} .



Watch Video Solution

17. If $A = [31 - 12]$, show that $A^2 - 5A + 7I = O$. Hence, find A^{-1} .



Watch Video Solution

18. For the matrix $A = \begin{bmatrix} 2 & 1 \\ 3 & 0 \end{bmatrix}$, find the numbers 'a' and 'b' such that $A^2 + aA + bI = O$. Hence, find A^{-1} .



Watch Video Solution

19. If $A = \begin{bmatrix} 2 & -3 \\ -4 & 7 \end{bmatrix}$, compute A^{-1} and show that $2A^{-1} + A - 9I = O$.



Watch Video Solution

Exercise 4 G Long Answer Type Questions

1. Find the inverse of the matrix $\begin{bmatrix} -3 & 2 \\ 5 & -3 \end{bmatrix}$ Hence, find the matrix P satisfying the matrix equation :

$$P \begin{bmatrix} -3 & 2 \\ 5 & -3 \end{bmatrix} = \begin{bmatrix} 1 & 2 \\ 2 & -1 \end{bmatrix}$$



[Watch Video Solution](#)

2. Find the inverse the matrix (if it exists)given in $[1 \ -1 \ 2 \ 0 \ 2 \ -3 \ 3 \ -2 \ 4]$



[Watch Video Solution](#)

3. Find the inverse of each of the following :

$$\begin{bmatrix} 1 & 2 & 3 \\ 0 & 2 & 4 \\ 0 & 0 & 5 \end{bmatrix}$$



[Watch Video Solution](#)

4. Find the inverse of each of the following :

$$\begin{bmatrix} 2 & 1 & 3 \\ 4 & -1 & 0 \\ -7 & 2 & 1 \end{bmatrix}$$



Watch Video Solution

5. Find the inverse of each of the following :

$$\begin{bmatrix} 1 & 0 & 0 \\ 3 & 3 & 0 \\ 5 & 2 & -1 \end{bmatrix}$$



Watch Video Solution

6. If $P = \begin{bmatrix} 1 & 2 & -2 \\ -1 & 3 & 0 \\ 0 & -2 & 1 \end{bmatrix}$, find P^{-1} . Verify that $PP^{-1} = I$



Watch Video Solution

7. Given $A = \begin{bmatrix} 5 & 0 & 4 \\ 2 & 3 & 2 \\ 1 & 2 & 1 \end{bmatrix}$, $B^{-1} = \begin{bmatrix} 1 & 2 & 3 \\ 1 & 4 & 3 \\ 1 & 3 & 4 \end{bmatrix}$. Compute $(AB)^{-1}$.



[Watch Video Solution](#)

8. If $A = \begin{bmatrix} 1 & 2 & 2 \\ 2 & 1 & 2 \\ 2 & 2 & 1 \end{bmatrix}$, prove that $A^2 - 4A - 5I = O$ and hence, obtain A^{-1} .



[Watch Video Solution](#)

9. If $A = \begin{bmatrix} 2 & 0 & 1 \\ 2 & 1 & 3 \\ 1 & -1 & 0 \end{bmatrix}$. find $A^2 - 5A + 6I$ and hence , find a matrix X such that $A^2 - 5A + 6I + X = O$.



[Watch Video Solution](#)

10. Find AB , if $A = \begin{bmatrix} 1 & 2 & 3 \\ 1 & -2 & 3 \end{bmatrix}$ and $B = \begin{bmatrix} 1 & -1 \\ 1 & 2 \\ 1 & -2 \end{bmatrix}$. Examine whether $(AB)^{-1}$ exist or not.



Watch Video Solution

11. Verify : $A(\text{adj. } A) = (\text{adj. } A)A = |A|I$ when :

$$A = \begin{bmatrix} 1 & -2 & 2 \\ 2 & 3 & 5 \\ -2 & 0 & 1 \end{bmatrix}$$



Watch Video Solution

12. Verify : $A(\text{adj. } A) = (\text{adj. } A)A = |A|I$ when :

$$A = \begin{bmatrix} 1 & -1 & 2 \\ 3 & 0 & -2 \\ 1 & 0 & 3 \end{bmatrix}$$



Watch Video Solution

13. Verify : $A(\text{adj. } A) = (\text{adj. } A)A = |A|I$ when :

$$A = \begin{bmatrix} 2 & 1 & 5 \\ 3 & -2 & -4 \\ -3 & 1 & -2 \end{bmatrix}$$



Watch Video Solution

14. Verify : $A(\text{adj. } A) = (\text{adj. } A)A = |A|I$ when :

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 0 & 5 & 0 \\ 2 & 4 & 3 \end{bmatrix}$$



Watch Video Solution

15. Find the inverse of the matrix $A = \begin{bmatrix} abc & 1+bc \\ a & a \end{bmatrix}$ and show that
 $aA^{-1} = (a^2 + bc + 1)I - aA$.



Watch Video Solution

Exercise 4 H Short Answer Type Questions

1. Classify the following system of equations as consistent or inconsistent

:

$$x + 2y = 2$$

$$2x + 3y = 3$$



Watch Video Solution

2. Classify the following system of equations as consistent or inconsistent

:

$$2x - y = 5$$

$$x + y = 4$$



Watch Video Solution

3. Classify the following system of equations as consistent or inconsistent

:

$$x + 3y = 5$$

$$2x + 6y = 8$$



Watch Video Solution

4. Solve the following equations using inverse of a matrix.

$$2x - y = -2$$

$$3x + 4y = 3$$



Watch Video Solution

5. Solve the following equations using inverse of a matrix.

$$5x + 2y = 3$$

$$3x + 2y = 5$$



Watch Video Solution

6. Solve the following equations using inverse of a matrix.

$$2x + 3y = 4$$

$$4x + 5y = 6$$



Watch Video Solution

7. Solve the following equations using inverse of a matrix.

$$2x + 5y = 1$$

$$3x + 2y = 7$$



Watch Video Solution

8. Solve system of linear equations, using matrix method, $5x + 2y = 4$

$$7x + 3y = 5$$



Watch Video Solution

9. Solve system of linear equations, using matrix method, $4x - 3y = 3$

$$3x - 5y = 7$$



Watch Video Solution

10. Solve the following equations using inverse of a matrix.

$$x + y = 5, y + z = 3, z + x = 4$$



Watch Video Solution

1. Classify the following system of equations as consistent or inconsistent

:

$$x + y + z = 1$$

$$2x + 3y + 2z = 2$$

$$ax + ay + 2az = 4$$



Watch Video Solution

2. Show that following system of linear equations is inconsistent:

$$3x - y - 2z = 2, \quad 2y - z = -1, \quad 3x - 5y = 3$$



Watch Video Solution

3. Examine the consistency of the system of equations in questions 1 to 6.

$$5x - y + 4z = 5, \quad 2x + 3y + 5z = 2, \quad 5x - 2y + 6z = -1$$



Watch Video Solution

4. Classify the following system of equations as consistent or inconsistent

:

$$5x - 6y + 4z = 15$$

$$7x + 4y - 3z = 19$$

$$2x + y + z = 46$$



Watch Video Solution

5. Solve the following system of equations by matrix method :

$$x + y - z = 3, 2x + 3y + z = 10, 3x - y - 7z = 1$$



Watch Video Solution

6. Using matrices, solve the following system of equations:

$$x - y + z = 4; 2x + y - 3z = 0; x + y + z = 2$$



Watch Video Solution

7. Solve the following system of equations, using matrix method.

$$x + 2y + z = 7, x + 3z = 11, 2x - 3y = 1$$



Watch Video Solution

8. Using matrices, solve the following system of equations for x,y and z.

$$x + 2y + z = 8$$

$$2x + y - z = 1$$

$$x - y + z = 2$$



Watch Video Solution

9. Using matrices, solve the following system of equations:

$$2x = 3y + 5z = 11 \quad 3x + 2y - 4z = -5$$



Watch Video Solution

10. Solve the following system of linear equations by matrix method:

$$x + y + z = 3, 2x - y + z = 2, x - 2y = 3z = 2$$



Watch Video Solution

11. Using matrices, solve the following system of equations for x,y and z.

$$x + y + z = 3, y + 3z = 4, x - 2y + z = 0$$



Watch Video Solution

12. Using matrices, solve the following system of equations for x,y and z.

$$x + y - z = -1, 3x + y - 2z = 3, x - y - z = -1$$



Watch Video Solution

13. Using matrices, solve the following system of equations for x,y and z.

$$2x + 3y + 3z = 5, x - 2y + z = -4, 3x - y - 2z = 3$$



Watch Video Solution

14. Using matrices, solve the following system of equations for x,y and z.

$$3x - 2y + 3z = 8, 2x + y - z = 1, 4x - 3y + 2z = 4$$



Watch Video Solution

15. Using matrices, solve the following system of equations for x,y and z.

$$2x + y + z = 1, x - 2y - z = \frac{3}{2}, 3y - 5z = 9.$$



Watch Video Solution

16. Solve system of linear equations, using matrix method, in questions 7

to 14.

$$x-y+2z=7, 3x+4y-5z=-5, 2x-y+3z=12$$



Watch Video Solution

17. Solve the following equations, using inverse of a matrix :

$$x - 2y + 3z = -5$$

$$3x + y + z = 8$$

$$2x - y + 2z = 1$$



Watch Video Solution

18. Solve the following equations, using inverse of a matrix :

$$x + 2y = 5$$

$$y + 2z = 8$$

$$z + 2x = 5$$



Watch Video Solution

19. Find inverse of matrix, solve the equation $x - y + z = 4$,

$$2x + y - 3z = 0, x + y + z = 2$$



Watch Video Solution

20. Solve the following equations, using inverse of a matrix :

$$x + 2y - 3z = -4$$

$$2x + 3y + 2z = 2$$

$$3x - 3y - 4z = 11$$



Watch Video Solution

21. $2x+3y+5z=16,$

$3x+ 2y-4z= 4,$

$x + y - 2z = -3.$



Watch Video Solution

22. Solve the following equations, using inverse of a matrix :

$$2x + y + z = 1$$

$$2x - 4y - 2z = 3$$

$$3y - 5z = 9$$



View Text Solution

23. Solve the following equations, using inverse of a matrix :

$$2x + 3y + 3z = 5$$

$$x - 2y + z = 4$$

$$3x - y - 2z = 3$$



Watch Video Solution

24. Solve the following equations, using inverse of a matrix :

$$3x + 4y + 7z = 4$$

$$2x - y + 3z = -3$$

$$x + 2y - 3z = 8$$



Watch Video Solution

25. Solve the following equations, using inverse of a matrix :

$$8x + 4y + 3z = 18$$

$$2x + y + z = 5$$

$$x + 2y + z = 5$$



Watch Video Solution

26. Solve the following equations, using inverse of a matrix :

$$x + y + z = 6$$

$$y + 3z = 11$$

$$x - 2y + z = 0$$



Watch Video Solution

27. Solve the following equations, using inverse of a matrix :

$$5x - y + z = 4$$

$$3x + 2y - 5z = 2$$

$$x + 3y - 2z = 5$$



Watch Video Solution

28. Solve the following equations, using inverse of a matrix :

$$3x - 2y + 3z = 8$$

$$2x + y - z = 1$$

$$4x - 3y + 2z = 4$$



Watch Video Solution

29. Solve the following equations, using inverse of a matrix :

$$3x - y + z = 5$$

$$2x - 2y + 3z = 7$$

$$x + y - z = -1$$



Watch Video Solution

30. Solve the following equations, using inverse of a matrix :

$$4x + 3y + z = 10$$

$$3x - y + 2z = 8$$

$$x - 2y - 3z = -10$$



Watch Video Solution

31. $\frac{2}{x} + \frac{3}{y} + \frac{10}{z} = 4, \frac{4}{x} - \frac{6}{y} + \frac{5}{z} = 1, \frac{6}{x} + \frac{9}{y} - \frac{20}{z} = 2$



Watch Video Solution

32. Solve the following system of equations :

$$\frac{1}{x} - \frac{1}{y} + \frac{2}{z} = 7$$

$$\frac{3}{x} + \frac{4}{y} - \frac{5}{z} = -5$$

$$\frac{2}{x} - \frac{1}{y} + \frac{3}{z} = 12$$



Watch Video Solution

33. Solve the following system of equations :

$$\left(\frac{1}{x} - \frac{1}{y} + \frac{2}{z} = 4, , \right), \left(\frac{2}{x} + \frac{1}{y} - \frac{3}{z} = 0, , \right), \left(\frac{1}{x} + \frac{1}{y} + \frac{1}{z} = 2, x, y, z \right)$$



Watch Video Solution

34. If $A = \begin{pmatrix} 2 & -3 & 5 \\ 3 & 2 & -4 \\ 1 & -2 & -2 \end{pmatrix}$, find A^{-1} . Using A^{-1} solve the following system of equations:

$$2x - 3y + 5z = 16; \quad 3x + 2y - 4z = -4; \quad x + y - 2z = -3$$



Watch Video Solution

35. If $A = \begin{pmatrix} 2 & -3 & 5 \\ 3 & 2 & -4 \\ 1 & 1 & -2 \end{pmatrix}$ find A^{-1} . Use it to solve the system of equations $2x - 3y + 5z = 11$, $3x + 2y - 4z = -5$ and $x + y - 2z = -3$



Watch Video Solution

36. If $A = \begin{bmatrix} 3 & 2 & 1 \\ 4 & -1 & 2 \\ 7 & 3 & -3 \end{bmatrix}$ then find A^{-1} and hence solve the following system of equations :

$$3x + 4y + 7z = 14, 2x - y + 3z = 4 \text{ and } x + 2y - 3z = 0.$$



Watch Video Solution

37. If $A = \begin{bmatrix} 3 & 1 & 2 \\ 3 & 2 & -3 \\ 2 & 0 & -1 \end{bmatrix}$, find A^{-1} . Hence, solve the system of equations :

$$3x + 3y + 2z = 1, x + 2y = 4, 2x - 3y - z = 5$$



Watch Video Solution

38. If $A = \begin{bmatrix} 1 & 1 & 1 \\ 1 & 0 & 2 \\ 3 & 1 & 1 \end{bmatrix}$, find A^{-1} . Hence, solve the system of equations :

$$x + y + z = 6, x + 2z = 7, 3x + y + z = 12$$



Watch Video Solution

39. If $A = \begin{pmatrix} 2 & -3 & 5 \\ 3 & 2 & -4 \\ 1 & 1 & -2 \end{pmatrix}$ find A^{-1} . Use it to solve the system of

equations $2x - 3y + 5z = 11$, $3x + 2y - 4z = -5$ and

$$x + y - 2z = -3$$



Watch Video Solution

40. Given that $A = \begin{bmatrix} 1 & -1 & 0 \\ 2 & 3 & 4 \\ 0 & 1 & 2 \end{bmatrix}$ and $B = \begin{bmatrix} 2 & 2 & -4 \\ -4 & 2 & -4 \\ 2 & -1 & 5 \end{bmatrix}$. Find AB .

Use this to solve that following system of equations :

$$x - y = 3, 2x + 3y + 4z = 17, y + 2z = 7.$$



Watch Video Solution

41. Given $A = \begin{bmatrix} 1 & -1 & 1 \\ 1 & -2 & -2 \\ 2 & 1 & 3 \end{bmatrix}$ and $B = \begin{bmatrix} -4 & 4 & 4 \\ -7 & 1 & 3 \\ 5 & -3 & -1 \end{bmatrix}$,

find AB and use this result in solving the following system of equation

$$x-y+z=4,$$

$$x-2y-2z=9$$

$$2x+2y+3z=1$$



Watch Video Solution

42. Use product $[1 - 1202 - 33 - 24] [- 20192 - 361 - 2]$ to solve the system of equation: $x - y + 2z = 1$ $2y - 3z = 1$ $3x - 2y + 4z = 2$



Watch Video Solution

43. Solve the following system of homogeneous equations:

$$2x + 3y - z = 0 \quad x - y - 2z = 0 \quad 3x + y + 3z = 0$$



Watch Video Solution

44. Solve the following system of homogeneous linear equations by matrix method: $3x + y - 2z = 0$, $x + y + z = 0$, $x - 2y + z = 0$



Watch Video Solution

45. Solve following system of homogeneous linear equations:
 $x + y - 2z = 0$, $2x + y - 3z = 0$, $5x + 4y - 9z = 0$



Watch Video Solution

46. Solve the following system of homogeneous equations:
 $x + y + z = 0$ $x - 2y + z = 0$ $3x + 6y - 5z = 0$



Watch Video Solution

47. A school wants to award its students for the value of honesty, regularity and hard work will total cash award of Rs. 6000. Three times the award money for hard work added to that added to that given for honesty amounts to Rs. 11000. The award money given for honesty and hard work together is double the one given for regularity. Represent the above situation algebraically and find the award money for each value, using matrix method.



[Watch Video Solution](#)

48. Two schools A and B want to award their selected students on the values of sincerity, truthfulness and helpfulness. The school A wants to award $\text{₹}x$ each, $\text{₹}y$ each and $\text{₹}z$ each for the three respective values to 3, 2 and 1 students respectively with a total award money of $\text{₹}1600$. School B wants to spend $\text{₹}2300$ to award its 4, 1 and 3 students on the respective values (by giving the same award money to the three values as before). If the total amount of award for one prize on each value is $\text{₹}900$, using matrices, find the award money for each value.



Watch Video Solution

49. Two schools P and Q want to award their selected students on the values of Discipline, politeness and punctuality. The school P wants to awards Rs. x each, Rs. y each and Rs. z each for the three respective values to its 3, 2 and 1 students with a total award money of Rs. 1000. School Q wants to spend Rs 1500 to award its 4, 1 and 3 students on the respective values (by giving the same award money for the three values before) If the total amount of awards for one prize on each value is Rs. 600, using matrices, find the award money for each value. Apart from the above three values suggest one more value for awards.



Watch Video Solution

50. The sum of three numbers is 6. If we multiply third number by 3 and add second number to it, we get 11. By adding first and third numbers, we get double of the second number. Represent it algebraically and find the numbers using matrix method.



Watch Video Solution

51. The sum of three numbers is 6. Twice the third number when added to the first number gives 7. On adding the sum of the second and the third numbers to thrice the first number, we get 12. Find the numbers by using matrix method.



Watch Video Solution

52. The cost of 4 kg onion, 3 kg wheat and 2 kg rice is Rs 60. The cost of 2 kg onion, 4 kg wheat and 6 kg rice is Rs 90. The cost of 6 kg onion 2 kg wheat and 3 kg rice is Rs 70. Find cost of each item per kg by matrix method.



Watch Video Solution

Objective Type Questions Multiple Choice Question

1. Evaluate the determinants in questions 1 and 2 :

If $\begin{vmatrix} x & 2 \\ 18 & x \end{vmatrix} = \begin{vmatrix} 6 & 2 \\ 18 & 6 \end{vmatrix}$, then x is equal to :

(a) 6

(b) ± 6

(c) -6

0

A. 6

B. ± 6

C. - 6

D. 6, 6

Answer: A



Watch Video Solution

2. Let A be a square matrix of order 3×3 , then $|kA|$ is equal to (A) $k|A|$ (B)

$k^2|A|$ (C) $K^3|A|$ (D) $3k|A|$

A. $k|A|$

B. $k^2|A|$

C. $k^3|A|$

D. $3k|A|$

Answer: C



Watch Video Solution

3. Which of the following is correct ?

A. Determinant is a square matrix

B. Determinant is a number associated to a matrix

C. Determinant is a number associated to a square matrix

D. None of these

Answer: C



Watch Video Solution

4. If area of triangle is 35 sq units with vertices $(2, -6)$, $(5, 4)$ and $(k, 4)$.

Then k is (A) 12 (B) -2 (C) 12, 2 (D) 12, -2

A. 12

B. -2

C. -12, -2

D. 12, -2

Answer: D



Watch Video Solution

5. If $A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$ and A_{ij} is co-factors of a_{ij} , then A is given by :

A. $a_{11}A_{31} + a_{12}A_{32} + a_{13}A_{33}$

B. $a_{11}A_{11} + a_{12}A_{21} + a_{13}A_{33}$

C. $a_{21}A_{11} + a_{22}A_{12} + a_{23}A_{13}$

D. $a_{11}A_{11} + a_{21}A_{21} + a_{31}A_{31}$

Answer: B



Watch Video Solution

6. Let A be a non-singular square matrix of order 3×3 . Then $|\text{adj } A|$ is equal to (a) $|A|$ (B) $|A|^2$ (C) $|A|^3$ (D) $3|A|$

A. $|A|$

B. $|A|^2$

C. $|A|^3$

D. $3|A|$

Answer: B



Watch Video Solution

7. If A is an invertible matrix of order 2, then $\det(A^{-1})$ is equal to (a) \det

(A) (B) $\frac{1}{\det(A)}$ (C) 1 (D) 0

A. $\det(A)$

B. $\frac{1}{\det(A)}$

C. 1

D. 0

Answer: A



Watch Video Solution

8. Choose the correct answer in questions 17 to 19:

If a, b, c are in A.P., then the determinant $\begin{bmatrix} x+2 & x+3 & x+2a \\ x+3 & x+4 & x+3b \\ x+4 & x+5 & x+2c \end{bmatrix}$ is :

A. 0

B. 1

C. x

D. $2x$

Answer: A



Watch Video Solution

9. Choose the correct answer

If x, y, z are nonzero real numbers then the inverse of matrix

$$A = \begin{bmatrix} x & 0 & 0 \\ 0 & y & 0 \\ 0 & 0 & z \end{bmatrix} \text{ is :}$$

A. $\begin{bmatrix} x^{-1} & 0 & 0 \\ 0 & y^{-1} & 0 \\ 0 & 0 & z^{-1} \end{bmatrix}$

B. $xyz \begin{bmatrix} x^{-1} & 0 & 0 \\ 0 & y^{-1} & 0 \\ 0 & 0 & z^{-1} \end{bmatrix}$

C. $\frac{1}{xyz} \begin{bmatrix} x^{-1} & 0 & 0 \\ 0 & y^{-1} & 0 \\ 0 & 0 & z^{-1} \end{bmatrix}$

D. $\frac{1}{xyz} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$

Answer: A



Watch Video Solution

10. Let $A = \begin{bmatrix} 1 & \sin \theta & 1 \\ -\sin \theta & 1 & \sin \theta \\ -1 & -\sin \theta & 1 \end{bmatrix}$, where $0 \leq \theta < 2\pi$ then (A)

$|A| = 0$ (B) $|A|\varepsilon[2, 4]$ (C) $|A|\varepsilon(0, \infty)$ (D) $|A|\varepsilon(2, \infty)$

A. $\text{Det}(A) = 0$

B. $\text{Det}(A)\varepsilon(2, \infty)$

C. $\text{Det}(A)\varepsilon(2, 4)$

D. $\text{Det}(A)\varepsilon[2, 4]$

Answer: D



Watch Video Solution

11. If $\begin{vmatrix} 2x & 5 \\ 8 & x \end{vmatrix} = \begin{vmatrix} 6 & -2 \\ 7 & 3 \end{vmatrix}$ then the value of x is

A. 3

B. ± 3

C. ± 6

D. 6

Answer: C



Watch Video Solution

12. Let $\Delta_1 = \begin{vmatrix} Ax & x^2 & 1 \\ By & y^2 & 1 \\ Cz & z^2 & 1 \end{vmatrix}$ and $\Delta_2 = \begin{vmatrix} A & B & C \\ x & y & z \\ yz & zx & xy \end{vmatrix}$, then :

A. $\Delta_1 = -\Delta$

B. $\Delta \neq \Delta_1$

C. $\Delta - \Delta_1 = 0$

D. None of these

Answer: C



13. If $x, y \in R$, then the determinant

$= |\cos x - \sin x| \sin x \cos x | \cos(x+y) - \sin(x+y)|$ lies in the interval $[-\sqrt{2}, \sqrt{2}]$ (b) $[-1, 1]$ (c) $[-\sqrt{2}, 1]$ (d) $[-1, -\sqrt{2}]$

A. $[-\sqrt{2}, \sqrt{2}]$

B. $[-1, 1]$

C. $[-\sqrt{2}, 1]$

D. $[-1, -\sqrt{2}]$

Answer: B



Watch Video Solution

14. If the area of a triangle with vertices $(-3, 0)$, $(3, 0)$ and $(0, k)$ is 9 sq. units. Then the value of k will be

A. 9

B. 3

C. -9

D. 6

Answer: B



Watch Video Solution

15. If A , B and C are angles of a triangle then the determinant

$$\begin{vmatrix} -1 & \cos C & \cos B \\ \cos C & -1 & \cos A \\ \cos B & \cos A & -1 \end{vmatrix}$$
 is equal to

A. 0

B. -1

C. 1

D. None of these

Answer: A



Watch Video Solution

16. If A is a square matrix of order 3×3 and $|A| = 5$, then $|adj. A|$ is

- A. 5
- B. 125
- C. 15
- D. 25

Answer: D



Watch Video Solution

17. The value of 'x' for which $\begin{vmatrix} 3 & x \\ x & 1 \end{vmatrix} = \begin{vmatrix} 3 & 2 \\ 4 & 1 \end{vmatrix}$ is

- A. $\pm 2\sqrt{3}$

B. $\pm 3\sqrt{3}$

C. $\pm 2\sqrt{2}$

D. None of these

Answer: C



Watch Video Solution

18. Evaluate the determinants in questions 1 and 2 :

If $\begin{vmatrix} x & 2 \\ 18 & x \end{vmatrix} = \begin{vmatrix} 6 & 2 \\ 18 & 6 \end{vmatrix}$, then x is equal to :

(a) 6

(b) ± 6

(c) -6

0

A. 6

B. ± 6

C. -6

D. 0

Answer: B



Watch Video Solution

19. If A is a matrix of order 3×3 and $|A| = 10$, then $|adj. A|$ is

A. 0

B. 10

C. 100

D. 1000

Answer: C



Watch Video Solution

20. If A is an invertible matrix of order 2, then $\det(A^{-1})$ is equal to (a) \det

(A) (B) $\frac{1}{\det(A)}$ (C) 1 (D) 0

A. $\det A$

B. $\frac{1}{\det(A)}$

C. 1

D. 0

Answer: A



Watch Video Solution

21. If $\begin{vmatrix} 2 & 3-x \\ 1 & 4 \end{vmatrix} = 0$, then value of 'x' is

A. 3

B. -3

C. 5

Answer: D



Watch Video Solution

22. If $\begin{vmatrix} x & 12 \\ 3 & x \end{vmatrix} = \begin{vmatrix} 6 & 18 \\ 2 & 6 \end{vmatrix}$, then value of 'x' is

A. ± 4

B. ± 6

C. ± 8

D. None of these

Answer: B



Watch Video Solution

23. Let A be a non-singular square matrix of order 3×3 . Then $|\text{adj } A|$ is equal to (a) $|A|$ (B) $|A|^2$ (C) $|A|^3$ (D) $3|A|$

A. $|A|^3$

B. $|A|$

C. $3|A|$

D. $|A|^2$

Answer: D



Watch Video Solution

24. If A and B are invertible matrices of the same order, then $(AB)'$ is equal to

A. $A'B'$

B. $B'A'$

C. $-A'B'$

D. $-B'A'$

Answer: B



Watch Video Solution

25. If determinant A is order 2×2 and $|A| = 3$, then the value of $|2A|$ is

A. 6

B. 12

C. - 6

D. None of these

Answer: B



Watch Video Solution

Objective Type Questions Fill In The Blanks

1. If $\begin{vmatrix} x & 2 \\ 8 & x \end{vmatrix} = \begin{vmatrix} 3 & 2 \\ 9 & 6 \end{vmatrix}$, then the value of 'x' is



Watch Video Solution

2. Let A be a 3×3 determinant and $|A| = 7$. Then the value of $|2A|$ is.....



Watch Video Solution

3. If $A = \begin{bmatrix} 1 & 2 \\ 4 & 2 \end{bmatrix}$, then the value of $k = \text{_____}$ if $|2A| = k|A|$.



Watch Video Solution

4. If A is a skew-symmetric matrix of order 3, then $\det A = \text{_____}$



Watch Video Solution

5. The value of $\begin{bmatrix} 102 & 18 & 36 \\ 1 & 3 & 4 \\ 17 & 3 & 6 \end{bmatrix}$ is _____



Watch Video Solution

6. If $\Delta = \begin{vmatrix} 1 & a \\ 1 & b \end{vmatrix}$, then minor of 'b' is _____



Watch Video Solution

7. Minor of 'd' in $\begin{vmatrix} a & c \\ b & d \end{vmatrix}$ is _____



Watch Video Solution

8. Co-factor of 'b' in $\begin{vmatrix} a & c \\ b & d \end{vmatrix}$ is _____



Watch Video Solution

9. If $\Delta = \begin{vmatrix} 1 & 2 & 3 \\ 2 & 0 & 1 \\ 5 & 3 & 8 \end{vmatrix}$, then minor of a_{22} =_____



Watch Video Solution

10. A square matrix A has inverse if and only if A is



Watch Video Solution

Objective Type Questions True Or False

1. If $\begin{bmatrix} x & 2 \\ 18 & x \end{bmatrix} = \begin{bmatrix} 6 & 2 \\ 18 & 6 \end{bmatrix}$, then 'x' is equal to ± 6 .



Watch Video Solution

2. If $A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$, then $|2A| = .$



Watch Video Solution

3. Let A be a square matrix of order 3×3 , then $|kA|$ is equal to (A) $k|A|$ (B) $k^2|A|$ (C) $K^3|A|$ (D) $3k|A|$

 Watch Video Solution

4. If any two rows (or columns) of a determinant are identical, the value of the determinant is zero.

 Watch Video Solution

5. If A; B are invertible matrices of the same order; then show that $(AB)^{-1} = B^{-1}A^{-1}$

 Watch Video Solution

Objective Type Questions Very Short Answer Type Question

1. Evaluate :
$$\begin{vmatrix} 2 & 4 \\ -1 & 2 \end{vmatrix}$$



[Watch Video Solution](#)

2. Evaluate :
$$\begin{vmatrix} -2 & 3 \\ 4 & -9 \end{vmatrix}$$



[Watch Video Solution](#)

3.
$$\begin{vmatrix} \cos 70^\circ & \sin 20^\circ \\ \sin 70^\circ & \cos 20^\circ \end{vmatrix} = ?$$



[Watch Video Solution](#)

4. Evaluate
$$\begin{vmatrix} \cos 15^\circ & \sin 15^\circ \\ \sin 75^\circ & \cos 75^\circ \end{vmatrix}$$



[Watch Video Solution](#)

5. Evaluate :
$$\begin{vmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{vmatrix}$$



Watch Video Solution

6. Evaluate :
$$\begin{vmatrix} x & x + 1 \\ x - 1 & x \end{vmatrix}$$



Watch Video Solution

7.
$$\begin{vmatrix} x^2 - x + 1 & x - 1 \\ x + 1 & x + 1 \end{vmatrix}$$



Watch Video Solution

8. If A is a 3×3 matrix, $|A| \neq 0$ and $|3A| = k|A|$, then write the value of k .



Watch Video Solution

9. Let A be a square matrix of order 3×3 . Write the value of $2A$, where $A = 4$.



Watch Video Solution

10. If A and B are square matrices of same order 3 , such that $|A| = 2$ and $AB = 2I$, write the value of $|B|$



Watch Video Solution

11. Find the co-factor of the element a_{23} of the determinant $\begin{vmatrix} 5 & 3 & 5 \\ 2 & 0 & 1 \\ 1 & 2 & 3 \end{vmatrix}$



Watch Video Solution

12. Write the minor of 6 in $\begin{vmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{vmatrix}$



Watch Video Solution

13. Write the co-factor of 7 in

$$\begin{vmatrix} 4 & 5 & 6 \\ 5 & 6 & 7 \\ 13 & 15 & 17 \end{vmatrix}$$


Watch Video Solution

14. For what value of x , is the following matrix singular ?

$$\begin{bmatrix} 3 - 2x & x + 1 \\ 2 & 4 \end{bmatrix}$$



Watch Video Solution

15. For what value of x , the matrix $[5 - x + 124]$ is singular?



Watch Video Solution

16. For what value of 'x' the matrix $\begin{bmatrix} 2 - x & 3 \\ -5 & -1 \end{bmatrix}$ is not invertible ?



Watch Video Solution

17. Find the adjoint of the following matrices :

$$\begin{bmatrix} 2 & -1 \\ 4 & 3 \end{bmatrix}$$



Watch Video Solution

18. Find the adjoint of each of the matrices in questions 1 and 2.

$$\begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$$



Watch Video Solution

19. Find the adjoint of the following matrices :

$$\begin{bmatrix} 2 & 3 \\ 1 & 4 \end{bmatrix}$$



Watch Video Solution

20. If $A = \begin{bmatrix} 3 & 1 \\ 2 & -3 \end{bmatrix}$, then find $|adj. A|$.



Watch Video Solution

21. If $A = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$ then $A^{-1} = ?$



[Watch Video Solution](#)

22. If A is a square matrix of order 3 with $|A| = 9$, then write the value of $|adj. A|$.



[Watch Video Solution](#)

23. If A is a square matrix with $|A| = 8$, then find the value of $|AA'|$.



[Watch Video Solution](#)

24. If A is a 3×3 invertible matrix, then what will be the value of k if $\det(A-1) = (\det A)k$.



[Watch Video Solution](#)

25. Write A^{-1} for $A = \begin{bmatrix} 2 & 5 \\ 1 & 3 \end{bmatrix}$.



Watch Video Solution

26. if for any 2×2 square matrix A , $A(adjA) = \begin{bmatrix} 8 & 0 \\ 0 & 8 \end{bmatrix}$ then write the value of $\det(A)$



Watch Video Solution

Ncert File Exercise 4 1

1. Evaluate the determinants in $|24 - 5 - 1|$



Watch Video Solution

2. Evaluate the determinants in $\begin{vmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{vmatrix}$



Watch Video Solution

3.
$$\begin{vmatrix} x^2 - x + 1 & x - 1 \\ x + 1 & x + 1 \end{vmatrix}$$



Watch Video Solution

4. Evaluate the determinants in questions 1 and 2 :

If $A = \begin{bmatrix} 1 & 2 \\ 4 & 2 \end{bmatrix}$, then show that $|2A|=4|A|$.



Watch Video Solution

5. If $A = [101012004]$, then show that $|3A| = 27|A|$



Watch Video Solution

6. Evaluate the determinants :

(i)
$$\begin{vmatrix} 3 & -1 & -2 \\ 0 & 0 & 1 \\ 3 & -5 & 0 \end{vmatrix}$$

(ii)
$$\begin{vmatrix} 3 & -4 & 5 \\ 1 & 1 & -2 \\ 2 & 3 & 1 \end{vmatrix}$$

(iii)
$$\begin{vmatrix} 0 & 1 & 2 \\ -1 & 0 & -3 \\ -2 & 3 & 0 \end{vmatrix}$$

(iv)
$$\begin{vmatrix} 2 & -1 & -2 \\ 0 & 2 & -1 \\ 3 & -5 & 0 \end{vmatrix}$$



Watch Video Solution

7. Evaluate the determinants :

(i)
$$\begin{vmatrix} 3 & -1 & -2 \\ 0 & 0 & 1 \\ 3 & -5 & 0 \end{vmatrix}$$

(ii)
$$\begin{vmatrix} 3 & -4 & 5 \\ 1 & 1 & -2 \\ 2 & 3 & 1 \end{vmatrix}$$

$$\begin{array}{l} \text{(iii)} \left| \begin{array}{ccc} 0 & 1 & 2 \\ -1 & 0 & -3 \\ -2 & 3 & 0 \end{array} \right| \\ \text{(iv)} \left| \begin{array}{ccc} 2 & -1 & -2 \\ 0 & 2 & -1 \\ 3 & -5 & 0 \end{array} \right| \end{array}$$



Watch Video Solution

8. Evaluate the determinants :

$$\begin{array}{l} \text{(i)} \left| \begin{array}{ccc} 3 & -1 & -2 \\ 0 & 0 & 1 \\ 3 & -5 & 0 \end{array} \right| \\ \text{(ii)} \left| \begin{array}{ccc} 3 & -4 & 5 \\ 1 & 1 & -2 \\ 2 & 3 & 1 \end{array} \right| \\ \text{(iii)} \left| \begin{array}{ccc} 0 & 1 & 2 \\ -1 & 0 & -3 \\ -2 & 3 & 0 \end{array} \right| \\ \text{(iv)} \left| \begin{array}{ccc} 2 & -1 & -2 \\ 0 & 2 & -1 \\ 3 & -5 & 0 \end{array} \right| \end{array}$$



Watch Video Solution

9. Evaluate the determinants :

(i)
$$\begin{vmatrix} 3 & -1 & -2 \\ 0 & 0 & 1 \\ 3 & -5 & 0 \end{vmatrix}$$

(ii)
$$\begin{vmatrix} 3 & -4 & 5 \\ 1 & 1 & -2 \\ 2 & 3 & 1 \end{vmatrix}$$

(iii)
$$\begin{vmatrix} 0 & 1 & 2 \\ -1 & 0 & -3 \\ -2 & 3 & 0 \end{vmatrix}$$

(iv)
$$\begin{vmatrix} 2 & -1 & -2 \\ 0 & 2 & -1 \\ 3 & -5 & 0 \end{vmatrix}$$



Watch Video Solution

10. Evaluate the determinants in questions 1 and 2 :

If $A = \begin{vmatrix} 1 & 1 & -2 \\ 2 & 1 & -3 \\ 5 & 4 & -9 \end{vmatrix}$, find $|A|$.



Watch Video Solution

11. Evaluate the determinants in questions 1 and 2 :

Find the values of x, if

$$(i) \begin{vmatrix} 2 & 4 \\ 5 & 1 \end{vmatrix} = \begin{vmatrix} 2x & 4 \\ 6 & x \end{vmatrix}$$
$$(ii) \begin{vmatrix} 2 & 3 \\ 4 & 5 \end{vmatrix} = \begin{vmatrix} x & 3 \\ 2x & 5 \end{vmatrix}$$



Watch Video Solution

12. Evaluate the determinants in questions 1 and 2 :

Find the values of x, if

$$(i) \begin{vmatrix} 2 & 4 \\ 5 & 1 \end{vmatrix} = \begin{vmatrix} 2x & 4 \\ 6 & x \end{vmatrix}$$
$$(ii) \begin{vmatrix} 2 & 3 \\ 4 & 5 \end{vmatrix} = \begin{vmatrix} x & 3 \\ 2x & 5 \end{vmatrix}$$



Watch Video Solution

13. Evaluate the determinants in questions 1 and 2 :

If $\begin{vmatrix} x & 2 \\ 18 & x \end{vmatrix} = \begin{vmatrix} 6 & 2 \\ 18 & 6 \end{vmatrix}$, then x is equal to :

(a) 6

(b) ± 6

(c) -6

0

A. 6

B. ± 6

C. -6

D. 0

Answer: B



Watch Video Solution

Ncert File Exercise 4 2

1. Using the property of determinants and without expanding, prove that:

$$|xax + ayby + bzc z + c| = 0$$



Watch Video Solution

2. Using the property of determinants and without expanding, prove that:

$$|a - ab - aa - bc - aa - c| = 0$$



Watch Video Solution

3. Using the property of determinants and without expanding, prove that

$$\begin{vmatrix} 2 & 7 & 65 \\ 3 & 8 & 75 \\ 5 & 9 & 86 \end{vmatrix} = 0$$



Watch Video Solution

4. Using the property of determinants and without expanding, prove that:

$$|1bca(b + c)1cab(c + a)1abx(a + b)| = 0$$



Watch Video Solution

5. Use the properties of determinant and without expanding prove that

$$\begin{vmatrix} b+c & q+r & y+z \\ c+a & r+p & z+x \\ a+b & p+q & x+y \end{vmatrix}.$$



Watch Video Solution

6. By using properties of determinants in

$$\begin{vmatrix} 0 & a & -b \\ -a & 0 & -c \\ b & c & 0 \end{vmatrix} = 0$$



Watch Video Solution

7. Using properties of determinants, prove that

$$|-a^2abacba - b^2bacb - c^2| = 4a^2b^2c^2$$



Watch Video Solution

8. Prove that

$$\begin{vmatrix} 1 & a & a^2 \\ 1 & b & b^2 \\ 1 & c & c^2 \end{vmatrix} = (a-b)(b-c)(c-a)$$



Watch Video Solution

9. Prove that:

$$(i) \begin{vmatrix} 1 & 1 & 1 \\ a & b & c \\ a^3 & b^3 & c^3 \end{vmatrix} = (a - b)(b - c)(c - a)(a + b + c)$$

$$(ii) \begin{vmatrix} a & b + c & a^2 \\ b & c + a & b^2 \\ c & a + b & c^2 \end{vmatrix} = -(a + b + c)(a - b)(b - c)(c - a)$$

$$(iii) \begin{vmatrix} b + c & a & a \\ b & c + a & b \\ c & c & a + b \end{vmatrix} = 4abc$$

$$(iv) \text{ If } \begin{vmatrix} 1 & a^2 & a^4 \\ 1 & b^2 & b^4 \\ 1 & c^2 & c^4 \end{vmatrix} = (a + b)(b + c)(c + a) = \begin{vmatrix} 1 & 1 & 1 \\ a & b & c \\ a^2 & b^2 & c^2 \end{vmatrix}$$



Watch Video Solution

$$10. \begin{bmatrix} x & x^2 & yz \\ y & y^2 & zx \\ z & z^2 & xy \end{bmatrix} = (x - y)(y - z)(z - x)(xy + yz + zx)$$



Watch Video Solution

11. By using properties of determinants. Show that: (i)

$$|x + 42x^2 - 2x^2 \times + 42x^2 - 2x^2 \times + 4| = (5x - 4)(4 - x)^2 \quad (\text{ii})$$

$$|y + kyyyy + kyyyy + k| = k^2 (2yk)^2$$



[Watch Video Solution](#)

12. Prove, using properties of determinants:

$$|y + kyyyy + kyyyy + k| = k^2(3y + k)$$



[Watch Video Solution](#)

13. Prove that: $\begin{vmatrix} a - b - c & 2a & 2a \\ 2b & b - c - a & 2b \\ 2c & 2c & c - a - b \end{vmatrix} = (a + b + c)^3$



[Watch Video Solution](#)

14.

Prove

that

$$\text{Det} \begin{bmatrix} x + y + 2z & x & y \\ z & y + z + 2x & y \\ z & x & z + x + 2y \end{bmatrix} = 2(x + y + z)^3$$



Watch Video Solution

15. By using properties of determinants. Show that:

$$|1 \times^2 x^2 1 \times x^2 1| = (1 - x^3)^2$$



Watch Video Solution

16. Show that

$$\begin{vmatrix} 1 + a^2 - b^2 & 2ab & -2b \\ 2ab & 1 - a^2 + b^2 & 2a \\ 2b & -2a & 1 - a^2 - b^2 \end{vmatrix} = (1 + a^2 + b^2)^3$$



Watch Video Solution

17. Using properties of determinants, prove the following:

$$|a^2 abaca \wedge 2 + 1 bacbc^2 + 1| = 1 + a^2 + b^2 + c^2.$$



Watch Video Solution

18. Let A be a square matrix of order 3×3 , then $|kA|$ is equal to (A) $k|A|$

(B) $k^2|A|$ (C) $K^3|A|$ (D) $3k|A|$

A. $k|A|$

B. $k^2|A|$

C. $k^3|A|$

D. $3k|A|$

Answer: C



Watch Video Solution

19. Which of the following is correct ?

A. Determinant is a square matrix

B. Determinant is a number associated to a matrix

C. Determinant is a number associated to a square matrix

D. None of these

Answer: C



Watch Video Solution

Ncert File Exercise 4 3

1. Find area of the triangle with vertices at the point given in each of the following :

(i) (1,0), (6,0), (4,3)

(ii) (2,7), (1,1), (10,8)

(iii) (-2,-3), (3,2), (-1,-8)



Watch Video Solution

2. Find area of the triangle with vertices at the point given in each of the following :

(i) $(1,0), (6,0), (4,3)$

(ii) $(2,7), (1,1), (10,8)$

(iii) $(-2,-3), (3,2), (-1,-8)$



Watch Video Solution

3. Find area of the triangle with vertices at the points given in each of the following

$(-2, -3), (3, 2), (-1, -8)$



Watch Video Solution

4. Show that points $A(a, b + c), B(b, c + a), C(c, a + b)$ are collinear.



Watch Video Solution

5. Find the values of k if area of triangle is 4 sq. units and vertices are :

(i) $(k,0), (4,0), (0,2)$

(ii) $(-2,0), (0,4), (0,k)$



Watch Video Solution

6. Find the values of k if area of triangle is 4 sq. units and vertices are:

$(-2,0), (0,4), (0,k)$



Watch Video Solution

7. (i) Find equation of line joining $(1,2)$ and $(3,6)$ using determinants, **(ii)**

Find equation of line joining $(3, 1)$ and $(9,3)$ using determinants.



Watch Video Solution

8. (i) Find equation of line joining $(1,2)$ and $(3,6)$ using determinants, **(ii)**

Find equation of line joining $(3, 1)$ and $(9,3)$ using determinants.



Watch Video Solution

9. If area of triangle is 35 sq units with vertices $(2, -6)$, $(5, 4)$ and $(k, 4)$.

Then k is

- A. 12
- B. -2
- C. $-12, -2$
- D. $12, -2$

Answer: D



Watch Video Solution

Ncert File Exercise 4 4

1. Write Minors and Cofactors of the elements of following determinants:

- (i) $|2 - 403|$ (ii) $|acbd|$



Watch Video Solution

2. Write Minors and Cofactors of the elements of following determinants:

- (i) $|2 - 403|$ (ii) $|acbd|$



Watch Video Solution

3. Write the minor and cofactor of each element of the following

determinants and also evaluate the determinant in each case:

$$\begin{vmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{vmatrix}$$



Watch Video Solution

4. Write minors and cofactors of the elements of the following

determinants:

$$\begin{vmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{vmatrix}$$

$$(ii) \begin{vmatrix} 1 & 0 & 4 \\ 3 & 5 & -1 \\ 0 & 1 & 2 \end{vmatrix}$$



Watch Video Solution

5. Using Cofactors of elements of second row, evaluate $\Delta = |538201123|$



Watch Video Solution

6. Using Cofactors of elements of third column, evaluate $\Delta = |1xyz1yzx1zxy|$



Watch Video Solution

7. If $\Delta = |a_{11}a_{12}a_{13}a_{21}a_{22}a_{23}a_{31}a_{32}a_{33}|$ and A_{ij} is cofactors of a_{ij} , then value of Δ is given by (A) $a_{11} + A_{31} + a_{12}A_{32} + a_{13}A_{33}$ (B) $a_{11}A_{11} + a_{12}A_{21} + a_{13}A_{31}$ (C) $a_{11}A_{11} + a_{12}A_{21} + a_{13}A_{31}$ (D) $a_{11} + A_{31} + a_{12}A_{32} + a_{13}A_{33}$

A. $a_{11}A_{31} + a_{12} + A_{32} + a_{13}A_{33}$

B. $a_{11}A_{11} + a_{12} + A_{21} + a_{13}A_{31}$

C. $a_{21}A_{11} + a_{22} + A_{12} + a_{23}A_{13}$

D. $a_{11}A_{11} + a_{21} + A_{21} + a_{31}A_{31}$

Answer: D



Watch Video Solution

Ncert File Exercise 4 5

1. Find the adjoint of each of the matrices in questions 1 and 2.

$$\begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$$



Watch Video Solution

2. Find the adjoint of the matrices

$$\begin{bmatrix} 1 & -1 & 2 \\ 2 & 3 & 5 \\ -2 & 0 & 1 \end{bmatrix}$$



Watch Video Solution

3. Verify $A (\text{adj}A) = (\text{adj}A)A = |A| I$ in questions 3 and 4.

$$\begin{bmatrix} 2 & 3 \\ -4 & -6 \end{bmatrix}$$



Watch Video Solution

4. Find the adjoint of the given matrix and verify in each case that

A. $(\text{adj}A) = (\text{adj}A) \cdot A = |A| \cdot I$.

$$\begin{bmatrix} 1 & -1 & 2 \\ 3 & 1 & -2 \\ 1 & 0 & 3 \end{bmatrix}$$



Watch Video Solution

5. Find the inverse the matrix (if it exists) given in $[2 - 243]$



Watch Video Solution

6. Find the inverse of each of the matrices (if it exists) given in $\begin{bmatrix} -1 & 5 \\ -3 & 2 \end{bmatrix}$



Watch Video Solution

7. Find the inverse the matrix (if it exists) given in $[123024005]$



Watch Video Solution

8. Find the inverse the matrix (if it exists) given in $[10033052 - 1]$



Watch Video Solution

9. Find the inverse the matrix (if it exists) given in $[2134 - 10 - 721]$



Watch Video Solution

10. Find the inverse the matrix (if it exists) given in $[1 - 1 | 202 - 33 - 24]$



Watch Video Solution

11. Find the inverse of each of the matrices (if it exists) given in

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & \cos \alpha & \sin \alpha \\ 0 & \sin \alpha & -\cos \alpha \end{bmatrix}$$



Watch Video Solution

12. Let $A = [37|25]$ and $B = [68|79]$. Verify that $(AB)^{-1} = B^{-1}A^{-1}$.



Watch Video Solution

13. If $A = [31 - 12]$, show that $A^2 - 5A + 7I = O$. Hence, find A^{-1} .



Watch Video Solution



Watch Video Solution

14. Solve system of linear equations, using matrix method,

$$xy + 2z = 7 \quad 3x + 4y + 5z = 5$$

$$2xy + 3z = 12$$



Watch Video Solution

15. For the matrix $A = \begin{vmatrix} 1 & 1 & 1 \\ 1 & 2 & -3 \\ 2 & -1 & 3 \end{vmatrix}$, then that

$$A^3 - 6A^2 + 5A + 11I = 0. \text{ find } A^{-1}$$



Watch Video Solution

16. If $A = [2 - 11 - 12 - 11 - 12]$. Verify that

$$A^3 - 6A^2 + 9A - 4I = O \text{ and hence find } A^{-1}.$$



Watch Video Solution

17. Let A be a non-singular square matrix of order 3×3 . Then $|\text{adj } A|$ is equal to

A. $|A|$

B. $|A|^2$

C. $|A|^3$

D. $3|A|$

Answer: B



Watch Video Solution

18. If A is an invertible matrix of order 2, then $\det(A^{-1})$ is equal to (a) \det

(A) (B) $\frac{1}{\det(A)}$ (C) 1 (D) 0

A. $\det(A)$

B. $\frac{1}{\det(A)}$

C. 1

D. 0

Answer: B



Watch Video Solution

Ncert File Exercise 4 6

1. Examine the consistency of the system of equations $x + 2y = 2$
 $2x + 3y = 3$



Watch Video Solution

2. Examine the consistency of the system of equations $2x - y = 5$
 $x + y = 4$



Watch Video Solution

3. Examine the consistency of the system of equations $x + 2y = 2$

$$2x + 3y = 3$$



Watch Video Solution

4. Examine the consistency of the system of equations

$$x + y + z = 1 \quad 2x + 3y + 2z = 2$$

$$ax + ay + 2a - z = 4$$



Watch Video Solution

5. Examine the consistency of the system of equations $3xy - 2z = 2$

$$3x - 5y = 3$$



Watch Video Solution

6. Examine the consistency of the system of equations in

$$5x - y - 4z = 5$$

$$2x + 3y + 5z = 2$$

$$5x - 2y + 6z = -1$$



Watch Video Solution

7. Solve system of linear equations, using matrix method, $5x + 2y = 4$

$$7x + 3y = 5$$



Watch Video Solution

8. Solve system of linear equations, using matrix method,

$$2. \begin{matrix} x & y & = & 2 \\ 3x & + & 4y & = & 3 \end{matrix}$$



Watch Video Solution

9. Solve system of linear equations, using matrix method, $4x - 3y = 3$

$$3x - 5y = 7$$



Watch Video Solution

10. Solve system of linear equations, using matrix method,
 $5x + 2y = 4$
 $7x + 3y = 5$



Watch Video Solution

11. Solve system of linear equations, using matrix method,
 $2x + y + z = 1$
 $x - 2y - z = \frac{3}{2}$
 $3y - 5z = 9$



Watch Video Solution

12. Solve system of linear equations, using matrix method,
- $$xy + z = 4 \qquad \qquad \qquad 2x + y - 3z = 0$$
- $$x + y + z = 2$$



Watch Video Solution

13. Solve system of linear equations, using matrix method,

$$2x + 3y + 3z = 5 \\ x - 2y + z = -4 \\ 3x - y - 2z = 3$$



[Watch Video Solution](#)

14. Solve system of linear equations, using matrix method, in

$$x - y + 2z = 7$$

$$3x + 4y - 5z = -5$$

$$2x - y + 3z = 12$$



[Watch Video Solution](#)

15. If $A = \begin{pmatrix} 2 & -3 & 5 \\ 3 & 2 & 3 \\ 1 & -4 & 2 \end{pmatrix}$, find A^{-1} . Using A^{-1} solve the

following system of equations:

$$2x - 3y + 5z = 16; \quad 3x + 2y - 4z = -4; \quad x + y - 2z = -3$$



[Watch Video Solution](#)

16. The cost of 4 kg onion, 3 kg wheat and 2 kg rice is Rs 60. The cost of 2 kg onion, 4 kg wheat and 6 kg rice is Rs 90. The cost of 6 kg onion 2 kg wheat and 3 kg rice is Rs 70. Find cost of each item per kg by matrix method.



[Watch Video Solution](#)

Miscellaneous Exercise On Chapter 4

1. Prove that the determinant $\begin{vmatrix} x & \sin \theta & \cos \theta \\ -\sin \theta & -x & 1 \\ \cos \theta & 1 & x \end{vmatrix}$ is independent of θ .



[Watch Video Solution](#)

2. Without expanding the determinant , prove that

$$\begin{vmatrix} a & a^2 & bc \\ b & b^2 & ca \\ c & c^2 & ab \end{vmatrix} = \begin{vmatrix} 1 & a^2 & a^3 \\ 1 & b^2 & b^3 \\ 1 & c^2 & c^3 \end{vmatrix}$$



[Watch Video Solution](#)

3. Evaluate

$$\begin{bmatrix} \cos \alpha \cos \beta & \cos \alpha \sin \beta & -\sin \alpha \\ -\sin \beta & \cos \beta & 0 \\ \sin \alpha \cos \beta & \sin \alpha \sin \beta & \cos \alpha \end{bmatrix}$$



Watch Video Solution

4. If a , b and c are real numbers, and

$\Delta = |b + aa + bc + aa + ca + + + a| = 0$. Show that either $a + b + c = 0$ or $a = b = c$.



Watch Video Solution

5. Solve the equation $|x + a \times \times + a \times \times + a| = 0, a \neq 0$



Watch Video Solution

6. Prove that $|a^2bcac + c^2a^2 + a^2aca^2 + b^2| = 4a^2b^2c^2$.



Watch Video Solution

7. If $A^{-1} = [3 - 11 - 156 - 55 - 22]$ and $B = [12 - 2 - 1300 - 21]$, find $(AB)^{-1}$.



Watch Video Solution

8. Let $A = \begin{bmatrix} 1 & -2 & 1 \\ -2 & 3 & 1 \\ 1 & 1 & 5 \end{bmatrix}$. Verify that
(i) $[adjA]^{-1} = adj(A^{-1})$
(ii) $(A^{-1})^{-1} = A$



Watch Video Solution

9. Evaluate: $\begin{bmatrix} x & y & x+y \\ y & x+y & x \\ x+y & x & y \end{bmatrix}$



Watch Video Solution

10. Evaluate the following:

$$\begin{vmatrix} 1 & x & y \\ 1 & x+y & y \\ 1 & x & x+y \end{vmatrix}$$



Watch Video Solution

11. Using properties of determinants in questions 11 to 15, prove that :

$$\begin{vmatrix} \alpha & \alpha^2 & \beta + \gamma \\ \beta & \beta^2 & \gamma + \alpha \\ \gamma & \gamma^2 & \alpha + \beta \end{vmatrix} = (\beta - \gamma)(\gamma - \alpha)(\alpha - \beta + \gamma)$$



Watch Video Solution

12. Using properties of determinants. Prove that

$$\begin{vmatrix} x^2 & 1 + px^3 & yy^2 & 1 + py^3 & zz^2 & 1 + pz^3 \end{vmatrix} = (1 + pxyz)(x - y)(y - z)(z - x)$$
, where p is any scalar.



Watch Video Solution

13. Using properties of determinants, prove the following:

$$|3a - a + b - a + ca - b3bc - ba - cb - c3c| = 3(a + b + c)(ab + bc +$$



Watch Video Solution

14.

Show

that

$$|11 + p1 + p + q23 + 2p1 + 3p + 2q36 + 3p106p + 3q| = 1.$$



Watch Video Solution

15. Show that $\begin{vmatrix} \sin \alpha & \cos \alpha & \cos(\alpha + \delta) \\ \sin \beta & \cos \beta & \cos(\beta + \delta) \\ \sin \gamma & \cos \gamma & \cos(\gamma + \delta) \end{vmatrix} = 0$



Watch Video Solution

16. $\frac{2}{x} + \frac{3}{y} + \frac{10}{z} = 4, \frac{4}{x} - \frac{6}{y} + \frac{5}{z} = 1, \frac{6}{x} + \frac{9}{y} - \frac{20}{z} = 2$



Watch Video Solution

17. Choose the correct answer in questions 17 to 19:

If a, b, c are in $A. P$, then the determinant $\begin{bmatrix} x+2 & x+3 & x+2a \\ x+3 & x+4 & x+2b \\ x+4 & x+5 & x+2c \end{bmatrix}$ is :

A. 0

B. 1

C. x

D. $2x$

Answer: A



Watch Video Solution

18. Choose the correct answer in questions 17 to 19:

If x, y, z are nonzero real numbers then the inverse of metrix

$$A = \begin{bmatrix} x & 0 & 0 \\ 0 & y & 0 \\ 0 & 0 & z \end{bmatrix} \text{ is :}$$

(a) $\begin{bmatrix} x^{-1} & 0 & 0 \\ 0 & y^1 & 0 \\ 0 & 0 & z^1 \end{bmatrix}$

(b) $xyz \begin{bmatrix} x^{-1} & 0 & 0 \\ 0 & y^1 & 0 \\ 0 & 0 & z^1 \end{bmatrix}$

(c) $\frac{1}{xyz} \begin{bmatrix} x & 0 & 0 \\ 0 & y & 0 \\ 0 & 0 & z \end{bmatrix}$

(d) $\frac{1}{xyz} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$

A. $\begin{bmatrix} x^{-1} & 0 & 0 \\ 0 & y^{-1} & 0 \\ 0 & 0 & z^{-1} \end{bmatrix}$

B. $xyz \begin{bmatrix} x^{-1} & 0 & 0 \\ 0 & y^{-1} & 0 \\ 0 & 0 & z^{-1} \end{bmatrix}$

C. $\frac{1}{xyz} \begin{bmatrix} x & 0 & 0 \\ 0 & y & 0 \\ 0 & 0 & z \end{bmatrix}$

D. $\frac{1}{xyz} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$

Answer: A



Watch Video Solution

19. Let $A = \begin{bmatrix} 1 & \sin \theta & 1 \\ -\sin \theta & 1 & \sin \theta \\ -1 & -\sin \theta & 1 \end{bmatrix}$, where $0 \leq \theta < 2\pi$ then (A)

$|A| = 0$ (B) $|A|\varepsilon[2, 4]$ (C) $|A|\varepsilon(0, \infty)$ (D) $|A|\varepsilon(2, \infty)$

A. $\text{Det}(A) = 0$

B. $\text{Det}(A)\varepsilon(2, \infty)$

C. $\text{Det}(A)\varepsilon(2, 4)$

D. $\text{Det}(A)\varepsilon[2, 4]$

Answer: D



Watch Video Solution

Exercise

1. $\begin{vmatrix} x & p & q \\ p & x & q \\ p & q & x \end{vmatrix} =$



Watch Video Solution

2. Show that $\begin{vmatrix} 0 & b-a & c-a \\ a-b & 0 & c-b \\ a-c & b-c & 0 \end{vmatrix} = 0$



Watch Video Solution

3. If $x = -4$ is a root of $\Delta = \begin{vmatrix} x & 2 & 3 \\ 1 & x & 1 \\ 3 & 2 & x \end{vmatrix} = 0$, then find the other two roots.



Watch Video Solution

4. Show that the points $(a + 5, a - 4)$, $(a - 2, a + 3)$ and (a, a) do not lie on a straight line of any value of a .



Watch Video Solution

Revision Exercise

1. For the matrix $A = \begin{bmatrix} 3 & 1 \\ 7 & 5 \end{bmatrix}$. Find 'x' and 'y' so that $A^2 + xI = yA$.

Hence find A^{-1} .



[Watch Video Solution](#)

2. $\begin{vmatrix} 1 & \cos(\beta - \alpha) & \cos(\gamma - \alpha) \\ \cos(\alpha - \beta) & 1 & \cos(\gamma - \beta) \\ \cos(\beta - \alpha) & \cos(\beta - \gamma) & 1 \end{vmatrix} =$



[Watch Video Solution](#)

3. Prove tha following

$$\begin{vmatrix} a + b + nc & na - a & nb - b \\ nc - c & b + c + na & nb - b \\ nc - c & na - a & c + a + nb \end{vmatrix} = n(a + b + c)^3.$$



[Watch Video Solution](#)

4. Prove: $\left| \frac{a^2 + b^2}{c} a \frac{b^2 + c^2}{a} a \frac{\mathbf{c}^2 + \mathbf{a}^2}{b} \right| = 4abc$



[Watch Video Solution](#)

5.

Prove

that

$$\begin{vmatrix} a^2 & a^2 - (b - c)^2 & bc \\ b^2 & b^2 - (c - a)^2 & ca \\ c^2 & c^2 - (a - b)^2 & ab \end{vmatrix} = (a - b)(b - c)(c - a)(a + b + c)(a^2 + b^2 + c^2)$$



Watch Video Solution

6.

Show

that

$$\begin{vmatrix} -a(b^2 + c^2 - a^2) & 2b^3 & 2c^3 \\ 2a^3 & -b(c^2 + a^2 - b^2) & 2c^3 \\ 2a^3 & 2b^3 & -c(a^2 + b^2 - C^2) \end{vmatrix} = abc(a^2 + b^2 + c^2)^3$$



Watch Video Solution

7. Prove that $\Delta =$

$$\begin{vmatrix} 1 & bc + ad & b^2c^2 + a^2d^2 \\ 1 & ca + bd & c^2a^2 + b^2d^2 \\ 1 & ab + cd & a^2b^2 + c^2d^2 \end{vmatrix} = (a - b)(a - c)(a - d)(b - c)(b - d)(c - d).$$



Watch Video Solution

8.
$$\begin{vmatrix} (a+1)(a+2) & a+2 & 1 \\ (a+2)(a+3) & a+3 & 1 \\ (a+3)(a+4) & a+4 & 1 \end{vmatrix} = -2$$



Watch Video Solution

9. Evaluate
$$\begin{vmatrix} .^x C_1 & .^x C_2 & .^x C_3 \\ .^y C_1 & .^y C_2 & .^y C_3 \\ .^z C_1 & .^z C_2 & .^z C_3 \end{vmatrix}$$



Watch Video Solution

10. Prove that

$$\Delta = [a + bxc + dxp + qxax + bcx + dpx + quvw] = (1 - x^2)|acpbqdqucw|$$



Watch Video Solution

11. Evaluate

$$\begin{bmatrix} \cos \alpha \cos \beta & \cos \alpha \sin \beta & -\sin \alpha \\ -\sin \beta & \cos \beta & 0 \\ \sin \alpha \cos \beta & \sin \alpha \sin \beta & \cos \alpha \end{bmatrix}$$



Watch Video Solution

12. If $\begin{vmatrix} 1 & 1 & 1 \\ a & b & c \\ a^3 & b^3 & c^3 \end{vmatrix} = (a - b)(b - c)(c - a)(a + b + c)$

where a, b, c are all different, then the determinant

$$\begin{vmatrix} 1 & 1 & 1 \\ (x - a)^2 & (x - b)^2 & (x - c)^2 \\ (x - b)(x - c) & (x - c)(x - a) & (x - a)(x - b) \end{vmatrix} \text{ vanishes when}$$



Watch Video Solution

13. Solve for $x \in R$: $\begin{vmatrix} (x + a)(x - a) & (x + b)(x - b) & (x + c)(x - c) \\ (x - a)^3 & (x - b)^3 & (x - c)^3 \\ (x + a)^3 & (x + b)^3 & (x + c)^3 \end{vmatrix} = 0$, a, b and c being distinct real numbers.



Watch Video Solution

14. If a,b,c are in A.P. find the value of:

$$| | 2y + 4, 5y + 7, 8y + a], [3y + 5, 6y + 8, 9y + b], [4y + 6, 7y + 9, 10y + c]$$



Watch Video Solution

15.

If

$$ax^2 + 2hxy + by^2 + 2gx + 2fy + c \equiv (lx + my + n)(l'x + m'y + n')$$

, then prove that :

$$\begin{vmatrix} a & h & g \\ h & b & f \\ g & f & c \end{vmatrix} = 0$$



Watch Video Solution

16. If $a + b + c = 0$ and $\begin{vmatrix} a - x & c & b \\ c & b - x & a \\ b & a & c - x \end{vmatrix} = 0$ then $x =$



Watch Video Solution

17. If $A + B + C = \pi$, then value of $\begin{vmatrix} \sin(A + B + C) & \sin B & \cos C \\ -\sin B & 0 & \tan A \\ \cos(A + B) & -\tan A & 0 \end{vmatrix}$

is



Watch Video Solution

18. Using properties of determinants. Prove that

$$\left| \begin{matrix} x^2 & 1 + px^3y & y^2 \\ 1 + py^3z & z^2 & 1 + pz^3 \end{matrix} \right| = (1 + pxyz)(x - y)(y - z)(z - x)$$

, where p is any scalar.



Watch Video Solution

19. If $A = \begin{bmatrix} 3 & -3 & 4 \\ 2 & -3 & 4 \\ 0 & -1 & 1 \end{bmatrix}$, then



Watch Video Solution

20. If $A = \begin{bmatrix} 3 & -3 & 4 \\ 2 & -3 & 4 \\ 0 & -1 & 1 \end{bmatrix}$, then show that $A^3 = A^{-1}$.



Watch Video Solution

21. If $A = [1 \tan x - \tan x 1]$, show that
 $A^T A^{-1} = [\cos 2x - \sin 2x \sin 2x \cos 2x]$.



Watch Video Solution

22. If $A = \begin{bmatrix} 2 & -3 \\ 4 & 6 \end{bmatrix}$ verify that $(adjA)^{-1} = (adjA^{-1})$.



Watch Video Solution

23. Prove that :

$$adj. I_n = I_n$$



Watch Video Solution

24. Prove that :

$$\text{adj. } O = O$$



Watch Video Solution

25. Prove that :

$$I_n^{-1} = I_n$$



Watch Video Solution

26. Find the inverse of each of the matrices given below :

Let $D = \text{diag}[d_1, d_2, d_3]$ where none of d_1, d_2, d_3 is 0, prove that

$$D^{-1} = \text{diag}[d_1^{-1}, d_2^{-1}, d_3^{-1}].$$



Watch Video Solution

27. Let $F(\alpha) = \begin{bmatrix} \cos \alpha & -\sin \alpha & 0 \\ \sin \alpha & \cos \alpha & 0 \\ 0 & 0 & 1 \end{bmatrix}$ and $G(\beta) = \begin{bmatrix} \cos \beta & 0 & \sin \beta \\ 0 & 1 & 0 \\ -\sin \beta & 0 & \cos \beta \end{bmatrix}$.

Show that $[F(\alpha) \cdot G(\beta)]^{-1} = G(-\beta) \cdot F(-\alpha)$.



[Watch Video Solution](#)

28. Find the inverse of each of the matrices given below :

Obtain the inverse of the matrices $\begin{bmatrix} 1 & p & 0 \\ 0 & 1 & p \\ 0 & 0 & 1 \end{bmatrix}$ and $\begin{bmatrix} 1 & 0 & 0 \\ q & 1 & 0 \\ 0 & q & 1 \end{bmatrix}$. And, hence find the inverse of the matrix $\begin{bmatrix} (1+pq) & p & 0 \\ q & (1+pq) & p \\ 0 & q & 1 \end{bmatrix}$.

Let the first two matrices be A and B. Then, the third matrix is AB. Now,

$$(AB)^{-1} = (B^{-1}A^{-1})$$



[Watch Video Solution](#)

29. Use product $[1 \ 1202 \ -33 \ -24] \ [-20192 \ -361 \ -2]$ to solve the system of equation: $x - y + 2z = 1$ $2y - 3z = 1$ $3x - 2y + 4z = 2$



[Watch Video Solution](#)



Watch Video Solution

30. If $a \neq p, b \neq q, c \neq r$ and $|pbcaqcabr| = 0$, then find the value of

$$\frac{p}{p-a} + \frac{q}{q-b} + \frac{r}{r-c}.$$



Watch Video Solution

31. Suppose that digit numbers A28,3B9 and 62 C, where A,B and C are integers between 0 and 9 are divisible by a fixed integer k, prove that the

determinant $\begin{vmatrix} A & 3 & 6 \\ 8 & 9 & C \\ 2 & B & 2 \end{vmatrix}$ is also divisible by k.



Watch Video Solution

32. let $a > 0, d > 0$ find the value of the determinant

$$\begin{vmatrix} \frac{1}{a} & \frac{1}{a(a+d)} & \frac{1}{(a+d)(a+2d)} \\ \frac{1}{a+d} & \frac{1}{(a+d)(a+2d)} & \frac{1}{(a+2d)(a+3d)} \\ \frac{1}{a+2d} & \frac{1}{(a+2d)(a+3d)} & \frac{1}{(a+3d)(a+4d)} \end{vmatrix}$$



Watch Video Solution

Check Your Understanding

1. Evaluate the determinants in questions 1 and 2 :

If $A = \begin{bmatrix} 1 & 2 \\ 4 & 2 \end{bmatrix}$, then show that $|2A|=4|A|$.



[Watch Video Solution](#)

2. If $\begin{vmatrix} x & 2 \\ 8 & x \end{vmatrix} = \begin{vmatrix} 3 & 2 \\ 9 & 6 \end{vmatrix}$, then the value of 'x' is



[Watch Video Solution](#)

3. Answer in one word

$$\begin{vmatrix} 3 & 1 & 9 \\ 5 & 2 & 15 \\ 7 & 4 & 21 \end{vmatrix}.$$



[Watch Video Solution](#)

4. If any two rows (or columns) of a determinant are identical, then the value of the determinant is zero.

 Watch Video Solution

5. Is $\left(\frac{1}{a} + \frac{1}{b} + \frac{1}{c} - 1\right)$ a factor of $\begin{vmatrix} 1+a & 1 & 1 \\ 1 & 1+b & 1 \\ 1 & 1 & 1+c \end{vmatrix}$?

 Watch Video Solution

6. If A is an invertible square matrix of order 4, then $|adjA| = \dots$

 Watch Video Solution

7. If A is square matrix of order n then $adj(adjA) =$ (1) $|A|^{n-1}A$ (2) $|A|^{n-2}A$ (3) $|A|^{n-2}$ (4) $|A|^nA$

 Watch Video Solution

8. Find the inverse of $A = \begin{bmatrix} 2 & -6 \\ 1 & -2 \end{bmatrix}$.



Watch Video Solution

9. If A is a square matrix satisfying $A^2 = 1$, then what is the inverse of A ?



Watch Video Solution

10. Is the matrix $\begin{bmatrix} \pi & 22 \\ \frac{1}{7} & 1 \end{bmatrix}$ singular?



Watch Video Solution

Competition File

1. Let a, b and c be such that $(b+c) \neq 0$. If

$$\begin{vmatrix} a & a+1 & a-1 \\ -b & b+1 & b-1 \\ c & c-1 & c+1 \end{vmatrix} + \begin{vmatrix} a+1 & b+1 & c-1 \\ a-1 & b-1 & c+1 \\ (-1)^{n+2}a & (-1)^{n+1}b & (-1)^nc \end{vmatrix} = 0, \text{ then}$$

the value of 'n' is :

- A. zero
- B. any integer
- C. any odd integer
- D. any integer

Answer: C



Watch Video Solution

2. The number of values of k for which the linear equations $4x + ky + 2z = 0$, $kx + 4y + z = 0$, $2x + 2y + z = 0$ posses a non-zero solution is : (1) 3 (2) 2 (3) 1 (4) zero

- A. 3
- B. 2
- C. 1

D. zero

Answer: B



Watch Video Solution

3. The number of values of k , for which the system of equations:

$$(k + 1)x + 8y = 4k$$

$$kx + (k + 3)y = 3k - 1$$

has no solution is,

A. 1

B. 2

C. 3

D. infinite.

Answer: A



Watch Video Solution

4. If $P = \begin{bmatrix} 1 & \alpha & 3 \\ 1 & 3 & 3 \\ 2 & 4 & 4 \end{bmatrix}$ is the adjoint of 3×3 matrix A and $|A| = 4$, then

$$\alpha =$$

A. 11

B. 5

C. 0

D. 4

Answer: A



Watch Video Solution

5. If $\alpha, \beta \neq 0$, and $f(n) = \alpha^n + \beta^n$ and

$|31 + f(1)1 + f(2)1 + f(1)1 + f(2)1 + f(3)1 + f(2)1 + f(3)1 + f(4)| = K$, then K is equal to (1) $\alpha\beta$ (2) $\frac{1}{\alpha\beta}$ (3) 1 (4) -1

A. $\frac{1}{\alpha\beta}$

B. 1

C. -1

D. $\alpha\beta$

Answer: B



Watch Video Solution

6. The set of all values of λ for which the system of linear equations :

$2x_1 - 2x_2 + x_3 = \lambda x_1$ $2x_1 - 3x_2 + 2x_3 = \lambda x_2$ $-x_1 + 2x_2 = \lambda x_3$ has a non-trivial solution, (1) is an empty set (2) is a singleton (3) contains two elements (4) contains more than two elements

A. is an empty set

B. is a singleton

C. contains 2 elements

D. contains more than 2 elements

Answer: C



Watch Video Solution

7. The system of linear equations $x + \lambda y - z = 0$ $\lambda x - y - z = 0$ $x + y - \lambda z = 0$ has a non-trivial solution for : (1) infinitely many values of λ . (2) exactly one value of λ . (3) exactly two values of λ . (4) exactly three values of λ .

A. Exactly one value of λ

B. Exactly two values of λ

C. Exactly three values of λ

D. Infinitely many values of λ .

Answer: C



Watch Video Solution

8. Let ω be a complex number such that $2\omega + 1 = z$ where $z = \sqrt{-3}$. If

$|((1, 1, 1), (1, -\omega^2 - 1, \omega^2), (1, \omega^2, \omega^7))| = 3k$, then k is equal to

A. -1

B. 1

C. $-z$

D. z

Answer: C



Watch Video Solution

9. If S is the set of distinct values of ' b ' for which the following system of linear equations $x + y + z = 1$ $x + ay + z = 1$ $ax + by + z = 0$ has no solution, then S is :
a finite set containing two or more elements (2) a singleton
an empty set (4) an infinite set

A. a finite set containing two or more elements

B. a singleton

C. an empty set

D. an infinite set

Answer: B



Watch Video Solution

10. If $A = \begin{vmatrix} 2 & -3 \\ -4 & 1 \end{vmatrix}$ then $\text{adj}(3A^2 + 12A)$ is equal to

A. $\begin{bmatrix} 51 & 84 \\ 63 & 72 \end{bmatrix}$

B. $\begin{bmatrix} 72 & -63 \\ -84 & 51 \end{bmatrix}$

C. $\begin{bmatrix} 72 & -84 \\ -63 & 51 \end{bmatrix}$

D. $\begin{bmatrix} 51 & 63 \\ 84 & 72 \end{bmatrix}$

Answer: D



Watch Video Solution

11. If $\begin{vmatrix} x - 4 & 2x & 2x \\ 2x & x - 4 & 2x \\ 2x & 2x & x - 4 \end{vmatrix} = (A + Bx)(x - A)^2$ then the ordered pair (A,B) is equal to

A. (-4, -5)

B. (-4, 3)

C. (-4, 5)

D. (4, 5)

Answer: C



Watch Video Solution

12. If the system of linear equations $x+ky+3z=0$ $3x+ky-2z=0$ $2x+4y-3z=0$ has a non-zero solution (x,y,z) then $\frac{xz}{y^2}$ is equal to

A. -10

B. 10

C. -30

D. 30

Answer: B



Watch Video Solution

13. An ordered pair (α, β) for which the system of linear equations

$$(1 + \alpha)x + \beta y + z = 2$$

$$\alpha x + (1 + \beta)y + z = 3$$

$$\alpha x + \beta y + 2z = 2$$

has a unique solution, is

A. $(1, -3)$

B. $(2, 4)$

C. $(-3, 1)$

D. $(-4, 2)$

Answer: B



Watch Video Solution

14. If α, β are the roots of $x^2 + x + 1 = 0$ then

$$\begin{vmatrix} Y+1 & \beta & \alpha \\ \beta & y+\alpha & 1 \\ \alpha & 1 & y+\beta \end{vmatrix}$$

A. $y^2 - 1$

B. $y(y^2 - 1)$

C. $y^2 - y$

D. y^3

Answer: D



Watch Video Solution

Chapter Test 4

1. If $\Delta = |a_{11}a_{12}a_{13}a_{21}a_{22}a_{23}a_{31}a_{32}a_{33}|$ and A_{ij} is cofactors of a_{ij} , then value of Δ is given by (A) $a_{11} + A_{31} + a_{12}A_{32} + a_{13}A_{33}$ (B) $a_{11}A_{11} + a_{12}A_{21} + a_{13}A_{31}$ (C) $a_{11}A_{31} + a_{12} + A_{32} + a_{12}A_{33}$

A. $a_{11}A_{31} + a_{12} + A_{32} + a_{12}A_{33}$

B. $a_{11}A_{11} + a_{12} + A_{21} + a_{13}A_{33}$

C. $a_{21}A_{11} + a_{22} + A_{12} + a_{23}A_{13}$

D. $a_{11}A_{11} + a_{21} + A_{21} + a_{31}A_{31}$

Answer: B



Watch Video Solution

2. Let $A = \begin{bmatrix} 1 & \sin \theta & 1 \\ -\sin \theta & 1 & \sin \theta \\ -1 & -\sin \theta & 1 \end{bmatrix}$, where $0 \leq \theta < 2\pi$ then (A)

|A| = 0 (B) |A| \in [2, 4] (C) |A| \in (0, \infty) (D) |A| \in (2, \infty)

A. $\text{Det}(A) = 0$

B. $\text{Det}(A)\varepsilon(2, \infty)$

C. $\text{Det}(A)\varepsilon(2, 4)$

D. $\text{Det}(A)\varepsilon[2, 4]$

Answer: D



Watch Video Solution

3. If A is a square matrix of order 3 and $|3A| = k|A|$ then find the value of k,



Watch Video Solution

4. If $| \begin{matrix} 1 & x \\ 3 & 4 \end{matrix} | = |3412|$, write the positive value of x.



Watch Video Solution

5. If A is a square matrix of order 3 such that $|\text{adj } A|=225$ find $|A'|$



Watch Video Solution

6. Using properties of determinants, prove that following:

$$|a + b + 2cabcb + c + 2abcac + a + 2b| = 2(a + b + c)^3$$



Watch Video Solution

7. Prove that

$$\begin{vmatrix} a^2 + 1 & ab & ac \\ ab & b^2 + 1 & bc \\ ac & bc & c^2 + 1 \end{vmatrix} = 1 + a^2 + b^2 + c^2.$$



Watch Video Solution

8. Using Cofactors of elements of third column, evaluate

$$\Delta = |1xyz\ 1yzx\ 1zxy|$$



Watch Video Solution

9. Find the values of k if area of triangle is 4 sq. units and vertices are:
 $(-2,0), (0,4), (0,k)$

 Watch Video Solution

10. Find the inverse of the matrix $A = \begin{bmatrix} (1, b), \left(c, \frac{1+bc}{a}\right) \end{bmatrix}$ and show that $aA^{-1} = (a^2 + bc + 1)I - aA$

 Watch Video Solution

11. Using properties of determinants, prove that
 $|aa + ba + b + c| = a^3$

 Watch Video Solution

12. If $A = \begin{pmatrix} 2 & -3 & 5 \\ 3 & 2 & -4 \\ 1 & 1 & -2 \end{pmatrix}$, find A^{-1} . Using A^{-1} solve the following system of equations:

$$2x - 3y + 5z = 16; \quad 3x + 2y - 4z = -4; \quad x + y - 2z = -3$$



Watch Video Solution