



PHYSICS

BOOKS - HC VERMA PHYSICS (ENGLISH)

WORK AND ENERGY

Example

1. A spring of spring constant 50 N/m is compressed from its natural position through

1 cm. Find the work done by the work done by the spring force on the agency compressing the spring.




[Watch Video Solution](#)

2. A particle of mass 20 g is thrown vertically upwards with a speed of 10 m/s. Find the work done by the force of gravity during the time the particle goes up.



[Watch Video Solution](#)

3. Two charged particles A and B repel each other by a force $\frac{k}{r^2}$ where k is a constant and r is the separation between them. Particle A is clamped to a fixed point in the lab and particle B which has a mass m , is released from rest with an initial separation r_0 from A. Find the change in the potential energy of the two particle system as the separation increases to a large value. What will be the speed of the particle B in this situation?



Watch Video Solution

4. A block of mass m slides along a frictionless surface as shown in the figure. If it is released from rest from A what is its speed at B?




Figure 8.10



[Watch Video Solution](#)

5. A pendulum bob has a speed 3m/s while passing through its lowest position. What is its speed when it makes an angle of 60° with the vertical? The length of the pendulum is 0.5m Take $g = 10\frac{\text{m}}{\text{s}^2}$.



Watch Video Solution

6. A block of mass m , attached to a spring of spring constant k , oscillates on a smooth horizontal table. The other end of the spring is

fixed to a wall. If it has speed v when the spring is at its natural length, how far will it move on the table before coming to an instantaneous rest?



[Watch Video Solution](#)

7. A block of mass m is suspended through a spring of spring constant k and is in equilibrium. A sharp blow gives the block an initial downward velocity v . How far below the

equilibrium position, the block comes to an instantaneous rest?



[Watch Video Solution](#)

Worked Out Examples

1. A porter lifts a suitcase weighing 20 kg from the platform and puts it on his head 2.0 m above the platform. Calculate the work done by the porter on the suitcase.



[Watch Video Solution](#)

2. An elevator weighing 500 kg is to be lifted up at a constant velocity of 0.20 m/s. What would be the minimum horsepower of the motor to be used?



[Watch Video Solution](#)

3. A block of mass 2.0 kg is pulled up on a smooth incline of angle 30° with the horizontal. If the block moves with an acceleration of $1.0 \frac{m}{s^2}$, find the power

delivered by the pulling force at a time 4.0 s after the motion starts. What is the average power delivered during the 4.0 s after the motion starts?



[Watch Video Solution](#)

4. A force $F = (10 + 0.50x)$ acts on a particle in the x direction, where F is in newton and x in meter./find the work done by this force during a displacement from $x=0$ to $x=2.0\text{m}$



[Watch Video Solution](#)

5. A body dropped from a height H reaches the ground with a speed of $1.2 \sqrt{gH}$. Calculate the work done by air friction.



[Watch Video Solution](#)

6. A block of mass M is pulled along a horizontal surface by applying a force at angle θ with the horizontal. The friction coefficient between the block and the surface is μ . If the block travels at a uniform velocity, find the

work done by this applied force during a displacement d of the block.



[Watch Video Solution](#)

7. Two identical cylindrical vessels with their bases at the same level each contain a liquid of density ρ . The height of the liquid in one vessel is h_1 and in the other is h_2 the area of either base is A . What is the work done by gravity in equalising the levels when the two vessels are connected?



[Watch Video Solution](#)

8. What minimum horizontal speed should be given to the bob of a simple pendulum of length l so that it describes a complete circle?



[Watch Video Solution](#)

9. A uniform chain of length l and mass m overhangs a smooth table with its two third part lying on the table. Find the kinetic energy of the chain as it completely slips off the table.



Watch Video Solution

10. A block of mass m is pushed against a spring of spring constant k fixed at the end to a wall. The block can slide on a frictionless table as shown in figure. The natural length of the spring is L_0 and it is compressed to half its natural length when the block is released. Find the velocity of the block as a function of its

distance x from the wall .




Figure 8-W6



[Watch Video Solution](#)

11. A particle is placed at the point A of a frictionless track ABC as shown in figure. It is pushed slightly towards right. Find its speed

when it reaches the point B. Take $g = 10 \frac{m}{s^2}$.




Figure 8-W7



[Watch Video Solution](#)

12. Figure shows a smooth curved track terminating in a smooth horizontal part. A spring of spring constant 400 N/m is attached at one end to a wedge fixed rigidly with the horizontal part. A 40 g mass is

released from rest at a height of 4.9 m on the curved track. Find the maximum compression of the spring.




Figure 8-W8



[Watch Video Solution](#)

13. Figure shows a loop the loop track of radius R . A car (without engine) starts from a

platform at a distance h above the top of the loop and goes around the loop without falling off the track. Find the minimum value of h for a successful looping. Neglect friction.




Figure 8-W9



Watch Video Solution

14. A heavy particle is suspended by a string of length l . The particle is given a horizontal

velocity v_0 . The string becomes slack at some angle and the particle proceeds on a parabola. find the value of v_0 if the particle passes through the point of suspension




Figure 8-W10



Watch Video Solution

Objective 1

1. A heavy stone is thrown from a cliff of height h with a speed v . The stone will hit the ground with maximum speed if it is thrown

A. vertically downward

B. vertically upward

C. horizontally

D. the speed does not depend on the initial direction.

Answer: D



Watch Video Solution

2. Two springs A and B ($k_A = 2k_B$) are stretched by applying forces of equal magnitudes at the force ends. If the energy stored in A is E, that in B is

A. $\frac{E}{2}$

B. $2E$

C. E

D. $\frac{E}{4}$

Answer: B



Watch Video Solution

3. Two equal masses are attached to the two ends of a spring of spring constant k . The masses are pulled out symmetrically to stretch

the spring by a length x over its natural length. The work done by the spring on each mass is

A. $\frac{1}{2}kx^2$

B. $-\frac{1}{2}kx^2$

C. $\frac{1}{4}kx^2$

D. $-\frac{1}{4}kx^2$

Answer: D



Watch Video Solution

4. The negative of the work done by the conservative internal forces on a system equals the change in

- A. total energy
- B. kinetic energy
- C. potential energy
- D. none of these

Answer: C



Watch Video Solution

5. The work done by the external forces on a system equals the change in

- A. total energy
- B. kinetic energy
- C. potential energy
- D. none of these

Answer: A



Watch Video Solution

6. The work done by all the forces (external and internal) on a system equals the change in

- A. total energy
- B. kinetic energy
- C. potential energy
- D. none of these

Answer: B



Watch Video Solution

7. _____ of a two article system depends only on the separation between the two particles. The most appropriate choice for the blank space in the above sentence is

- A. kinetic energy
- B. total mechanical energy
- C. potential energy
- D. total enerty

Answer: C



Watch Video Solution

8. A small block of mass m is kept on a rough inclined surface of inclination θ fixed in an elevator. The elevator goes up with a uniform velocity v and the block does not slide on the wedge. The work done by the force of friction on the block in time t will be

A. zero

B. $mgvt \cos^2 \theta$

C. $mgvt \sin^2 \theta$

$D. mgvtsi2\theta$

Answer: C



Watch Video Solution

9. A block of mass m slides down a smooth vertical circular track. During the motion, the block is in

A. vertical equilibrium

B. horizontal equilibrium

C. radia equilibrium

D. none of these

Answer: D



Watch Video Solution

10. A particle is rotated in vertical circle by connecting it to string fixed. The minimum speed of the particle when the string is horizontal for which the particle will complete the circle is

A. \sqrt{gl}

B. $\sqrt{2ghl}$

C. $\sqrt{3gl}$

D. $\sqrt{5gl}$

Answer: C



Watch Video Solution

Objective 2

1. A heavy stone is thrown from a cliff of height h in a given direction. The speed with which it hits the ground

A. must depend on the speed of projection

B. must be larger than the speed of projection

C. must be independent of the speed of projection

D. may be smaller than the speed of projection

Answer: A::B



Watch Video Solution

2. The total work done on a particle is equal to the change in its kinetic energy

A. always

B. only if the forces acting on it are conservative

C. only if gravitational force alone acts on it

D. only if elastic force alone acts on it

Answer: A



Watch Video Solution

3. A particle is acted upon by a force of constant magnitude which is always perpendicular to the velocity of the particle.

The motion of the particle takes place in a plane. It follows that

- A. its velocity is constant
- B. its acceleration is constant
- C. its kinetic energy is constant
- D. it moves in a circular path

Answer: C::D



Watch Video Solution

4. Consider two observers moving with respect to each other at a speed v along a straight line. They observe a block of mass m moving a distance l on a rough surface. The following quantities will be same as observed by the two observers

A. kinetic energy of the block at time t

B. work done by friction

C. total work done on the block

D. acceleration of the block

Answer: D



Watch Video Solution

5. You lift a suitcase from the floor and keep it on a table. The work done by you on the suitcase does not depend on

- A. the path taken by the suitcase
- B. the time taken by you in doing so
- C. the weight of the suitcase
- D. your weight

Answer: A::B::D



Watch Video Solution

6. The kinetic energy of a particle continuously increases with time

A. the resultant force on the particle must

be parallel to the velocity at all instants.

B. the resultant force on the particle must

be at an angle less than 90^0 all the time

C. Its height above the ground level must continuously decrease

D. the magnitude of its linear momentum is increasing continuously.

Answer: B::D



Watch Video Solution

7. One end of a light spring of spring constant k is fixed to a wall and the other end is tied to a block placed on a smooth horizontal surface.

In a displacement, the work by the spring is

$\frac{1}{2}kx^2$. The possible cases are

- A. the spring was initially compressed by a distance x and was finally in its natural length
- B. it was initially in its natural length and finally was in its natural length
- C. it was initially in its natural length and finally in a compressed position.

D. it was initially in its natural length and finally in a stretched position.

Answer: A::B



Watch Video Solution

8. A block of mass M is hanging over a smooth and light pulley through a light string. The other end of the string is pulled by a constant force F . The kinetic energy of the block increases by 20J in 1s .

A. the tension in the string is Mg

B. The tension in the string is F

C. The work done by the tension on the block is 20 J in the above 1 s .

D. the work done by the force of gravity is -20 J in the above 1 s .

Answer: B



Watch Video Solution

Exercises

1. The mass of cyclist together with the bike is 90 kg. Calculate the increase in kinetic energy if the speed increases from 6.0 km/h to 12 km/h.



[Watch Video Solution](#)

2. A block of mass 2.00 kg moving at a speed of 10.0 m/s accelerates at $3.0 \frac{m}{s^2}$ for 5.00 s. Compute its final kinetic energy.



[Watch Video Solution](#)

3. A box is pushed through 4.0 m across floor offering 100 N resistance. How much work is done by the resisting force?



[Watch Video Solution](#)

4. A block of mass 5.0 kg slides down an incline of inclination 30° and length 10 m. Find the work done by the force of gravity.



Watch Video Solution

5. A constant force of 2.50 N accelerates a stationary particle of mass 15 g through a displacement of 2.50 m. Find the work done and the average power delivered.



Watch Video Solution

6. A particle moves from a point

$\vec{r}_1 = (2m)\vec{i} + (3m)\vec{j}$ to another point

$\vec{r}_2 = (3m)\vec{i} + (2m)\vec{j}$ during which a

certain force $\vec{F} = (5N)\vec{i} + (5N)\vec{j}$ acts on it. Find the work done by the force on the particle during the displacement.



[Watch Video Solution](#)

7. A man moves on a straight horizontal road with a block of mass 2 kg in his hand. If he covers a distance of 340 m with an acceleration of $0.5\frac{m}{s^2}$ find the work done by the man on the block during the motion.



[Watch Video Solution](#)

8. A force $F=a+bx$ acts on a particle in the x -direction, where a and b are constants. Find the work done by this force during a displacement from $x = 0 \rightarrow x = d$.



Watch Video Solution

9. A block of mass 250 g slides down an incline of inclination 37° with a uniform speed. Find the work done against the friction as the block slides through 1.0 m.



Watch Video Solution

10. A block of mass m is kept over another block of mass M and the system rests on a horizontal force F acting on the lower block produces an acceleration $\frac{F}{2(m + M)}$ in the system the two blocks always move together.

A. Find the coefficient of kinetic friction between the bigger block and the horizontal surface. b. find the frictional force acting on the smaller block. c. Find the work done by the force of friction on the smaller block by the

bigger block during a displacement d of the system.




Figure 8-E1



[Watch Video Solution](#)

11. A box weighing 2000 N is to be slowly slid through 20 m on a straight track having friction coefficient 0.2 with the box. A find the

work done by the person pulling the box with a chain at angle θ with the horizontal. B. find the work when the person has chosen a value of θ which ensures him the minimum magnitude of the force.



[Watch Video Solution](#)

12. A block of weight 100 N is slowly slide up on a smooth incline of inclination 37° by as person. Calculate the work done by the person in moving the block through a distance of 2.0

m, if the driving force is a. parallel to the incline and b. in the horizontal direction.



Watch Video Solution

13. Find the average frictional force needed to stop a a car weighing 500 kg in a distance of 25 m (the initial speed is 72 km/h.)



Watch Video Solution

14. Find the average force needed to accelerate a car weighing 500 kg from rest to 72 km/h in a distance of 25 m.



Watch Video Solution

15. A particle of mass m moves on a straight line with its velocity varying with the distance travelled according to the equation $v = a\sqrt{x}$, where a is a constant. Find the total work

done by all the forces during a displacement from $x = 0 \rightarrow x = d$.



Watch Video Solution

16. A block of mass 2.0 kg kept at rest on an inclined plane of inclination 37° is pulled up the plane by applying a constant force of 20 N parallel to the incline. The force acts for one second. A. show that the work done by the applied force does not exceed 40 J. b. find the work done by the applied force is 40 J. c.

Find the kinetic energy of the block at the instant the force ceases to act. Take $g = 10 \frac{m}{s^2}$.



[Watch Video Solution](#)

17. A block of mass 2.0 kg is pushed down an inclined plane of inclination 37° with a force of 20N acting parallel to the incline. It is found that the block moves on the incline with an acceleration of $10 \frac{m}{s^2}$. If the block started from rest, find the work done a. by the applied force

in the first second b. by the weight of the block in the first second and c. by the frictional force acting on the block in the first second.

Take $g = 10 \frac{m}{s^2}$.



[Watch Video Solution](#)

18. A 250 g block slides on a smooth horizontal table. Find the work done by the frictional force in bringing the block to rest if it is initially moving at a speed of 40 cm/s. If the friction coefficient between the table and the

block is 0.1 how far does the block move before coming to rest?



[Watch Video Solution](#)

19. Water falling from a 50 m high fall is to be used for generating electric energy. If $1.8 \times 10^5 \text{ kg}$ of water falls per hour and half the gravitational potential energy can be converted into electric energy, how many 100 W lamps can be it?



[Watch Video Solution](#)

20. A person is painting his house walls. He stands on a ladder with a bucket containing paint in one hand and a brush in other. Suddenly the bucket slips from his hand and falls down on the floor. If the bucket with the paint had a mass of 6.0 kg and was at a height of 2.0 m at the time it slipped, how much gravitational potential energy is lost together with the paint?



[Watch Video Solution](#)

21. A projectile is fired from the top of a 40 m high cliff with an initial speed of 50 m/s at an unknown angle. Find its speed when it hits the ground.



[Watch Video Solution](#)

22. The 200 m free style women's swimming gold medal at Seol Olympic 1988 went to Heike Friedrich of East Germany when she set a new Olympic record of 1 minute and 57.56 seconds. Assume that she covered most of the

distance with a uniform speed and had to exert 460 W to maintain her speed. Calculate the average force of resistance offered by the water during the swim.



[Watch Video Solution](#)

23. The US athlete Florence Griffith Joyner won the 100 m sprint gold medal at Seoul Olympic 1988 setting a new Olympic record of 10.54s. Assume that she achieved her maximum speed in a very short time and then ran the race with

that speed till she crossed the line. Take her mass to be 50 kg. a. Calculate the kinetic energy of Griffith Joyner at her full speed. b. Assuming that the track the wind etc. offered an average resistance of one tenth of her weight, calculate the work done by the resistance during the run. c. What power Griffith Joyner had to exert to maintain uniform speed?



Watch Video Solution

24. A water pump lifts water from a level 10 m below the ground. Water is pumped at a rate of 30 kg /minute with negligible velocity. Calculate the minimum horsepower the engine should have to do this.



Watch Video Solution

25. An unruly demonstrator lifts a stone of mass 200 g from the ground and throws it at his opponent. At the time of projection, the

stone is 150 cm above the ground and has a speed of 3.00 m/s. Calculate the work done by the demonstrator during the process. If it takes one second for the demonstrator to lift the stone and throw, what horsepower does he use?



[Watch Video Solution](#)

26. In a factory it is desired to lift 2000 kg of metal through a distance of 12 m in 1 minute.

Find the minimum horsepower of the engine to be used.



[Watch Video Solution](#)

27. A scooter company gives the following specifications about its product.

Weight of the scooter -95 kg

Maximum speed -60 km/h

Maximum engine power -3.5 hp

Pick up time to get the maximum speed -5s

Check the validity of these specifications.



[Watch Video Solution](#)

28. A block of mass 30.0 kg is being brought down by a chain. If the block acquires a speed of $40.0 \frac{cm}{s}$ in dropping down 2.00 m, find the work done by the chain during the process.



[Watch Video Solution](#)

29. The heavier block in an Atwood machine has a mass twice that of the lighter one. The tension in the string is 16.0 N when the system

is set into motion. Find the decrease in the gravitational potential energy during the first second after the system is released from rest.



[Watch Video Solution](#)

30. The two blocks in an Atwood machine have masses 2.0 kg and 3.0 kg. Find the work done by gravity during the fourth second after the system is released from rest.



[Watch Video Solution](#)

31. Consider the situation shown in figure. The system is released from rest and the block of mass 1.0 kg is found to have a speed 0.3 m/s after it has descended through a distance of 1 m . Find the coefficient of kinetic friction between the block and the table.




Figure 8-E2



Watch Video Solution

32. A block of mass 100g is moved with a speed of 5.0 m/s at the highest point in a closed circular tube of radius 10 cm kept in a vertical plane. The cross section of the tube is such that the block just fits in it. The block makes several oscillations inside the tube and finally stops at the lowest point. Find the work done by the tube on the block during the process



Watch Video Solution

33. A car weighing 1400 kg is moving at speed of 54 km/h up a hill when the motor stops. If it is just able to reach the destination which is at a height of 10 m above the point calculate the work done against friction (negative of the work done by the friction).



Watch Video Solution

34. small block of mass 200 g is kept at the top of a frictionless incline which is 10 m long and

3.2 m high. How much work was required a. to lift the block from the ground and put it at the top b. to slide the block up the incline? What will be the speed of the block when it reaches the ground if, c. it falls off the incline and drops vertically on the ground d. it slides down the incline? Take $g = 10 \frac{m}{s^2}$.



[Watch Video Solution](#)

35. In a children's park, there is a slide which has a total length of 10 m and a height of 8.0

m figure. Vertical ladder are provided to reach the top. A boy weighting 200 N climbs up the ladder to the top of the slide and slides down to the ground. The averages friction offered by the slide is three tenth of his weight. Find a. the work done by the ladder on the boy as he goes up. b. the work done by the slide on the boy as he comes down. Neglect any work done by forces inside the body of the boy.



[Watch Video Solution](#)

36. Figure shows a particle sliding on a frictionless track which terminates in straight horizontal section. If the particle starts slipping from the point. A, how far way from the track will the particle hit the ground ?




Figure 8-E3



View Text Solution

37. A block weighing 10 N travels down a smooth curved track AB joined to a rough horizontal surface. The rough surface has a friction coefficient of 0.20 with the block. If the block starts slipping on the track from a point 1.0 m above the horizontal surface, how far will it move on the rough surface?




Figure 8-E4



[Watch Video Solution](#)

38. A uniform chain of mass m and length l overhangs a table with its two third part on the table. Find the work to be done by a person to put the hanging part back on the table.



Watch Video Solution

39. A uniform chain of length L and mass M overhangs a horizontal table with its two third part on the table. The friction coefficient

between the table and the chain is μ . Find the work done by the friction during the period the chain slips off the table.



[Watch Video Solution](#)

40. A block of mass m slides along a frictionless surface as shown in the figure. If it is released from rest from A what is its speed

at B?




Figure 8.10



[Watch Video Solution](#)

41. A block of mass 5.0 kg is suspended from the end of a vertical spring which is stretched by 10 cm under the load of the block. The block is given a sharp impulse from below so that it

acquires an upward speed of 2.0 m/s. How high will it rise? Take $g = 10 \frac{m}{s^2}$



[Watch Video Solution](#)

42. A block of mass 250g is kept on a vertical spring of spring constant 100N/m fixed from below. The spring is now compressed to have a length 10cm shorter than its natural length and the system is released from this position. How high does the block rise? take $g = 10 \frac{m}{s^2}$.



[Watch Video Solution](#)

43. Figure shows a spring fixed at the bottom end of an incline of inclination 37° . A small block of mass 2 kg starts slipping down the incline from a point 4.8 m away from the spring. The block compresses the spring by 20 cm, stops momentarily and then rebounds through a distance of 1 m up the incline. Find

- the friction coefficient between the plane and the block and
- the spring constant of

the spring. Take $g = 10 \frac{m}{s^2}$.




Figure 8-E7



[View Text Solution](#)

44. A block of mass m moving at a speed ' v ' compresses as spring through a distance ' x ' before its speed is halved. Find the spring constant of the spring.



Watch Video Solution

45. Consider the situation shown in figure. Initially the spring is unstretched when the system is released from rest. Assuming no friction in the pulley, find the maximum

elongation of the spring.




Figure 8-E8



Watch Video Solution

46. A block of mass m is attached to two unstretched springs of spring constants k_1 and k_2 as shown in figure. The block is displaced towards right through a distance x and is released. Find the speed of the block as it passes through the mean position shown.




Figure 8-E9



Watch Video Solution

47. A block of mass m sliding on a smooth horizontal surface with velocity \vec{v} meets a long horizontal spring fixed at one end and having spring constant k as shown in figure. Find the maximum compression of the spring. Will the velocity of the block be the same as v when it comes back to the original position shown?



Figure 8-E10



Watch Video Solution

48. A small block of mass 100 g is pressed against a horizontal spring fixed at one end to compress the spring through 5.0 cm. The spring constant is 100 N/m. When released, the block moves horizontally till it leaves the spring. Where will it hit the ground 2 m below the spring?




Figure 8-E11

 [Watch Video Solution](#)


49. A small heavy block is attached to the lower end of a light rod of length l which can be rotated about its clamped upper end. What minimum horizontal velocity should the block be given so that it moves in a complete vertical circle?



 [View Text Solution](#)

50. Figure shows two block A and B , each having a mass of 320 g connected by a light string passing over a smooth light pulley. The horizontal surface on which the block A can slide is smooth. The block A is attached to spring constant 40 N/m whose other end is fixed to a support 40 cm above the horizontal surface. Initially, the spring is vertical and unstretched when the system is released to move. Find the velocity of the block A at the instant it breaks off the surface below it. Take

$$g = 10 \text{ m/s}^2 .$$



[Watch Video Solution](#)

51. one end of a spring of natural length h and spring constant k is fixed at the ground and the other is fitted with a smooth ring of

mass m which is allowed to slide on a horizontal rod fixed at a height h figure. Initially, the spring makes an angle of 37° with the vertical when the system is released from rest. find the speed of the ring when the spring becomes vertical.




Figure 8-E13



[Watch Video Solution](#)

52. Figure shows a light rod of length l rigidly attached to a small heavy block at one end and a hook at the other end. The system is released from rest with the rod in a horizontal position. There is a fixed smooth ring at a depth h below the initial position of the hook and the hook gets into the ring as it reaches there. What should be the minimum value of h so that the block moves in a complete circle

about the ring?




Figure 8-E14



Watch Video Solution

53. The bob of a pendulum at rest is given a sharp hit to impart a horizontal velocity $\sqrt{10gl}$ where l is the length of the pendulum. Find the tension in the string when a. the

string is horizontal. B. The bob is at its highest point and c. the string makes an angle of 60° with the upward vertical.



[Watch Video Solution](#)

54. A simple pendulum consists of a 50 cm long string connected to a 100 g ball. The ball is pulled aside so that the string makes an angle of 37° with the vertical and is then released. Find the tension in the string when

the bob is at its lowest position.



[Watch Video Solution](#)

55. Figure shows a smooth track, a part of which is a circle of radius R . A block of mass m is pushed against a spring of spring constant k fixed at the left end and is then released. Find the initial compression of the spring so that the block presses the track with a force

mg when it reaches the point P, where the radius of the track is horizontal.



[Watch Video Solution](#)

56. The bob of a stationary pendulum is given a sharp hit to impart it a horizontal speed of $\sqrt{3gl}$. Find the angle rotated by the string before it becomes slack.



[Watch Video Solution](#)

57. A heavy particle is suspended by a 1.5 m long string . It is given a horizontal velocity of $\sqrt{57} \frac{m}{s}$. a. Find the angle made by the string with the upward vertical, when it becomes slack. B. Find the speed of the particle at this instant. c. Find the maximum height reached by the particle over the point of suspension. Take

$$g = 10 \frac{m}{s^2}$$



[Watch Video Solution](#)


58. A simple pendulum of length L having a bob of mass m is deflected from its rest position by an angle θ and released figure. The string hits a peg which is fixed at distance x below the point of suspension and the bob starts going in a circle centred at the peg. a. Assuming that initially the bob has a height less than the peg, show that the maximum height reached by the bob equals its initial height. b. If the pendulum is released with $\theta = 90^\circ$ and $m = \frac{L}{2}$ find the maximum height reached by the bob above its lowest

position before the string becomes slack. c.

Find the minimum value of x/L for which the

bob goes in a complete circle about the pet

when the pendulum is released from $\theta = 90^\circ$.



Watch Video Solution

59. A particle slides on the surface of a fixed smooth sphere starting from the topmost point. Find the angle rotated by the radius through the particle, when it leaves contact with the sphere.



Watch Video Solution

60. A particle of mass m is kept on a fixed, smooth sphere of radius R at a position, where the radius through the particle makes an angle of 30° with the vertical. The particle is

released from this position. (a) What is the force exerted by the sphere on the particle just after the release? (b) Find the distance traveled by the particle before it leaves contact with the sphere.



[Watch Video Solution](#)

61. A particle of mass m is kept on the top of a smooth sphere of radius R . It is given a sharp impulse which imparts it a horizontal speed v .
a. find the normal force between the sphere


and the particle just after the impulse. B. What should be the minimum value of v for which the particle does not slip on the sphere? c. Assuming the velocity v to be half the minimum calculated in part, d. find the angle made by the radius through the particle with the vertical when it leaves the sphere.



Watch Video Solution

62. Figure shows a smooth track which consists of a straight inclined part of length l joining smoothly with the circular part. A particle of mass m is projected up the incline from its bottom. a. Find the minimum projected speed v_0 for which the particle reaches the top of the track. b. Assuming that the projection speed is $2v_0$ and that the block does not lose contact with the track before reaching its top, find the force acting on it when it reaches the top. c. Assuming that the projection speed is only slightly greater than

v_0 where will the block lose contact with the track?



[▶ Watch Video Solution](#)

63. A chain of length l and mass m lies on the surface of a smooth sphere of radius $R > l$ with one end tied to the top of the sphere.

a. Find the gravitational potential energy of the chain with reference level at the centre of the sphere. B. suppose the chain is released and slides down the sphere. Find the kinetic energy of the chain, when it has slid through an angle θ c. find the tangential acceleration $\frac{dv}{dt}$ of the chain when the chain starts sliding down.



Watch Video Solution

64. A smooth sphere of radius R is made to translate line with a constant acceleration $a=g$. A particle kept on the top of the sphere is released from there at zero velocity with respect to the sphere. Find the speed of the particle with respect to the sphere as a function of angle θ as it slides down.



Watch Video Solution

Question For Short Answer

1. When you lift a box from the floor and put it on an almirah the potential energy of the box increases, but there is no change in its kinetic energy. Is it a violation of conservation of energy?



Watch Video Solution

2. A particle is released from the top of an incline of height h . Does the kinetic energy of the particle at the bottom of the incline depend on the angle of incline? Do you need

any more information to answer this question
in Yes or No?



[Watch Video Solution](#)

3. Can the work by kinetic friction on an object
be positive? Zero?



[Watch Video Solution](#)

4. Can static friction do nonzero work on an
object? If yes give an example.If no, give

reason.



Watch Video Solution

5. Can normal force do a nonzero work on an object. If yes, give an example. If no, give reason.



Watch Video Solution

6. Can kinetic energy of a system be increased without applying any external force on the

system?



[Watch Video Solution](#)

7. Is work energy theorem valid in noninertial frames?



[Watch Video Solution](#)

8. A heavy box is kept on a smooth inclined plane and is pushed up by a force F acting parallel to the plane. Does the work done by

the force F as the box goes from A to B depend on how fast the box was moving at A and B? does the work by the force of gravity depend on this?



[Watch Video Solution](#)

9. One person says that the potential energy of a particular book kept in an almirah is 20 J and the other says it is 30J. Is one of them necessarily wrong?



[Watch Video Solution](#)

10. A book is lifted from the floor and is kept in an almirah. One person says that the potential energy of the book is increased by 20 J and the other says it is increased by 30 J. Is one of them necessarily wrong?



Watch Video Solution

11. In one of the exercises to strengthen the wrist and fingers, a person squeezes and releases a soft rubber ball. Is the work done

on the ball positive, negative or zero during compression? During expansion?



[Watch Video Solution](#)

12. When an apple falls from a tree what happens to its gravitational potential energy just as it reaches the ground? After it strikes the ground?



[Watch Video Solution](#)

13. When you push your bicycle up on an incline the potential energy of the bicycle and yourself increases. Where does this energy come from?



Watch Video Solution

14. The magnetic force on a charged particle is always perpendicular to its velocity. Can the magnetic force change the velocity of the particle? Speed of the particle?





[Watch Video Solution](#)

15. A ball is given a speed v on a rough horizontal surface. The ball travels through a distance l on the surface and stops. A. What are the initial and final kinetic energies of the ball? b. What is the work done by the kinetic friction?



[Watch Video Solution](#)

16. Consider the situation of the previous question from a frame moving with a speed v_0 parallel to the initial velocity of the block? b. What is the work done by the kinetic friction?



Watch Video Solution