



PHYSICS

BOOKS - HC VERMA PHYSICS (ENGLISH)

DISPERSION AND SPECTRA



1. Find the dispersive power of flint glass. The refractive indices of flint glass for red, yellow

and violet light are 1.613, 1.620 and 1.632 respectively.



2. The focal lengths of a thin lens for red and violet light are 90.0 cm and 86.4cm respectively. Find the dispersive power of the material of the lens. Make appropriate assumptions.



1. The refractive indices of filnt glass for red and violet light are 1.613 and 1.632 respectively. Find the angular dispersion prouduced by a thin prism of flint glass having refracting angle 5°

Watch Video Solution

2. A crown glass prism of angle 5° is to be combined with a flint glass prism in such a way

that the mean ray passes undeviated. Find (a) the angle of the flint glass prism needed and (b) the angular dispersion produced by the combination when white light goes through it. Refractive indices for red, yellow and viloet light are 1.514, 1.517 and 1.523 respectively for crown glass and 1.613 1.620 and 1.632 for flint glass.



3. The dispersive powers of crown and flint glasses are 0.03 and 0.05 respectively, The refractive indices for yellow light for these glasses are 1.517 and 1.621 respectively. It is desired to from an achromatic combination of prisms of crown and flint glasses which can produce a deviation of 1° in the yellow ray. Find the refracting angles of the two prisms needed.



1. The angular dispersion produced by a prism

A. increases if the average refractive index

increases

B. increases if the averge refractive index

decrease

C. remains constant whether the average

refractive index increases or decreases

D. has no relation with average refractive

index.

Answer: A



2. If a glass prism in dipped in water its dispersive power

A. increases

B. decreases

C. does not change

D. may increase or decrease depedig on

whether the angle of the prism is less

than or greater than 60°

Answer: B

Watch Video Solution

3. A prism can produce a minmum deviation δ in a the beam. If three such prisms are

combined, the minimum deviation that can be

produced in this beam is

A. 0

 $\mathsf{B.}\,2\delta$

 $\mathsf{C}.\,\delta$

D. `3 delta

Answer: B



4. Consider the following two statemets
Line spectra contain information about atoms.
Band spectra contain information about
molecules

A. Both A and B are wrong

B. A is correct but B is wrong

C. B is correct but A wrong

D. Both A and B are correct

Answer: D



5. The focal length of a converging lens are f_v and f_r for violet and red light respectively.

A. a.
$$f_v > f_r$$

- $\mathsf{B}.\,\mathsf{b}.f_v=f_r$
- C. c. $f_v \, < \, f_r$

D. d.Any of the three is possible depending

on the value of the average refractive index μ .







1. A narrow beam of white light goes through a slab having parallel faces.

A. a.The light never splits in different colours.

B. b.The emergent beam is white

C. c.the light inside the slab is split into

different colours

D. d.The light inside the slab is white

Answer: B::C

Watch Video Solution

2. By properly combining two prisms made of

different materials, it is possible

A. a.have	dispersior	n with	out	average
deviatio	on			
B. b.have deviation without dispersion				
C. c.have	both disp	ersion	and	average
deviatio	on			
D. d.have	neither dis	spersion	nor	average
deviatio	on.			

Answer: A::B::C

Watch Video Solution

3. In producing a pure spectrum, the incident light is passed through a narrow slit placed in the focal plane of an chromatic lens because a narrow slit

A. 1.a produces less diffraction

B. 2. increases intensity

C. 3.allows onlyone colour at a time

D. 4.allows more parallel beam when it

passes thorugh the lens

Answer: D



4. Which of the following quantities related to a lens depend on the wavelength or wavelengths of the incident light?

(Choose the incorrect option)

A. a.power

B. b.focal length

C. c.chromatic aberration

D. d.radii of curvature

Answer: A::B::C



5. Which of the following quantities increaes when wavelength is increased? Consider only the magnitudes

- A. 1.the power of a converging lens
- B. 2.the focal length of a converging lens
- C. 3.the power of a diverging lens
- D. 4.the focal length of a diverging lens.

Answer: B::D



Exercises

1. A flint glass prism and a crown glass prism are to be combined in such a way that the deviation of the mean ray is zero. The refractive index of flint and crown glasses for the mean ray are 1.620 and 1.518 respectively. If the refracting angle of the flint prism is 6.0° , what would be the refracting angle of the

crown prism?



2. A certain material has refractive indices 1.56,
1.60 and 1.68 for red, yellow and violet light
respectively. (a) Calculate the dispersive power.
(b) Find the angular dispersion produced by a
thin prism of angle 6° made of this material.



3. The focal lengths of a convex lens for red, yellow and violet rays are 100 cm, 98 cm and 96 cm respectively. Find the dispersive power of

the material of the lens.



4. The refractive index of a material changes by 0.014 as the colour of the light changes from red to violet. A rectangular slab of height 2.00 cm made of this material is placed on a newspaper. When viewed normally in yellow light, the letters appear 1.32 cm below the top

surface of the slab. Calculate the dispersive

power of the material.

Watch Video Solution

5. A thin prism is made of a material having refractive indices 1.61 and 1.65 for red and violet light. The dispersive power of the material is 0.07. It is found that a beam of yellow light passing through the prism suffers

a minimum deviation of 4.0° in favourable

conditions. Calculate the angle of the prism.



6. The minimum deviations suffered by red, yellow and violet beam passing through an equilateral transparent prism are 38.4° , 38.7° and 39.2° respectively. Calculate the dispersive power of the medium.



7. Two prisms of identical geometrical shape are combined with their refracting angles oppositely directed. The materials of the prisms have refractive indices 1.52 and 1.62 for violet light. A violet ray is deviated by 1.0° when passes symmetrically through this combination. What is the angle of the prisms?

Watch Video Solution

8. Three thin prisms are combined as shown in figure. The refractive indices of the crown

glass for red, yellow and violet rays are μ_r , μ_y and μ_v respectively and those for the flint glass are μ'_r , μ'_y and μ'_v respectively. Find the ratio A'/A for which (a) there is no net angular dispersion, and (b) there is no net deviation in the yellow ray.





9. A thin prism of crown glass $(\mu_r = 1.515, \mu_v = 1.525)$ and a thin prism of flint glass $(\mu_r = 1.612, \mu_v = 1.632)$ are placed in contact with each other. Their refracting angles are 5.0° each and are similarly directed. Calculate the angular dispersion produced by the combination.

Watch Video Solution

10. A thin prism of angle $6.0^{\,\circ}\,,\,\omega^{\,\prime}\,=\,0.07\, ext{ and }\,\mu_y^{\,\prime}\,=\,1.50$ is combined with another thin prism having $\omega=0.08$ and $\mu_y=1.60$. The combination produces no deviation in the mean ray. (a) Find the angle of the second prism. (b) Find the net angular dispersion produced by the combination when a beam of white light passes through it. (c) If the prisms are similarly directed, what will be the deviation in the mean ray ? (d) Find the angular dispersion in the situation described in (c).

11. The refractive index of a material M_1 changes by 0.014 and that of another material M_2 changes by 0.024 as the colour of the light is changed from red to violet. Two thin prisms one made of $M_1(A=5.3^\circ)$ and other made of $M_2(A=3.7^\circ)$ are combined with their refracting angles oppositely directed.

(a) Find the angular dispesion produced by the combination.

(b) the prisms are now combined with their

refracting angles similarly directed. Find the angular dispersion produced by the combination.

Watch Video Solution

Question For Short Answer

1. The equation $\omega = \frac{\mu_v - \mu_r}{\mu - 1}$ was derived for a prism having small refracting angle. Is it also valid for a prism of large refracting angle? Is it also valid for a glass slab or a glass sphere?



Watch Video Solution

2. Can the dispersive power $\omega = \frac{\mu_v - \mu_r}{\mu - 1}$ be negative? What is the sign of ω if a hollow prism is immersed into water?

3. If three identical prisms are combined is it possible to pass a beam that emerges undeviated? Undispersed?





4. Monochromatic light shold be used to produce pure specturm. Comment on this statement.

Watch Video Solution

5. Does focal length of a lens depend on the colour often light used? Does focal length of a mirror depend on the colour?

Watch Video Solution

6. Suggest a method to produce a rainbow in

your house.

Watch Video Solution