



PHYSICS

NCERT - NCERT PHYSICS(TELUGU)

KINETIC THEORY

Example

1. The density of water is 1000kgm^{-3} . The density of water vapour at 100°C and 1 atm pressure is 0.6kgm^{-3} . The volume of a

molecule multiplied by the total number gives ,what is called, molecular volume. Estimate the ratio (or fraction) of the molecular volume to the total volume occupied by the water vapour under the above conditions of temperature and pressure.



Watch Video Solution

2. Estimate the volume of a water molecule using the data in Example 13.1.



View Text Solution

3. What is the average distance between atoms (interatomic distance) in water? Use the data given in Examples 13.1 and 13.2.



[View Text Solution](#)

4. A vessel contains two non reactive gases : neon (monatomic) and oxygen (diatomic). The ratio of their partial pressures is 3:2. Estimate the ratio of (i) number of molecules and (ii) mass density of neon and oxygen in the vessel.

Atomic mass of Ne = 20.2 u, molecular mass of

$$O_2 = 32.0u.$$



[Watch Video Solution](#)

5. A flask contains argon and chlorine in the ratio of 2:1 by mass. The temperature of the mixture is $27^\circ C$. Obtain the ratio of (i) average kinetic energy per molecule, and (ii) root mean square speed v_{rms} of the molecules of the two gases. Atomic mass of argon = 39.9 u, Molecular mass of chlorine = 70.9 u.



[Watch Video Solution](#)

6. Uranium has two isotopes of masses 235 and 238 units. If both are present in Uranium hexafluoride gas which would have the larger average speed ? If atomic mass of fluorine is 19 units, estimate the percentage difference in speeds at any temperature.



[Watch Video Solution](#)

7. (a) When a molecule (or an elastic ball) hits a (massive) wall, it rebounds with the same speed. When a ball hits a massive bat held firmly, the same thing happens. However, when the bat is moving towards the ball, the ball rebounds with a different speed. Does the ball move faster or slower?

(b) When gas in a cylinder is compressed by pushing in a piston, its temperature rises. Guess at an explanation of this in terms of kinetic theory using (a) above.

(c) What happens when a compressed gas

pushes a piston out and expands. What would you observe ?

(d) Sachin Tendulkar used a heavy cricket bat while playing. Did it help him in anyway ?



[View Text Solution](#)

8. A cylinder of fixed capacity 44.8 litres contains helium gas at standard temperature and pressure. What is the amount of heat needed to raise the temperature of the gas in

the cylinder by $15.0^\circ C$? (

$$R = 8.31 J mol^{-1} K^{-1})$$



[Watch Video Solution](#)

9. Estimate the mean free path for a water molecule in water vapour at 373 K.



[Watch Video Solution](#)

Exercises

1. Estimate the fraction of molecular volume to the actual volume occupied by oxygen gas at STP. Take the diameter of an oxygen molecule to be 3\AA .




[Watch Video Solution](#)

2. Molar volume is the volume occupied by 1 mole of any (ideal) gas at standard temperature and pressure (STP : 1 atmospheric pressure , $0^\circ C$) . Show that it is 22.4 litres.



3. Figure Show plot of PV/T versus P for 1.00×10^{-3} kg of oxygen gas at two different temperatures.



(a) What does the dotted plot signify?

(b) Which is true: $T_1 > T_2$ or $T_1 < T_2$?

(c) What is the value of PV/T where the

curves meet on the y-axis?

(d) If we obtained similar plots for 1.00×10^{-3} kg of hydrogen, would we get the same value of PV/T at the point where the curves meet on the y-axis? If not, what mass of hydrogen yields the same value of PV/T (for low pressure high temperature region of the plot) ? (Molecular mass of $\text{H}_2 = 2.02$ u, of $\text{O}_2 = 32.0$ u, $R = 8.31 \text{ J mol}^{-1} \text{ K}^{-1}$.)



[Watch Video Solution](#)

4. An oxygen cylinder of volume 30 litres has an initial gauge pressure of 15 atm and a temperature of $27^{\circ}C$. After some oxygen is withdrawn from the cylinder, the gauge pressure drops to 11 atm and its temperature drops to $17^{\circ}C$. Estimate the mass of oxygen taken out of the cylinder ($R = 8.31 J mol^{-1} K^{-1}$. molecular mass of $O_2 = 32u$).



[Watch Video Solution](#)

5. An air bubble of volume 1.0cm^3 rises from the bottom of a lake 40m deep at a temperature of 12°C . To what volume does it grow when it reaches the surface, which is at a temperature of 35°C ?



[Watch Video Solution](#)

6. Estimate the total number of air molecules (inclusive of oxygen, nitrogen, water vapour and other constituents) in a room of capacity

25.0m^3 at a temperature of 27°C and 1 atm pressure.



[Watch Video Solution](#)

7. Estimate the average thermal energy of a helium atom at (i) room temperature (27°C), (ii) the temperature on the surface of the Sun (6000 K), (iii) the temperature of 10 million kelvin (the typical core temperature in the case of a star).



[Watch Video Solution](#)

8. Three vessels of equal capacity have gases at the same temperature and pressure. The first vessel contains neon (monatomic), the second contains chlorine (diatomic), and the third contains uranium hexafluoride (polyatomic). Do the vessels contain equal number of respective molecules? Is the root mean square speed of molecules the same in the three cases? If not, in which case is v_{rms} the largest?



Watch Video Solution

9. At what temperature is the root mean square speed of an atom in an argon gas cylinder equal to the rms speed of a helium gas atom at -20°C ? (atomic mass of $\text{Ar} = 39.9u$, of $\text{He} = 4.0u$).



[Watch Video Solution](#)

10. Estimate the mean free path and collision frequency of a nitrogen molecule in a cylinder containing nitrogen at 2.0 atm and

temperature 17°C . Take the radius of a nitrogen molecule to be roughly 1.0 \AA . Compare the collision time with the time the molecule moves freely between two successive collisions (Molecular mass of $\text{N}_2 = 28.0\text{ u}$).



[Watch Video Solution](#)

Additional Exercise

1. A metre long narrow bore held horizontally (and closed at one end) contains a 76 cm long

mercury thread, which traps a 15 cm column of air. What happens if the tube is held vertically with the open end at the bottom ?



[View Text Solution](#)

2. From a certain apparatus, the diffusion rate of hydrogen has an average value of $28.7\text{cm}^3\text{s}^{-1}$. The diffusion of another gas under the same conditions is measured to have an average rate of $7.2\text{cm}^3\text{s}^{-1}$. Identify the gas. [Hint : Use Graham's law of diffusion:

$R_1 / R_2 = (M_2 / M_1)^{1/2}$, where R_1, R_2 , are diffusion rates of gases 1 and 2, and M_1 and M_2 their respective molecular masses. The law is a simple consequence of kinetic theory.]



[Watch Video Solution](#)

3. Given below are densities of some solids and liquids. Give rough estimates of the size of their atoms :

Substance	Atomic Mass (u)	Density (10^3 Kg m^{-3})
Carbon (diamond)	12.01	2.22
Gold	197.00	19.32
Nitrogen (liquid)	14.01	1.00
Lithium	6.94	0.53
Fluorine (liquid)	19.00	1.14

[Hint : Assume the atoms to be 'tightly packed' in a solid or liquid phase, and use the known value of Avogadro's number. You should, however, not take the actual numbers you obtain for various atomic sizes too literally. Because of the crudeness of the tight packing approximation, the results only indicate that atomic sizes are in the range of a few Å].



Watch Video Solution