



## PHYSICS

### NCERT - NCERT PHYSICS(TELUGU)

#### THERMAL PROPERTIME OF MATTER

##### Example

1. Show that the coefficient of area expansion ,  $(\Delta A / A) / \Delta T$  , of a rectangular sheet of the solid is twice its linear expansivity ,  $\alpha_1$ .

 [Watch Video Solution](#)

2. A blacksmith fixes Iron ring on the rim of the wooden wheel of a horse cart. The diameter of the rim and the fron ring are

5.243m and 5.231m, respectively at  $27^{\circ}C$ . To what temperature should the ring be heated so as to fit the rim of the wheel?

 [Watch Video Solution](#)

3. A sphere of 0.047 kg aluminium is placed for sufficient time in a vessel containing boiling water, so that the sphere is at  $100^{\circ}C$ . It is then immediately transferred to 0.14 kg copper calorimeter containing 0.25 kg water at  $20^{\circ}C$ . The temperature of water rises and attains a steady state at  $23^{\circ}C$ . Calculate the specific heat capacity of aluminium.

 [Watch Video Solution](#)

4. When 0.15kg of ice at  $0^{\circ}C$  is mixed with 0.30 kg of water at  $50^{\circ}C$  in a container, the resulting temperature is  $6.7^{\circ}C$ . Calculate the heat of fusion of ice. ( $S_{\text{water}} = 4186 J kg^{-1} K^{-1}$ )




Watch Video Solution

5. Calculate the heat required to convert 3 kg of ice at  $-12^{\circ}C$  kept in a calorimeter to steam at  $100^{\circ}C$  at atmospheric pressure. Given specific heat capacity of ice  $= 2100 Jkg^{-1}K^{-1}$ , specific heat capacity of water  $= 4186 Jkg^{-1}K^{-1}$ , latent heat of fusion of ice  $= 3.35 \times 10^5 Jkg^{-1}$  and latent heat of steam  $= 2.256 \times 10^8 Jkg^{-1}$ .



Watch Video Solution

6. What is the temperature of the steel - copper junction in the steady state of the system shown in Fig.11.15.




Length of the steel rod = 15.0 cm, length of the copper rod = 10.0 cm, temperature of the furnace =  $300^{\circ}\text{C}$ , temperature of the other end =  $0^{\circ}\text{C}$ . The area of cross section of the steel rod is twice that of the copper rod. (Thermal conductivity of steel =  $50.2\text{Js}^{-1}\text{m}^{-1}\text{K}^{-1}$ , and of copper =  $385\text{Js}^{-1}\text{m}^{-1}\text{K}^{-1}$ ).



[Watch Video Solution](#)

7. An iron bar ( $L_1 = 0.1\text{m}$ ,  $A_1 = 0.02\text{m}^2$ ,  $K_1 = 79\text{Wm}^{-1}\text{K}^{-1}$ ) and a brass bar ( $L_2 = 0.1\text{m}$ ,  $A_2 = 0.02\text{m}^2$ ,  $K_2 = 109\text{Wm}^{-1}\text{K}^{-1}$ ) are soldered end to end as shown in Fig. 11.16. The free ends of the iron bar and brass bar are maintained at 373 K and 273 K respectively. Obtain expressions for and hence compute (i) the temperature of the junction of the two bars, (ii) the equivalent thermal conductivity of the compound bar, and (iii) the heat current through the

compound bar.



[Watch Video Solution](#)

8. A pan filled with hot food cools from  $94^\circ\text{C}$  to  $86^\circ\text{C}$  in 2 minutes when the room temperature is at  $20^\circ\text{C}$ . How long will it take to cool from  $71^\circ\text{C}$  to  $69^\circ\text{C}$ ?

[Watch Video Solution](#)

## Exercises

1. The triple points of neon and carbon dioxide are  $24.57\text{ K}$  and  $216.55\text{ K}$  respectively. Express these temperatures

on the Celsius and Fahrenheit scales.



[Watch Video Solution](#)

2. Two absolute scales A and B have triple points of water defined to be 200 A and 350 B. What is the relation between  $T_A$  and  $T_B$ ?



[Watch Video Solution](#)

3. The electrical resistance in ohms of a certain thermometer varies with temperature according to the approximate law:

$$R = R_0[1 + \alpha(T - T_0)]$$

The resistance is  $101.6\Omega$  at the triple-point of water  $273.16K$ , and  $165.5\Omega$  at the normal melting point of lead ( $600.5K$ ). What is the temperature when the resistance is  $123.4\Omega$ ?



[Watch Video Solution](#)

4. The triple-point of water is a standard fixed point in modern thermometry.

Why? What is wrong in taking the melting point of Ice and the boiling point of water as standard fixed points as was originally done in the Celsius scale) ?

 [Watch Video Solution](#)

5. There were two fixed points in the original Celsius scale as mentioned above which were assigned the number  $0^{\circ}C$  and  $100^{\circ}C$  respectively. On the absolute scale, one of the fixed points is the triple-point of water, which on the Kelvin absolute scale is assigned the number 273.16K What is the other fixed point on this (Kelvin) scale ?

 [Watch Video Solution](#)

6. The absolute temperature (Kelvin scale)  $T$  is related to the temperature  $t_c$  on the Celsius scale by  $t_c = T - 273.15$

Why do we have 273.15 in this relation, and not 273.16?

 [Watch Video Solution](#)

7. What is the temperature of the triple-point of water on an absolute scale whose unit interval size is equal to that of the Fahrenheit scale ?

 [Watch Video Solution](#)

8. Two ideal gas thermometers A and B use oxygen and hydrogen respectively. The following observations are made:

Temperature	Pressure thermometer A	Pressure thermometer B
Triple - point water	$1.250 \times 10^5 Pa$	$0.200 \times 10^5 Pa$
Normal melting point of sulphur	$1.797 \times 10^5 Pa$	$0.287 \times 10^5 Pa$



(a) What is the absolute temperature of normal melting point of sulphur as read by thermometers A and B ?

(b) What do you think is the reason behind the slight difference in answers of thermometers A and B ? (The thermometers are not faulty). What further procedure is needed in the experiment to reduce the discrepancy between the two readings ?

 [View Text Solution](#)

9. A steel tape 1m long is correctly calibrated for a temperature of  $27.0^{\circ}C$ . The length of a steel rod measured by this tape is found to be 63.0 cm on a hot day when the temperature is  $45.0^{\circ}C$ . What is the actual length of the steel rod on that day? What is the length of the same steel rod on a day when the temperature is  $27.0^{\circ}C$  ? Coefficient of linear expansion of steel  $= 1.20 \times 10^{-5} K^{-1}$ .

 [Watch Video Solution](#)

**10.** A large steel wheel is to be fitted on to a shaft of the same material. At  $27^{\circ}C$ , the outer diameter of the shaft is 8.70 cm and the diameter of the central hole in the wheel is 8.69cm. The shaft is cooled using dry Icel. At what temperature of the shaft does the wheel slip on the shaft? Assume coefficient of linear expansion of the steel to be constant over the required temperature range:

$$\alpha_{\text{steel}} = 1.20 \times 10^{-5} K^{-1}.$$

 [Watch Video Solution](#)

**11.** A hole is drilled in a copper sheet. The diameter of the hole is 4.24 cm at  $27.0^{\circ}C$ . What is the change in the diameter of the hole when the sheet is heated to  $227^{\circ}C$ ? Coefficient of linear expansion of copper =  $1.70 \times 10^{-5} K^{-1}$ .

 [Watch Video Solution](#)

**12.** A brass wire 1.8 m long at  $27^\circ C$  is held taut with little tension between two rigid supports. If the wire is cooled to a temperature of  $-39^\circ C$ . what is the tension developed in the wire, if its diameter is 2.0 mm ? Co-efficient of linear expansion of brass  $= 2.0 \times 10^{-5} K^{-1}$ , Young's modulus of brass  $= 0.91 \times 10^{11}$  Pa.



[Watch Video Solution](#)

**13.** A brass rod of length 50 cm and diameter 3.0 mm is joined to a steel rod of the same length and diameter. What is the change in length of the combined rod at  $250^\circ C$ . if the original lengths are at  $40.0^\circ C$ ? Is there a "thermal stress developed at the junction ? The ends of the rod are free to expand (Co-efficient of linear expansion of brass  $= 2.0 \times 10^{-5} K^{-1}$ , steel  $= 1.2 \times 10^{-5} K^{-1}$ ).



[Watch Video Solution](#)

14. The coefficient of volume expansion of glycerine is  $49 \times 10^{-5} K^{-1}$ . What is the fractional change in its density for a  $30^\circ C$  rise in temperature ?

 [Watch Video Solution](#)

15. A 10 kW drilling machine is used to drill a bore in a small aluminium block of mass 8.0kg. How much is the rise in temperature of the block in 2.5 minutes, assuming 50% of power is used up in heating the machine itself or lost to the surroundings. Specific heat of aluminium =  $0.91 J g^{-1} K^{-1}$ .

 [Watch Video Solution](#)

16. A copper block of mass 2.5 kg is heated in a furnace to a temperature of  $500^\circ C$  and then placed on a large ice block. What

is the maximum amount of ice that can melt? (Specific heat of copper =  $0.39 J g^{-1} K^{-1}$ , heat of fusion of water =  $335 J g^{-1}$ ).

 [Watch Video Solution](#)

**17.** In an experiment on the specific heat of a metal, a  $0.20 \text{ kg}$  block of the metal at  $150^\circ C$  is dropped in a copper calorimeter (of water equivalent  $0.025 \text{ kg}$  containing  $150 \text{ cm}^3$  of water at  $27^\circ C$ ). The final temperature is  $40^\circ C$ . Compute the specific heat of the metal. If heat losses to the surroundings are not negligible. Is your answer greater or smaller than the actual value for specific heat of the metal ?

 [Watch Video Solution](#)

**18.** Given below are observations on molar specific heats at room temperature of some common gases.

Gas	Molar specific heat ( $C_v$ ) ( $\text{cal mol}^{-1} \text{K}^{-1}$ )
Hydrogen	4.87
Nitrogen	4.97
Oxygen	5.02
Nitric oxide	4.99
Carbon monoxide	5.01
Chlorine	6.17

The measured molar specific heats of these gases are markedly different from those for monatomic gases. Typically, molar specific heat of a monatomic gas is  $2.52 \text{ cal/mol K}$ . Explain this difference.

What can you infer from the somewhat larger than the rest) value for chlorine ?

 [Watch Video Solution](#)

**19.** A child running a temperature of  $101^\circ \text{F}$  is given an antipyrim (1.e. a medicine that lowers fever) which causes an increase in the rate of evaporation of sweat from his body. If the fever is brought down to  $98^\circ \text{F}$  in 20 minutes, what is the average rate of extra

evaporation caused by the drug. Assume the evaporation mechanism to be the only way by which heat is lost. The mass of the child is 30 kg. The specific heat of human body is approximately the same as that of water, and latent heat of evaporation of water at that temperature is about  $580 \text{ cal g}^{-1}$ .

 [Watch Video Solution](#)

**20.** A 'thermacole' icebox is a cheap and an efficient method for storing small quantities of cooked food in summer in particular. A cubical icebox of side 30 cm has a thickness of 5.0 cm. If 4.0 kg of ice is put in the box, estimate the amount of ice remaining after 6 h. The outside temperature is  $45^\circ \text{C}$  and coefficient of thermal conductivity of thermacole is  $0.01 \text{ Js}^{-1} \text{ m}^{-1} \text{ K}^{-1}$ . [Heat of fusion of water =  $335 \times 10^3 \text{ J kg}^{-1}$ ]

 [Watch Video Solution](#)

21. A brass boiler has a base area of  $0.15\text{m}^2$  and thickness  $1.0\text{ cm}$ . It boils water at the rate of  $6.0\text{ kg/min}$  when placed on a gas stove. Estimate the temperature of the part of the flame in contact with the boiler. Thermal conductivity of brass  $= 109\text{Js}^{-1}\text{m}^{-1}\text{K}^{-1}$ , Heat of vaporisation of water  $= 2256 \times 10^3\text{Jkg}^{-1}$ .

 [Watch Video Solution](#)

22. A body cools from  $80^\circ\text{C}$  to  $50^\circ\text{C}$  in 5 minutes. Calculate the time it takes to cool from  $60^\circ\text{C}$  to  $30^\circ\text{C}$ . The temperature of the surroundings is  $20^\circ\text{C}$ .

 [Watch Video Solution](#)

23. Answer the following questions based on the P-T phase diagram of carbon dioxide:



At what temperature and pressure can the solid, liquid and vapour phases of  $CO_2$  co-exist in equilibrium ?

 [Watch Video Solution](#)

24. Answer the following questions based on the P-T phase diagram of carbon dioxide:

What is the effect of decrease of pressure on the fusion and boiling point of  $CO_2$  ?

 [Watch Video Solution](#)

25. Answer the following questions based on the P-T phase diagram of carbon dioxide:

What are the critical temperature and pressure for  $CO_2$ ? What is their significance ?

 [Watch Video Solution](#)

**26.** Answer the following questions based on the P-T phase diagram of carbon dioxide:

Is  $CO_2$  solid, liquid or gas at (a)  $-70^\circ C$  under 1 atm, (b)  $-60^\circ C$  under 10 atm, (c)  $15^\circ C$  under 56 atm?

 [Watch Video Solution](#)

**27.** Answer the following questions based on the P-T phase diagram of  $CO_2$ :

$CO_2$  at 1 atm pressure and temperature  $-60^\circ C$  is compressed isothermally Does it go through a liquid phase ?

 [Watch Video Solution](#)

**28.** Answer the following questions based on the P-T phase diagram of  $CO_2$ :

What happens when  $CO_2$  at 4 atm pressure is cooled from room temperature at constant pressure ?

 [Watch Video Solution](#)

**29.** Answer the following questions based on the P-T phase diagram of  $CO_2$ :

Describe qualitatively the changes in a given mass of solid  $CO_2$  at 10 atm pressure and temperature  $-65^\circ C$  as it is heated up to room temperature at constant pressure.

 [Watch Video Solution](#)

**30.** Answer the following questions based on the P-T phase diagram of  $CO_2$ :

$CO_2$  is heated to a temperature  $70^\circ C$  and compressed isothermally. What changes in its properties do you expect to observe ?



**Watch Video Solution**