



PHYSICS

NCERT - NCERT PHYSICS(TELUGU)

UNITS AND MEASUREMENT

Example

1. Calculate the angle of (a) 1° (degree) (b) $1'$ (minute of arc or arcmin) and (c) $1''$ (second of arc or arc second) in radians. Use $360^\circ = 2\pi$ rad, $1^\circ = 60'$ and $1' = 60''$.




[Watch Video Solution](#)

2. A man wishes to estimate the distance of a nearby tower from him. He stands at a point A in front of the tower C and spots a very distant object in line with AC. He then walks perpendicular to AC up to B, a distance of 100 m, and looks at O and C again. Since O is very distant, the direction BO is practically the same as AO, but he finds the line of sight of C shifted from the original line of sight by an angle = 40° (θ is known as 'parallax') estimate the distance of the tower C from his

original position A.

D is tant t
object O



 Watch Video Solution

3. From two diametrically opposite points A and B on earth, moon is observed. The angle θ subtended at the moon by the two directions of observation is $1^\circ 54'$. If radius of earth is 0.638×10^7 m, find the distance of the moon from the earth.

 [Watch Video Solution](#)

4. The Sun's angular diameter is measured to be $1920''$. The distance D of the Sun from the Earth is 1.496×10^{11} m. What is the diameter of the Sun ?

 [Watch Video Solution](#)

5. If the size of a nucleus (in the range of $10^{-15} \text{ m} \rightarrow 10^{-14} \text{ m}$) is scaled up to the tip of a sharp pin, what roughly is the size of an atom? Assume tip of the pin to be in the range 10^{-5} to 10^{-4} m .



[Watch Video Solution](#)

6. Two clocks are being tested against a standard clock located in a national laboratory. At 12:00:00 noon by the standard clock, the readings of the two clocks are:

	Clock 1	Clock 2
Monday	12:00:05	10:15:06
Tuesday	12:01:15	10:14:59
Wednesday	11:59:08	10:15:18
Thursday	12:01:50	10:15:07
Friday	11:59:15	10:14:53
Saturday	12:01:30	10:15:24
Sunday	12:01:19	10:15:11

If you are doing an experiment that requires precision time interval measurements, which of the two clocks will you prefer ?

 [View Text Solution](#)

7. The readings of a length come out to be 2.63 m , 2.56 m , 2.42 m , 2.71 m and 2.80 m . Calculate the absolute errors and relative errors or percentage errors. What do you think of the actual value of the length and its limits ?

 [Watch Video Solution](#)

8. The temperature of two bodies measured by a thermometer are $t_1 = 20^\circ C \pm 0.5^\circ C$ and $t_2 = 50^\circ C \pm 0.5^\circ C$. Calculate the temperature difference and the error therein.

 [Watch Video Solution](#)

9. The resistance $R=V/I$ where $V=(100 \pm 5)V$ and $I=(10 \pm 0.2)A$. Find the percentage error in R .

 [Watch Video Solution](#)

10. Two resistors of resistances $R_1 = 100 \pm 3$ ohm and $R_2 = 200 \pm 4$ ohm are connected (a) in series , (b) in parallel. Find the equivalent resistance of the (a) series combination , (b) parallel combination . Use for (a) the

relation $R = R_1 + R_2$ and $\frac{\Delta R'}{R'^2} = \frac{\Delta R_1}{R_1^2} + \frac{\Delta R_2}{R_2^2}$ for (b)

$$\frac{1}{R} = \frac{1}{R_1} + \frac{1}{R_2} \text{ and } \frac{\Delta R'}{R'^2} = \frac{\Delta R_1}{R_1^2} + \frac{\Delta R_2}{R_2^2}$$

 [Watch Video Solution](#)

11. Find the relative error in Z, if $Z = A^4 \frac{B^{\frac{1}{3}}}{C} D^{\frac{3}{2}}$.

 [Watch Video Solution](#)

12. The period of oscillation of a simple pendulum is $T =$

$$2\pi\sqrt{\frac{L}{g}}.$$

Measured value of L is 20.0 cm known to 1mm

accuracy and time for 100 oscillations of the pendulum

is found to be 90s using a wrist watch of 1 s resolution.

What is the accuracy in the determination of g ?



[Watch Video Solution](#)

13. Each side of a cube is measured to be 7.203m. What

is (i) the total surface area and (ii) the volume of the

cube to appropriate significant figures ?



[Watch Video Solution](#)

14. Find the number of significant figures in the 6.0023

 [Watch Video Solution](#)

15. Let us consider an equation $\frac{1}{2}mv^2 = mgh$

Where m is the mass of the body. v its velocity, g is the acceleration due to gravity and h is the height. Check whether this equation is dimensionally correct.

 [Watch Video Solution](#)

16. The SI unit of energy is $J = kgm^2s^{-2}$, that of speed v is ms^{-1} and of acceleration a is ms^{-2} . What of the

formulae for kinetic energy (k) given below can you rule out on the basis of dimensional arguments (m stands for the mass of the body) :

(a) $K = m^2v^2$

(b) $K = \left(\frac{1}{2}\right)mv^2$

(c) $K = ma$

(d) $K = \left(\frac{3}{16}\right)mv^2$

(e) $K = \left(\frac{1}{2}\right)mv^2 + ma$



[Watch Video Solution](#)

17. Consider a simple pendulum, having a bob attached to a string, that oscillates under the action of the force of gravity. Suppose that the period of oscillation of the

simple pendulum depends on its length (l), mass of the bob (m) and acceleration due to gravity (g). Derive the expression for its time period using method of dimensions.



[Watch Video Solution](#)

Exercise Fill In The Blanks

1. The volume of a cube of side 1 cm is equal to m^3 .



[Watch Video Solution](#)

2. The surface area of a solid cylinder of radius 2.0 cm and height 10.0 cm is equal to $(mm)^2$

 [Watch Video Solution](#)

3. A vehicle moving with a speed of $18kmh^{-1}$ covers.....m in 1 s.

 [Watch Video Solution](#)

4. The relative density of lead is 11.3. Its density is gcm^{-3} or kgm^{-3} .

 [Watch Video Solution](#)

$$5. 1kgm^2s^{-2} = \dots\dots\dots gcm^2s^{-2}$$

 [Watch Video Solution](#)

$$6. 1 m \dots\dots\dots ly$$

 [Watch Video Solution](#)

$$7. 3.0ms^{-2} = \dots\dots\dots Kmh^{-2}$$

 [Watch Video Solution](#)

8.

$$G = 6.67 \times 10^{-11} Nm^2(kg)^{-2} = \dots\dots\dots (cm)^3 s^{-2} g^{-1}$$

.



Watch Video Solution

Exercise

1. 1 calorie=4.2 J where $1J = 1 \text{ kg } m^2 s^{-2}$. Suppose we employ a system of units in which the unit of mass is α kg, the unit of length is β m and the unit of time γ s, show that a calorie has a magnitude $4.2\alpha^{-1}\beta^{-2}\gamma^{-2}$ in the new system.



Watch Video Solution



2. Explain this statement clearly :

"To call a dimensional quantity 'large' or 'small' is meaningless without specifying a standard for comparison". In view of this, reframe the following statements wherever necessary:

(a) atoms are very small objects

(b) a jet plane moves with great speed

(c) the mass of jupiter is very large

(d) the air inside this room contains a large number of molecules

(e) a proton is much more massive than an electron

(f) the speed of sound is much smaller than the speed of light.



[Watch Video Solution](#)

3. A new unit of length is chosen so that the speed of light in vacuum is 1ms^{-2} . If light takes 8 min and 20 s to cover this distance, what is the distance between the Sun and Earth in terms of the new unit?



[Watch Video Solution](#)

4. Which of the following is the most precise device for measuring length :

(a) a vernier callipers with 20 divisions on the sliding scale

(b) a screw gauge of pitch 1 mm and 100 divisions on the circular scale.

(c) an optical instrument that can measure length to within a wavelength of light ?



[Watch Video Solution](#)

5. A student measures the thickness of a human hair using a microscope of magnification 100. He makes 20 observations and find that the average thickness (as viewed in the microscope) is 3.5 mm. What is the estimate of the thickness of hair?



[Watch Video Solution](#)



[Watch Video Solution](#)

6. A screw gauge has a pitch of 1.0mm and 200 divisions on the circular scale. Do you think it is possible to increase the accuracy of the screw gauge arbitrarily by increasing the number of divisions on the circular scale ?



[Watch Video Solution](#)

7. The mean diameter of a thin brass rod is to be measured by vernier callipers. Why is a set of 100 measurements of the diameter expected to yield a

more reliable estimate than a set of 5 measurements only?



[Watch Video Solution](#)

8. The photograph of a house occupies an area of 1.75cm^2 on a 35 mm side. The slide is projected on to a screen, and the area of the house on the screen is 1.55m^2 . What is the linear magnification of the projector-screen arrangement.



[Watch Video Solution](#)

9. State the number of significant figures in the 0.007m^2

 [Watch Video Solution](#)

10. State the number of significant

$$0.0006032m^2$$

 [Watch Video Solution](#)

11. State the number of significant figures in the

$$0.2370gcm^{-3}$$

 [Watch Video Solution](#)

12. State the number of significant figures in the $6.320J$



Watch Video Solution

13. State the number of significant figures in the following

6.032



Watch Video Solution

14. State the number of significant

$0.0006032m^2$



Watch Video Solution

15. The length , breadth and thickness of a rectangular sheet of metal are 4.234 m, 1.005 m, and 2.01 cm respectively. Give the area and volume of the sheet to correct significant figures.



Watch Video Solution

16. The mass of a box measured by a grocer's balance is 2.30 kg. Two gold pieces of masses 20.15 g and 20.17 g are added to the box. What is (a) the total mass of the box, (b) the difference in the masses of the pieces to correct significant figures ?



Watch Video Solution

17. A physical quantity P is related to four observables a,b,c and d as follows:

$$P = a^3 \frac{b^2}{\sqrt{cd}}$$

The percentage errors of measurement in a,b, c and d are 1%, 3% , 4% and 2% respectively. What is the percentage error in the quantity P ? If the value of P calculated using the above relation turns out to be 3.763, to what value should you round off the result ?



[Watch Video Solution](#)

18. A book with many printing errors contains four different formulas for the displacement y of a particle

undergoing a certain periodic motion :

(a) $y = a \sin 2\pi t / T$

(b) $y = a \sin vt$

(c) $y = (a / T) \sin t / a$

(d) $y = (a / \sqrt{2}) (\sin 2\pi t / T + \cos 2\pi t / T)$

(a = maximum displacement of the particle, v=speed of the particle. T=time-period of motion). Rule out the wrong formulas on dimensional grounds)



[Watch Video Solution](#)

19. A famous relation in physics relates 'moving mass' m to the 'rest mass' m_0 of a particle in terms of its speed v and the speed of light, c . (This relation first arose as a

consequence of special relativity due to Albert Einstein). A boy recalls the relation almost correctly but forgets where to put the constant c . He writes:

$$m = \frac{m_0}{(1 - v^2)^{1/2}}$$

Guess where to put the missing c .

 [Watch Video Solution](#)

20. The unit of length convenient on the atomic scale is known as an angstrom and is denoted by \AA : $1\text{\AA} = 10^{-10}m$. The size of a hydrogen atom is about 0.5\AA . What is the total atomic volume in m^{-3} of a mole of hydrogen atoms ?

 [Watch Video Solution](#)

21. One mole of an ideal gas at standard temperature and pressure occupies 22.4 L (molar volume). IF the size of the hydrogen molecule is about 1A. What is the ratio of molar volume to the atomic volume of a mole of hydrogen?



[Watch Video Solution](#)

22. Explain this common observation clearly: If you look out of the window of a fast moving train, the nearby trees, houses etc. seem to move rapidly in a direction opposite to the train's motion, but the distant objects (hills tops, the Moons, the stars etc) seem to be

stationary . (In fact , since you are aware that you are moving , these distant objects seem to move with you.)



[Watch Video Solution](#)

23. The principle of 'parallax' in section 2.3.1 is used in the determination of distances of very distant stars. The baseline AB is the line joining the Earth's two locations six months apart in its orbit around the Sun. That is, the baseline is about the diameter of the Earth's orbit $= 3 \times 10^{11} m$. However, even the nearest stars are so distant that with such a long baseline, they show parallax only of the order of 1" (second) of arc or so. A parsec is a convenient unit of length on the

astronomical scale. It is the distance of an object that will show a parallax of 1" (second of arc) from opposite ends of a baseline equal to the distance from the Earth to the Sun. How much is a parsec in terms of metres ?



[View Text Solution](#)

24. The nearest star to our solar system is 4.29 light years away. How much is this distance in terms of parsecs ? How much parallax would this star (named Alpha Centauri) show when viewed from two locations of the Earth six months apart in its orbit around the Sun ?



[View Text Solution](#)

25. The Sun is a hot plasma (ionized matter) with its inner core at a temperature exceeding $10^7 K$, and its outer surface at a temperature of about 6000K. At these high temperatures, no substance remains in a solid or liquid phase. In what range do you expect the mass density of the Sun to be, in the range of densities of solids and liquids or gases? Check if your guess is correct from the following data: mass of the Sun = $2.0 \times 10^{30} Kg$, radius of the Sun = $7.0 \times 10^8 m$.



[View Text Solution](#)

26. When the planet Jupiter is at a distance of 824.7 million kilometers from the Earth, its angular diameter is measured to be 35.72" of arc. Calculate the diameter of Jupiter.



[Watch Video Solution](#)

Exercise Additional Exercise

1. A man walking briskly in rain with speed v must slant his umbrella forward making an angle θ with the vertical. A student derives the following relation between θ and v : $\tan \theta = v$ and checks that the relation

has a correct limit : as $v \rightarrow 0$, $\theta \rightarrow 0$, as expected , (we are assuming there is no strong wind and that the rain falls vertically for a stationary man.) Do you think this relation can be correct ? If you, guess the correct relation.

 [Watch Video Solution](#)

2. It is claimed that two atomic clocks if allowed to run for 100 years, free from any disturbance, may differ by only about 0.02 second. In measuring a time interval of second what is the accuracy?

 [Watch Video Solution](#)

3. Estimate the average mass density of a sodium atom assuming its size to be about 2.5\AA . (Use the known values of Avagardo's number and the atomic mass of sodium). Compare it with the mass density of sodium in its crystalline phase : 970kgm^{-3} . Are the two densities of the same order of magnitude ? If so, why ?



[Watch Video Solution](#)

4. The unit of length conventent on the nuclear scale is a fermi : $1f = 10^{-15}\text{m}$. Nuclear sized obey roughly the following empirical relation:

$$r = r_0 A^{1/3}$$

Where r is the radius of the nucleus, A its mass number,

and r_0 is a constant equal to about, 1.2 f. Show that the rule implies that nuclear mass density is nearly constant for different nuclei. Estimate the mass density of sodium nucleus. Compare it with the average mass density of a sodium atom obtained in Exercise . 2.27.



[View Text Solution](#)

5. A LASER is a source of very intense, monochromatic, and unidirectional beam of light. These properties of a laser light can be exploited to measure long distance. The distance of the Moon from the Earth has been already determined very precisely using a laser as a source of light . A laser light beamed at the Moon takes

2.56 s to return after reflection at the Moon's surface.

How much is the radius of the lunar orbit around the Earth ?



[Watch Video Solution](#)

6. A SONAR (sound navigation and ranging) uses ultrasonic waves to detect and locate objects under water. In a submarine equipped with a SONAR , the time delay between generation of a probe wave and the reception of its echo after reflection from an enemy submarine is found to be 77.0s. What is the distance of the enemy submarine ? (Speed of sound in water = 1450ms^{-1}).



[Watch Video Solution](#)



[Watch Video Solution](#)

7. The farthest objects in our universe discovered by modern astronomers are so distant that light emitted by them takes billions of years to reach the earth. What is the distance in km of a quasar from which light takes 3 billion years to reach us?



[Watch Video Solution](#)

8. It is a well known fact that during a total solar eclipse the disk of the moon almost completely covers the disk of the Sun. From this fact and from the information you

can gather from examples 2.3 and 2.4, determine the approximate diameter of the moon.



[View Text Solution](#)

9. A great physicist of this century (P.A.M. Dirac) loved playing with numerical values of Fundamental constants of nature. This led him to an interesting observation. Dirac found that from the basic constants of atomic physics (c, e , mass of electron, mass of proton) and the gravitational constant G , he could arrive at a number with the dimension of time. Further, it was a very large number, its magnitude being close to the present estimate on the age of the universe (~ 15

billion years). From the table of fundamental constants in this book, try to see if you too can construct this number (or any other interesting number you can think of). If its coincidence with the age of the universe were significant, what would this imply for the constancy of fundamental constants?



[View Text Solution](#)