



## MATHS

### NCERT - NCERT MATHEMATICS(TELUGU)

#### DETERMINANTS

#### Example

1. Evaluate  $\begin{bmatrix} 2 & 4 \\ -1 & 2 \end{bmatrix}$

 [Watch Video Solution](#)

2. Evaluate  $\begin{bmatrix} x & x + 1 \\ x - 1 & x \end{bmatrix}$

 [Watch Video Solution](#)

3. Evaluate the determinant  $\Delta = \begin{bmatrix} 1 & 2 & 4 \\ -1 & 3 & 0 \\ 4 & 1 & 0 \end{bmatrix}$

 [Watch Video Solution](#)

4. Evaluate  $\Delta = \begin{bmatrix} 0 & \sin \alpha & -\cos \alpha \\ -\sin \alpha & 0 & \sin \beta \\ \cos \alpha & -\sin \beta & 0 \end{bmatrix}$

 [Watch Video Solution](#)

5. Find values of x for which  $\begin{bmatrix} 3 & x \\ x & 1 \end{bmatrix} = \begin{bmatrix} 3 & 2 \\ 4 & 1 \end{bmatrix}$

 [Watch Video Solution](#)

6. Verify property 1 for  $\Delta = \begin{bmatrix} 2 & -3 & 5 \\ 6 & 0 & 4 \\ 1 & 5 & -7 \end{bmatrix}$

 [View Text Solution](#)

7. Verify Property 2 for  $\Delta = \begin{bmatrix} 2 & -3 & 5 \\ 6 & 0 & 4 \\ 1 & 5 & -7 \end{bmatrix}$

 [Watch Video Solution](#)

8. Evaluate  $\Delta = \begin{bmatrix} 3 & 2 & 3 \\ 2 & 2 & 3 \\ 3 & 2 & 3 \end{bmatrix}$

 [Watch Video Solution](#)

9. Evaluate  $\begin{bmatrix} 102 & 18 & 36 \\ 1 & 3 & 4 \\ 17 & 3 & 6 \end{bmatrix}$

 [Watch Video Solution](#)

10. Show that  $\begin{vmatrix} a + b + 2c & a & b \\ c & b + c + 2a & b \\ c & a & c + a + 2b \end{vmatrix} = 2(a + b + c)^3$

 [Watch Video Solution](#)

11. Prove that 
$$\begin{bmatrix} a & a+b & a+b+c \\ 2a & 3a+2b & 4a+3b+2c \\ 3a & 6a+3b & 10a+6b+3c \end{bmatrix} = a^3$$

 [View Text Solution](#)

12. Without expanding prove that 
$$\Delta = \begin{bmatrix} x+y & y+z & z+x \\ z & x & y \\ 1 & 1 & 1 \end{bmatrix} = 0$$

 [Watch Video Solution](#)

13. Evaluate 
$$\Delta = \begin{bmatrix} 1 & a & bc \\ 1 & b & ca \\ 1 & c & ab \end{bmatrix}$$

 [Watch Video Solution](#)

14. Prove that 
$$\begin{bmatrix} b+c & a & a \\ b & c+a & b \\ c & c & b+a \end{bmatrix}$$



[View Text Solution](#)

15. If  $x, y, z$  are different and  $\Delta = \begin{bmatrix} x & x^2 & 1 + x^3 \\ y & y^2 & 1 + y^3 \\ z & z^2 & 1 + z^3 \end{bmatrix}$  then prove that

$$1 + XYZ = 0$$

 Watch Video Solution

16. Show that

$$\begin{bmatrix} 1 + a & 1 & 1 \\ 1 & 1 + b & 1 \\ 1 & 1 & 1 + c \end{bmatrix} = abc \left( 1 + \frac{1}{a} + \frac{1}{b} + \frac{1}{c} \right) = abc + bc + ca + ab$$

 Watch Video Solution

17. Find the area of the triangle whose vertices are  $(3, 8)$ ,  $(-4, 2)$  and  $(5, 1)$

 Watch Video Solution

18. Find the equation of the line joining A(1,3) and B(0,0) using determinants and find k if D(k,0) is a point such that area of triangle ABD is  $3\text{sq units}$ .

 [Watch Video Solution](#)

19. Find the minor of elements 6 in the determinants  $\Delta = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix}$

 [Watch Video Solution](#)

20. Find the minor and cofactors of all the elements of the determinants

$$\begin{bmatrix} 1 & -2 \\ 4 & 3 \end{bmatrix}$$

 [Watch Video Solution](#)

21. Find minors and cofactors of the elements  $a_{11}, a_{21}$  in the

$$\text{determinant } \Delta = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$$

 [Watch Video Solution](#)

22. Find  $\text{adj } A$  for  $A = \begin{bmatrix} 2 & 3 \\ 1 & 4 \end{bmatrix}$

 [Watch Video Solution](#)

23. If  $A = \begin{bmatrix} 1 & 3 & 3 \\ 1 & 4 & 3 \\ 1 & 3 & 4 \end{bmatrix}$ , then verify that  $A \text{ adj } A = |A| I$ . Also find  $A^{-1}$

 [Watch Video Solution](#)

24. If  $A = \begin{bmatrix} 2 & 3 \\ 1 & -4 \end{bmatrix}$  and  $B = \begin{bmatrix} 1 & -2 \\ -1 & 3 \end{bmatrix}$ , then verify that  $(AB)^{-1} = B^{-1}A^{-1}$

[Watch Video Solution](#)

25. Show that the matrix  $A = \begin{bmatrix} 2 & 3 \\ 1 & 2 \end{bmatrix}$  satisfies equation  $A^2 - 4A + I = 0$  where  $I$  is  $2 \times 2$  identity matrix and  $O$  is  $2 \times 2$  Zero matrix. Using this equation, Find  $A^{-1}$

[Watch Video Solution](#)

26. Solve the system of equation

$$2x + 5y = 1$$

$$3x + 2y = 7$$

[Watch Video Solution](#)

27. The sum of three numbers is 6. If we multiply third numbers by 3. and add second numbers to it . We get 11. by adding first and third numbers ,



we get double of the second number. Represent it algebraically and find the numbers using matrix method.

A.  $x = 1, y = 2, z = 3$

B.

C.

D.

**Answer:**

 [Watch Video Solution](#)

**28.** If,  $a, b, c$  are positive and unequal, show that value of the determinant

$$\Delta = \begin{vmatrix} a & b & c \\ b & c & a \\ c & a & b \end{vmatrix} \text{ is negative}$$

 [Watch Video Solution](#)

29. Use product  $\begin{bmatrix} 1 & -1 & 2 \\ 0 & 2 & -3 \\ 3 & -2 & 4 \end{bmatrix} \begin{bmatrix} -2 & 0 & 1 \\ 9 & 2 & -3 \\ 6 & 1 & -2 \end{bmatrix}$  to solve the system of equations

$$x - y + 2z = 1$$

$$2y - 3z = 1$$

$$3x - 2y + 4z = 2$$

A.  $z = 3$

B.

C.

D.

**Answer:**



**Watch Video Solution**

30. Prove that

$$\Delta = \begin{bmatrix} a + bx & c + dx & p + qx \\ ax + b & cx + d & px + q \\ u & v & w \end{bmatrix} = (1 - x^2) \begin{bmatrix} a & c & p \\ b & d & q \\ u & v & w \end{bmatrix}$$

 [Watch Video Solution](#)

### Exercise 4 1

1. Evaluate the determinants in Exercises 1 and 2.

$$\begin{bmatrix} 2 & 4 \\ -5 & -1 \end{bmatrix}$$

 [Watch Video Solution](#)

2. (i)  $\begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$

(ii)  $\begin{bmatrix} x^2 - x + 1 & x - 1 \\ x + 1 & x + 1 \end{bmatrix}$

 [Watch Video Solution](#)

3. If  $A = \begin{bmatrix} 1 & 2 \\ 4 & 2 \end{bmatrix}$ , then show that  $|2A| = 4|A|$

 [Watch Video Solution](#)

4. If  $A = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 2 \\ 0 & 0 & 4 \end{bmatrix}$ , then show that  $|3A| = 27|A|$

 [Watch Video Solution](#)

5. Evaluate the determinants

(i)  $(3, -1, -2), (0, 0, -1), (3, -5, 0)$

 [Watch Video Solution](#)

6. If  $A = \begin{bmatrix} 1 & 1 & -2 \\ 2 & 1 & -3 \\ 5 & 4 & -9 \end{bmatrix}$ , find  $|A|$

 [Watch Video Solution](#)

7. Find values of  $x$ , if

$$(i) \begin{bmatrix} 2 & 4 \\ 5 & 1 \end{bmatrix} = \begin{bmatrix} 2x & 4 \\ 6 & x \end{bmatrix}$$

$$(ii) \begin{bmatrix} 2 & 3 \\ 4 & 5 \end{bmatrix} = \begin{bmatrix} x & 3 \\ 2x & 5 \end{bmatrix}$$



Watch Video Solution

8. If  $\begin{bmatrix} x & 2 \\ 18 & x \end{bmatrix} = \begin{bmatrix} 6 & 2 \\ 18 & 6 \end{bmatrix}$ , then  $x$  is equal to

A. 6

B.  $\pm 6$

C.  $-6$

D. 0

Answer: B



Watch Video Solution

1. Using the property of determinants and without expanding

$$\begin{bmatrix} x & a & x+a \\ y & b & y+b \\ z & c & z+c \end{bmatrix} = 0$$

 [Watch Video Solution](#)

2. 
$$\begin{vmatrix} a-b & b-c & c-a \\ b-c & c-a & a-b \\ c-a & a-b & b-c \end{vmatrix} =$$

 [Watch Video Solution](#)

3. Using the property of determinants and without expanding

$$\begin{bmatrix} 2 & 7 & 65 \\ 3 & 8 & 75 \\ 5 & 9 & 86 \end{bmatrix} = 0$$

 [Watch Video Solution](#)

$$4. \begin{vmatrix} 1 & bc & a(b+c) \\ 1 & ca & b(c+a) \\ 1 & ab & c(a+b) \end{vmatrix} =$$



[Watch Video Solution](#)

5. Using the property of determinants and without expanding

$$\begin{bmatrix} b+c & q+r & y+z \\ c+a & r+p & z+x \\ a+b & p+q & x+y \end{bmatrix} = 2 \begin{bmatrix} a & p & x \\ b & q & y \\ c & r & z \end{bmatrix}$$



[Watch Video Solution](#)

6. Using the property of determinants and without expanding

$$\begin{bmatrix} 0 & a & -b \\ -a & 0 & -c \\ b & c & 0 \end{bmatrix} = 0$$



[Watch Video Solution](#)

$$7. \begin{vmatrix} -a^2 & ab & ac \\ ab & -b^2 & bc \\ ac & bc & -c^2 \end{vmatrix} =$$



Watch Video Solution

$$8. \begin{vmatrix} 1 & a & a^2 \\ 1 & b & b^2 \\ 1 & c & c^2 \end{vmatrix} =$$



Watch Video Solution

9. By using properties of determinants, show that :

$$\begin{bmatrix} x & x^2 & yz \\ y & y^2 & zx \\ z & z^2 & xy \end{bmatrix} = (x - y)(y - z)(z - x)(xy + yz + zx)$$



Watch Video Solution



10. By using properties of determinants , show that : (i)

$$\begin{bmatrix} x+4 & 2x & 2x \\ 2x & x+4 & 2x \\ 2x & 2x & x+4 \end{bmatrix} = (5x+4)(4-x)^2$$

(ii)  $\begin{bmatrix} y+k & y & y \\ y & y+k & y \\ y & y & y+k \end{bmatrix} = k^2(3y+k)$

 [Watch Video Solution](#)

11. Show that  $\begin{vmatrix} a-b-c & 2a & 2a \\ 2b & b-c-a & 2b \\ 2c & 2c & c-a-b \end{vmatrix} = (a+b+c)^3$

 [Watch Video Solution](#)

12. By using properties of determinants , show that :

$$\begin{bmatrix} 1 & x & x^2 \\ x^2 & 1 & x \\ x & x^2 & 1 \end{bmatrix} = (1-x^3)^2$$

 [Watch Video Solution](#)

13. Prove that

$$\begin{vmatrix} 1 + a^2 - b^2 & 2ab & -2b \\ 2ab & 1 - a^2 + b^2 & 2a \\ 2b & -2a & 1 - a^2 - b^2 \end{vmatrix} = (1 + a^2 + b^2)^3$$

 [Watch Video Solution](#)

14.  $\begin{vmatrix} a^2 + 1 & ab & ac \\ ab & b^2 + 1 & bc \\ ac & bc & c^2 + 1 \end{vmatrix} =$

 [Watch Video Solution](#)

15. Let A be a square matrix of order  $3 \times 3$  then  $|kA|$  is equal to

A.  $k|A|$

B.  $k^2|A|$

C.  $k^3|A|$

D.  $3k|A|$

**Answer: C**



[Watch Video Solution](#)

**16.** Which of the following is correct

- A. Determinant is a square matrix.
- B. Determinant is a number associated to a matrix
- C. Determinant is a number associated to a square matrix
- D. None of these

**Answer: C**



[Watch Video Solution](#)

**Exercise 4 3**

1. Find area of the triangle with vertices at the point given in each of the following:

$(1, 0), (6, 0), (4, 3)$



[Watch Video Solution](#)

2. Find area of the triangle with vertices at the point given in each of the following:

$(2, 7), (1, 1), (10, 8)$



[Watch Video Solution](#)

3. Find area of the triangle with vertices at the point given in each of the following:

$(-2, -3), (3, 2), (-1, -8)$



[Watch Video Solution](#)

4. Prove that the points  $(a, b + c)$ ,  $(b, c + a)$  and  $(c, a + b)$  are collinear and find the equation of the straight line containing them .



[Watch Video Solution](#)

5. Find values of  $k$  if area of triangle is 4 sq. units and vertices are

(i)  $(k, 0)$ ,  $(4, 0)$ ,  $(0, 2)$        $(-2, 0)$ ,  $(0, 4)$ ,  $(0, k)$



[Watch Video Solution](#)

6. (i) Find equation of line joining  $(1,2)$  and  $(3,6)$  using determinants .

(ii) Find equation of line joining  $(3,1)$  and  $(9,3)$  using determinants.



[View Text Solution](#)

7. If area of triangle is 35sq units with vertices  $(2,-6)$ ,  $(5,4)$  and  $(k,4)$  Then  $k$  is

A. 12

B.  $-2$

C.  $-12, -2$

D.  $12, -2$

**Answer: D**



[Watch Video Solution](#)

### Exercise 4 4

1. Write Minors and Cofactors of the elements of following determinants:

(i)  $\begin{bmatrix} 2 & -4 \\ 0 & 3 \end{bmatrix}$

$$\begin{bmatrix} a & c \\ b & d \end{bmatrix}$$



[Watch Video Solution](#)

2. The sum of the values of  $x$  so that the matrix

$$\begin{pmatrix} 2 & 2 & 1 \\ 1 & 3 & 1 \\ 1 & 2 & 2 \end{pmatrix} - x \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \text{ is singular, is}$$

 [Watch Video Solution](#)

3. Using Cofactors of elements of second row, evaluate  $\Delta = \begin{bmatrix} 5 & 3 & 8 \\ 2 & 0 & 1 \\ 1 & 2 & 3 \end{bmatrix}$

 [Watch Video Solution](#)

4. Using Cofactors of elements of third column, evaluate,

$$\Delta = \begin{bmatrix} 1 & x & yz \\ 1 & y & zx \\ 1 & z & xy \end{bmatrix}$$

 [Watch Video Solution](#)

5. If  $\Delta = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$  and  $A_{ij}$  is Cofactors of  $a_{ij}$  then value of  $\Delta$  is given by

A.  $a_{11}A_{31} + a_{12}A_{32} + a_{13}A_{33}$

B.  $a_{11}A_{11} + a_{12}A_{21} + a_{13}A_{31}$

C.  $a_{21}A_{11} + a_{22}A_{12} + a_{23}A_{13}$

D.  $a_{11}A_{11} + a_{21}A_{21} + a_{31}A_{31}$

**Answer: D**

 [Watch Video Solution](#)

## Exercise 4 5

1. Find adjoint of each of the matrices

$$\begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$$

 [Watch Video Solution](#)



[Watch Video Solution](#)

2. Find adjoint of each of the matrices

$$\begin{bmatrix} 1 & -1 & 2 \\ 2 & 3 & 5 \\ -2 & 0 & 1 \end{bmatrix}$$



[Watch Video Solution](#)

3. Verify  $A (\text{adj } A) = (\text{adj } A) A = |A| I$   $\begin{bmatrix} 2 & 3 \\ -4 & -6 \end{bmatrix}$



[Watch Video Solution](#)

4. Find the discriminant of the matrix.  $\begin{bmatrix} 1 & -1 & 2 \\ 3 & 0 & -2 \\ 1 & 0 & 3 \end{bmatrix}$



[Watch Video Solution](#)

5. Find the inverse of each of the matrices (if it exists )

$$\begin{bmatrix} 2 & -2 \\ 4 & 3 \end{bmatrix}$$

 [Watch Video Solution](#)

6. Find the inverse of each of the matrices (if it exists )

$$\begin{bmatrix} 1 & 2 & 3 \\ 0 & 2 & 4 \\ 0 & 0 & 5 \end{bmatrix}$$

 [View Text Solution](#)

7. Find the inverse of each of the matrices (if it exists )

$$\begin{bmatrix} 1 & 0 & 0 \\ 3 & 3 & 0 \\ 5 & 2 & -1 \end{bmatrix}$$

 [View Text Solution](#)

8. Find the inverse of each of the matrices (if it exists )

$$\begin{bmatrix} 2 & 1 & 3 \\ 4 & -1 & 0 \\ -7 & 2 & 1 \end{bmatrix}$$



Watch Video Solution

9. Find the inverse of each of the matrices (if it exists )

$$\begin{bmatrix} 1 & -1 & 2 \\ 0 & 2 & -3 \\ 3 & -2 & 4 \end{bmatrix}$$



Watch Video Solution

10. Find the inverse of each of the matrices (if it exists )

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & \cos \alpha & \sin \alpha \\ 0 & \sin \alpha & -\cos \alpha \end{bmatrix}$$



Watch Video Solution

11. Let  $A = \begin{bmatrix} 3 & 7 \\ 2 & 5 \end{bmatrix}$  and  $B = \begin{bmatrix} 6 & 8 \\ 7 & 9 \end{bmatrix}$ . Verify that  $(AB)^{-1} = B^{-1}A^{-1}$

 [Watch Video Solution](#)

12. If  $A = \begin{bmatrix} 3 & 1 \\ -1 & 2 \end{bmatrix}$  Show that  $A^2 - 5A + 7I = O$ . Hence find  $A^{-1}$

 [Watch Video Solution](#)

13. For the matrix  $A = \begin{bmatrix} 3 & 2 \\ -1 & 2 \end{bmatrix}$ . Find the numbers  $a$  and  $b$  such that  $A^2 + aA + bI = O$ .

 [Watch Video Solution](#)

14.  $A = \begin{bmatrix} 1 & 1 & 1 \\ 1 & 2 & -3 \\ 2 & -1 & 3 \end{bmatrix}$

Show that  $A^3 - 6A^2 + 5A + 11I = O$ . Hence, find  $A^{-1}$

 [Watch Video Solution](#)

15. If  $A = \begin{bmatrix} 2 & -1 & 1 \\ -1 & 2 & -1 \\ 1 & -1 & 2 \end{bmatrix}$

Verify that  $A^3 - 6A^2 + 9A - 4I = O$  and hence find  $A^{-1}$

 [Watch Video Solution](#)

16. Let  $A$  be a nonsingular square matrix of order  $3 \times 3$ . Then  $|\text{adj } A|$  is equal to

A.  $|A|$

B.  $|A|^2$

C.  $|A|^3$

D.  $3|A|$

**Answer: B**

 [Watch Video Solution](#)

17. If  $A$  is an invertible matrix of order 2, then  $\det (A^{-1})$  is equal to

A.  $\det (A)$

B.  $\frac{1}{\det(A)}$

C. 1

D. 0

**Answer: B**



[Watch Video Solution](#)

### Exercise 4 6

1. Examine the consistency of the system of equations

$$x + 2y = 2$$

$$2x + 3y = 3$$



[Watch Video Solution](#)

2. Examine the consistency of the system of equations

$$2x - y = 5$$

$$x + y = 4$$



Watch Video Solution

3. Examine the consistency of the system of equations

$$x + 3y = 5$$

$$2x + 6y = 8$$



Watch Video Solution

4. Examine the consistency of the system of equations

$$x + y + z = 1$$

$$2x + 3y + 2z = 2$$

$$ax + ay + 2az = 4$$



Watch Video Solution

5. Examine the consistency of the system of equations

$$3x - y - 2z = 2$$

$$2y - z = -1$$

$$3x - 5y = 3$$



Watch Video Solution

6. Examine the consistency of the system of equations

$$5x - y + 4z = 5$$

$$2x + 3y + 5z = 2$$

$$5x - 2y + 6z = -1$$



Watch Video Solution

7. Solve system of linear equations , using matrix method

$$5x + 2y = 4$$



$$7x + 3y = 5$$



[View Text Solution](#)

**8.** Solve system of linear equations , using matrix method

$$2x - y = -2$$

$$3x + 4y = 3$$



[View Text Solution](#)

**9.** Solve system of linear equations , using matrix method

$$4x - 3y = 3$$

$$3x - 5y = 7$$



[Watch Video Solution](#)

**10.** Solve system of linear equations , using matrix method

$$5x + 2y = 3$$

$$3x + 2y = 5$$



[View Text Solution](#)

11. Solve system of linear equations , using matrix method

$$2x + y + z = 1$$

$$x - 2y - z = \frac{3}{2}$$

$$3y - 5z = 9$$



[View Text Solution](#)

12. Solve system of linear equations , using matrix method

$$x - y + z = 4$$

$$2x + y - 3z = 0$$

$$x + y + z = 2$$



[View Text Solution](#)

13. Solve system of linear equations , using matrix method

$$2x + 3y + 3z = 5$$

$$x - 2y + z = -4$$

$$3x - y - 2z = 3$$



[View Text Solution](#)

14. Solve system of linear equations , using matrix method

$$x - y + 2z = 7$$

$$3x + 4y - 5z = -5$$

$$2x - y + 3z = 12$$



[View Text Solution](#)

15. If  $A = \begin{bmatrix} 2 & -3 & 5 \\ 3 & 2 & -4 \\ 1 & 1 & -2 \end{bmatrix}$ , find  $A^{-1}$  Using  $A^{-1}$  solve the system of equations

$$2x - 3y + 5z = 11$$

$$3x + 2y - 4z = -5$$

$$x + y - 2z = -3$$

 [Watch Video Solution](#)

**16.** The cost of 4 kg. onion ,3kg wheat and 2kg rice is rupees 60. The cost of 2 kg onion , 4kg wheat and 6kg rice is rupees 90. The cost of 6kg onion 2kg wheat and 3 kg rice is rupees 70. Find cost of each item per kg by matrix method.

 [Watch Video Solution](#)

## Miscellaneous Exercises On Chapter 4

**1.** Prove that the determinant  $\begin{bmatrix} x & \sin \theta & \cos \theta \\ -\sin \theta & -x & 1 \\ \cos \theta & 1 & x \end{bmatrix}$  is independent of  $\theta$ .

 [Watch Video Solution](#)

2. Without expanding the determinant, prove that

$$\begin{bmatrix} a & a^2 & bc \\ b & b^2 & ca \\ c & c^2 & ab \end{bmatrix} = \begin{bmatrix} 1 & a^2 & a^3 \\ 1 & b^2 & b^3 \\ 1 & c^2 & c^3 \end{bmatrix}$$

 [Watch Video Solution](#)

3. Evaluate  $\begin{bmatrix} \cos \alpha \cos \beta & \cos \alpha \sin \beta & -\sin \alpha \\ -\sin \beta & \cos \beta & 0 \\ \sin \alpha \cos \beta & \sin \alpha \sin \beta & \cos \alpha \end{bmatrix} = 0$

 [Watch Video Solution](#)

4. If  $a, b$  and  $c$  are real numbers, and

$$\Delta = \begin{bmatrix} b+c & c+a & a+b \\ c+a & a+b & b+c \\ a+b & b+c & c+a \end{bmatrix} = 0$$

Show that either  $a + b + c = 0$  or  $a = b = c$ .

 [Watch Video Solution](#)

5. Solve the equation  $\begin{bmatrix} x + a & x & x \\ x & x + a & x \\ x & x & x + a \end{bmatrix} = 0, a \neq 0$

 [Watch Video Solution](#)

6. Prove that  $\begin{bmatrix} a^2 & bc & ac + c^2 \\ a^2 + ab & b^2 & ac \\ ab & b^2 + bc & c^2 \end{bmatrix} = 4a^2b^2c^2$

 [Watch Video Solution](#)

7. If  $A^{-1} = \begin{bmatrix} 3 & -1 & 1 \\ -15 & 6 & -5 \\ 5 & -2 & 2 \end{bmatrix}$  and  $B = \begin{bmatrix} 1 & 2 & -2 \\ -1 & 3 & 0 \\ 0 & -2 & 1 \end{bmatrix}$ , find  $(AB)^{-1}$

 [Watch Video Solution](#)

8. Evaluate  $\begin{bmatrix} x & y & x + y \\ y & x + y & x \\ x + y & x & y \end{bmatrix}$



[Watch Video Solution](#)

9. Evaluate 
$$\begin{bmatrix} 1 & x & y \\ 1 & x + y & y \\ 1 & x & x + y \end{bmatrix}$$

[Watch Video Solution](#)

10. Using properties of determinants in Exercises prove that :

$$\begin{bmatrix} \alpha & \alpha^2 & \beta + \gamma \\ \beta & \beta^2 & \gamma + \alpha \\ \gamma & \gamma^2 & \alpha + \beta \end{bmatrix} = (\beta - \gamma)(\gamma - \alpha)(\alpha - \beta)(\alpha + \beta + \gamma)$$

[Watch Video Solution](#)

11. 
$$\begin{bmatrix} x & x^2 & 1 + px^3 \\ y & y^2 & 1 + py^3 \\ z & z^2 & 1 + pz^3 \end{bmatrix} = (1 + pxyz)(x - y)(y - z)(z - x),$$

[Watch Video Solution](#)

12. Prove 
$$\begin{bmatrix} 3a & -a + b & -a + c \\ -b + a & 3b & -b + c \\ -c + a & -c + b & 3c \end{bmatrix} = 3(a + b + c)(ab + bc + ca)$$

 [Watch Video Solution](#)

13. 
$$\begin{bmatrix} 1 & 1 + p & 1 + p + q \\ 2 & 3 + 2p & 1 + 3p + 2q \\ 3 & 6 + 3p & 1 + 6p + 3q \end{bmatrix} = 1$$

 [Watch Video Solution](#)

14. Prove that 
$$\begin{bmatrix} \sin \alpha & \cos \alpha & \cos(\alpha + \delta) \\ \sin \beta & \cos \beta & \cos(\beta + \delta) \\ \sin \gamma & \cos \gamma & \cos(\gamma + \delta) \end{bmatrix} = 0$$

 [Watch Video Solution](#)

15. If a,b,c are in A.P. then the determinant

$$\begin{bmatrix} x + 2 & x + 3 & x + 2a \\ x + 3 & x + 4 & x + 2b \\ x + 4 & x + 5 & x + 2c \end{bmatrix} \text{ is}$$



A. 0

B. 1

C.  $x$

D.  $2x$

**Answer: A**



**Watch Video Solution**

16. If  $x, y, z$  are nonzero real number , then the inverse of matrix

$$A = \begin{bmatrix} x & 0 & 0 \\ 0 & y & 0 \\ 0 & 0 & z \end{bmatrix} \text{ is}$$

A.  $\begin{bmatrix} x^{-1} & 0 & 0 \\ 0 & y^{-1} & 0 \\ 0 & 0 & z^{-1} \end{bmatrix}$

B.  $xyz \begin{bmatrix} x^{-1} & 0 & 0 \\ 0 & y^{-1} & 0 \\ 0 & 0 & z^{-1} \end{bmatrix}$

C.  $\frac{1}{xyz} \begin{bmatrix} x & 0 & 0 \\ 0 & y & 0 \\ 0 & 0 & z \end{bmatrix}$

$$D. \frac{1}{xyz} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Answer: A



Watch Video Solution

17. Let  $A = \begin{bmatrix} 1 & \sin \theta & 1 \\ -\sin \theta & 1 & \sin \theta \\ -1 & -\sin \theta & 1 \end{bmatrix}$ , where  $0 \leq \theta \leq 2\pi$ , Then

A.  $\text{Det}(A) = 0$

B.  $\text{Det}(A) \in (2, \infty)$

C.  $\text{Det}(A) \in (2, 4)$

D.  $\text{Det}(A) \in [2, 4]$

Answer: D



Watch Video Solution