



MATHS

NCERT - NCERT MATHEMATICS(TELUGU)

DIFFERENTIAL EQUATIONS

Example

1. Find the order and degree, if defined, of each of the following differential equations:

$$(i) \frac{dy}{dx} - \cos x = 0$$

$$(ii) xy \frac{d^2y}{dx^2} + x \left(\frac{dy}{dx} \right)^2 - y \frac{dy}{dx} = 0$$

$$(iii) y'''' + y^2 + e^{y'} = 0$$



Watch Video Solution

2. Verify that the function $y = e^{-3x}$ is a solution of the differential equation

$$\frac{d^2y}{dx^2} + \frac{dy}{dx} - 6y = 0$$

 [Watch Video Solution](#)

3. Verify that the function $y = a \cos x + b \sin x$, where $a, b \in \mathbb{R}$ is a solution of the differential equation $\frac{d^2y}{dx^2} + y = 0$

 [Watch Video Solution](#)

4. Form the differential equation representing the family of curves $y = mx$ where, m is arbitrary constant.

 [Watch Video Solution](#)

5. Form the differential equation representing the family of curves

$y = a \sin(x + b)$, where a, b are arbitrary constants.



[Watch Video Solution](#)

6. Form the differential equation representing the family of ellipse

having foci on x-axis and centre at the origin.



[Watch Video Solution](#)

7. Form the differential equation of the family of circles touching

the x-axis at origin.



[Watch Video Solution](#)

8. Form the differential equation representing the family of parabolas having vertex at origin and axis along positive direction of x-axis.

 [Watch Video Solution](#)

9. Find the general solution of the differential equation

$$\frac{dy}{dx} = \frac{x + 1}{2 - y}, (y \neq 2).$$

 [Watch Video Solution](#)

10. Find the general solution of the differential equation

$$\frac{dy}{dx} = \frac{1 + y^2}{1 + x^2}.$$

 [Watch Video Solution](#)

11. Find the particular solution of the differential equation

$$\frac{dy}{dx} = -4xy^2 \text{ given that } y = 1, \text{ when } x = 0.$$

 [Watch Video Solution](#)

12. Find the equation of the curve passing through the point (1,1)

whose differential equation is $x dy = (2x^2 + 1) dx (x \neq 0)$.

 [Watch Video Solution](#)

13. Find the equation of a curve passing through the point (-2,3),

given that the slope of the tangent to the curve at any point (x,y)

is $\frac{2x}{y^2}$.

 [Watch Video Solution](#)

14. In a bank, principal increases continuously at the rate of 5% per year. In how many years Rs 1000 double itself?

 [Watch Video Solution](#)

15. Show that the differential equation $(x - y) \frac{dy}{dx} = x + 2y$ is homogeneous and solve it.

 [View Text Solution](#)

16. Show that the differential equation $x \cos\left(\frac{y}{x}\right) \frac{dy}{dx} = y \cos\left(\frac{y}{x}\right) + x$ is homogeneous and solve it.

 [View Text Solution](#)

17. Show that the differential equation

$$2y \frac{e^x}{y} dx + \left(y - 2x \frac{e^x}{y} \right) dy = 0$$
 is homogeneous and find its

particular solution, given that, $x = 0$ when $y = 1$.



[View Text Solution](#)

18. Show that the family of curves for which the slope of the tangent at any point (x,y) on its $\frac{x^2 + y^2}{2xy}$, is given by $x^2 - y^2 = cx$.



[View Text Solution](#)

19. Find the general solution of the differential equation

$$\frac{dy}{dx} - y = \cos x.$$



[View Text Solution](#)

20. Find the general solution of the differential equation

$$x \frac{dy}{dx} + 2y = x^2 (x \neq 0).$$

 [Watch Video Solution](#)

21. Find the general solution of the differential equation

$$ydx - (x + 2y^2)dy = 0.$$

 [View Text Solution](#)

22. Find the particular solution of the differential equation

$$\frac{dy}{dx} + y \cot x = 2x + x^2 \cot x (x \neq 0) \text{ given that } y = 0 \text{ when } x = \frac{\pi}{2}.$$

 [View Text Solution](#)

23. Find the equation of a curve passing through the point (0,1). If the slope of the tangent to the curve at any point (x,y) is equal to the sum of the x coordinate (abscissa) and the product of the x coordinate and y coordinate (ordinate) of that point.



[View Text Solution](#)

Miscellaneous Examples

1. Verify that the function $y = c_1 e^{ax} \cos bx + c_2 e^{ax} \sin bx$, where c_1, c_2 are arbitrary constants is a solution of the differential equation

$$\frac{d^2 y}{dx^2} - 2a \frac{dy}{dx} + (a^2 + b^2)y = 0$$



[View Text Solution](#)

2. Form the differential equation of the family of circles in the second quadrant and touching the coordinate axes.



[View Text Solution](#)

3. Find the particular solution of the differential equations log

$$\left(\frac{dy}{dx}\right) = 3x + 4y \text{ given that } y = 0 \text{ when } x = 0.$$



[View Text Solution](#)

4. Solve the differential equation

$$(x dy - y dx) y \sin\left(\frac{y}{x}\right) = (y dx + x dy) x \cos\left(\frac{y}{x}\right).$$



[View Text Solution](#)

5. Solve the differential equation $(\tan^{-1} y - x)dy = (1 + y^2)dx$.



[View Text Solution](#)

Exercise 9 1

1. Determine order and degree (if defined) of differential equations given in Exercises 1 to 10

$$(1) \frac{d^4 y}{dx^4} + \sin(y''') = 0$$



[View Text Solution](#)

$$2. y' + 5y = 0$$



[View Text Solution](#)

$$3. \left(\frac{ds}{dt}\right)^4 + 3s\frac{d^2s}{dt^2} = 0$$

 [View Text Solution](#)

$$4. \left(\frac{d^2y}{dx^2}\right)^2 + \cos\left(\frac{dy}{dx}\right) = 0$$

 [View Text Solution](#)

$$5. \frac{d^2y}{dx^2} = \cos 3x + \sin 3x$$

 [View Text Solution](#)

$$6. (y''')^2 + (y'')^3 + (y')^4 + y^5 = 0$$

 [View Text Solution](#)

$$7. y' + 2y'' + y' = 0$$



[View Text Solution](#)

$$8. y' + y = e^x$$



[View Text Solution](#)

$$9. y + (y')^2 + 2y = 0$$



[View Text Solution](#)

$$10. y + 2y' + \sin y = 0$$



[View Text Solution](#)

11. The degree of the differential equation

$$\frac{d^2y}{(dx^2)^3} + \left(\frac{dy}{dx}\right)^2 + \sin\left(\frac{dy}{dx}\right) + 1 = 0 \text{ is}$$

- A. 3
- B. 2
- C. 1
- D. not defined

Answer: D



[View Text Solution](#)

12. The order of the differential equation

$$2x^2 \frac{d^2y}{dx^2} - 3 \frac{dy}{dx} + y = 0 \text{ is}$$

- A. 2

B. 1

C. 0

D. not defined

Answer: A



[View Text Solution](#)

Exercise 9 2

1. $y = \cos x + C : y' + \sin x = 0$



[Watch Video Solution](#)

2. $y = \sqrt{1 + x^2} : y' = \frac{xy}{1 + x^2}$



[Watch Video Solution](#)

3. $y = Ax : xy' = y(x \neq 0)$

 [Watch Video Solution](#)

4.

$y = x \sin x : xy' = y + x\sqrt{x^2 - y^2} (x \neq 0 \text{ and } x > y \text{ or } x < -y)$

 [Watch Video Solution](#)

5. $xy = \log y + C : y' = \frac{y^2}{1 - xy} (xy \neq 1)$

 [Watch Video Solution](#)

6. $y - \cos y = x : (y \sin y + \cos y + x)y' = y$



 Watch Video Solution

7. $x + y = \tan^{-1} y: y^2 y' + y^2 + 1 = 0$

 Watch Video Solution

8. $y = \sqrt{a^2 - x^2} x \neq (-a, a): x + y \frac{dy}{dx} = 0 (y \neq 0)$

 Watch Video Solution

9. The number of arbitrary constant in the general solution of a differential equation of fourth order are

A. 0

B. 2

C. 3

D. 4

Answer: D



[Watch Video Solution](#)

10. The number of arbitrary constants in the particular solution of a differential equation of third order are

A. 3

B. 2

C. 1

D. 0

Answer: D



[Watch Video Solution](#)

Exercise 9 3

1. In each of the Exercises 1 to 5, form a differential equation representing the given family of curves by eliminating arbitrary constants a and b .

$$1. \frac{x}{a} + \frac{y}{b} = 1$$



Watch Video Solution

2. find the general solution of the differential equation

$$y^2 = a(b^2 - x^2)$$



Watch Video Solution

$$3. y = ae^{3x} + be^{-2x}$$



Watch Video Solution

4. $y = e^{2x}(a + bx)$

 [Watch Video Solution](#)

5. find the general solution for the differential equations

$$y = e^x(a \cos x + b \sin x)$$

 [Watch Video Solution](#)

6. Form the differential equation of the family of circles touching the y-axis at origin.

 [Watch Video Solution](#)

7. The differential equation of the family of parabolas having vertices at the origin and foci on y-axis is

 [Watch Video Solution](#)

8. Form the differential equation of the family of ellipse having foci on y - axis and centre at origin.

 [Watch Video Solution](#)

9. Form the differential equation of the family of hyperbolas having foci on x-axis and centre at origin.

 [Watch Video Solution](#)

10. Form the differential equation of the family of circles having centre on y - axis and radius 3 units.

 [Watch Video Solution](#)

11. Which of the following differential equations has $y = c_1e^x + c_2e^{-x}$ as the general solution?

A. $\frac{d^2y}{dx^2} + y = 0$

B. $\frac{d^2y}{dx^2} - y = 0$

C. $\frac{d^2y}{dx^2} + 1 = 0$

D. $\frac{d^2y}{dx^2} - 1 = 0$

Answer: B

 [Watch Video Solution](#)

12. Which of the following differential equations has $y = x$ as one of its particular solution?

A. $\frac{d^2y}{dx^2} - x^2 \frac{dy}{dx} + xy = x$

B. $\frac{d^2y}{dx^2} + x \frac{dy}{dx} + xy = x$

C. $\frac{d^2y}{dx^2} - x^2 \frac{dy}{dx} + xy = 0$

D. $\frac{d^2y}{dx^2} + x \frac{dy}{dx} + xy = 0$

Answer: C



Watch Video Solution

Exercise 9 4

1. For each of the differential equations in Exercises 1 to 10, find the general solution:

$$1. \frac{dy}{dx} = \frac{1 - \cos x}{1 + \cos x}$$

 [Watch Video Solution](#)

2. find the general solutions for the differential equation

$$\frac{dy}{dx} = \sqrt{4 - y^2} \quad (-2 < y < 2)$$

 [Watch Video Solution](#)

3. find the general solutions for the differential equation

$$\frac{dy}{dx} + y = 1 \quad (y \neq 1)$$

 [Watch Video Solution](#)

4. The general solution of $\tan y \sec^2 x dx + \tan x \sec^2 y dy = 0$ is

 [Watch Video Solution](#)

5. $(e^x + e^{-x})dy - (e^x - e^{-x})dx = 0$

 [Watch Video Solution](#)

6. find the general solution for the differential equation

$$\frac{dy}{dx} = (1 + x^2)(1 + y^2)$$

 [Watch Video Solution](#)

7. find the general solutions for the differential equation

$$y \log y dx - x dy = 0$$

 [Watch Video Solution](#)

8. find the general solutions for the differential equation

$$x^5 \frac{dy}{dx} = -y^5$$

 [Watch Video Solution](#)

9. $\frac{dy}{dx} = \sin^{-1} x$

 [Watch Video Solution](#)

10. The solution of $e^x \tan y dx + (1 - e^x) \sec^2 y dy = 0$ is

 [Watch Video Solution](#)

11. For each of the differential equations in Exercises 11 to 14, find a particular solution satisfying the given condition:

11. $(x^3 + x^2 + x + 1) \frac{dy}{dx} = 2x^2 + x, y = 1$ when $x = 0$.



[View Text Solution](#)

12. $x(x^2 - 1) \frac{dy}{dx} = 1, y = 0$ when $x = 2$.



[Watch Video Solution](#)

13. $\cos\left(\frac{dy}{dx}\right) = a (a \neq R), y = 1$ when $x = 0$



[View Text Solution](#)

14. $\frac{dy}{dx} = y \tan x, y = 1$ when $x = 0$



[Watch Video Solution](#)

15. Find the equation of a curve passing through the point (0,0) and whose differential equation is $y' = e^x \sin x$.

 [Watch Video Solution](#)

16. For the differential equation $xy \frac{dy}{dx} = (x + 2)(y + 2)$, find the solution curve passing through the point (1, -1).

 [Watch Video Solution](#)

17. Find the equation of a curve passing through the point (0, -2) given that at any point (x,y) on the curve, the product of the slope of its tangent and y coordinate of the point is equal to the x coordinate of the point.

 [Watch Video Solution](#)

18. At any point (x,y) of a curve, the slope of the tangent is twice the slope of the line segment joining the point of contact to the point $(-4, -3)$. Find the equation of the curve given that it passes through $(-2, 1)$.



[View Text Solution](#)

19. The volume of spherical ballon being inflated changes at a constant rate. If initially its radius is 3 units and after 3 seconds it is 6 units. Find the radius of ballon after t seconds.



[Watch Video Solution](#)

20. In a bank, principal increases continuously at the rate of $r\%$ per yeat. Find the value of r if Rs 100 double itself in 10 years
($\log_e 2 = 0.6931$)



[Watch Video Solution](#)

21. In a bank, principal increases continuously at the rate of 5% per year. An amount of Rs 1000 is deposited with this bank, how much will it worth after 10 years $\left(e^{\frac{1}{2}} = 1.648\right)$



[Watch Video Solution](#)

22. In a culture, the bacteria count is 1,00,000. The number is increased by 10% in 2 hours. In how many hours will the count reach 2,00,000, if the rate of growth of bacteria is proportional to the number present ?



[Watch Video Solution](#)

23. The general solution of the differential equation $\frac{dy}{dx} = e^{x+y}$ is

A. $e^x - e^{-y} = C$

B. $e^x + e^y = C$

C. $e^{-x} + e^y = C$

D. $e^{-x} + e^{-y} = C$

Answer: A



[Watch Video Solution](#)

Exercise 9 5

1. In each of the Exercises 1 to 10, show that the given differential equation is homogeneous and solve each of them.

1. $(x^2 + xy)dy = (x^2 + y^2)dx.$



[View Text Solution](#)

2. $y' = \frac{x + y}{x}$



[View Text Solution](#)

3. $(x - y)dy - (x + y)dx = 0$



[View Text Solution](#)

4. The solution of $(x^2 - y^2)dx + 2xydy = 0$ is



[Watch Video Solution](#)

5. $x^2 \frac{dy}{dx} = x^2 - 2y^2 + xy$



[View Text Solution](#)

6. The solution of $xdy - ydx = \sqrt{x^2 + y^2}dx$ is



[Watch Video Solution](#)

7.

$$\left\{ x \cos\left(\frac{y}{x}\right) + y \sin\left(\frac{y}{x}\right) \right\} y dx = \left\{ y \sin\left(\frac{y}{x}\right) - x \cos\left(\frac{y}{x}\right) \right\} x dy$$



[View Text Solution](#)

8. $x \frac{dy}{dx} - y + x \sin\left(\frac{y}{x}\right) = 0$



[View Text Solution](#)

9. $ydx + x \log\left(\frac{y}{x}\right)dy - 2xdy = 0$

 [View Text Solution](#)

10. Solve $\left(1 + e^{\frac{x}{y}}\right)dx + e^{\frac{x}{y}}\left(1 - \frac{x}{y}\right)dy = 0$

 [Watch Video Solution](#)

11. For each of the differential equations in Exercises from 11 to 15, find the particular solution satisfying the given condition :

11. $(x + y)dy + (x - y)dx = 0, y = 1$ when $x = 1$

 [View Text Solution](#)

12. $x^2dy + (xy + y^2)dx = 0, y = 1$ when $x = 1$



[View Text Solution](#)

13. Give the solution of $x \sin^2 \frac{y}{x} dx = y dx - x dy$ which passes through the point $\left(1, \frac{\pi}{4}\right)$.



[Watch Video Solution](#)

14. $\frac{dy}{dx} - \frac{y}{x} + \operatorname{cosec}\left(\frac{y}{x}\right) = 0, y = 0$ when $x = 1$.



[View Text Solution](#)

15. $2xy + y^2 - 2x^2 \frac{dy}{dx} = 0, y = 2$ when $x = 1$



[View Text Solution](#)

16. A homogeneous differential equation of the form

$\frac{dx}{dy} = h\left(\frac{x}{y}\right)$ can be solved by making the substitution.

A. $y = vx$

B. $v = yx$

C. $x = vy$

D. $x = v$

Answer: C



Watch Video Solution

17. Which of the following is a homogeneous differential equation?

A. $(4x + 6y + 5)dy - (3y + 2x + 4)dx = 0$

B. $(xy)dx - (x^3 + y^3)dy = 0$

$$C. (x^3 + 2y^2)dx + 2xydy = 0$$

$$D. y^2dx + (x^2 - xy - y^2)dy = 0$$

Answer: D

 [Watch Video Solution](#)

Exercise 9 6

1. For each of the differential equations given in Exercises 1 to 12, find the general solution :

$$1. \frac{dy}{dx} + 2y = \sin x$$

 [Watch Video Solution](#)

$$2. \text{ solve } \frac{dy}{dx} + 3y = e^{-2x}$$

 [Watch Video Solution](#)

3. solve $\frac{dy}{dx} + \frac{y}{x} = x^2$

 [Watch Video Solution](#)

4. solve $\frac{dy}{dx} + (\sec x)y = \tan x \left(0 \leq x < \frac{\pi}{2}\right)$

 [Watch Video Solution](#)

5. solve $\cos^2 x \frac{dy}{dx} + y = \tan x \left(0 \leq x \leq \frac{\pi}{2}\right)$

 [Watch Video Solution](#)

6. The solution of $x \frac{dy}{dx} + 2y = x^2 \log x$ is

 [Watch Video Solution](#)

 Watch Video Solution

7. $x \log x \frac{dy}{dx} + y = 2 \log x$

 Watch Video Solution

8. solve $(1 + x^2)dy + 2xydx = \cot x dx (x \neq 0)$

 Watch Video Solution

9. Solve $x \frac{dy}{dx} + y - x + xy \cot x = 0 (x \neq 0)$

 Watch Video Solution

10. Solve $(x + y) \frac{dy}{dx} = 1$

 Watch Video Solution

11. Solve $ydx + (x - y^2)dy = 0$

 [Watch Video Solution](#)

12. $(x + 3y^2) \frac{dy}{dx} = y$

 [View Text Solution](#)

13. The solution of $\frac{dy}{dx} + 2y \tan x = \sin x$, given that $y = 0, x = \frac{\pi}{3}$ is

 [Watch Video Solution](#)

14. Solve $(1 + x^2) \frac{dy}{dx} + 2xy = \frac{1}{1 + x^2}, y = 0$ when $x = 1$

 [View Text Solution](#)

15. $\frac{dy}{dx} = 3y \cot x = \sin 2x, y = 2$ when $x = \frac{\pi}{2}$

 [View Text Solution](#)

16. Find the equation of a curve passing through the origin given that the slope of the tangent to the curve at any point (x, y) is equal to the sum of the coordinates of the point.

 [Watch Video Solution](#)

17. Find the equation of a curve passing through the point $(0,2)$ given that the sum of the coordinates of any point on the curve exceeds the magnitude of the slope of the tangent to the curve at that point by 5.



[View Text Solution](#)

18. The Integrating Factor of the differential equation

$$x \frac{dy}{dx} - y = 2x^2 \text{ is}$$

A. e^{-x}

B. e^{-y}

C. $\frac{1}{x}$

D. x

Answer: C



[Watch Video Solution](#)

19. The Integrating Factor of the differential equation

$$(1 - y^2) \frac{dx}{dy} + yx = ay \quad (-1 < y < 1) \text{ is}$$

A. $\frac{1}{y^2 - 1}$

B. $\frac{1}{\sqrt{y^2 - 1}}$

C. $\frac{1}{1 - y^2}$

D. $\frac{1}{\sqrt{1 - y^2}}$

Answer: D

 [Watch Video Solution](#)

Miscellaneous Exercise

1. For each of the differential equations given below, indicate its order and degree(if defined).

(i) $\frac{d^2y}{dx^2} + 5x \left(\frac{dy}{dx} \right)^2 - 6y = \log x$

(ii) $\left(\frac{dy}{dx} \right)^3 - 4 \left(\frac{dy}{dx} \right)^2 + 7y = \sin x$

(iii) $\frac{d^4y}{dx^4} - \sin \left(\frac{d^3y}{dx^3} \right) = 0$



[Watch Video Solution](#)

2. For each of the exercises given below, verify that the given function (implicit or explicit) is a solution of the corresponding differential equation.

$$(i) \quad xy = ae^x + be^{-x} + x^2 : x \frac{d^2y}{dx^2} + 2 \frac{dy}{dx} - xy + x^2 - 2 = 0$$

$$(ii) \quad y = e^x(a \cos x + b \sin x) : \frac{d^2y}{dx^2} - 2 \frac{dy}{dx} + 2y = 0$$

$$(iii) \quad y = x \sin 3x : \frac{d^2y}{dx^2} + 9y - 6 \cos 3x = 0$$

$$(iv) \quad x^2 = 2y^2 \log y : \left(x^2 + y^2 \frac{dy}{dx} - xy \right) = 0$$



[View Text Solution](#)

3. Form the differential equation representing the family of curves given by $(x - aa)^2 + 2y^2 = a^2$, where a is an arbitrary constant.



[View Text Solution](#)

4. Prove the $x^2 - y^2 = c(x^2 + y^2)^2$ is the general solution of differential equation $(x^3 - 3xy^2)dx = (y^3 - 3x^2y)dy$, where c is a parameter.



[View Text Solution](#)

5. Form the differential equation of the family of circles in the first quadrant which touch the coordinate axes.



[View Text Solution](#)

6. Find the general solution of the differential equation

$$\frac{dy}{dx} + \sqrt{\frac{1-y^2}{1-x^2}} = 0.$$



[Watch Video Solution](#)

7. Show that the general solution of the differential equation

$$\frac{dy}{dx} + \frac{y^2 + y + 1}{x^2 + x + 1} = 0 \quad \text{is} \quad \text{given} \quad \text{by}$$

$(x + y + 1) = A(1 - x - y - 2xy)$, where A is parameter.



[View Text Solution](#)

8. Find the equation of the curve passing through the point

$\left(0, \frac{\pi}{4}\right)$ whose differential equation is $\sin x \cos y \, dx + \cos x \sin y \, dy = 0$.



[View Text Solution](#)

9. Find the particular solution of the differential equation

$(1 + e^{2x}) \, dy + (1 + y^2) \, e^x \, dx = 0$, given that $y = 1$ when $x = 0$



[Watch Video Solution](#)

10. Solve the differential equation

$$y \frac{e^x}{y} dx = \left(x \frac{e^x}{y} + y^2 \right) dy (y \neq 0).$$



[View Text Solution](#)

11. Find a particular solution of the differential equation

$$(x - y)(dx + dy) = dx - dy \text{ given that } y = -1, \text{ when } x = 0. (\text{Hint :}$$

put $x - y = t$)



[Watch Video Solution](#)

12. Solve the differential equation

$$\left[\frac{e^{-2\sqrt{x}}}{\sqrt{x}} - \frac{y}{\sqrt{x}} \right] \frac{dx}{dy} = 1 (x \neq 0).$$



[Watch Video Solution](#)

13. The solution of $\frac{dy}{dx} + y \cot x = 4x \cos ecx$, given that $y = 0$,
 $x = \frac{\pi}{2}$ is

 [Watch Video Solution](#)

14. Find a particular solution of the differential equation
 $(x + 1) \frac{dy}{dx} = 2e^{-y} - 1$, given that $y = 0$ when $x = 0$.

 [Watch Video Solution](#)

15. The population of a village increases continuously at the rate proportional to the number of its inhabitants present at any time. If the population of the village was 0,000 in 1999 and 25000 in the year 2004, what will be the population of the village in 2009 ?

 [Watch Video Solution](#)

16. The general solution of the differential equation

$$\frac{ydx - xdy}{y} = 0 \text{ is}$$

A. $xy = C$

B. $x = Cy^2$

C. $y = Cx$

D. $y = Cx^2$

Answer: C

 [Watch Video Solution](#)

17. The general solution of a differential equation of the type

$$\frac{dy}{dx} + P_1x = Q_1 \text{ is}$$

A. $y \cdot e^{\int P_1 dy} = \int (Q_1 e^{\int P_1 dy}) dy + C$

B. $y \cdot e^{\int P_1 dx} = \int (Q_1 e^{\int P_1 dx}) dx + C$

$$C. xe^{\int P_1 dx} = \int(Q_1 e^{\int P_1 dx}) dx + C$$

$$D. xe^{\int P_1 dx} = \int(Q_1 e^{\int P_1 dx}) dx + C$$

Answer: C

 [View Text Solution](#)

18. The general solution of the differential equation $e^x dy + (ye^x + 2x)dx = 0$ is

$$A. xe^y + x^2 = C$$

$$B. xe^y + y^2 = C$$

$$C. ye^x + x^2 = C$$

$$D. ye^y + x^2 = C$$

Answer: C

 [Watch Video Solution](#)

