



# MATHS

## BOOKS - RD SHARMA MATHS

### (ENGLISH)

# BRIEF REVIEW OF CARTESIAN SYSTEM OF RECTANGULAR COORDINATES

Others

1. If the two vertices of an equilateral triangle be  $(0, 0)$ ,  $(3, \sqrt{3})$ , find the third vertex.



**Watch Video Solution**

2. Find the coordinates of the circumcentre of the triangle whose vertices are  $(8, 6)$ ,  $(8 - 2)$  and  $(2, - 2)$  Also, find its circum-radius.



**Watch Video Solution**

3. Find the distance between the points :

$$(at_1^2, 2at_1) \text{ and } (at_2^2, 2at_2)$$



[Watch Video Solution](#)

4. Find the area of the quadrilateral  $ABCD$

whose vertices are respectively

$$A(1, 1), B(7, -3), C(12, 2) \text{ and } D(7, 21)$$

.



[Watch Video Solution](#)

5. Prove that the points  $(a, b + c)$ ,  $(b, c + a)$  and  $(c, a + b)$  are collinear.



[Watch Video Solution](#)

6. Let the opposite angular points of a square be  $(3, 4)$  and  $(1, -1)$ . Find the coordinates of the remaining angular points.



[Watch Video Solution](#)

7. Prove that the area of the triangle whose vertices are  $(t, t - 2)$ ,  $(t + 2, t + 2)$  and  $(t + 3, t)$  is independent of  $t$ .



[Watch Video Solution](#)

8. If the axes are shifted to the point  $(1, -2)$  without rotation, what do the following equations become?  $2x^2 + y^2 - 4x + 4y = 0$   
 $y^2 - 4x + 4y + 8 = 0$



[Watch Video Solution](#)

9. Shift the origin to a suitable point so that the equation  $y^2 + 4y + 8x - 2 = 0$  will not contain term of  $y$  and the constant term.



[Watch Video Solution](#)

10. For what value of  $k$  are the points  $(k, 2 - 2k)$   $(k + 1, 2k)$  and  $(-4 - k, 6 - 2k)$  are collinear?



[Watch Video Solution](#)

**11.** If the coordinates of the mid-points of the sides of a triangle are  $(1, 1)$ ,  $(2, -3)$  and  $(3, 4)$ . Find its (i) centroid (ii) in-centre.



**Watch Video Solution**

**12. STATEMENT-1 :** Two bodies of mass  $M$  and  $2M$  released from rest and they move towards each other due to their mutual gravitational force of attraction and collide at mid point.

STATEMENT-2 : If the total mechanical energy of a body is zero than its linear momentum is necessarily non zero at any point other than reference point.

STATEMENT-3 : The linear momentum of the bob of a simple pendulum suspended by the roof of a car accelerating on a horizontal road increases due to tension in the string of the pendulum.



[Watch Video Solution](#)



**13.** Prove that the area of a triangle is invariant under the translation of the axes.



**Watch Video Solution**

**14.** Find the point to which the origin should be shifted so that the equation  $y^2 - 6y - 4x + 13 = 0$  is transformed to the form  $y^2 + Ax = 0$ .



**Watch Video Solution**

**15.**  $AB$  is a variable line sliding between the coordinate axes in such a way that  $A$  lies on the  $x$ -axis and  $B$  lies on the  $y$ -axis. If  $P$  is a variable point on  $AB$  such that  $PA = b$ ,  $Pb = a$ , and  $AB = a + b$ , find the equation of the locus of  $P$ .



**Watch Video Solution**

**16.** A point moves so that the sum of its distances from  $(ae, 0)$  and  $(-ae, 0)$  is  $2a$ ,

prove that the equation to its locus is

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1, \text{ where } b^2 = a^2(1 - e^2).$$



[Watch Video Solution](#)

17. A point moves so that the sum of its distances from  $(ae, 0)$  and  $(-ae, 0)$  is  $2a$ ,

prove that the equation to its locus is

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1, \text{ where } b^2 = a^2(1 - e^2).$$



[Watch Video Solution](#)

**18.** Find the equation of the locus of a point which moves such that the ratio of its distances from  $(2, 0)$  and  $(1, 3)$  is  $5:4$ .



**Watch Video Solution**

**19.** Find the equation to the locus of a point equidistant from the points  $A(1, 3)$  and  $B(-2, 1)$ .



**Watch Video Solution**

**20.** The sum of the squares of the distances of a moving point from two fixed points  $(a, 0)$  and  $(-a, 0)$  is equal to a constant quantity  $2c^2$ . Find the equation to its locus.



**Watch Video Solution**

**21.** Find the locus of a point, such that the join of  $(-5, 1)$  and  $(3, 2)$  subtends a right angle at the moving point.



**Watch Video Solution**

**22.** Find the locus of a point such that the sum of its distances from the points  $(0, 2)$  and  $(0, -2)$  is 6.



[Watch Video Solution](#)

**23.** Find the locus of the mid-point of the portion of the line  $x \cos \alpha + y \sin \alpha = p$  which is intercepted between the axes.



[Watch Video Solution](#)

24. A common test to find the genotype of a hybrid is by



[Watch Video Solution](#)

25. Show that four points  $(0, -1)$ ,  $(6, 7)$ ,  $(-2, 3)$  and  $(8, 3)$  are the vertices of a rectangle. Also, find its area.



[Watch Video Solution](#)

26. If the segments joining the points  $A(a, b)$  and  $B(c, d)$  subtends an angle  $\theta$  at the origin, prove that :

$$\cos \theta = \frac{ac + bd}{\sqrt{(a^2 + b^2)(c^2 + d^2)}}$$



[Watch Video Solution](#)

27. The vertices of a triangle are  $A(1, 1)$ ,  $B(4, 5)$  and  $C(6, 13)$ . Find  $\cos A$ .



[Watch Video Solution](#)



**28.** If the vertices of a triangle having integral coordinates . Prove that triangle can't be equilateral .



**Watch Video Solution**

**29.** If the coordinates of two points  $A$  and  $B$  are  $(3, 4)$  and  $(5, -2)$  , respectively, find the coordinates of any point  $P$  if  $PA = PB$ . Area of  $PAB$  is 10 sq. units.



**Watch Video Solution**

**30.** The coordinates of  $A, B, C$  are  $(6, 3), (-3, 5), (4, -2)$ , respectively, and  $P$  is any point  $(x, y)$ . Show that the ratio of the area of  $PBC$  to that of  $ABC$  is  $\frac{|x + y - 2|}{7}$ .



**Watch Video Solution**

**31.** Find the coordinates of points lying on the line joining  $P(3, -4)$  and  $Q(-2, 5)$  that is twice as far from  $P$  as  $Q$



**Watch Video Solution**

**32.** Determine the ratio in which the line  $3x + y - 9 = 0$  divides the segment joining the points  $(1,3)$  and  $(2, 7)$ .



**Watch Video Solution**

**33.** Prove that:  $(4, -1)$ ,  $(6, 0)$ ,  $(7, 2)$  and  $(5, 1)$  are the vertices of a rhombus. Is it a square?



**Watch Video Solution**

**34.** if the coordinates of the mid points of the Sides of a triangle are  $(1, 2)$ ,  $(0, -1)$  and  $(2, -1)$ . Find the coordinates of its vertices :



**Watch Video Solution**

**35.** Two vertices of a triangle are  $(3, -5)$  and  $(-7, 4)$ . If its centroid is  $(2, -1)$ , find the third vertex.



**Watch Video Solution**

**36.** If the line segment joining the points  $P(x_1, y_1)$  and  $Q(x_2, y_2)$  subtends an angle  $\alpha$  at the origin  $O$ , prove that :

$$OP \cdot OQ \cos \alpha = x_1 x_2 + y_1 y_2.$$

 [Watch Video Solution](#)

**37.** Four points  $A(6, 3)$ ,  $B(-3, 5)$ ,  $C(4, -2)$  and  $D(x, 3x)$  are given in such a way that  $\frac{\Delta DBC}{\Delta ABC} = \frac{1}{2}$ , find  $x$

 [Watch Video Solution](#)

**38.** The points  $A(2, 0)$ ,  $B(9, 1)$ ,  $C(11, 6)$  and  $D(4, 4)$  are the vertices of a quadrilateral  $ABCD$ . Determine whether  $ABCD$  is a rhombus or not.



**Watch Video Solution**

**39.** Find the coordinates of the centre of the circle inscribed in a triangle whose angular points are  $(-36, 7)$ ,  $(20, 7)$  and  $(0, -8)$ .



Watch Video Solution

**40.** The base of an equilateral triangle with side  $2a$  lies along the  $y$ -axis such that the mid point of the base is at the origin. Find the vertices of the triangle.



Watch Video Solution

**41.** Find the distance between  $P(x_1, y_1)$  and  $Q(x_2, y_2)$  when i.  $PQ$  is

parallel to the y-axis ii. PQ is parallel to the x-axis.



[Watch Video Solution](#)

**42.** Find a point on the x-axis, which is equidistant from the point (7,6) and (3,4).



[Watch Video Solution](#)

**43.** Find the equation to the locus of a point which moves so that the sum of its distances



from (3,0) and (-3,0) is less than 9.



[Watch Video Solution](#)

**44.** Find the locus of the point of intersection of lines  $x \cos \alpha + y \sin \alpha = a$  and  $x \sin \alpha - y \cos \alpha = b$  ( $\alpha$  is a variable).



[Watch Video Solution](#)

**45.** A rod of length  $l$  slides with its ends on two perpendicular lines. Find the locus of its

midpoint.



[Watch Video Solution](#)

**46.** If  $O$  is the origin and  $Q$  is a variable points on  $x^2 = 4y$ . Find the locus of the mid pint of  $OQ$ .



[Watch Video Solution](#)

**47.** Find the locus of a point equidistant from the point  $(2,4)$  and the  $y$ -axis.



[Watch Video Solution](#)

**48.** Find the locus of a point such that the sum of its distances from the points  $(0, 2)$  and  $(0, -2)$  is 6.



[Watch Video Solution](#)

**49.** Find the locus of a point which is equidistant from  $(1, 3)$  and x-axis.



[Watch Video Solution](#)

**50.** Find the locus of a point which moves such that its distance from the origin is three times its distance from x-axis.



**Watch Video Solution**

**51.** Find the locus of a point such that the line segments having end points  $(2,0)$  and  $(-2,0)$  subtend a right angle at that point.



**Watch Video Solution**

52. If  $A(-1, 1)$  and  $B(2, 3)$  are two fixed points, find the locus of a point  $P$  so that the area of  $\Delta PAB = 8$  sq. units.



Watch Video Solution

53. If  $O$  is the origin and  $Q$  is a variable point on  $y^2 = x$ . Find the locus of the mid point of  $OQ$ .



Watch Video Solution

54. At what point should the origin be shifted if the coordinates of a point  $(4, 5)$  become  $(-3, 9)$ ?



[Watch Video Solution](#)

55. What does the equation  $(x - a)^2 + (y - b)^2 = r^2$  become when the axes are transferred to parallel axes through the point  $(a - c, b)$ ?



[Watch Video Solution](#)

56. What does the equation  $(a - b)(x^2 + y^2) - 2abx = 0$  become if the origin is shifted to the point  $\left(\frac{ab}{a - b}, 0\right)$  without rotation?



[Watch Video Solution](#)

57. Find what the following equation become when the origin is shifted to the point (1,1):

$$x^2 + xy - 3x - y + 2 = 0$$



[Watch Video Solution](#)

**58.** Find what the following equation become when the origin is shifted to the point (1,1):

$$xy - x - y + 1 = 0$$



**Watch Video Solution**

**59.** Find what the following equation become when the origin is shifted to the point (1,1):

$$x^2 - y^2 - 2x + 2y = 0$$



**Watch Video Solution**



**60.** Find what the following equation become when the origin is shifted to the point (1,1):

$$xy - y^2 - x + y = 0$$



[Watch Video Solution](#)

**61.** Find the point to which the origin should be shifted so that the equation  $y^2 - 6y - 4x + 13 = 0$  is transferred to the form  $y^2 + Ax = 0$



[Watch Video Solution](#)

**62.** Find what the following equations become when the origin is shifted to the point (1,1):

$$x^2 + xy - 3y^2 - y + 2 = 0$$



[Watch Video Solution](#)

**63.** Find what the following equations become when the origin is shifted to the point (1,1):

$$xy - x - y + 1 = 0$$



[Watch Video Solution](#)

**64.** Find what the following equations become when the origin is shifted to the point (1,1):

$$xy - y^2 - x + y = 0$$



[Watch Video Solution](#)

**65.** Find what the following equations become when the origin is shifted to the point (1,1):

$$x^2 - y^2 - 2x + 2y = 0$$



[Watch Video Solution](#)

**66.** Find the point to which the origin should be shifted after a translation of axes so that the following equations will have no first degree term:  $y^2 + x^2 - 4x - 8y + 3 = 0$



**Watch Video Solution**

**67.** Find the point to which the origin should be shifted after a translation of axes so that the following equations will have no first degree term:  $x^2 + y^2 - 5x + 2y - 5 = 0$



**Watch Video Solution**

**68.** Find the point to which the origin should be shifted after a translation of axes so that the following equations will have no first degree term:  $x^2 - 12x + 4 = 0$



**Watch Video Solution**

**69.** Verify that the area of the triangle with vertices (4,6), (7,10) and (1,-2) remains invariant

under the translation of axes when the origin is shifted to the point  $(-2,1)$ .



[Watch Video Solution](#)

**70.** The vertices of a triangle are  $O(0, 0)$ ,  $A(a, 0)$  and  $B(0, b)$ . Write the coordinates of its circumcentre.



[Watch Video Solution](#)

71. Write the coordinates of the orthocentre of the triangle formed by points  $(8,0)$ ,  $(4,6)$  and  $(0,0)$



[Watch Video Solution](#)

72. Three vertices of a parallelogram, taken in order, are  $(-1, -6)$ ,  $(2,-5)$  and  $(7,2)$ . Write the coordinates of its fourth vertex.



[Watch Video Solution](#)

73. If the points  $(a, 0)$ ,  $(at_1^2, 2at_1)$  and  $(at_2^2, 2at_2)$  are collinear, write the value of  $t_1 t_2$ .



[Watch Video Solution](#)

74. Write the co ordinates of the circumcentre of a triangle whose centroid and orthocenter are at  $(3, 3)$  and  $(-3, 5)$  respectively



[Watch Video Solution](#)



**75.** Write the coordinates of the incentre of the triangle having its vertices at  $(0,0)$ ,  $(5,0)$  and  $(0,12)$ .



**Watch Video Solution**

**76.** If the points  $(1,-1)$ ,  $(2,-1)$  and  $(4,-3)$  are the mid points of the sides of a triangle then write the coordinates of its centroid.



**Watch Video Solution**

77. Write the area of the triangle having vertices at  $(a, b + c)$ ,  $(b, c + a)$ ,  $(c, a + b)$ .



**Watch Video Solution**