



MATHS

BOOKS - RD SHARMA MATHS (ENGLISH)

STATISTICS

Others

1. From the data given below state which group is more variable G_1 or G_2 ? Marks 10-20 20-30 30-40 40-50 50-60 60-70 70-80 Group G_1 9 17 32 33 40 10 9 Group G_2 10 20 30 25 43 15 7

[Watch Video Solution](#)

2. From the prices of shares X and Y given below: find out which is more stable value: X: 35 54 52 53 56 58 52 50 50 49
Y: 108 107 105 105 106 107 104 103 104 101



Watch Video Solution

3. The following are some particulars of the distribution of weights of boys and girls in a class:

	Boys	Girls
Number	100	50
Mean	60kg	45kg
Variance	9	4

Which of the distributions is more variable?



Watch Video Solution

4. For a frequency distribution mean deviation from mean is

computed by: (a) $M.D. = \frac{\sum f}{\sum f|d|}$ (b) $M.D. = \frac{\sum d}{\sum f}$ (c) $M.D. = \frac{\sum fd}{\sum f}$ (d) $M.D. = \frac{\sum f|d|}{\sum f}$



Watch Video Solution

5. Calculate mean deviation about mean from the following

data: x_i : 3 9 17 23 27 f_i : 8 10 12 9 5



Watch Video Solution

6. Calculate the mean deviation about median age for the

age distribution of 100 persons given below: Age 16-20

21-25 26-30 31-35 36-40 41-45 46-50 51-55

Number 5 6 12 14 26 12

16 9



Watch Video Solution

7. Calculate mean deviation from the median of the following data: Class interval: 0-6 6-12 12-18 18-24 24-30
Frequency 4 5 3 6 2



Watch Video Solution

8. Find the mean deviation about the mean for the following data: Marks obtained 10-20 20-30 30-40 40-50 50-60 60-70
70-80 Number of students 2 3 8 14 8 3 2



Watch Video Solution

9. The age distribution of 100 life-insurance policy holders is as follows: Age (on nearest birthday) 17-19.5 20-25.5 26-35.5 36-40.5 41-50.5 51-55.5 56-60.5 61-70.5 No. of persons 5 16 12 26 14 12 6 5 Calculate the mean deviation from median age.



Watch Video Solution

10. Calculate the median deviation from the median of the following data: Wages per week (in Rs.), 10-20, 20-30, 30-40, 40-50, 50-60, 60-70, 70-80 No. of workers, 4, 6, 10, 20, 10, 6, 4



Watch Video Solution

11. Find the mean deviation from the mean of the following data: Classes:, 10-20, 20-30, 30-40, 40-50, 50-60, 60-70, 70-80
frequency, 2, 3, 8, 14, 8, 3, 2



Watch Video Solution

12. Find the mean deviation from the median for the following data: Size:, 10, 11, 12, 14, 15 Frequency:, 2, 3, 8, 3, 4



Watch Video Solution

13. Find the mean deviation about the median of the following frequency distribution: Class:, 0-6, 6-12, 12-18, 18-24, 24-30 Frequency:, 8, 10, 12, 9, 5



Watch Video Solution

[Watch Video Solution](#)

14. Find the mean deviation from the mean for the following data: Size:, 20, 21, 22, 23, 24 frequency, 6, 4, 5, 1, 4

[Watch Video Solution](#)

15. Find the mean deviation from the mean for the following data: Size: 1 3 5 7 9 11 13 15 Frequency: 3 3 4 14 7 4 3 4

[Watch Video Solution](#)

16. The following table gives the number of finished articles turned out per day by different number of workers in a factory. Find the standard deviation of the daily output of

finished articles. Find the variance and standard deviation of the following frequency distribution: Number of article:, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27 No. of workers:, 3, 7, 11, 14, 18, 17, 13, 8, 5, 4



Watch Video Solution

17. If a is positive integer and the frequency distribution: x : , $A, 2a, 3a, 4a, 5a, 6a$ f : , 2, 1, 1, 1, 1, 1 and variance is 160 then find the value of a .



Watch Video Solution

18. There are 60 students in a class. The following is the frequency distribution of marks obtained by the students in

a test: Marks: 0 1 2 3 4 5 Frequency $x - 2$, x , x^2 , $(x + 1)^2$, $2x$, $x + 1$ Determine the mean and standard deviation.



Watch Video Solution

19. Find the mean, and standard deviation for the following data: (i) Year render: 10 20 30 40 50 60 No. of persons (cumulative): 15 32 51 78 97 109



Watch Video Solution

20. Find the mean and standard deviation of first n terms of an A.P. whose first term is a and common difference is d .



Watch Video Solution

21. Find the mean of the following frequency distribution:

Variable (x_i) , 2, 4, 6, 8, 10, 12, 14, 16 Frequency (f_i) , 4, 4, 5,

15, 8, 5, 4, 5



Watch Video Solution

22. Calculate the variance and standard deviation from the

data given below: Size of item, 3.5, 4.5, 5.5, 6.5, 7.5, 8.5, 9.5

Frequency , 3, 7, 22, 60, 85, 32, 8



Watch Video Solution

23. Find the variance and standard deviation for the

following distribution: X : 4.5, 14.5, 24.5, 34.5, 44.5, 54.5, 64.5

f : , 1, 5, 12, 22, 17, 9, 4



[Watch Video Solution](#)

24. Find the standard deviation for the following data: (i) x :
3, 8, 13, 18, 23 f : 7, 10, 15, 10, 6



[Watch Video Solution](#)

25. Calculate the mean and standard deviation for the following distribution: Marks: 20-30, 30-40, 40-50, 50-60, 60-70, 70-80, 80-90 No. of students: 3, 6, 13, 15, 14, 5, 4



[Watch Video Solution](#)

26. Let x_1, x_2, \dots, x_n be x_n observations. Let $y_i = ax_i + b$ for $i=1,2,\dots,n$ where a and b are constants. If the mean of x_i 's is 48 and their standard deviation is 12, the mean of y_i 's is 55 and standard deviation of y_i 's is 15, the values of a and b are (a) $a = 1.25, b = -5$ (b) $a = -1.25, b = 5$ (c) $a = 2.5, b = -5$ (d) $a = 2.5, b = 5$



Watch Video Solution

27. For a frequency distribution standard deviation is computed by applying the formula (a)

$$\sigma = \sqrt{\frac{\sum fd^2}{\sum f} - \left(\frac{\sum fd}{\sum f}\right)^2} \quad (b)$$

$$\sigma = \sqrt{\left(\frac{\sum fd}{\sum f}\right)^2 - \frac{\sum fd^2}{\sum f}} \quad (c) \quad \sigma = \sqrt{\frac{\sum fd^2}{\sum f} - \frac{\sum fd}{\sum f}}$$

$$(d) \quad \sigma = \sqrt{\left(\frac{\sum fd}{\sum f}\right)^2 - \frac{\sum fd^2}{\sum f}}$$



[Watch Video Solution](#)

28. The mean of 100 numbers observations is 50 and their standards deviation is 5. The sum of all squares of all the observations is (a) 50,000 (b) 250,000 (c) 252500 (d) 255000



[Watch Video Solution](#)

29. If $n = 10$, $X = 12$ and $\sum x_i^2 = 1530$, then the coefficient of variation is (a) 36% (b) 41% (c) 25% (d) none of these



[Watch Video Solution](#)

30. Life of bulbs produced by two factories A and B are given below: Length of life (in hours): 550-650 650-750 750-850 850-950 950-1050
 Factory A: (Number of bulbs) 10 22 52 20 16
 Factory B: (Number of bulbs) 8 60 24 16 12



Watch Video Solution

31. If the standard deviation of a variable X is σ , then standard deviation of variable $\frac{aX + b}{c}$ is (a) $a\sigma$ (b) $\frac{a}{c}\sigma$ (c) $\left|\frac{a}{c}\right|\sigma$ (d) $\frac{a\sigma + b}{c}$



Watch Video Solution

32. Following are the marks obtained, out of 100, by two students Ravi and Hashina in 10 life? Ravi: 25 50 45 30 70 42 36 48 35 60 Hashina: 10 70 50 20 95 55 42 60 48 80 .Who is more intelligent and who is more consistent?



Watch Video Solution

33. Let a, b, c, d, e , be the observations with m and standard deviation s . The standard deviation of the observations $a+k, b+k, c+k, d+k, e+k$ is (a) s (b) ks (c) $s + k$ (d) $\frac{s}{k}$



Watch Video Solution

34. The lengths (in cm) of 10 rods in a shop are given below:
40.0, 52.3, 55.2, 72.9, 52.8, 79.0, 32.5, 15.2, 27.9, 30.2 Find mean deviation from median (ii) find mean deviation from the mean also.



Watch Video Solution

35. If the S.D. of a set of observations is 8 and if each observation is divided by -2, the S.D. of the new set of observations will be (a) -4 (b) -8 (c) 8 (d) 4



Watch Video Solution

36. The standard deviation of the observations 6,5,9,13,12,8,10 is 6 (b) $\sqrt{6}$ (c) $\frac{52}{7}$ (d) $\sqrt{\frac{52}{7}}$



Watch Video Solution

37. The standard deviation of first 10 natural numbers is (a) 8.25 (b) 6.5 (c) 3.87 (d) 2.87



Watch Video Solution

38. The mean deviation of the data 3,10,10,4,7,10,5 (a) 2 (b) 2.57 (c) 3 (d) 3.57



Watch Video Solution

39. The mean deviation from the median is (a) equal to that measured from another value (b) maximum if all observations are positive (c) greater than that measured from any other value. (d) less than that measured from any other value.



Watch Video Solution

40. The weight of coffee in 70 jars is shown in the following table: Weight (in grams): 200 – 201 201 – 202 202 – 203 203 – 204 204-205 205-206 *Frequency* 13 27 18 10 1 1
Determine the variance and standard deviation of the above distribution.



Watch Video Solution

41. Find the mean and variance of frequency distribution given below: x_i : $1 \leq x < 3$ $3 \leq x < 5$ $5 \leq x < 7$ $7 \leq x < 10$ f_i : 6 4 5 1



Watch Video Solution

42. While calculating the mean and variance of 10 readings, a student wrongly used the reading of 52 for the correct reading 25. He obtained the mean and variance as 45 and 16 respectively. Find the correct mean and the variance.



Watch Video Solution

43. Mean and standard deviation of 100 observations were found to be 40 and 10 respectively. If at the time of

calculation two observations were wrongly taken as 30 and 70 in place of 3 and 27 respectively. Find the correct standard deviation.



[Watch Video Solution](#)

44. Calculate the mean and standard deviation for the following data: Wages upto (into Rs.) 15 30 45 60 75 90 105 120
No. of workers 12 30 65 107 157 202 222 230



[Watch Video Solution](#)

45. The following table gives the distribution of income of 100 families in a village. Calculate the standard deviation:

Income Rs.	0-1000	1000-2000	2000-3000	3000-4000	4000-5000	5000-6000	No. of Families	18	26	30	12	10	4
------------	--------	-----------	-----------	-----------	-----------	-----------	-----------------	----	----	----	----	----	---


[Watch Video Solution](#)

46. Calculate the mean and standard deviation for the following table given the age distribution of a group of people:

Age:	20-30	30-40	40-50	50-60	60-70	70-80	80-90	No.
of persons:	3	51	122	141	130	51	2	


[Watch Video Solution](#)

47. The measurements of the diameters (in mm) of the heads of 107 screws are given below:

Diameter (in mm)	33-35
------------------	-------

36-38 39-41 42-44 45-47 No. of screws 10 19 23 21 27. calculate mean diameter of head of screw.



Watch Video Solution

48. The mean deviation of the numbers 3, 4, 5, 6, 7 from the mean is (a) 25 (b) 5 (c) 1.2 (d) 0



Watch Video Solution

49. A batsman scores run in 10 innings as 38,70,48,34,42,55,63,46,54 and 44. The mean deviation about mean is (a) 8.6 (b) 6.4 (c) 10.6 (d) 7.6



Watch Video Solution

50. The mean deviation of the series $a, a + d, a + 2d, \dots, a + 2n$ from its mean is (a) $\frac{(n+1)d}{2n+1}$ (b) $\frac{nd}{2n+1}$ (c) $\frac{n(n+1)d}{2n+1}$ (d) $\frac{(2n+1)d}{n(n+1)}$



Watch Video Solution

51. The following the record of goals scored by team a in football session. Number of goals scored: 0 1 2 3 4 Number of matches: 1 9 7 5 3 For the team B, mean number of goals scored per match was 2 with a standard deviation 1.25 goals. Find which team may be considered more consistent?



Watch Video Solution

52. An analysis of monthly wages paid to the works of two firms A and B belonging to the same industry gives the following results:

	Firm A	Firm B
Number of workers	1000	1200
Average monthly wages	Rs. 2800	Rs. 2800
Variance of distribution of wages	100	169

In which firm, A or B is there greater variability in individual wages?



Watch Video Solution

53. Let x_1, x_2, \dots, x_n be values taken by a variable X and y_1, y_2, \dots, y_n be the values taken by a variable Y such that $y_i = ax_i + b, i=1,2,3,\dots$ Then, (a) $\text{Var.}(Y) = a^2 \text{Var.}(X)$ (b) $\text{Var.}(X) = a^2 \text{Var.}(Y)$ (c) $\text{Var.}(X) = \text{Var.}(X) + b$ (d) none of these



Watch Video Solution

54. The algebraic sum of deviations of 10 observations measured from 16 is 7. The mean is (a) 105 (b) 70 (c) 15.7 (d) None of these



Watch Video Solution

55. Calculate the mean deviation about median from the following data: 340, 150, 210, 240, 300, 310, 320.



Watch Video Solution

56. For a group of 200 candidates the mean and S.D. were found to be 40 and 15 respectively. Later on it was found

that the score 43 was misread as 34. Find the correct mean and correct S.D.



Watch Video Solution

57. Find the mean deviation from the mean for the data:
6,7,10,12,13,4,8,20.



Watch Video Solution

58. The scores of a batsman in ten innings are :
38,70,48,34,42,55,63,46,54,44. Find the mean deviation about the median.



Watch Video Solution

59. Calculate the mean deviation about the mean of the set of first n natural numbers when n is even natural number.



Watch Video Solution

60. Calculate the mean deviation about the mean of the set of first n natural numbers when n is odd natural number.



Watch Video Solution

61. Find the variance and standard deviation of the following data 5,12,3,18,6,8,2,10



Watch Video Solution

62. Compute the variance and standard deviation of the following observations of marks of 5 students of a tutorial group: Marks out of 25: 8, 12, 13, 15, 22



Watch Video Solution

63. The number of telephone calls received at an exchange in 245 successive one-minute intervals are shown in the following frequency distribution:

Number of calls	0	1	2	3	4	5	6	7
Frequency	14	21	25	43	51	40	39	12



Watch Video Solution

64. Calculate mean deviation from the median for the following distribution:

x_i	10	15	20	25	30	35	40	45
f_i	7	3	8					

[Watch Video Solution](#)

65. The mean deviation for n observations x_1, x_2, \dots, x_n from their mean X is given by (a) $\sum_{i=1}^n (x_i - X)$ (b) $\frac{1}{n} \sum_{i=1}^n (x_i - X)$ (c) $\sum_{i=1}^n (x_i - X)^2$ (d) $\frac{1}{n} \sum_{i=1}^n (x_i - X)^2$

[Watch Video Solution](#)

66. Let x_1, x_2, \dots, x_n be n observations, and let \bar{x} be their arithmetic mean and σ^2 be their variance. Statement 1: Variance of $2x_1, 2x_2, \dots, 2x_n$ is $4\sigma^2$. Statement 2: Arithmetic mean of $2x_1, 2x_2, \dots, 2x_n$ is $4\bar{x}$. (1) Statement 1 is false, statement 2 is true (2) Statement 1 is

true, statement 2 is true; statement 2 is a correct explanation for statement 1 (3) Statement 1 is true, statement 2 is true; statement 2 is not a correct explanation for statement 1 (4) Statement 1 is true, statement 2 is false



Watch Video Solution

67. The scores of a batsman in 10 matches were as follows: 38,70,48,34,42,55,63,46,54,44 compute the variance and standard deviation.



Watch Video Solution

68. The mean and variance of 7 observation is 8 and 16. If five of the observations are 2, 4, 10, 12, 14. Find the remaining

two observations.



Watch Video Solution

69. If for a sample of size 60, we have the following information $\sum x_i^2 = 18000$ and $\sum x_i = 960$, then the variance is (a) 6.63 (b) 16 (c) 22 (d) 44



Watch Video Solution

70. The mean and standard deviation of 20 observations are found to be 10 and 2, respectively. One rechecking, it was found that an observation 8 was incorrect. Calculate the correct mean and standard deviation in each of the

following cases. (i) If wrong item is omitted (ii) If it is replaced by 12.



Watch Video Solution

71. Consider the numbers 1,2,3,4,5,6,7,8,9,10. If 1 is added to each number, the variance of the numbers so obtained is (a) 6.5 (b) 2.87 (c) 3.87 (d) 8.25



Watch Video Solution

72. If for distribution of 18 observations $\sum (x_i - 5) = 3$ and $\sum (x_i - 5)^2 = 43$, find the mean and standard deviation.



Watch Video Solution

73. If the mean and standard deviation of 100 observations are 50 and 4 respectively. Find the sum of all the observations and the sum of their squares.



Watch Video Solution

74. Let x_1, x_2, \dots, x_n values of variable X and let ' a ' be non zero real number. Then prove that the variance of the observations ax_1, ax_2, \dots, ax_n is $a^2 \text{Var}(X)$. Also, find their standard deviation.



Watch Video Solution

75. If two variates X and Y are connected by the relation

$$Y = \frac{aX + b}{c}, \text{ where } a, b, c \text{ are constants such that}$$

$$ac < 0, \text{ then (a) } \sigma_Y = \frac{a}{c} \sigma_X \quad \text{(b) } \sigma_Y = -\frac{a}{c} \sigma_X \quad \text{(c)}$$

$$\sigma_Y = \frac{a}{c} \sigma_X + b \quad \text{(d) none of these}$$



Watch Video Solution

76. Let $x_1, x_2, x_3, \dots, x_n$ be n values of a variable X . If these

values are changed to

$x_1 + a, x_2 + a, \dots, x_n + a$, where $a \in R$, show that the

variance remains unchanged.



Watch Video Solution

77. If for a sample of size 60, we have the following information $\sum x_i^2 = 18000$ and $\sum x_i = 960$, then the variance is (a) 6.63 (b) 16 (c) 22 (d) 44



Watch Video Solution

78. The weight of coffee in 70 jars is shown in the following table: Weight (in grams):, 200 – 201 , 201 – 202 , 202 – 203 , 203 – 204 , 204-205, 205-206 *Frequency* , 13, 27, 18, 10, 1, 1
Determine the variance and standard deviation of the above distribution.



Watch Video Solution

79. Suppose that samples of polythene bags from two manufacturers, A and B, are tested by a prospective buyer for bursting pressure, with the following results. Bursting Pressure in kg, Number of bags manufactured by manufacturer

Pressure in kg	Number of bags (A)	Number of bags (B)
5-10	2	9
10-15	9	11
15-20	29	18
20-25	54	32
25-30	11	27
30-35	5	13

Which set of the bags has the highest average bursting pressure?



Watch Video Solution

80. Let x_1, x_2, \dots, x_n be values taken by a variable X and y_1, y_2, \dots, y_n be the values taken by a variable Y such that $y_i = ax_i + b, i = 1, 2, \dots, n$. Then, (a)

$Var(Y) = a^2 Var(X)$ (b) $Var(X) = a^2 Var(Y)$ (c)

$Var(X) = Var(X) + b$ (d) none of these



[Watch Video Solution](#)

81. The sum of the squares of deviation of 10 observations from their mean 50 is 250, then coefficient of variation is



[Watch Video Solution](#)

82. If the S.D. of a set of observations is 8 and if each observation is divided by -2, the S.D. of the new set of observations will be (a) -4 (b) -8 (c) 8 (d)

4



[Watch Video Solution](#)

83. Find the mean and variance of frequency distribution given below: x_i : , $1 \leq x < 3$, $3 \leq x < 5$, $5 \leq x < 7$, $7 \leq x < 10$ f_i : , 6, 4, 5, 1



Watch Video Solution

84. The mean deviation of the series $a, a + d, a + 2d, \dots, a + 2n$ from its mean is $\frac{(n+1)d}{2n+1}$ (b) $\frac{nd}{2n+1}$ (c) $\frac{n(n+1)d}{2n+1}$ (d) $\frac{(2n+1)d}{n(n+1)}$



Watch Video Solution

85. The standard deviation of the data: x : , $1, a, a^2, \dots, a^n$ f : , nC_0 , nC_1 , nC_2 , ..., nC_n is

$$\left(\frac{1+a^2}{2}\right)^n - \left(\frac{1+a}{2}\right)^{2n} \quad (b)$$

$$\left(\frac{1+a^2}{2}\right)^{2n} - \left(\frac{1+a^2}{2}\right)^{2n} \quad (c)$$

$$\left(\frac{1+a}{2}\right)^{2n} - \left(\frac{1+a^2}{2}\right)^n \quad (d) \text{ none of these}$$



Watch Video Solution

86. The mean deviation of the numbers 3,4,5,6,7 from the mean is (a) 25 (b) 5 (c) 1.2 (d) 0



Watch Video Solution

87. A batsman scores run in 10 innings as 38,70,48,34,42,55,63,46,54 and 44. The mean deviation about mean is 8.6 (b) 6.4 (c) 10.6 (d) 7.6



Watch Video Solution

88. The scores of a batsman in 10 matches were as follows:
38,70,48,34,42,55,63,46,54,44 compute the variance and standard deviation.



Watch Video Solution

89. Calculate mean for the following data: x_i : 3 9 17 23 27
 f_i : 8 10 12 9 5



Watch Video Solution

90. Calculate the mean and standard deviation for the following data: Wages upto (into Rs.), 15, 30, 45, 60, 75, 90,

105, 120 No. of workers, 12, 30, 65, 107, 157, 202, 222, 230



[Watch Video Solution](#)

91. The following table gives the distribution of income of 100 families in a village. Calculate the standard deviation:

Income Rs.,	0-1000,	1000-2000,	2000-3000,	3000-4000,	4000-5000,	5000-6000
No. of Families,	18,	26,	30,	12,	10,	4



[Watch Video Solution](#)

92. While calculating mean and variance of 10 readings, a student wrongly used the reading of 52 for the correct reading 25. He obtained the mean and variance as 45 and 16 respectively. Find the correct mean and the variance.

[Watch Video Solution](#)

93. Mean and standard deviation of 100 observations were found to be 40 and 10 respectively. If at the time of calculation two observations were wrongly taken as 30 and 70 in place of 3 and 27 respectively. Find the correct standard deviation.

[Watch Video Solution](#)

94. The number of telephone calls received at an exchange in 245 successive one-minute intervals are shown in the following frequency distribution: Number of calls, 0, 1, 2, 3, 4, 5, 6, 7 Frequency, 14, 21, 25, 43, 51, 40, 39, 12 Calculate mean deviation about Median.

[Watch Video Solution](#)

95. Calculate mean deviation from the median for the following distribution: x_i : , 10, 15, 20, 25, 30, 35, 40, 45 f_i : , 7, 3, 8, 5, 6, 8, 4, 9

[Watch Video Solution](#)

96. Find the variance and standard deviation for the following data: 4,6,10,12,14,18,20

[Watch Video Solution](#)

97. Complete the variance and standard deviation of the following observations of marks of 5 students of a tutorial

group: Marks out of 25:8,12,13,15,22



Watch Video Solution

98. Calculate the mean deviation about the mean of the set of first n natural numbers when n is even natural number.



Watch Video Solution

99. Let $x_1, x_2, x_3, \dots, x_n$ be n values of a variable X . If these values are changed to $x_1 + a, x_2 + a, \dots, x_n + a$, where $a \in R$, show that the variance remains unchanged.



Watch Video Solution

100. Find the mean deviation from the mean for the data:
6,7,10,12,13,4,8,20.



Watch Video Solution

101. The scores of a batsman in ten innings are :
38,70,48,34,42,55,63,46,54,44. Find the mean deviation about
the median.



Watch Video Solution

102. Calculate the mean deviation about median from the
following data: 340, 150, 210, 240, 300, 310, 320.



Watch Video Solution

103. For a group of 200 candidates the mean and S.D. were found to be 40 and 15 respectively. Later on it was found that the score 43 was misread as 34. Find the correct mean and correct S.D.



Watch Video Solution

104. The mean deviation from e median is a.equal to that measured from another value b.maximum if all observations are positive c,greater than that measured from any other value.d,less than that measured form any other value.



Watch Video Solution

105. Calculate the mean deviation about the mean of the set of first n natural numbers when n is odd natural number.



Watch Video Solution

106. Let x_1, x_2, \dots, x_n values of variable X and let ' a ' be non zero real number. Then prove that the variance of the observations ax_1, ax_2, \dots, ax_n is $a^2 \text{Var}(X)$. Also, find their standard deviation.



Watch Video Solution

107. Life of bulbs produced by two factories A and B are given below: Length of life (in hours);, 550-650, 650-750, 750-

850, 850-950, 950-1050 Factory A: (Number of bulbs), 10, 22, 52, 20, 16 Factory B (Number of bulbs), 8, 60, 24, 16, 12 The bulbs of which factory are more consistent from the point of view of length of life?



[Watch Video Solution](#)

108. If for distribution of 18 observations $\sum (x_i - 5) = 3$ and $\sum (x_i - 5)^2 = 43$, find the mean and standard deviation.



[Watch Video Solution](#)

109. If the mean and standard deviation of 100 observations are 50 and 4 respectively. Find the sum of all the

observations and the sum of their squares.



[Watch Video Solution](#)

110. Find the standard deviation for the following data: (i)

$x: , 3, 8, 13, 18, 23$ $f: , 7, 10, 15, 10, 6$ (ii) $x: , 2, 3, 4, 5, 6, 7, 7$ $f:$

$, 4, 9, 16, 14, 11, 6, 6$



[Watch Video Solution](#)

111. The mean and standard deviation of 20 observations are found to be 10 and 2, respectively. One rechecking, it was found that an observation 8 was incorrect. Calculate the correct mean and standard deviation in each of the

following cases. (i) If wrong item is omitted (ii) If it is replaced by 12.

 [Watch Video Solution](#)

112. For a frequency distribution mean deviation from mean

in computed by (a) $\dot{MD} = \frac{\sum f}{\sum f|d|}$ (b) $\dot{MD} = \frac{\sum d}{\sum f}$ (c) $\dot{MD} = \frac{\sum fd}{\sum f}$ (d) $\dot{MD} = \frac{\sum f|d|}{\sum f}$

 [Watch Video Solution](#)

113. Calculate the mean and standard deviation for the following distribution: Marks:, 20-30, 30-40, 40-50, 50-60, 60-70, 70-80, 80-90 No. of students:, 3, 6, 13, 15, 14, 5, 4

 [Watch Video Solution](#)

114. The measurements of the diameters (in mm) of the heads of 107 screws are given below: Diameter (in mm), 33-35, 36-38, 39-41, 42-44, 45-47 No. of screws, 10, 19, 23, 21, 27. Find the mean diameter.



Watch Video Solution

115. Let a, b, c, d, e , be the observations with m and standard deviation s . The standard deviation of the observations $a+k, b+k, c+k, d+k, e+k$ is s (b) ks (c) $s + k$ (d) $\frac{s}{k}$



Watch Video Solution

116. If $n = 10$, $X = 12$ and $\sum x^2 = 1530$, then the coefficient of variation is (a) 36% (b) 41% (c) 25% (d) none of these



Watch Video Solution

117. The mean deviation for n observations x_1, x_2, \dots, x_n from their mean X is given by (a)

$\sum_{i=1}^n (x_i - X)$ (b) $\frac{1}{n} \sum_{i=1}^n |x_i - X|$ (c) $\sum_{i=1}^n (x_i - X)^2$ (d)

$\frac{1}{n} \sum_{i=1}^n (x_i - X)^2$



Watch Video Solution

118. The standard deviation of the observations 6,5,9,13,12,8,10 is (a) 6 (b) $\sqrt{6}$ (c) $\frac{52}{7}$ (d) $\sqrt{\frac{52}{7}}$



Watch Video Solution

119. If the standard deviation of a variable X is σ , then standard deviation of variable $\frac{aX + b}{c}$ is (a) $a\sigma$ (b) $\frac{a}{c}\sigma$ (c) $\left|\frac{a}{c}\right|\sigma$ (d) $\frac{a\sigma + b}{c}$



Watch Video Solution

120. Following are the marks obtained, out of 100, by two students Ravi and Hashina in 10 life? Ravi: 25 50 45 30 70 42

36 48 35 60 Hashina: 10 70 50 20 95 55 42 60 48 80 .Who is more intelligent and who is more consistent?



Watch Video Solution

121. Consider the numbers 1,2,3,4,5,6,7,8,9,10. If 1 is added to each number, the variance of the numbers so obtained is

A. 6.5

B. 2.87

C. 3.87

D. 4.87

Answer: null



Watch Video Solution

122. The mean of 100 numbers observations is 50 and their standards deviation is 5. The sum of all squares of all the observations is (a) 50,000 (b) 250,000 (c) 252500 (d) 255000



Watch Video Solution

123. The following is ht record of goals scored by team A in a football session For the team b, mean number of goals scored per match was 2 with a standard deviation 1.25 goals. Find which team may be considered more consistent?



Watch Video Solution

- 124.** The standard deviation of first 10 natural numbers is (a) 8.25 (b) 6.5 (c) 3.87 (d) 2.87



Watch Video Solution

- 125.** Let x_1, x_2, \dots, x_n be x_n observations. Let $y_i = ax_i + b$ for $i=1,2,\dots,n$ where a and b are constants. If the mean of x_i 's is 48 and their standard deviation is 12, the mean of y_i 's is 55 and standard deviation of y_i 's is 15, the values of a and b are (a) $a = 1.25, b = -5$ (b) $a = -1.25, b = 5$ (c) $a = 2.5, b = -5$ (d) $a = 2.5, b = 5$



Watch Video Solution

126. Let x_1, x_2, \dots, x_n be n observations and \bar{X} be their arithmetic mean. The standard deviation is given by (a)

$$\sum_{i=1}^n (x_i - \bar{X})^2 \quad (b) \quad \frac{1}{n} \sum_{i=1}^n (x_i - \bar{X})^2 \quad (c)$$

$$\sqrt{\frac{1}{n} \sum_{i=1}^n (x_i - \bar{X})^2} \quad (c) \quad \frac{1}{n} \sum_{i=1}^n x_i^2 - \bar{X}^2$$



Watch Video Solution

127. The following are some particulars of the distribution of weights of boys and girls in a class:

	Boys	Girls
Number	100	50
Mean	60kg	45kg
Variance	9	4

Which of the distributions is more variable?



Watch Video Solution

128. The mean deviation of the data 3,10,10,4,7,10,5 from the mean is (a) 2 (b) 2.57 (c) 3 (d) 3.57



Watch Video Solution

129. There are 60 students in a class. The following is the frequency distribution of marks obtained by the students in a test: Marks:, 0, 1, 2, 3, 4, 5 Frequency , $x - 2$, x , x^2 , $(x + 1)^2$, $2x$, $x + 1$. Determine mean and standard deviation of the marks.



Watch Video Solution

130. Find the mean deviation about median for the following

data : Marks		0-10	10-20	20-30	30-40	40-50
50-60		Number of	6	8	14	16
4		2 Girls				

 [Watch Video Solution](#)

131. The following table gives the number of finished articles turned out per day by different number of workers in a factory. Find the standard deviation of the daily output of finished articles. Find the variance and standard deviation of the following frequency distribution: Number of article:, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27 No. of workers:, 3, 7, 11, 14, 18, 17, 13, 8, 5, 4

 [Watch Video Solution](#)

132. If a is positive integer and the frequency distribution:
 $x: , A, 2a, 3a, 4a, 5a, 6a$ $f: , 2, 1, 1, 1, 1, 1$ and variance is 160
 then find the value of a .


[Watch Video Solution](#)

133. Calculate the mean deviation about median age for the
 age distribution of 100 persons given below:

Age	16-20	21-25	26-30	31-35	36-40	41-45	46-50	51-55
Number	5	6	12	14	26	12	16	9


[Watch Video Solution](#)

134. Calculate mean deviation from the median of the following data: Class interval: 0-6 6-12 12-18 18-24 24-30
Frequency 4 5 3 6 2



Watch Video Solution

135. Find the mean of first n terms of an A.P. whose first term is a and common difference is d .



Watch Video Solution

136. Find the variance and standard deviation of the following frequency distribution: Variable (x_i) , 2, 4, 6, 8, 10, 12, 14, 16 Frequency (f_i) , 4, 4, 5, 15, 8, 5, 4, 5



Watch Video Solution

137. Calculate the mean deviation from the median of the following data: Wages per week (in Rs.), 10-20, 20-30, 30-40, 40-50, 50-60, 60-70, 70-80 No. of workers, 4, 6, 10, 20, 10, 6, 4



Watch Video Solution

138. Find the mean deviation from the mean of the following data: Classes:, 10-20, 20-30, 30-40, 40-50, 50-60, 60-70, 70-80 frequency, 2, 3, 8, 14, 8, 3, 2



Watch Video Solution

139. Find the mean deviation about the mean for the following data: Marks obtained , 10-20, 20-30, 30-40, 40-50, 50-60, 60-70, 70-80 Number of students, 2, 3, 8, 14, 8, 3, 2



Watch Video Solution

140. The age distribution of 100 life-insurance policy holders is as follows: Age (on nearest birthday), 17-19.5, 20-25.5, 26-35.5, 36-40.5, 41-50.5, 51-55.5, 56-60.5, 61-70.5 No. of persons, 5, 16, 12, 26, 14, 12, 6, 5



Watch Video Solution

141. Find the mean deviation from the mean for the following data: Size:, 20, 21, 22, 23, 24 frequency, 6, 4, 5, 1, 4



Watch Video Solution

142. Find the mean deviation from the mean for the following data: Size: 1 3 5 7 9 11 13 15 Frequency: 3 3 4 14 7 4 3 4



Watch Video Solution

143. Find the mean deviation from the median for the following data: Size:, 10, 11, 12, 14, 15 Frequency:, 2, 3, 8, 3, 4



Watch Video Solution

144. Find the mean deviation about the median of the following frequency distribution: Class:, 0-6, 6-12, 12-18, 18-24, 24-30 Frequency:, 8, 10, 12, 9, 5



Watch Video Solution

145. Calculate the mean and standard deviation for the following table given the age distribution of a group of people: Age: 20-30 30-40 40-50 50-60 60-70 70-80 80-90 No. of persons: 3 51 122 141 130 51 2



Watch Video Solution

146. The lengths (in cm) of 10 rods in a shop are given below:
40.0, 52.3, 55.2, 72.9, 52.8, 79.0, 32.5, 15.2, 27.9, 30.2 Find mean deviation from median (ii) find mean deviation from the mean also.



Watch Video Solution

147. The mean and variance of 7 observations are 8 and 16, respectively. If five of the observations are 2, 4, 10, 12, 14. Find the remaining two observations.



Watch Video Solution

148. An analysis of monthly wages paid to workers in two firms A and B, belonging to the same industry, gives the following result and no. of workers of firm A and B are 586

and 648 respectively		No. of wage earners	Firm
A	Firm B	Mean of monthly wages	Rs
5253	Rs 5253		Variance of the
distribution	100	121	of wages (i)

Which firm A or B pays larger amount as monthly wages?

(ii) Which firm, A or B, show greater variability in individual wages?


[Watch Video Solution](#)

149. From the data given below state which group is more variable G_1 or G_2 ? Marks, 10-20, 20-30, 30-40, 40-50, 50-60,

60-70, 70-80 Group G_1 , 9, 17, 32, 33, 40, 10, 9 Group G_2 , 10, 20, 30, 25, 43, 15, 7



Watch Video Solution

150. From the prices of shares X and Y below, find out which is more stable in value X 35 54 52 53 56 58 52 50 51 49 . Y 108 107 105 105 106 107 104 103 104 101



Watch Video Solution

151. If two variates X and Y are connected by the relation $Y = \frac{aX + b}{c}$, where a, b, c are constants such that $ac < 0$, then (a) $\sigma_Y = \frac{a}{c}\sigma_X$ (b) $\sigma_Y = -\frac{a}{c}\sigma_X$ (c) $\sigma_Y = \frac{a}{c}\sigma_X + b$ (d) none of these

[Watch Video Solution](#)

152. Calculate the variance and standard deviation from the data given below: Size of item, 3.5, 4.5, 5.5, 6.5, 7.5, 8.5, 9.5
Frequency , 3, 7, 22, 60, 85, 32, 8

[Watch Video Solution](#)

153. Find the variance and standard deviation for the following distribution: X:, 4.5, 14.5, 24.5, 34.5, 44.5, 54.5, 64.5
 f : , 1, 5, 12, 22, 17, 9, 4

[Watch Video Solution](#)

154. Calculate the mean deviation about the median of the following observation: 3011,2780,3020,2354,3541,4150,5000



Watch Video Solution

155. The scores of a batsman in ten innings are : 38,70,48,34,42,55,63,46,54,44. Find the mean deviation about the median.



Watch Video Solution

156. Calculate the mean deviation about median of the following data: 34, 66, 30, 38, 44, 50, 40, 60, 42, 51



Watch Video Solution

157. Calculate the mean deviation about the median of the following observation: 22, 24, 30, 27, 29, 31, 25, 28, 41, 42



Watch Video Solution

158. Calculate the mean deviation about the median of the following observation: 38, 70, 48, 34, 63, 42, 55, 44, 53, 47



Watch Video Solution

159. Calculate the mean deviation about mean of the following data. (i) 1, 4, 7, 8, 9, 10, 12, 13, 17



Watch Video Solution

160. Calculate the mean deviation from the mean for the following data: 13, 17, 16, 14, 11, 13, 10, 16, 11, 18, 12, 17



Watch Video Solution

161. Calculate the mean deviation from the mean for the following data: 38, 70, 48, 40, 42, 55, 63, 46, 54, 44



Watch Video Solution

162. Calculate the mean deviation from the mean for the following data: 36, 72, 46, 42, 60, 45, 53, 46, 51, 49



Watch Video Solution

163. Calculate the mean deviation from the mean for the following data: 57, 64, 43, 67, 49, 59, 44, 47, 61, 59



Watch Video Solution

164. Calculate the mean deviation of the following income groups of five and seven members from their medians: 1st income group (in Rs.): 4000 4200 4400 4600 4800 2nd income group (in Rs.): 3800 4000 4200 4400 4600 4800 5800



Watch Video Solution

165. Calculate the mean deviation from the median of the following frequency distribution: 5 7 9 11 13 15 17 2 4 6 8 10 12

[Watch Video Solution](#)

166. Calculate the mean deviation about the median of the following frequency distribution:

	5	7	9	11	13	15	17
Having							
Frequency	2	4	6	8	10	12	8

[Watch Video Solution](#)

167. Find the mean deviation from the mean for the following data: 5 2 9 10 12 15 8 6 2 2 2 6

[Watch Video Solution](#)

168. Find the mean deviation from the median of the following data: 15 21 27 30 3 5 6 7



Watch Video Solution

169. Compute the mean deviation from the median of the following distribution:

Class	0-10	10-20	20-30	30-40	40-50
Frequency	5	10	20	5	10



Watch Video Solution

170. Find the mean deviation from the mean for the following data:

Classes	0-100	100-200	200-300	300-400	400-500	500-600	600-700	700-800
Frequencies	4	8	9	10	7	5	4	3



Watch Video Solution

 [Watch Video Solution](#)

171. Compute mean deviation from mean of the following distribution: Marks 10-20 20-30 30-40 40-50 50-60 60-70 70-80 80-90 No. of students 8 10 15 2 20 18 9 5

 [Watch Video Solution](#)

172. Find the deviation from the mean and from median of the following distribution: Marks 0-10 10-20 20-30 30-40 40-50 No. of Students 5 8 15 16 6

 [Watch Video Solution](#)

173. Calculate the mean deviation about the mean for the following frequency distribution

Class interval	Frequency
0 – 4, 4 – 8, 8 – 12, 12 – 16, 16 – 20	4, 6, 8, 5, 2



Watch Video Solution

174. Let x_1, x_2, \dots, x_n values of variable X and let ' a ' be non zero real number. Then prove that the variance of the observations ax_1, ax_2, \dots, ax_n is $a^2 \text{Var}(X)$. Also, find their standard deviation.



Watch Video Solution

175. Calculate the mean and standard deviation of first n natural numbers.



Watch Video Solution

176. Find the mean variance and standard deviation for the following data: 2,4,5,6,8,17



Watch Video Solution

177. Find the mean, variance and standard deviation for the following data: 6,7,10,12,13,4,8,12



Watch Video Solution

178. Find the mean, variance and standard deviation for the following data: 227,235,255,269,292,299,312,321,333,348



Watch Video Solution

179. The variance of 20 observations is 5. If each observation is multiplied by 2 find the variance of the resulting observations.



Watch Video Solution

180. The variance of 15 observations is 4. If each observations increased by 9 find the variance of the resulting observations.



Watch Video Solution

181. The mean of 5 observations is 4.4 and their variance is 8.24. If three of the observations are 1, 2 and 6, find the other two observations.



Watch Video Solution

182. The mean and standard deviation of six observations are 8 and 4, respectively. If each observation is multiplied by 3, find the new mean and new standard deviation of the resulting observations.



Watch Video Solution

183. The mean and variance of eight observations are 9 and 9.25, respectively. If six of the observations are 6, 7, 10, 12, 12 and 13, find the remaining two observations.



Watch Video Solution

184. For a group of 200 candidates, mean was found to 40 but it was discovered later that the scores of 43 and 35 were misread as 34 and 53 respectively. Then the correct mean is



Watch Video Solution

185. The mean and standard deviation of 100 observations were calculated as 40 and 5.1 respectively by a student who

took by mistake 50 instead of 40 for one observation. What are the correct mean and standard deviation?



Watch Video Solution

186. The mean and standard deviation of 20 observations are found to be 10 and 2, respectively. One rechecking, it was found that an observation 8 was incorrect. Calculate the correct mean and standard deviation in each of the following cases. (i) If wrong item is omitted (ii) If it is replaced by 12.



Watch Video Solution

187. The mean and standard deviation of a group of 100 observations were found to be 20 and 3, respectively. Later on it was found that three observations were incorrect, which are recorded as 21, 21 and 18. Find the mean and standard deviation if the incorrect observations are omitted.



Watch Video Solution

188. Show that the two formulae for the standard deviation of ungrouped data:

$$\sigma = \sqrt{\frac{1}{n} \sum (x_i - \bar{X})^2}$$

and $\sigma' = \sqrt{\frac{1}{n} \sum x_i^2 - \bar{X}^2}$ are equivalent where

$$\bar{X} = \frac{1}{n} \sum x_i.$$



Watch Video Solution

189. Find the variance and standard deviation for the following distribution: X : 4.5, 14.5, 24.5, 34.5, 44.5, 54.5, 64.5
 f : 1, 5, 12, 22, 17, 9, 4



Watch Video Solution

190. Table below shows the frequency f with which x alpha particles were radiated from a diskette x : 0 1 2 3 4 5 6 7 8 9
 10 11 12 f : 51 203 383 525 532 408 273 139 43 27 10 4 2.
 Calculate the mean and variance



Watch Video Solution

191. Calculate the mean and S.D for the following data:
 Expenditure (in Rs.) 0-10 10-20 20-30 30-40 40-50 Frequency:

14 13 27 21 15



[Watch Video Solution](#)

192. Calculate the standard deviation for the following data:

Class	0-30	30-60	60-90	90-120	120-150	150-180	180-210
-------	------	-------	-------	--------	---------	---------	---------

Frequency:	9	17	43	82	81	44	24
------------	---	----	----	----	----	----	----



[Watch Video Solution](#)

193. The mean and standard deviation of 100 observations were calculated as 40 and 5.1 respectively by a student who took by mistake 50 instead of 40 for one observation. What are the correct mean and standard deviation?



[Watch Video Solution](#)

194. Calculate the mean, median and standard deviation of the following distribution: Class interval 31-35 36-40 41-45 46-50 51-55 56-60 61-65 66-70 Frequency: 2 3 8 12 16 5 2 3



Watch Video Solution

195. Calculate mean, variance and standard deviation of the following frequency distribution: Class 1-10 10-20 20-30 30-40 40-50 50-60 Frequency: 11 29 18 4 5 3



Watch Video Solution

196. An analysis of monthly wages paid to workers in two firms A and B, belonging to the same industry, gives the

following results: Firm A Firm B No. of wages earners 586 684 Mean of monthly wages Rs. 5253 Rs. 5253 Variance of the distribution of wages 100 121 Which firm A or B pays to larger mount as monthly wages? Which firm A or B shows greater variability in individual wages?



[Watch Video Solution](#)

197. An analysis of monthly wages paid to workers in two firms A and B, belonging to the same industry, gives the following results: Firm A Firm B No. of wage earners 586 648 Mean of monthly wages Rs 5253 Rs 5253 Variance of the distribution of wages 100 121 (i) Which firm A or B pays larger amount as monthly wages? (ii) Which firm, A or B, shows greater variability in individual wages?



[Watch Video Solution](#)

198. The following values are calculated in respect of heights and weights of the students of a section of class XI:

Height Mean 162.6 cm Weight Mean 52.36 kg Height Variance 127.69 cm^2 Weight Variance 23.1361 kg^2

Can we say that the weights show greater variation than the heights?

[Watch Video Solution](#)

199. If the heights of 5 persons are 140 cm, 150cm, 158cm, 152 cm and 161 cm respectively. Find the mean height.

[Watch Video Solution](#)

200. The sum and sum of squares corresponding to length x (in cm) and weight y (in gm) of 50 plant products are given below :

$$\sum_{i=1}^{50} x_i = 212, \sum_{i=1}^{50} x_i^2 = 902.8, \sum_{i=1}^{50} y_i = 261, \sum_{i=1}^{50} y_i^2 = 1457.6$$

Which is more varying, the length or weight?



Watch Video Solution

201. Two plants A and B of a factory show following results about the number of workers and the wages paid to them.

Plant A	Plant B	No. of workers	Average monthly wages	Rs.	Rs.	Rs.	Rs.
5000	6000	2500	2500	81	100		

In which plant A or B is there greater variability in individual wages?



Watch Video Solution

 [Watch Video Solution](#)

202. find the mean salary of 60 workers of a factory from the given table



[Watch Video Solution](#)

203. The means and standard deviations of heights and weights of 50 students of a class are as follows:

	Weights	Heights
Mean	63.2 kg	63.2 inch
Standard deviation	5.6 kg	11.5 inch

Which shows more variability, heights or weights?



[Watch Video Solution](#)

204. The means and standard deviations of heights and weights of 50 students of a class are as follows: Weights Mean 63.2 kg Standard deviation 5.6 kg
Heights Mean 63.2 inch Standard deviation 11.5 inch Which shows more variability, heights or weights?



Watch Video Solution

205. Coefficient of variation of two distributions are 60% and 70% and their standard deviations are 21 and 16 respectively. What are their arithmetic means?



Watch Video Solution

206. Calculate coefficient of variation from the following data: Income (in Rs.) 1000-1700 1700-2400 2400-3100 3100-3800 3800-4500 4500-5200 No. o families 12 18 20 25 35 10



Watch Video Solution

207. An analysis of the weekly wages paid to workers in two firms A and B, belonging to the same industry gives the following results; Firm A Firm B No. of wages earners 586 648 Average weekly wages Rs. 2. Rs. 47.5 Distribution of wages 100 121 Which firm A or B pays out larger amount as weekly wages? Which firm A or B has greater variability in individual wages?



Watch Video Solution

208. An analysis of monthly wages paid to workers in two firms A and B, belonging to the same industry, gives the following results:

	Firm A	Firm B
No. of wage earners	586	648
Mean of monthly wages	Rs 5253	Rs 5253
Variance of the distribution of wages	100	121

(i) Which firm A or B pays larger amount as monthly wages? (ii) Which firm, A or B, shows greater variability in individual wages?



Watch Video Solution

209. The mean and standard deviation of marks obtained by 50 students of a class in their subjects, mathematics, physics and chemistry are given below:

Subject	Mathematics	Physics	Chemistry
Mean	42	32	40.9
Standard Deviation	12	15	

20 Which of the three subjects shows the highest variability in marks and which shows the lowest?



Watch Video Solution

210. Find the coefficient of variation for the following data:

Size (in cms)	10-15	15-20	20-25	25-30	30-35	35-40	No. of items
---------------	-------	-------	-------	-------	-------	-------	--------------

	2	8	20	35	20	15	
--	---	---	----	----	----	----	--



Watch Video Solution

211. The variance of first n natural number is:



Watch Video Solution

212. If the sum of the squares of deviations for 10 observations taken from their means is 2.5, then write the value of standard deviation.



Watch Video Solution

213. If x_1, x_2, \dots, x_n are n values of a variable X and y_1, y_2, \dots, y_n are n values of variable Y such that $y_i = ax_i + b, i = 1, 2, \dots, n$ then write $Var(Y)$ in terms of $Var(X)$



Watch Video Solution

214. In a series of 20 observations, 10 observation each equal to k and each of the remaining half is equal to $-k$. If the standard deviation of the observations is 2, then write the value of k .



Watch Video Solution

215. If each observation of a dist., whose variance is σ^2 , is multiplied by λ , then the S.D. of the new observations is



Watch Video Solution

216. If a variable X takes values of $0, 1, 2, \dots, n$ with frequencies ${}^nC_0, {}^nC_1, {}^nC_2, \dots, {}^nC_n$, then variance of

distribution is

 **Watch Video Solution**