



# MATHS

## BOOKS - KCET PREVIOUS YEAR PAPERS

### SOLVED PAPER 04

#### Mathematics

1. If  $ax^2 - y^2 + 4x - y = 0$  represents a pair of lines then a ...

A.  $-16$

B.  $1/(16)$

C.  $4$

D.  $-4$

**Answer: B**



**Watch Video Solution**

2. What is the equation of the locus of a point which moves such that 4 times its distance

from the x axis is the square of its distance from the origin?

A.  $x^2 + y^2 - 4y = 0$

B.  $x^2 + y^2 - 4|y| = 0$

C.  $x^2 + y^2 - 4x = 0$

D.  $x^2 + y^2 - 4|x| = 0$

**Answer: D**



**Watch Video Solution**

3. Equation of the straight line making equal intercepts on the axes and passing through the point (2,4) is

A.  $4x - y - 4 = 0$

B.  $2x + y - 8 = 0$

C.  $x + y - 6 = 0$

D.  $x + 2y = 100$

**Answer: C**



**Watch Video Solution**

4. If the area of the triangle with vertices  $(x, 0)$ ,  $(1, 1)$  and  $(0, 2)$  is 4 square units then a value of

A.  $-2$

B.  $-4$

C.  $-6$

D.  $8$

**Answer: C**



**Watch Video Solution**

5.  $\lim_{\theta \rightarrow x/2} \frac{\frac{\pi}{2} - \theta}{\cot \theta}$

A. 0

B. -1

C. 1

D.  $\infty$

**Answer: C**



**Watch Video Solution**

6. The co-axial system of circles given by

$$x^2 + y^2 + 2gx + c = 0$$

- A. intersecting circles
- B. non intersecting circles
- C. touching circles
- D. touching or non-intersecting circles

**Answer: D**



**View Text Solution**

7. The radius of the circle passing through the point (6,2) and two of whose diameters are  $x + y = 6$  and  $x + 2y = 4$  is

A. 4

B. 6

C. 20

D.  $\sqrt{20}$

**Answer: D**



**Watch Video Solution**



8. If  $(0, 6)$  and  $(0, 3)$  are respectively the vertex & focus of a parabola then its equation is

A.  $x^2 + 12y = 72$

B.  $x^2 - 12y = 72$

C.  $y^2 + 12x = 72$

D.  $y^2 + 12x = 72$

**Answer: A**



**Watch Video Solution**

9. For the ellipse  
 $25x^2 + 9y^2 - 150x - 90y + 225 = 0$  the  
eccentricity  $e =$

A.  $\frac{2}{5}$

B.  $\frac{3}{5}$

C.  $\frac{4}{5}$

D.  $\frac{1}{5}$

**Answer: C**



**Watch Video Solution**

10. If the foci of the ellipse  $\frac{x^2}{16} + \frac{y^2}{b^2} = 1$  and the hyperbola  $\frac{x^2}{144} - \frac{y^2}{81} = \frac{1}{25}$  coincide then the value of  $b^2$  is

A. 1

B. 7

C. 5

D. 9

**Answer: B**



**Watch Video Solution**

11. If  $Q_1$  is the set of all rationals other than 1 with the binary operation defined by  $a \cdot b = a + b - ab$  for all  $a, b$  in  $Q_1$ , then the identity in  $Q_1$  with respect to is

A. 1

B. 0

C.  $-1$

D. 2

**Answer: B**



Watch Video Solution

12. The circle  $x^2 + y^2 - 8x + 4y + 4 = 0$  touches

A. x-axis

B. y-axis

C. both axes

D. neither x-axies nor y-axis

**Answer: B**



Watch Video Solution

13. Which of the following is true?

A. The set of all fourth roots of unity is a multiplicative group.

B. The set of all cube roots of unity is an additive group

C.  $(ab)^{-1} = a^{-1}b^{-1}$  for all  $a, b$  in any group  $G$ .

D. If  $(ab)^2 = a^2b^2$  for all  $a, b$  in any group

$G$ , then the group  $G$  is non abelian

**Answer: A**



**View Text Solution**

**14.** The set of all integral multiples of 5 is a subgroup

A. The set of all rational numbers under multiplication

B. The set of all integers under multiplication

C. The set of all non zero rational numbers under multiplication

D. The set of all integers under addition

**Answer: C**



**Watch Video Solution**



15. The value of  $k$  so that

$$x^2 + y^2 + kx + 4y + 2 = 0 \quad \text{and}$$

$$2(x^2 + y^2) - 4x - 3y + k = 0 \quad \text{cut}$$

orthogonally is

A.  $\frac{10}{3}$

B.  $\frac{-8}{3}$

C.  $\frac{-10}{3}$

D.  $\frac{8}{3}$

**Answer: B**



Watch Video Solution

16.  $\lim_{x \rightarrow \infty} \left(1 - \frac{4}{x-1}\right)^{3x-1} =$

A.  $e^{12}$

B.  $e^{-12}$

C.  $e^4$

D.  $e^3$

**Answer: B**



**Watch Video Solution**

17. If  $A+B+C = 180^\circ$  then  $\sum \frac{\tan(A)}{2} \frac{\tan(B)}{2} =$

A. 0

B. 1

C. 2

D. 3

**Answer: B**



**Watch Video Solution**

**18.** In a triangle ABC if  $b = 2, B = 30^\circ$  then the area of the circumcircle of triangle ABC in square units is

A.  $\pi$

B.  $2\pi$

C.  $4\pi$

D.  $6\pi$

**Answer: C**



**Watch Video Solution**

19. If  $\sin x + \sin^2 x = 1$  then

$$\cos^{12} 2x + 3 \cos^{10} x + 3 \cos^8 x + \cos^6 x =$$

A. 1

B. 2

C. 3

D. 0

**Answer: A**



**Watch Video Solution**

20. If  $R$  denotes the set of all real numbers then the function  $f: R \rightarrow R$  defined  $f(x) = |x|$

- A. one-one only
- B. onto only
- C. both one-one & onto
- D. neither one-one nor onto

**Answer: D**



**Watch Video Solution**

21. The equation of the director circle of the hyperbola  $\frac{x^2}{16} - \frac{y^2}{4} = 1$  is given by

A.  $x^2 + y^2 = 16$

B.  $x^2 + y^2 = 4$

C.  $x^2 + y^2 = 20$

D.  $x^2 + y^2 = 12$

**Answer: D**



**Watch Video Solution**

22. If  $0 \leq x \leq \pi$  and  $81^{\sin^2 x} + 81^{\cos^2 x} = 30$

then  $x =$

A.  $\frac{\pi}{6}$

B.  $\frac{\pi}{2}$

C.  $\frac{\pi}{4}$

D.  $\frac{3\pi}{4}$

**Answer: A**



**Watch Video Solution**



23. If  $\frac{\sin^{-1}(x)}{5} + \operatorname{cosec}^{-1} \frac{5}{4} = \frac{\pi}{2}$  then  $x =$

A. 1

B. 4

C. 3

D. 5

**Answer: C**



**Watch Video Solution**

24. If  $\cos^{-1} p + \cos^{-1} q + \cos^{-1} r = \pi$  then

$$p^2 + q^2 + r^2 + 2pqr =$$

A. 3

B. 1

C. 2

D. -1

**Answer: B**



**Watch Video Solution**

25. The smallest positive integer for which

$$(1 + i)^{2n} = (1 - i)^{2n} \text{ is}$$

A. 1

B. 2

C. 3

D. 4

**Answer: B**



**Watch Video Solution**

26. If  $\frac{\log x}{a - b} = \frac{\log y}{b - c} = \frac{\log z}{c - a}$  then  $xyz =$

A. 0

B. 1

C. -1

D. 2

**Answer: B**



**Watch Video Solution**

27. The last digit in  $7^{300}$  is

A. 7

B. 9

C. 1

D. 3

**Answer: C**



**Watch Video Solution**

28. How many numbers of 6 digits can be formed from the digits of the number 112233?

A. 30

B. 60

C. 90

D. 120

**Answer: C**



**Watch Video Solution**

29. The number of solutions for the equation

$$x^2 - 5x + 6 = 0 \text{ is..}$$

A. 4

B. 3

C. 2

D. 1

**Answer: A**



**Watch Video Solution**

30. 0.5737373 ...

A.  $\frac{284}{497}$

B.  $\frac{284}{495}$

C.  $\frac{568}{999}$

D.  $\frac{567}{990}$

**Answer: D**



**Watch Video Solution**



31.  $\sim p \wedge q$  is logically equivalent to ...

A.  $p \rightarrow q$

B.  $q \rightarrow p$

C.  $\sim(p \rightarrow q)$

D.  $\sim(q \rightarrow p)$

**Answer: D**



**Watch Video Solution**

**32.** Which of the following is the inverse of the proposition: "If a number is a prime then it is odd?"

A. If a number is not a prime then it is odd.

B. If a number is not a prime then it is not odd.

C. If a number is not odd then it is not a prime.

D. If a number is odd then it is a prime.

**Answer: B**



**Watch Video Solution**

**33.** What must be the matrix  $X$  if

$$2X + \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} = \begin{bmatrix} 3 & 8 \\ 7 & 2 \end{bmatrix}?$$

A.  $\begin{bmatrix} 1 & 3 \\ 2 & -1 \end{bmatrix}$

B.  $\begin{bmatrix} 1 & -3 \\ 2 & -1 \end{bmatrix}$

C.  $\begin{bmatrix} 2 & 6 \\ 4 & -2 \end{bmatrix}$

D.  $\begin{bmatrix} 2 & -6 \\ 4 & -2 \end{bmatrix}$

**Answer: A**



**Watch Video Solution**

34. The value of 
$$\begin{vmatrix} 1 & 1 & 1 \\ bc & ca & ab \\ b+c & c+a & a+b \end{vmatrix}$$

A. 1

B. 0

C.  $(a-b)(b-c)(c-a)$

D.  $(a+b)(b+c)(c+a)$

**Answer: C**



**Watch Video Solution**

35. The value of  $\begin{vmatrix} 441 & 442 & 443 \\ 445 & 446 & 447 \\ 449 & 450 & 451 \end{vmatrix}$  is

A.  $441 \times 446 \times 451$

B. 0

C.  $-1$

D. 1

**Answer: B**



**Watch Video Solution**

$$36. \left( \vec{a} \cdot \vec{i} \right) \vec{i} + \left( \vec{a} \cdot \vec{j} \right) \vec{j} + \left( \vec{a} \cdot \vec{k} \right) \vec{k}$$

=

A.  $\vec{a}$

B.  $2\vec{a}$

C.  $3\vec{a}$

D.  $\vec{0}$

**Answer: A**



**Watch Video Solution**

37. Inverse of the matrix  $\begin{bmatrix} \cos 2\theta & -\sin 2\theta \\ \sin 2\theta & \cos 2\theta \end{bmatrix}$  is

A.  $\begin{bmatrix} \cos 2\theta & -\sin 2\theta \\ \sin 2\theta & \cos 2\theta \end{bmatrix}$  is

B.  $\begin{bmatrix} \cos 2\theta & \sin 2\theta \\ \sin 2\theta & -\cos 2\theta \end{bmatrix}$  is

C.  $\begin{bmatrix} \cos 2\theta & \sin 2\theta \\ \sin 2\theta & \cos 2\theta \end{bmatrix}$  is

D.  $\begin{bmatrix} \cos 2\theta & \sin 2\theta \\ -\sin 2\theta & \cos 2\theta \end{bmatrix}$  is

**Answer: D**



Watch Video Solution

38. IF  $|\vec{a}| = 3, |\vec{b}| = 4$  then a value of  $\lambda$  for which  $\vec{a} + \lambda \vec{b}$  is perpendicular to  $\vec{a} - \lambda \vec{b}$

A.  $\frac{9}{16}$

B.  $\frac{3}{4}$

C.  $\frac{3}{2}$

D.  $\frac{4}{3}$

Answer: D





39. The projection of  $\vec{a} = 2i + 3j - 2k$  on  $\vec{b} = i + 2j + 3k$  is

A.  $\frac{1}{\sqrt{(14)}}$

B.  $\frac{2}{\sqrt{(14)}}$

C.  $\sqrt{14}$

D.  $\frac{-2}{\sqrt{(14)}}$

**Answer: B**



40. In the group  $\{1, 2, 3, 4, 5, 6\}$  under multiplication mod 7,  $2^{-1} \times 4 =$

A. 1

B. 4

C. 2

D. 3

**Answer: C**



**Watch Video Solution**

41. If a tangent to the curve  $y = 6x - x^2$  is parallel to the line  $4x - 2y - 1 = 0$ , then the point of tangency on the curve is

A. (2,8)

B. (8,2)

C. (6,1)

D. (4,2)

**Answer: A**



**View Text Solution**

42.  $\int \frac{dx}{x^2 + 2x + 2} =$

A.  $\sin^{-1}(x + 1) + c$

B.  $\sinh^{-1}(x + 1) + c$

C.  $\tanh^{-1}(x + 1) + c$

D.  $\tan^{-1}(x + 1) + c$

**Answer: D**



**Watch Video Solution**

43.  $\int \sqrt{x} e^{\sqrt{x}} dx =$

A.  $2\sqrt{x} - e^{\sqrt{x}} - 4\sqrt{x}e^{\sqrt{x}} + c$

B.  $(2x - 4\sqrt{x} + 4)e^{\sqrt{x}} + c$

C.  $(2x + 4\sqrt{x} + 4)e^{\sqrt{x}} + c$

D.  $(1 - 4\sqrt{x})e^{\sqrt{x}} + c$

**Answer: B**



**Watch Video Solution**

$$44.1 \int \frac{dx}{x(x^7 + 1)} =$$

A.  $\log\left(\frac{x^7}{x^7 + 1}\right) + c$

B.  $\frac{1}{7} \log\left(\frac{x^2}{x^7 + 1}\right) + c$

C.  $\log\left(\frac{x^{2 \log\left(\frac{x^7+1}{x^7}\right) + c}}{x^7 + 1}\right) + c$

D.  $\frac{1}{7} \log\left(\frac{x^7 + 1}{x^7}\right) + c$

**Answer: B**



**Watch Video Solution**

45.  $\int_0^1 |1 - x| dx =$

A.  $-2$

B.  $0$

C.  $2$

D.  $4$

**Answer: C**



**Watch Video Solution**

46. If  $x + \frac{1}{x} = 2 \cos \alpha$  then  $x^n + \frac{1}{x^n} =$

A.  $2^n \cos \alpha$

B.  $2^n \cos n\alpha$

C.  $2^n \sin n\alpha$

D.  $2^1 \cos n\alpha$

**Answer: D**



**Watch Video Solution**



47. If  $y = \tan^{-1}(\sec x + \tan x)$  then  $\frac{dy}{dx} =$

A. 2

B.  $-2$

C.  $\frac{1}{2}$

D.  $-\frac{1}{2}$

**Answer: C**



**Watch Video Solution**

48. If  $\omega = \frac{-1 + i\sqrt{3}}{2}$  then

$$(3 + \omega + 3\omega^2)^4 =$$

A. 16

B.  $-16$

C.  $16\omega$

D.  $16\omega^2$

**Answer: A**



**Watch Video Solution**

49. If  $f(x) = \begin{cases} \frac{1 - \cos x}{x} & x \neq 0 \\ k & x = 0 \end{cases}$  is continuous

at  $x=0$  then  $k=$

A. 0

B.  $\frac{1}{2}$

C.  $\frac{1}{4}$

D.  $-\frac{1}{2}$

**Answer: A**



**Watch Video Solution**

50. The differential coefficient of  $f(\sin x)$  with respect to  $x$  where  $f(x) = \log x$  is

A.  $\tan x$

B.  $\cot x$

C.  $f(\cos x)$

D.  $1/x$

**Answer: B**



**Watch Video Solution**

51.  $\int_0^{n/2} \frac{\cos x - \sin x}{1 + \cos x \sin x} dx$

A. 0

B.  $\frac{\pi}{2}$

C.  $\frac{\pi}{4}$

D.  $\frac{\pi}{6}$

**Answer: A**



**Watch Video Solution**

52.  $\int_0^{n/8} \cos^3 4\theta d\theta =$

A.  $\frac{2}{3}$

B.  $\frac{1}{4}$

C.  $\frac{1}{3}$

D.  $\frac{1}{6}$

**Answer: D**



**Watch Video Solution**

53. The area enclosed between the curves  $y = x^3$  and  $y = \sqrt{x}$  is, (in square units)

A.  $\frac{5}{3}$

B.  $\frac{5}{4}$

C.  $\frac{5}{12}$

D.  $\frac{12}{5}$

**Answer: C**



**Watch Video Solution**

54. The degree of the differential Equation

$$\left(1 + \left(\frac{dy}{dx}\right)^2\right)^{3/4} = \left(\frac{d^2y}{dx^2}\right)^{1/3}$$

A.  $\frac{1}{3}$

B. 4

C. 9

D.  $\frac{3}{4}$

**Answer: B**



**Watch Video Solution**



55. The general solution of the differential

equation  $\frac{dy}{dx} + \frac{1 - \cos 2x}{1 + \cos 2y} = 0$  is given by

A.  $\tan y + \cot x = c$

B.  $\tan y - \cot x = c$

C.  $\tan x - \cot y = c$

D.  $\tan x + \cot x = c$

**Answer: B**



**Watch Video Solution**

56. If  $x = a\left(t - \frac{1}{t}\right), y = a\left(t + \frac{1}{t}\right)$  then

$$\frac{dy}{dx} =$$

A.  $\frac{y}{x}$

B.  $\frac{-y}{x}$

C.  $\frac{x}{y}$

D.  $\frac{-x}{y}$

**Answer: C**



**Watch Video Solution**

57. If  $x = A \cos 4t + B \sin 4t$  then  $\frac{d^2x}{dt^2} =$

A.  $-16x$

B.  $16x$

C.  $x + y - 6 = 0$

D.  $-x$

**Answer: A**



**Watch Video Solution**

58. For the curve  $y^n \equiv a^{n-1}x$  if the subnormal at any point is a constant then  $n =$

A. 1

B. 2

C.  $-2$

D.  $-1$

**Answer: B**



**Watch Video Solution**

59. If the distance 's' metres traversed by a particle in 1 seconds is given by  $s = t^3 - 3t^2$ , then the velocity of the particle when the acceleration is zero, in metres/sec. is

A. 3

B. - 2

C. - 3

D. 2

**Answer: C**



**Watch Video Solution**

60. The maximum of the function  $3 \cos x - 4 \sin x$  is

A. 2

B. 3

C. 4

D. 5

**Answer: D**



**Watch Video Solution**

